Full Title: Extended hopanoid lipids promote bacterial motility, surface attachment, and root nodule

development in the *Bradyrhizobium diazoefficiens-Aeschynomene afraspera* symbiosis

Short Title: Extended hopanoids promote efficient *Bradyrhizobium-Aeschynomene* symbiosis

Belin, B.J.,[1] Tookmanian, E.T.,[1] de Anda, J.,[3] Wong, G. C. L.,[3] Newman, D.K.[1,2]

[1]Division of Biology & Bioengineering, [2]Division of Geological & Planetary Sciences, California Institute of

Technology, Pasadena, CA.

[3]Department of Bioengineering, Department of Chemistry and Biochemistry, and California NanoSystems

Institute, University of California Los Angeles, Los Angeles, CA 90095-1600, USA University of California

Los Angeles, Los Angeles, CA.

12   **Abstract**

Hopanoids are steroid-like bacterial lipids that enhance membrane rigidity and promote bacterial growth

under diverse stresses. Roughly 10% of bacteria contain genes involved in hopanoid biosynthesis, and

these genes are particularly conserved in plant-associated organisms. We previously found that

the extended class of hopanoids (C_{35}) in the nitrogen-fixing soil bacterium *Bradyrhizobium*

diazoefficiens promotes its root nodule symbiosis with the tropical legume *Aeschynomene afraspera*. By

quantitatively modeling root nodule development, we identify independent roles for hopanoids in the

initiation of root nodule formation and in determining the rate of root nodule maturation. *In vitro* studies

demonstrate that extended hopanoids support *B. diazoefficiens* motility and surface attachment, which

may correlate with stable root colonization *in planta*. Confocal microscopy of maturing root nodules

reveals that root nodules infected with extended hopanoid-deficient *B. diazoefficiens* contain unusually

low densities of bacterial symbionts, indicating that extended hopanoids are necessary for persistent, high

levels of host infection. This work identifies extended hopanoids as regulators of the efficiency

of *Bradyrhizobia* nitrogen-fixing symbioses, agriculturally and economically significant associations with

growing importance in a changing climate.

Introduction

29   Hopanoids are steroid-like lipids synthesized by metabolically and environmentally diverse

bacteria. Studies in multiple organisms have shown that hopanoids support bacterial survival in response

to stress, including growth under conditions of high osmolarity, low pH, elevated temperatures,

detergents, and antibiotics (reviewed in ref. [1]). It is thought that hopanoids primarily promote survival by

rigidifying and decreasing the permeability of membranes,[2],[3] providing a better barrier against external

insults. Hopanoids may also directly regulate proteins, chemical signals, and other lipids in the membrane

(as do eukaryotic sterols[4]), but such interactions are yet to be characterized.

36   Hopanoids consist of a 30-carbon pentacyclic core, generated by the squalene-hopene cyclase

(*shc*) family of enzymes,[5],[6] that can be modified to generate a medley of distinct molecules. The most

common hopanoid modifications are methylation by the enzyme HpnP[7] and addition of a ribose-derived

side chain by the enzyme HpnH (Fig. 1a)[8]. Downstream enzymes can further diversify side chains

added by HpnH,[9] and the resulting molecules are known collectively as the C_{35} or extended hopanoids.

Common extended hopanoid side chains include aminotriol-, polyol-, and adenosyl- moieties, although

organism-specific side chains have also been observed. For example, the *Bradyrhizobiaceae* family of

soil bacteria encode an extended hopanoid with a lipid A side chain, known as HoLA[10–12]. It is an open

question whether HoLA and other specific extended hopanoids perform non-overlapping functions in

cells, although evidence that methylated and extended hopanoids affect membrane biophysics and stress

tolerance to differing degrees supports this notion (reviewed in ref. [1]).

In the environment, hopanoid biosynthesis genes are enriched in plant-associated

metagenomes,[13] and they are highly conserved in the plant growth-promoting *Bacilli*, *Nostoc* and

Anabaena symbionts of various green plants, and *Frankia* and *Bradyrhizobia* nitrogen-fixing symbionts of

legumes[14]'[15]. We previously found that hopanoids are required for optimal symbiosis of two strains of

Bradyrhizobia, the photosynthetic *Bradyrhizobium* BTAi1 and the model *Bradyrhizobium diazoefficiens*,

with members of the *Aeschynomene* genus of tropical legumes[10,11]. The symbiotic properties of *B.*

diazoefficiens are of particular agricultural and economic significance because it natively forms symbioses

with soybean crops[16]. In a *B. diazoefficiens-Aeschynomene afraspera* symbiosis model, we found that

the bacterial extended hopanoids specifically are required for optimal rates of *in planta* nitrogen

fixation[11].

Numerous processes in bacteria and plant hosts determine the rate of nitrogen fixation in legume-

rhizobia symbioses. The most basic requirement is the inherent capacity of bacteria to perform the

nitrogen fixation reaction, which demands a high uptake of various trace metals, rapid consumption of

protons and ATP, and the absence of freely-diffusing oxygen. In *Frankia* legume symbionts, it has been

suggested that hopanoids help to maintain a low-oxygen environment by limiting oxygen diffusion across

membranes[17–19], although comparative measurements of free oxygen in wild type and hopanoid-

deficient strains have not been performed.

Another determinant of root nodule nitrogen fixation levels is the efficiency of root nodule

development, beginning with rhizobial colonization of the plant root surface. In *Aeschynomene* hosts root

colonization occurs on plant root "cracks", breaks in the outer epidermal layers formed naturally by lateral

root emergence[20]. *B. diazoefficiens* possesses two systems that guide motility towards these cracks, a

constitutive subpolar flagellum and inducible lateral flagella[21], which appear to fill complementary roles.

The subpolar flagellum primarily drives chemotaxis in open aquatic environments, whereas the lateral

flagella promote motility on surfaces or in viscous fluids[22]; presumably these activities correspond to

persistent, long-range movement towards hosts and clearing obstacles on the soil and root surface,

respectively[23,24]. In *A. afraspera,* root surface colonization also requires bacterial Nod factors,

immunosuppressant molecules that activate pro-symbiotic signaling[25].

After stable colonies are established at root cracks, rhizobial infections progress to the root

interior. The cytological details of this process have been characterized for *A. afraspera* only in

association with the *Bradyrhizobium* strain ORS285,[25] and in this context, rhizobia can be observed

invading roots via thread-like patterns in the intercellular spaces of the inner epidermal and cortical cell

layers. At the cortex, several host cells become intracellularly infected with bacteria. These infected cells

appear to internalize the bacteria through narrow membrane invaginations, each enclosing multiple

bacterial cells, that pinch off to form intracellular infection pockets[25]. In a few infected cells, these

infection pockets presumably break down into symbiosomes, an organelle-like structure consisting of

intracellular bacteria (now called "bacteroids") surrounded by a plant membrane, which is the basic

nitrogen-fixing unit and defining feature of the root nodule.

Two developmental programs follow symbiosome formation in *A. afraspera*. In the first, infected

host cells enter a proliferative state[25], forming the nodule primordium, and grow to form a large,

spherical mass of infected cells known as the central infection zone. As the infection zone develops it is

enveloped by epidermal tissue, and eventually, a vasculature system. This pathway for infection zone

formations is unusual among legumes. More commonly, the nodule primordium arises from a meristem

that is regulated independently and spatially separated from the bacterial invasion site; paradoxically,

bacterial internalization in meristem-dependent nodules often inhibits, rather than stimulates, further host

proliferation[26]. Another notable feature of *A. afraspera* is that two or more discrete infection zones may

develop in the same nodule, resulting in a multi-lobed morphology. Each infection zone is thought to arise

from a single infected cell, as co-inoculation of *A. afraspera* with a mixture of GFP- and mCherry-

expressing *Bradyrhizobia* strains generally results in infection zones expressing only one

fluorophore[25,27].

The second program initiated by symbiosome formation is the establishment of major genetic and

chemical changes for bacteroids, including a sharp reduction in pH[28] and free oxygen levels[29] in the

surrounding environment, high levels of bacteroid genome duplication, suppression of non-symbiotic

genes, extreme changes in bacterial morphology, and a loss of bacteroid viability. These genetic changes

are directed largely by antimicrobial NCR (nodule cysteine-rich) peptides produced by legume hosts

(reviewed in [30]). Mutant rhizobia with higher susceptibility to the NCR peptides *in vitro* tend to be

ineffective symbionts that trigger plant defense reactions or early senescence of root nodules, a process

in which the central nodule tissue is degraded and recycled by the host[31].

In this study, we sought to better understand how the extended hopanoids of *B. diazoefficiens*

support nitrogen fixation in the *A. afraspera* root nodule symbiosis. We found that the lower nitrogen

fixation of extended hopanoid mutants can be explained by a reduction in root nodule size and rhizobial

occupancy, indicating that the underlying defect is primarily developmental rather than relating to per-

bacteroid nitrogen fixation productivity. By novel application of a quantitative framework for modeling plant

tissue growth using single-nodule tracking, we find that extended hopanoids support root nodule

development through two independent pathways: by promoting bacterial motility towards and attachment

to root surfaces and by maintaining normal host and symbiosome proliferation rates in the infection zone.

In addition, our results reveal the dynamics of nodulation in *A. afraspera*, demonstrating that for these

plants nodulation occurs in bursts separated by fixed intervals that do not depend on net fixed nitrogen

production across the root. We further suggest that nodule growth rates are key determinants of an *A.*

afraspera nodule's nitrogen fixation output at maturity, in which growth rates may be "tuned" to stimulate

proliferation in nodules with more effective symbionts.

Results

Loss of extended hopanoids results in reduced nodule size

Previously, we observed a symbiotic defect for a mutant lacking extended hopanoids (ΔhpnH) in

the *B. diazoefficiens-Aeschynomene afraspera* symbiosis model[11]. To verify and better understand this

defect, we inoculated *A. afraspera* plants with ΔhpnH, ΔhpnP (lacking 2-Me hopanoids), or wild-type *B.*

diazoefficiens. At 24 days post-inoculation (dpi), plants inoculated with Δ*hpnH* were shorter than wild

type-inoculated plants, although both strains produced equivalent numbers of nodules (Fig. 1b). Δ*hpnH*-

inoculated plants also exhibited a roughly 50% decrease in the rate of reduction of acetylene gas (a proxy

for nitrogen fixation) compared to wild type-inoculated plants at this time point (Fig. 1c). In contrast, the

Δ*hpnP* mutant was similar to wild type (Fig. 1b-c). These results are consistent with our previous

findings[11].

We initially hypothesized that the Δ*hpnH* defect in nitrogen fixation *in planta* occurs because of

poor survival of the Δ*hpnH* mutant in hosts, given that *in vitro* the Δ*hpnH* strain is more susceptible to

conditions mimicking the nodule environment (*e.g.* low pH, low oxygen, high osmolarity, and NCR

peptides)[11]. To assess Δ*hpnH* viability within *A. afraspera* nodules, we performed morphological

analyses of nodule semi-thin sections using confocal fluorescent microscopy. Hand-sectioned nodule

slices (of roughly 200-500 µm thickness) were stained with a bacterial live:dead kit, consisting of the

SYTO9 DNA dye, which is cell-permeable and labels all cells, and propidium iodide (PI), a dye that

uniquely labels cells with compromised membranes, which are generally non-viable. Contrary to our

expectations, we did not observe an increase in PI staining in nodules containing Δ*hpnH* (Fig. 1d; Fig.

S1,S2). Signatures of plant defense reactions, which previously were associated with Δ*hpnH*[11], were

also rare, occurring prominently in only ~1.5% (1/67) of nodules (Fig. S2).

The most apparent difference in the morphology of Δ*hpnH* nodules compared to wild type was

the presence of relatively small nodules (Fig. S1,S2). To determine whether lower nodule size could

explain reduced nitrogen fixation in Δ*hpnH*-inoculated plants, we repeated acetylene reduction assays for

wild-type- and Δ*hpnH*-inoculated plants, harvested their nodules, and calculated the total nodule dry

mass for each plant at 24 dpi. We found a decrease in the nodule dry mass per plant for Δ*hpnH*-

inoculated plants that is commensurate with the decrease in acetylene reduction rates (Fig. 1e). After

normalization by nodule dry mass, no significant difference could be observed in acetylene reduction

between the wild-type- and Δ*hpnH*-inoculated plants (Fig. 1e). This suggests the *in planta* nitrogen

fixation defect of Δ*hpnH* at 24 dpi can be explained by smaller average nodule sizes. It also rules out the

possibility that nitrogenase functions ineffectively in the absence of extended hopanoids due to

inactivation by oxygen, as has been suggested in *Frankia*[17–19], as the per-mg nitrogen fixation rates

are unaffected in Δ*hpnH*.

*Δ*hpnH *nodules are more variable in size than wild-type nodules*

Curious about the origin of the smaller Δ*hpnH* nodule sizes, we devised three *a priori* hypotheses

for how small nodules could arise: first, that Δ*hpnH* nodule growth is terminated prematurely; second, that

Δ*hpnH* nodules grow at a slower rate; and third, that the initiation of Δ*hpnH* nodules is delayed. Of these

scenarios, the first possibility is unique in that Δ*hpnH* nodules would be *necessarily* smaller than wild

type; in other words, Δ*hpnH* nodules could not "catch up" to wild type given additional time. To test this

possibility, we measured acetylene reduction per plant across an extended 40 dpi period, and we

observed that the differences in both acetylene reduction rates and nodule dry masses between wild type

and Δ*hpnH* steadily decreased with time (Fig. 2a-b). By 40 dpi the overall symbiotic efficiencies of wild

type and Δ*hpnH* per plant were indistinguishable, in terms of the plants' qualitative appearance (Fig. 2c,d)

as well as their average shoot heights and acetylene reduction rates (Fig. S3). Total nodule counts per

plant also did not differ between wild type and Δ*hpnH* at 40 dpi, indicating that the increase in total nodule

mass reflects growing nodules rather than more frequent nodulation (Fig. S3).

These total dry mass measurements appeared consistent with Δ*hpnH* nodules "catching up" to

wild type, so we sought to independently verify this interpretation by measuring the radii of individual

nodules on ten plants for each strain at 40 dpi (Fig. 2e,f). From these data we determined that although

average nodule sizes (and, by extension, the total nodule mass) did become similar between strains by

this time point (0.73 vs. 0.88 mm average radii), their underlying distributions were markedly distinct.

Wild-type nodule radii appear to form a roughly normal distribution, whereas the Δ*hpnH* nodule radius

distribution is bimodal, consisting of a subpopulation of small nodules with small radii (<0.5 mm) that are

rarely observed in wild type, as well as a second, larger subpopulation that has a similar median radius as

wild type but is skewed towards larger radii (>1.5mm). These data demonstrate that the small-nodule

phenotype of Δ*hpnH* persists throughout a 40 dpi time course, but is compensated by greater size

heterogeneity, in which a handful of "mega" nodules offset smaller nodules over time.

ΔhpnH nodule size heterogeneity reflects variable nodule growth rates

The static measurements described above were insufficient to evaluate our hypothesis that

Δ*hpnH* nodule growth is aborted, because we could not infer from these assays whether the small

nodules observed at 24 (Fig. 1) and 40 (Fig. 2) dpi represent the same population, or whether they simply

began growing at different times. To better evaluate the possible mechanisms of the Δ*hpnH* nodule size

defect, we studied the kinetics of single nodule development. Beginning one week after inoculation of *A.*

afraspera plants with either wild type or Δ*hpnH*, we collected images of entire plant roots every 3-5 days

up to ~40 days post-inoculation (Fig. S4, S5). From these images, we identified nodules that were clearly

visible (*e.g.* not obscured by lateral roots or more recently emerged nodules) in at least five time points

(Fig. 3a) and measured their radii. We then calculated nodule volumes by approximating nodules as

spheres and plotted the volume of the tracked nodules over time. While we again observed that many

Δ*hpnH* nodules were smaller at 40 dpi than any of the wild type nodules, we also found that nodule

growth was highly variable both within and between strains (Fig. 3b,c).

To describe this heterogeneity, we employed a simple framework for quantifying nodule

development. We describe nodule growth by the following variables: the time (t_i) of the initial intracellular

infection event and the volume of the nascent nodule (V_i), equivalent to the volume of one infected *A.*

afraspera cortical cell; the time (t_{min}) and volume (V_{min}) at which a clearly visible, spherical nodule has

developed; the rate of growth of a nodule once it has become visible (dV/dt); and the time (t_{max}) and

volume (V_{max}) of a nodule when its growth has stopped (Fig. 3d). To calculate these variables, we fit each

nodule's growth over time to three different growth models: exponential, quadratic, and a generalized

logistic (*e.g.* sigmoidal) equation commonly used for plant growth[32,33] (see Methods for complete

details). Sigmoidal models generally provided the best fit to the experimental data, so these models were

used for growth parameter calculation (Fig. 3e; Fig. S6, S7).

The growth rates of Δ*hpnH* nodules were lower on average than wild-type nodules (Fig. 3f), with

roughly a third of tracked nodules exhibiting growth rates lower than observed for wild type by 40 dpi

(<0.1 mm^3/dpi). We noted that this was the same fraction of nodules as in the subpopulation of nodules

with smaller-than wild type radii at 40 dpi (Fig. 2e; Fig. 3g), and indeed, the subpopulation of nodules with

lower-than-wild-type growth rates and small nodule sizes are the same (Fig. 3h). Additionally, we found

that the growth rate of a nodule and its maximum size are positively linearly correlated for both strains,

with Pearson coefficients of ~0.64 ($p<10^{-9}$) for wild type and ~0.75 ($p<10^{-15}$) for Δ*hpnH*. We interpret these

data to suggest that host proliferation is slower in a subset of nodules infected with Δ*hpnH*, and that this

at least partially accounts for the low final volume of these nodules.

We also noted that the Δ*hpnH* nodule sizes at 40 dpi differed between these single-nodule

volume measurements (Fig. 3g) and our previous 40 dpi end-point measurements of nodule radii (Fig.

2e), in that we did not observe larger-than-wild-type "mega" nodules in the single-nodule dataset. This

discrepancy likely reflects the smaller sample size in our single-nodule tracking experiments (84

compared to 268 end-point nodules), and the low frequency of "mega" nodule formation. To verify this, we

selected 10,000 random subsets of 84 nodules from the 268 Δ*hpnH* nodules shown in Figure 2e,

converted the nodule radii to volumes, and found that there is no statistically significant difference

($p<0.05$) between a random subset of Fig. 2e and the Δ*hpnH* single-nodule tracking data in ~92%

(9184/10000) of cases. Thus the differences in nodule size distributions in Figure 3g and Figure 2e are

consistent with sampling error.

We also used nodule growth fitting to test our hypothesis that the growth period is shorter for

Δ*hpnH* nodules compared to wild type – *e.g.* that growth stops prematurely. We estimated the window of

maximum growth as the time required for a nodule to increase from 10% to 90% of its final volume.

Neither the time at which a nodule reaches 90% of its maximum volume, t_{max}, nor the window of

maximum growth differs significantly between Δ*hpnH* and wild type (Fig. S8a,b). The window of maximum

growth for each nodule is also uncorrelated with their final volume or growth rate, indicating that small

nodules are not prematurely aborted; rather, their growth periods are similar to larger nodules (Fig. S9a-

d).

To better understand the subpopulation of small, slow-growing Δ*hpnH* nodules, we sectioned

nodules from Δ*hpnH*-inoculated plants at 40 dpi and separated the small (<0.5 mm radius) subpopulation

of nodules from all others (>0.5 mm radius). We stained sections with SYTO9, PI and Calcofluor and

imaged the sections with confocal microscopy. We found that while most small Δ*hpnH* nodules contained

a single, continuous infection zone, a large fraction were un- or under-infected with bacteria, often

exhibiting disorganized central infection zones (~37%; 28/75) (Fig. 4a; Fig. S10). Of the fully infected

small Δ*hpnH* nodules, some contained primarily PI-stained, likely dead bacterial cells (~25%; 12/47) (Fig.

4a; Fig. S10). Similarly high proportions of under-infected nodules or nodules primarily occupied with

membrane-compromised bacteria did not occur in larger Δ*hpnH* nodules harvested at the same time

point; however, we did notice that many of these nodules contained noticeably fragmented infection

zones (Fig. 4b; Fig. S11).

We also compared the subpopulation of small Δ*hpnH* nodules at 40 dpi to two wild-type nodule

populations: similarly small nodules harvested at 10 and 25 dpi (Fig. 4a; Fig S12; Fig. S13), and nodules

harvested at the same 40 dpi time point (Fig. 4b; Fig. S14). Again, we found that higher proportions of

under-infected nodules and nodules containing primarily membrane-compromised bacteria were unique

to the Δ*hpnH* small-nodule subset. It is interesting that these phenotypes were not apparent for Δ*hpnH*

nodules in the previously analyzed 24 dpi time point (Fig. S1), in which nodules that are under-infected or

primarily occupied by dead bacteroids are rare. It is possible we did not capture these phenotypes

because they occur infrequently in the overall nodule population; infection zones with primarily dead

bacteroids occur in only 12/75 small Δ*hpnH* nodules in Fig. S10, and small (<0.5 mm radius) nodules

themselves make up only a third of all Δ*hpnH* nodules at 40 dpi. An alternate conclusion from these data

is that poor Δ*hpnH* survival or infection density is only apparent at later stages in development, indicating

inherently poor persistence of Δ*hpnH* within host cells and/or premature nodule senescence.

*Δ*hpnH *is delayed in early plant-symbiont association*

Our final hypothesis for the origin of small nodules, that the onset of nodule growth is delayed,

proved more difficult to address. The "true" beginning of nodule formation is the time when the first *A.*

afraspera cortical cell is infected, t_i (Fig. 3d). However, this initial infection event is not visible at the root

surface, and it is difficult to extrapolate from sigmoidal models in which the growth curves approach the

initial volume $V_i \sim 0$ mm^3 asymptotically. As a proxy for t_i, we defined three alternate t_{min} as the times at

which nodules reached three arbitrarily small volumes: $V = 0.05$ mm^3, $V = 0.1$ mm^3, and $V = 0.2$ mm^3.

When t_{min} is defined by $V = 0.05$ mm^3 or 0.1 mm^3, t_{min} could not be accurately calculated for all nodules, as

the sigmoidal models sometimes predicted $t_{min} < 0$ (an impossible value) (Fig. S8c). These nodule

volumes are also too small to be seen on the root surface, and we had no experimental means to

determine the accuracy of the calculations in this low-volume regime. When t_{min} is defined by V = 0.2 mm^3

(the smallest nodule volume that we could identify in our single-nodule tracking assays), there is a small

but statistically significant increase for Δ*hpnH* relative to wild type (Fig. S8c). To independently verify this

delay in visible nodule appearance, we inspected the roots of 20 wild type- and 20 Δ*hpnH*-inoculated

plants over 40 dpi and recorded the number of visible nodules per plant each day. We found a more even

distribution of observed t_{min} for Δ*hpnH* relative to wild type, with a 1-3 day shift in the most frequent dpi.

Surprisingly, we also found that the formation of new nodules is periodic, with a new "burst" of nodules

emerging roughly every 18 days (Fig. 3i). This periodicity of nodule emergence appears to be similar

between strains.

While the slight t_{min} delay for Δ*hpnH* is consistent with longer times required to initiate the

symbiosis (*e.g.* root surface colonization, invasion of the root epidermis and cortex, and intracellular

uptake), it is also possible that a delay in t_{min} simply reflects a lower rate of nodule growth immediately

after the first host cell is intracellularly infected. To address this, we compared the calculated value of t_{min}

(defined by V = 0.2 mm^3) to the maximum growth rates and volumes for each nodule (Fig. S9e,f). We did

not find that the subpopulation of nodules with lower growth rates and final volumes than wild type was

more likely to have a later t_{min}, supporting the interpretation that the delay in t_{min} of Δ*hpnH* could be due to

a separate initiation defect, although it is also possible that the growth rates of emergent nodules and

nascent nodule primordia are not strictly related. Interestingly, t_{min} is also not correlated with the period in

which maximum nodule growth occurs, such that later-emerging nodules have similar periods of growth

as nodules formed within a few dpi (Fig. S9g,h). This indicates that although the appearance of new

nodules is restricted to narrow, periodic time frames (Fig. 3i), once a nodule has entered its maximum

growth phase, its continued growth is comparatively unconstrained.

To further verify the slight delay in the initiation of the symbiosis for Δ*hpnH* relative to wild type,

we performed competition assays using a standard fluorescence labeling approach. We first generated

Δ*hpnH* and wild-type strains expressing chromosomally-integrated fluorescent proteins, and then we co-

inoculated *A. afraspera* with different ratios of these two strains. As control experiments, we also co-

inoculated each tagged strain with its untagged counterpart, in order to determine the effect of fluorescent

protein overexpression on each strain's competitiveness. After >40 dpi we measured the size of nodules

on plants inoculated with each strain combination and ratio, then sectioned and fixed nodules for imaging.

Although we expected each nodule to contain a clonal population based on previous work[25,27], the

majority of nodules appeared to contain a mixture of both strains (Fig. 5a; Fig. S15-S19). We instead

quantified the relative abundance of each strain in each nodule by fluorescence imaging; in our control

experiments, in which only one fluorophore-expressing strain was present, a DNA dye (either SYTO9 for

mCherry-expressing bacteria or PI, which stains all cells in fixed samples, for GFP-expressing bacteria)

was used to label all bacteria.

Both WT-YFP and Δ$hpnH$-mCherry were significantly out-competed by their corresponding

untagged strains, with higher proportions of tagged strains correlating with lower bacterial DNA

abundance and smaller nodule and/or infection zone sizes (Fig. 4b-c; Fig. S21-22). Additionally, plants

co-inoculated with untagged-Δ$hpnH$ and Δ$hpnH$-mCherry were significantly shorter than plants inoculated

with untagged-Δ$hpnH$ only, suggesting Δ$hpnH$-mCherry is symbiotically defective (Fig. S20). These

effects of fluorophore overexpression made it difficult to interpret our WT-YFP and Δ$hpnH$-mCherry

competition data (Fig. S23), so we developed an alternative, antibiotics-based method to study the timing

of early symbiotic initiation.

First we sought antibiotics that were effective against *B. diazoefficiens* but would minimally affect

A. afraspera growth. We selected three antibiotics (100 µg/ml streptomycin, 100 µg/ml kanamycin, and 20

µg/ml tetracycline) and treated non-inoculated plants with these antibiotics for two weeks, alone and in

combination. After this treatment, we found that neither kanamycin nor streptomycin, nor the combination

of the two, significantly affected plant appearance, shoot height, or root and shoot dry masses compared

to untreated controls (Fig. S24). Plants treated with tetracycline were noticeably more yellow in color,

indicating chlorosis, and the roots and plant medium became brown; these plants also had lower shoot

and root dry masses than untreated plants (Fig. S24).

Next, because the Δ$hpnH$ strain is more sensitive to antibiotics than wild type[11], we tested

various concentrations of the non-plant-perturbing antibiotics streptomycin and kanamycin to identify

concentrations that would result in the same rates of cell death for both strains. We inoculated plant

growth media with wild type or Δ$hpnH$ to the same concentrations and under the same environmental

conditions as in plant inoculation experiments. The wild-type culture was supplemented with 100 µg/ml

streptomycin plus 100 µg/ml kanamycin, and ΔhpnH cultures were supplemented with decreasing

concentrations of these antibiotics: 75, 50 and 25 µg/mL each. Samples of the cultures were then

collected, serially diluted and added to PSY plates to estimate colony-forming units (cfus) per mL over

time. At 50 µg/mL kanamycin plus 50 µg/mL streptomycin, the rate of decrease in cfus/mL for ΔhpnH was

equivalent to that of wild type treated with 100 µg/ml kanamycin plus streptomycin (Fig. 5d).

Finally, we inoculated 40 plants each with wild type or ΔhpnH and added streptomycin or

kanamycin to 100 µg/mL each or 50 µg/mL each, respectively, at various points post-inoculation. After 40

days we counted the number of nodules per plant, and found that antibiotics were able to block nodule

formation over a ~50% longer window in ΔhpnH compared to wild type (Fig. 5e). The decrease in nodules

formed at different antibiotic treatment time points was also evident in the overall appearance of the

plants (Fig. 5f). These results suggest that ΔhpnH requires more time on average to reach the

intracellular stage of the symbiosis, at which point we presume that the bacteria are protected from

antibiotic by the host cells. These data are consistent with ΔhpnH requiring more time to colonize the root

surface, invade the root epidermis, and/or be internalized by host cells.

Extended hopanoids support surface attachment and motility

Because we found that expression of genetic tags in wild type and ΔhpnH perturbed their

symbiosis with *A. afraspera*, and therefore we could not confidently track these strains *in planta*, we used

an *in vitro* approach to study two steps in the initiation of the symbiosis: (1) bacterial motility toward or

along the *A. afraspera* root, and (2) stable attachment of bacteria to the root surface[20]. To determine

whether ΔhpnH is less motile than wild type, we inoculated low-agar, PSY plates with ΔhpnH or wild type

and measured the rate of zone of swimming over time. We observed that diameter of motility was

reduced in ΔhpnH compared to wild type (Fig. 6a-b), consistent with a swimming motility defect; however,

because we have previously shown that ΔhpnH grows more slowly in this medium than wild type[11], we

could not rule out the possibility that slower zone expansion simply reflects a longer doubling time.

To investigate the nature of the plate motility defect, we studied the motility of single *B.*

diazoefficiens cells. We inoculated cells that have been incubated in PSY liquid medium into a glass

bottom steril flow cell with 100µL of each strain and recorded the trajectories of single cells near the glass

surface at 5 ms time resolution. The cells were tracked using an in-house tracking software in

MATLAB,[34] in which actively swimming cells are defined as having super-diffusive motion and a

trajectory radius of gyration greater than 2.5 µm, and we calculated mean swimming speeds for each cell

using a moving window average over the cell's full trajectory (see Methods for details). In agreement with

our low-agar swimming plates, we found a significantly lower ($p < 0.0001$) fraction of motile cells for

$\Delta hpnH$, $N_{\Delta hpnH}$ = 65 ± 29, compared to wild type, N_{wt}= 368 ± 60, in PSY medium (Fig. 6c), although the

average mean-speed among motile cells were similar between strains, $<V>_{\Delta hpnH}$ = 24.83 ± 7.0 µm/sec

and $<V>_{wt}$ = 22.75 ± 6.7 µm/sec (Fig. 6d). Because the composition of PSY differs greatly from that of the

plant growth medium (BNM), we repeated these assays in BNM supplemented with arabinose. Under this

condition, we again observed a lower fraction of motile $\Delta hpnH$ cells than wild type, ($N_{\Delta hpnH}$ = 54 ± 59, N_{wt}=

450 ± 310) (Fig. 6e), with similar mean speeds between strains and across media conditions (Fig. 6d,f).

These results are consistent with a delay in the initiation of the $\Delta hpnH$ symbiosis with *A. afraspera* due to

a lower occurrence of motile bacterial cells, indicating that the effect of extended hopanoids is likely

regulatory and upstream of the workings of the flagellum.

We next tested the surface attachment capabilities of $\Delta hpnH$ and wild type by incubating dense

bacterial cultures on glass coverslips and quantifying the fraction of the surface covered with stably

adhered cells after two hours. In PSY medium, both strains adhered poorly, and there was no significant

difference in their attachment efficiencies (Fig 6g; Fig. S25). In BNM supplemented with arabinose, both

strains adhered to glass better than in PSY, and $\Delta hpnH$ attachment levels were significantly lower than

wild type (Fig. 6g; Fig. S25). The origin of the differences in surface attachment between PSY and BNM is

unclear. While this result may indicate that the surface attachment machinery of *B. diazoefficiens*

functions better in BNM, we cannot exclude the interpretation that because PSY is more rich than BNM,

peptides and other components of PSY are more likely to interact with the glass surface to occlude or

disfavor bacterial attachment. Regardless, the lower adherence of $\Delta hpnH$ in BNM data suggest that

$\Delta hpnH$ may adhere to the *A. afraspera* root surface less efficiently than wild type, and that this also may

contribute to the delay in the initiation of the symbiosis.

Discussion

Hopanoids are well-established mediators of bacterial survival under stress, and previously we

showed that the capacity for hopanoid production is enriched in plant-associated environments[13] and

required for optimal *Bradyrhizobia-Aeschynomene* spp. symbioses[10,11]. Here we performed a detailed,

quantitative evaluation of the extended hopanoids' roles in the *Bradyrhizobium diazoefficiens-*

Aeschynomene afraspera symbiosis. Our results provide evidence that extended hopanoids' effects are

pleiotropic and span the stages of root nodule biogenesis. We identify specific extended-hopanoid

dependent processes that will provide a foundation for more detailed mechanistic work (Fig. 7). Our

description of the kinetics of nodule growth in *A. afraspera*, both for single nodules and across the root

system, more generally suggests new hypotheses about the cell and developmental biology of nodule

formation in this organism.

By tracking the development of individual root nodules, we found that the initial emergence of root

nodules is delayed in plants inoculated with extended hopanoid mutants. *In vitro*, extended hopanoids

appear to support both the efficiency of surface attachment and the proportion of motile cells in plant

growth medium, suggesting that delayed root nodule emergence may result from fewer cells swimming

and attaching to the root surface in the initial bacteria-host contact (Fig. 7a-b). At later stages in root

nodule development, roughly a third of nodules infected with extended hopanoid mutants grow at slower

rates than with wild type and consequently exhibit low volumes when fully mature. Many of these small

nodules contain reduced densities of symbionts throughout their infection zones; extended hopanoid

mutant nodules with wild-type nodule volumes can also have lower symbiont loads, due to infection zone

fragmentation. The origin of this under-infection phenotype is unclear from our experiments. One

explanation is that host cells in the nodule primordium either internalize fewer cells from the extracellular

environment, or that primordial symbiosomes duplicate or are segregated less effectively, and this

phenotype is simply propagated as nodules develop (Fig. 7d). Alternatively, low symbiont densities may

reflect symbiont degradation in a previously fully-infected nodule (Fig. 7e).

What mechanisms might underpin these extended hopanoid mutant phenotypes? Through a

biophysical lens, the effects of extended hopanoid loss in the *B. diazoefficiens-A. afraspera* symbiosis

could be viewed purely as consequences of less rigid *B. diazoefficiens* membranes. The fraction of motile

cells in *E. coli* populations has been suggested to be sensitive to changes to the mechanical properties of

the outer membrane[35], and membrane-based mechanotransduction is required by diverse bacteria to

stimulate extracellular matrix production and cement their attachment to surfaces[36,37]. Subsequent

progression of bacteria into the root interior through small (micron- to submicron-diameter) channels may

also be hindered by a less rigid *B. diazoefficiens* membrane, as growth rates under similarly confined

spaces *in vitro* have been shown to be positively correlated with membrane rigidity in *E. coli* and *B.*

subtilis models[38,39]. Though not much is known about the mechanical requirements of formation and

propagation of the symbiosome, bacterial membrane rigidity may be relevant in this context as well. For

example, the maintenance of single-bacteroid-containing symbiosomes within host cells (presumably

requiring coordination of bacteroid division with division of the surrounding host-derived membrane)

implies a mechanical linkage between the bacteroid and peribacteroid membranes, which may be

sensitive to changes in bacteroid membrane rigidity. NCR peptide-directed reprogramming of bacteroids

could also be less effective in *B. diazoefficiens* with less rigid membranes, increasing bacteroid

permeability to peptides or peptide-induced bacteroid lysis. While intriguing, these mechanics-based roles

for extended hopanoids are difficult to verify given how little information is available on the biophysical

aspects of legume-microbe interactions. Future work in this area would enrich our understanding of how

intracellular bacterial infections operate in native multicellular hosts.

Other potential extended hopanoid-mediated effects on the symbiosis include regulation of

membrane-based processes, such as motility, oxidative phosphorylation, the early legume-rhizobia

dialogue or transport across the symbiosome membranes. Specifically, the loss of HoLA from the outer

membrane in the *B. diazoefficiens* Δ*hpnH* mutant is likely to disrupt processes involving lipid A, which has

previously been shown to enhance bacterial fitness in *A. afraspera* hosts[40]. Identifying the enzyme(s)

responsible for the attachment of hopanoids to lipid A will be necessary to determine how much HoLA

contributes to the Δ*hpnH* phenotypes we observed in this work, as well as the universality of these

phenotypes to other host-bacteria interactions. While many hopanoid-producing bacteria are host-

associated, so far HoLA has only been found in the *Bradyrhizobiaceae*.

Regardless of the underlying mechanisms for extended hopanoids, it is curious that their absence

is not a death knell for the *B. diazoefficiens-A. afraspera* symbiosis at any stage. In our *in vitro* studies,

although fewer motile cells were present in the extended hopanoid mutants, mean speeds among motile

ΔhpnH cells were indistinguishable from wild type. While we cannot rule out more subtle defects in the

direction of the movement or in chemotaxis, the identical mean speeds suggest that once motility of

ΔhpnH cells is induced, their motility systems function properly. Similarly *in planta*, ΔhpnH nodules

developing at wild-type rates and reaching average wild-type volumes did occur – and, in the case of

"mega" nodules, some exceeded their wild-type counterparts. Why do ΔhpnH populations form two

distinct populations (wild-type-like or defective) rather than falling on a continuous distribution of

behavior? Bimodality in a population can reflect a switch-like, or threshold-based, regulation, and perhaps

in the ΔhpnH strain, a fraction of cells cannot support levels of signaling or biochemical events above the

appropriate threshold for proper function. Populations within ΔhpnH also may differ in the extent to which

extended hopanoid loss is compensated. In *Methylobacterium extorquens* and *Rhodopseudomonas*

palustris[41,42], hopanoid loss results in upregulation of other membrane-rigidifying lipids including

carotenoids and cardiolipins, and in other plant-microbe systems, lipid exchange between hosts and

microbes has been observed[43], suggesting that ΔhpnH mutant success at a given nodulation site may

relate to the local availability of exogenous plant steroids or structurally similar metabolites.

Beyond hopanoids, our results provide insight into the developmental control of nodule formation

by *A. afraspera* hosts. Although the time of emergence, growth rate and window of growth of any specific

nodule appears to be stochastic, the extent to which nodulation is permitted across the entire root system

is tightly determined. Host control of nodule numbers, known as the autoregulation of nodulation (AON),

is a well-established phenomenon[44], and a new, expanded model of AON derived from studies of *Lotus*

japonicum has recently been published[45]. In this model, the root-specific TOO MUCH LOVE (TML)

protein is a key negative regulator of nodule formation. In the absence of rhizobia, a microRNA (miR2111)

produced in plant leaves translocates to roots to suppress TML, poising uninfected plants in a nodulation-

susceptible ("ON") state. When rhizobia begin to infect the plants, a negative feedback loop is initiated,

beginning with the local production of CLE-family peptides. These CLE peptides then translocate to the

shoot and activate a receptor protein at the shoot apex known as HAR1. HAR1-mediated signaling blocks

the miR2111 shoot-root translocation, de-repressing TML to create a nodulation-resistant ("OFF") state.

The general features of this AON pathway are conserved across legumes (reviewed in [44]), and

although AON has not been specifically investigated in *A. afraspera*, our results suggest a similar

mechanism is at play. Our observation that the wild-type *B. diazoefficiens*-*A. afraspera* symbiosis is

characterized by an initial 2-3 day burst of nodulation followed by a sharp drop-off is consistent with an

initial "ON" state that is inhibited by a negative feedback loop with a several-day activation time. What is

most striking about nodule formation in *A. afraspera* is that nodulation-susceptible "ON" states appear to

be reactivated at fixed 18-day intervals, and that re-activation is unrelated to systemic nitrogen fixation

efficiency: at the end of the first 18 dpi cycle, Δ*hpnH* nodules fix considerably less nitrogen than wild type,

yet the timing and distribution of nodulation in the second cycle is unchanged. This both suggests that

there is a fixed oscillation between nodulation-susceptible and nodulation-resistant states in *A. afraspera*,

and that the timing of the oscillation is determined by the dynamics of the underlying regulatory

mechanisms (*e.g.* the rates of synthesis, activation/inhibition, and translocation of specific regulatory

molecules). Thus, we predict that a systems-level analysis of regulatory factor dynamics will be key to

elucidating the nodulation re-activation pathway, as will a more detailed analysis of nodulation timing in

other, model legumes. Specifically, it will be interesting to determine whether re-activation occurs simply

by dampening of the negative feedback pathway as inhibitory molecules reach the end of their lifetimes,

or if it involves an additional positive feedback mechanism that simply requires a longer activation time.

Since the systemic control of nodulation is insensitive to nitrogen fixation, it is possible that *A.*

afrapera plants maximize overall nitrogen fixation through the local regulation of nodule organogenesis. It

is already known that nitrogen-based signaling can locally regulate nodule formation in other legume

hosts (reviewed in [44]), albeit in the context of nodule growth inhibition by increased soil nitrogen

availability. We have found that the key variable that determines the final size of a nodule (and thus it

potential for symbiotic productivity) is the rate of nodule growth and, given that rates of growth are lower

in the less-productive Δ*hpnH* strain than wild-type, this could be coupled to the nodule's fixed nitrogen

output. Alternately, nodule growth rates may relate to the density and survival of symbionts present,

perhaps by indirect sensing of bacteroid-produced signals. An intriguing possibility is that bacteroids

secrete mitogenic compounds, such as phytohormones, to directly influence host cell division rates. *B.*

diazoefficiens has been shown to synthesize the majority of known phytohormones in pure culture[46],

and in *Frankia* symbionts, extended hopanoid levels may be correlated with amount of the auxin

phytohormome phenyl-acetic acid (PAA)[47]. It has also been shown that *B.* ORS285 mutants that cannot

produce cytokinin phytohormones exhibit slower nodulation kinetics and smaller nodule sizes in

Aeschynomene indica hosts[48]. The full extent to which primordia and infection zone growth are

dependent on bacteroid-secreted mitogens or nitrogen fixation productivity, and through what

mechanisms, will be an active area of future research.

On a technical level, our results underscore the importance of identifying the most informative,

least perturbing tools for interrogating legume-microbe symbiosis. Employing quantitative single-nodule

and single-cell approaches rather than bulk measurements were essential for uncovering the diverse

phenotypes of the *B. diazoefficiens* extended hopanoid mutants and yielded unexpected information on

regulation of nodule development by *A. afraspera*. We have also shown the limitations of introducing

overexpressed genetic tags into bacteria. While use of these tags has undoubtedly enhanced our

understanding of legume-microbe symbiosis[49], they may not fully capture the behavior of native

organisms, and in this work we provide a model for evaluating the extent of these genetic tags' effects.

Additionally, our work is one of many to emphasize the importance of appropriate culture models for

mimicking the host environment, as the Δ*hpnH* surface attachment defect was observed in plant growth

medium but not in a standard richer medium. Recent work has suggested that culturing rhizobia in the

latter is correlated with an increase in mutations of symbiosis-associated genes[50], raising questions

about the utility of rich medium cultures in symbiosis work overall. A more detailed analysis of the host

environment, including the full milieu of root exudates[51], available carbon sources[52] and trace metals

specific to each legume, will improve *in vitro* models of legume-bacteria interactions.

Finally, beyond opening up future work on the mechanism of hopanoids in motility and

attachment, this work highlights aspects of the development of *A. afraspera* symbioses most in need of

more detailed analysis. Key questions raised in this study include how relevant the biophysical

environment of the root invasion is for the mode and the efficiency of progressive bacterial invasion; how

is a high density of symbiosomes established and maintained within infected nodule cells; what factors

regulate the mitotic index of a nodule, and how dynamic are these indices; and how the answers to all of

these questions change in the presence of competition between multiple strains with different symbiotic

nitrogen-fixation capacities. Addressing these questions in the *Aeschynomene-Bradyrhizobia* symbiosis

will expand our knowledge of the diversity of approaches to establishing persistent, intracellular

infections, and perhaps provide a better "blueprint" for engineering novel beneficial plant-microbe

associations. These future studies may also prove to be ecologically significant, as African

Aeschynomene spp. are native to regions with the world's poorest food security and nitrogen fertilizer

accessibility and thus in greatest need of better symbiotic nitrogen fixation[53], and *Bradyrhizobia* are

dominant in harsh, warm soils predicted to be more prevalent in a changing global climate[54].

Methods

B. diazoefficiens *culture and strain generation*

B. diazoefficiens hopanoid biosynthesis mutants were generated previously[11]. For construction

of YFP- and mCherry-expressing strains, fluorophore expression vectors pRJPaph-YFP and pRJPaph-

mCherry[27] were provided as a gift from Prof. Dr. Hans-Martin Fischer (ETH Zurich). These vectors were

introduced into *B. diazoefficiens* by conjugation with the β2155 DAP auxotroph strain of *E.coli*, using the

following protocol: *B. diazoefficiens* wild type and Δ*hpnH* were grown in 5 mL PSY medium[55] at 30°C

and 250 rpm to an OD_{600} of ~1.0 (wild type) or of 0.5-0.8 (Δ*hpnH*). β2155 strains carrying pRJPaph

vectors were grown to an OD_{600} of 0.5-0.8 in 5 mL LB supplemented with 10 µg/mL tetracycline and 300

µm DAP at 37°C and 250 rpm. Both *B. diazoefficiens* and β2155 donor cultures were pelleted at 3250 x g

for 30 minutes, washed three times in 0.9% sterile saline, and resuspended in 0.9% sterile saline to a

final OD_{600} of 1.0. *B. diazoefficiens* strains and β2155 donor cells were combined at a 4:1 ratio,

respectively, and mixed by repeated pipetting. Aliquots (50 µl) of these 4:1 mixtures were dropped to PSY

plates supplemented with 300 µm DAP, dried in a biosafety cabinet, and incubated for 48 hours at 30°C.

Conjugation pastes were then removed from plates and resuspended in 5 mL sterile saline, pelleted at

3250 x g for 30 minutes and washed twice, in order to remove residual DAP. Washed cells were pelleted

a final time and resuspended to 200 µl in 0.9% sterile saline and plated onto PSY plates supplemented

with 20 µg/mL (wild type) or 10 µg/mL (Δ*hpnH*) tetracycline. Colonies appeared after 7-10 days (wild type)

or 10-14 days (Δ*hpnH*) and were streaked onto fresh PSY/tetracycline plates, then screened for

fluorescence using a Lumascope 720 fluorescent microscope (Etaluma). Fluorescent colonies were then

sequenced to verify insertion of the pRJPaph vectors into the *scoI* locus.

A. afraspera *cultivation and inoculation with* B. diazoefficiens

      A. afraspera seeds were obtained as a gift from the laboratory of Dr. Eric Giraud (LSTM/Cirad,

Montpelier, France). Seeds were sterilized and scarified by incubation in 95% sulfuric acid at RT for 45

minutes, followed by 5 washes in sterile-filtered nanopure water and a second incubation in 95% ethanol

for 5 minutes at RT. After ethanol treatment seeds were washed 5X and incubated overnight in sterile-

filtered nanopure water. Seeds were then transferred to freshly poured water/agar plates using sterile,

single-use forceps in a biosafety cabinet, and germinated for 24-72 hours in the dark at 28-32°C.

      Seedlings were then placed in clear glass test tubes containing 100 mL of sterile, nitrogen-free

Buffered Nodulation Medium (BNM)[56] and grown for 7-10 days in plant growth chambers (Percival)

under the following settings: 28°C, 80-90% humidity, and 16 hour photoperiod under photosynthetic light

bulbs (General Electric) emitting ~4000 lumens/ft^2. In parallel, *B. diazoefficiens* strains were grown in 5-10

mL PSY liquid culture at 30°C and 250 rpm to stationary phase (OD_{600} > 1.4). Stationary phase cultures

were diluted into PSY one day prior to plant inoculation to reach an OD_{600} of ~0.8 at the time of

inoculation. OD_{600} ~ 0.8 cultures were pelleted at 3250 x g for 30 minutes at RT, washed once in PSY,

then resuspended in PSY to a final OD_{600} of 1.0. Resuspended *B. diazoefficiens* cultures were directly

inoculated into the plant medium in a sterile biosafety cabinet; 1 mL of OD_{600}=1.0 culture was added per

plant. Inoculated plants were then returned to growth chambers and maintained for the times indicated for

each experiment. For longer experiments (lasting longer than ~30 days post-inoculation), plant growth

tubes were refilled with sterile-filered nanopure water as needed. To minimize cross-contamination,

inoculated plants and non-inoculated plants were cultivated in separate growth chambers, and growth

chambers were sterilized with 70% ethanol followed by UV irradiation for at least 24 hours between

experiments.

Acetylene reduction experiments

Individual plants were transferred to clear glass 150 mL Balch-type anaerobic culture bottles containing

15 mL BNM medium and sealed under a gas-tight septum. After sealing, 15 mL of headspace gas (10%

of the culture bottle volume) was removed and replaced with 15 mL of acetylene gas (Airgas). Plants in

culture bottles were incubated in the light at 28°C in growth chambers for 3-6 hours. A 100 µl sample of

the headspace gas was removed using a gas-tight syringe (Hamilton), and this sample was injected and

analyzed for ethylene signal intensities using a Hewlett Packard 5890 Series II GC with Hewlett Packard

5972 Mass Spectrometer with a 30mx0.320mm GasPro Column (Agilent Technologies) and a 2 mm ID

splitless liner (Restek Corporation). Following acetylene reduction measurements, plants were removed

from jars and plant shoot heights and number of nodules per plant were recorded. When nodule dry mass

measurements were performed, nodules were harvested with a razor blade, transferred into pre-weighed

Eppendorf tubes, dried at 50°C for a minimum of 48 hours, then weighed again.

Live:Dead staining and imaging of nodule cross-sections

Nodules were hand-sectioned with razor blades and immediately transferred into a fresh solution

of 5 µM SYTO9 (diluted 1:100 from a 500 uM stock in DMSO at -20°C; Thermo Fisher) and 0.02 mg/mL

(30 µM) propidium iodide (diluted 1:50 from a 1 mg/mL stock stored in water at 4°C; Thermo Fisher) in

PBS. Nodule sections were incubated in this SYTO9/propidium iodide solution at room temperature for 30

minutes in the dark with gentle shaking, washed 5X in PBS, and fixed in 4% paraformaldehyde (Electron

Microscopy Sciences) in PBS overnight in the dark at 4°C. Fixed sections were washed 5X in PBS and

transferred to a freshly prepared solution of 0.1 mg/mL Calcofluor White (Fluorescence Brightener 28;

Sigma) in PBS. The sections were incubated in the Calcofluor solution in the dark for 1 hour at RT with

gentle shaking and washed 5X in PBS to remove excess dye.

Prior to imaging, sections were transferred to 30 mm imaging dishes with 20 mm, #0 coverglass

bottoms (MatTek) and overlaid with sterile 50% glycerol. Nodule images were collected on a Leica TCS

SPE laser-scanning confocal (model DMI4000B-CS) using a 10X/0.3 NA APO ACS objective and solid-

state laser lines for fluorophore excitation at the following settings for each dye: Calcofluor, 405 nm

excitation/410-500 nm emission; SYTO9, 488 nm excitation/510-570 nm emission; PI, 532 nm

excitation/600-650 nm emission. These images were then processed to enhance brightness and contrast

in FIJI[57,58].

Nodule diameter and volume measurements

Inoculated *A. afraspera* root nodules were imaging using a high-definition Keyence VHX-600

digital microscope at 20X magnification. For end-point root nodule volume measurements at 40 days

post-inoculation, plants were removed from the growth chamber and imaged at RT on paper towels, then

discarded. Nodule diameters were measured using the line tool in FIJI and recorded using a custom FIJI

macro. For tracking nodule volumes over time, plants were serially removed from their growth chambers

and transferred to a plastic dish containing 150 mL of sterile BNM pre-warmed to 28°C. Images of

sections of the plant root were collected serially from the hypocotyl to the root tip. Following collection of

images, plants were immediately returned to their original growth tubes in the growth chamber. Plastic

dishes were sterilized for 10 minutes in 10% bleach, washed three times in sterile-filtered nanopure

water, sprayed with 70% ethanol/water, and air-dried before each new plant was imaged. A fresh aliquot

of sterile, pre-warmed BNM also was used for each plant. After the time course was completed, images of

entire plant root systems were reconstructed by eye for each plant at each time point. For nodules

appearing in at least five time points, nodule diameters were measured as described for the end-point

measurements and were converted to approximate volumes in R using the equation $V = 4/3\pi r^3$.

Nodule growth curve fitting and analysis

All analyses of nodule growth, and corresponding plots, were generated in R. For nodule growth

curve fitting, three model equations were used to identify the best fit, as follows:

(1) exponential function:

$$V = ae^{-bt} + c$$

620   (2) quadratic function:

$$V = at^2 + bt + c$$

622   (3) generalized logistic function (expressed as a Richard's function with a time shift):

$$V = \frac{a}{\left(1 + e^{-b(t-c)}\right)^{\left(\frac{1}{d}\right)}}$$

Calculation of the optimal parameter values for each equation (e.g. the values of **a**, **b**, **c**, and **d**) and the

standard error for each curve compared to the raw data were performed using the built-in function *nlm()* in

R. In some cases, *nlm()* could not produce a best-fit model without specifying initial values for the function

parameters. For exponential models, an equation of best fit could be successfully determined without

specification of initial values for parameters **a**, **b** and **c**. For quadratic models, initial parameter values

were required and were set to **a**=0, **b**=10 and **c**=0 for each nodule plot, after identifying these initial

parameter values as broadly optimal based on an initial parameter sweep of -50 to 50 for each plot. For

sigmoidal models, no broadly optimal initial values could be identified, so a parameter sweep was

performed for each plot with the initial value of **a** set to the maximum observed nodule volume (as **a**

describes the upper asymptote of the sigmoidal curve), **b** ranging from 0.1 to 1, **c** ranging from 0 to 10,

and **d** ranging from 0.01 to 1.0. In the sigmoidal plots, an initial point of (0,0) was added to the nodule

volume time series to improve fitting.

Because the sigmoidal model provided the best fits, extrapolation of nodule growth characteristics

was performed on sigmoidal models only. The maximum nodule volume, $\mathbf{V_{max}}$, is defined as the upper

asymptote of the sigmoidal growth curve, *e.g.* **a**. The nodule initiation time, $\mathbf{t_{min}}$, was defined in three

separate ways: the times at which the nodule volume is equal to 0.05, 0.1, or 0.2 mm^3 (*e.g.* through

solving 0.05, 0.1, or 0.2 = $\mathbf{a}/((1+e^{(-\mathbf{b}(t-\mathbf{c}))})^{(1/\mathbf{d})})$ for *t*). The maximum nodule growth rate, **dV/d***t*, was defined

as the average rate of growth (*e.g.* slope) between the time at which the volume is 10% of $\mathbf{V_{max}}$ and the

time at which the volume is 90% of $\mathbf{V_{max}}$. The time at which each nodule reaches its maximum size, $\mathbf{t_{max}}$,

was approximated as the time at which the volume is 90% of $\mathbf{V_{max}}$, since the "true" maximum volume is

asymptotic to the growth curve and is therefore never fully reached in the model.

Competition assays

mCherry-tagged Δ*hpnH* and YFP-tagged wild type *B. diazoefficiens* were grown to stationary

phase (OD$_{600}$ > 1.4) in 10 mL PSY cultures supplemented with 20 µg/mL (wild type) or 10 µg/mL (Δ*hpnH*)

tetracycline; untagged strains were grown in PSY. On the day prior to inoculation, all strains were diluted

into 50-150 mL tetracycline-free PSY to reach an OD_{600} of ~0.8 at the time of inoculation. *A. afraspera*

plants were cultivated pre-inoculation in test tubes as described above, with the addition of covering the

growth tubes in foil to minimize the production of chlorophyll in the plant roots, which spectrally overlaps

with mCherry. At the time of inoculation, all cultures were pelleted at 3250 x g for 30 minutes at RT,

washed three times, then resuspended in PSY to a final OD_{600} of 1.0. A 10 mL culture of each strain ratio

for inoculation was generated a sterile 15mL Falcon tube; for example, for a 50:50 mixture of mCherry-

tagged Δ*hpnH* and YFP-tagged wild type, 5 mL of each strain was combined. These cultures were mixed

thoroughly by gentle pipetting, and 1 mL of the mixtures was added to directly to the plant medium for 7-8

plants per strain mixture.

After 45-60 days, plants were harvested. First, plant heights and the number of nodules per plant

were recorded. Then, the roots were cut from the stem and images of all nodules for each plant were

collected on a high-definition Keyence VHX-600 digital microscope at 20X magnification. These nodules

were then cross-sectioned and immediately transferred to Eppendorfs containing 4% paraformaldehyde

(Electron Microscopy Sciences) in PBS. Fresh sections were fixed overnight in the dark at 4°C, washed

5X in PBS, and stored in PBS supplemented with 0.1% azide in the dark at 4°C until imaging.

Fixed sections were stained in Calcofluor (all strain combinations), SYTO9 (WT-YFP and WT co-

inoculation only) or propidium iodide (mCherry-Δ*hpnH* and Δ*hpnH* co-inoculation only) as described for

Live:Dead staining. Imaging was performed as described for Live:Dead staining using a 5X objective.

Given the high autofluorescence of these nodules and low mCherry and YFP signal intensities, the

following excitation/emission settings were used: Calcofluor, 405 nm excitation/410-460 nm emission;

YFP/SYTO9, 488 nm excitation/500-550 nm emisasion; mCherry, 532 nm excitation/600-650 nm

emission.

Quantification of nodule statistics (including nodule and infection zone areas, signal intensity of

YFP, mCherry, SYTO9 and propidium iodide) was performed on raw images using a custom FIJI macro.

Briefly, nodule images were opened at random, infection zones (IZs) and whole nodules were circled by

hand and saved as discrete regions of interest (ROIs), and the area and intensity in each channel were

measured automatically for all ROIs. These measurements were exported as a text table and various

parameters from these measurements were calculated using custom Python scripts, as indicated in the

Results. Plots of all parameters and statistical comparisons were generated using custom R scripts.

Antibiotic treatment of inoculated plants

A. afraspera plants were cultivated as described above and the following antibiotics were added

to non-inoculated plants 7 days after rooting in 100 mL BNM growth tubes: kanamycin to 100 µg/mL,

streptomycin to 100 µg/mL, tetracycline to 20 µg/mL, kanamycin plus tetracycline, kanamycin plus

streptomycin, streptomycin plus tetracycline. Plants were grown in antibiotics under normal plant growth

conditions for 14 days, after which plants were visually inspected. Plant heights were also recorded, and

the root and shoot systems were separated with a razor blade, transferred into pre-weighed 15 mL Falcon

tubes, dried at 50°C for a minimum of 48 hours, then weighed again.

Antibiotic treatments of Δ*hpnH* and wild-type *B. diazoefficiens* were performed by growing

antibiotic 5 mL PSY cultures of each strain to stationary phase (OD_{600} >1.4) and diluting strains in fresh

PSY to reach an OD_{600} of ~0.8 at the time of antibiotic treatment – *e.g.* as they would be grown prior to

plant inoculation. Cultures were pelleted at 3250 x g for 30 minutes at RT, washed three times, then

resuspended in PSY to a final OD_{600} of 1.0. Four 100 µl aliquots of these culture were diluted 1:00 into

separate 10 mL BNM cultures in clear glass tubes in plant growth chambers. Kanamycin (at 25, 50, 75,

and 100 µg/mL) and streptomycin (at 25, 50, 75, and 100 µg/mL) were added directly to the BNM

cultures, and 100 µl samples were taken immediately prior to antibiotic treatment and at 2, 4, 6, 8, and 10

hours post-antibiotic addition. These 100 µl samples were immediately diluted 1:10 in 900 µl and mixed

vigorously by repeated pipetting. Vortexing was avoided as we found that this method reduces Δ*hpnH*

viability. Ten serial 1:10 dilutions were performed, and three 10 µl samples of each dilution for each strain

were spotted and dripped across PSY plates. After 7 days (wild type) or 10 days (Δ*hpnH*), colonies were

counted manually and recorded for each dilution exhibiting discrete colonies. Log plots of colony counts

over time were generated in R.

Plants were then inoculated with Δ*hpnH* and wild-type *B. diazoefficiens* as described above, and

kanamycin and streptomycin were added to Δ*hpnH*-inoculated plants to 50 µg/mL each, and to wild type-

inoculated plants to 100 µg/mL at 12 hours and 36 hours and at 2, 2.5, 3, 3.5, 4, 4.5, 5, 6.5, 8.5, 9.5, 10.5,

and 12.5 days post-inoculation. Four plants were treated per time point per strain, with an additional four

plants each as an untreated control. At 40 dpi, the number of nodules per plant was recorded.

Bulk motility assays

Swimming motility assays were performed as previously described, with some modifications

(Althabegoiti et al. 2008). WT and Δ*hpnH* were grown to turbidity in 5 mL of PSY at 30°C and 250 rpm,

diluted to an OD_{600} of 0.02 in 5 mL of fresh PSY, and grown to exponential phase (OD_{600} = 0.3-0.5).

Exponential cultures then were diluted to an OD_{600} of 0.06 in fresh PSY and 2 µL of the adjusted cultures

into the center of swimming plate containing 0.3% agar/PSY. After inoculation, the plates were wrapped

with parafilm to prevent dehydration and incubated in a humidity-controlled environmental chamber

(Percival) at 30°C for 10 days total, with daily scans after 5 days. The resulting images were analyzed in

FIJI to measure the area of the swimming colony.

Surface attachment assays

Δ*hpnH* and wild-type *B. diazoefficiens* were grown in 5 mL PSY cultures to stationary phase

(OD_{600} >1.4) then diluted in fresh PSY to reach an OD_{600} of ~0.8 at the time of surface attachment

assays. Cultures were pelleted at 3250 x g for 30 minutes at RT, washed twice in the indicated

attachment medium, then resuspended in attachment medium to an OD_{600} of 1.0. These cultures were

mixed thoroughly by repeated pipetting, and 2 mL samples were added to sterile imaging dishes (30 mm

dishes with 20 mm, #1.5 coverglass bottoms; MatTek). Cultures were incubated on imaging dishes

without shaking at 30°C for two hours. To remove non-adhered cells, imaging dishes were immersed in

50 mL of attachment media in a 100 mL glass beaker on an orbital shaker and shaken gently at RT for 5

minutes; direct application of washing medium to the coverglass surface was avoided, as we found that

this creates a shear force sufficient to wash away adhered cells. Imaging dishes were then gently lifted

out of the washing medium and imaged with a 100X objective on a Lumascope 720 fluorescence

microscope (Etaluma). Forty fields of view were recorded for each strain and media combination. These

images were processed in FIJI using the Enhanced Local Contrast (CLAHE) plugin[59] and converted

into a binary image to determine the area of the imaging window covered with adhered cells. Calculation

of the fraction of the surface was performed in Excel and statistical analyses were conducted in R. Areas

of the surface containing groups of cells larger than 10 μm^2 in area were ignored in the calculations, as

these likely do not represent true attachment events rather than sedimentation of larger cell clumps. BNM

used for attachment assays was prepared as described above, with the addition of 1.0 g/mL arabinose.

Because BNM contains salt crystals that can sediment onto coverglass and occlude or obscure adhered

cells, this medium was passed through a 2 μm filter (Millipore) prior to the attachment experiments.

Single-cell motility assays and analysis

B. diazoefficiens wild-type and ΔhpnH were grown in 12.5 ml PSY medium at 30°C and 200 rpm

to an OD_{600} = 0.6-0.8 from an AG medium plate culture. Then, a 1:10 dilution of cell culture was

subcultured in PSY medium to a final volume of 12.5 ml and regrown to an OD_{600} of ~0.6. Two aliquots of

750 µL were sampled from the regrowth culture and pelleted at 3500 x g for 20 min (wild-type) or for 30

min (ΔhpnH) at RT. The supernatant was removed, and one pellet was resuspended in 500 µL PSY and

the other in 500 µL BNM medium. Because BNM contains salt crystals that can sediment onto coverglass

and occlude or obscure adhered cells, this medium was passed through a 2 µm filter (Millipore) prior to

usage for these experiments. The two medium conditions were then incubated for 2.5 hrs (wild-type) or

for 3.5 hrs (ΔhpnH) at 30°C; given the difference in growth time ΔhpnH incubated for longer. Right before

imaging, each culture was diluted at a 1:10 ratio with its respective medium. The bacteria were then

injected into a sterile flow cell (ibidi sticky-Slide VI0.4 with a glass coverslip). The flow cell was attached to

a heating stage set to 30°C.

The imaging protocol involved high-speed bright-field imaging for 5 min at a single XYZ location

per experimental repeat. High speed bright-field recordings used a Phantom V12.1 high speed camera

(Vision Research); images were taken with a 5 ms exposure at 200 fps and a resolution of 512×512

pixels (0.1 µm/pixel). This protocol was performed on an Olympus IX83 microscope equipped with a 100×

oil objective, a 2× multiplier lens, and a Zero Drift Correction autofocus system. The recorded movies

were extracted into single frames from the .cine files using PCC 2.8 (Phantom Software). Image

processing and cell tracking algorithms are adapted from previous work[34] and written in MATLAB

R2015a (Mathworks).

We identified cells swimming near the surface as cells with a trajectory radius of gyration greater

than 2.5 µm and a mean-squared displacement (MSD) slope greater than 1.5. Setting a minimum radius

of gyration selects for cells with a minimum net translation on the across the surface, while a minimum

MSD slope threshold ensured the cells are moving super-diffusively (MSD slope $\cong$ 1, diffusive motion;

MSD slope $\cong$ 2, super-diffusive motion). For each tracked cell, the mean-speed, $\mathbf{v}$, was calculated by

averaging a moving window, $\mathbf{w}$, of the displacement over the cell's full trajectory, using the following

equation:

$$< v > = \ Avg \left(\sum_{t=1}^{N-w} \frac{\sqrt{(x_{t+w} \ - \ x_t)^2 \ + \ (y_{t+w} \ - \ y_t)^2}}{w} * f * p \right)$$

where $\mathbf{N}$ is the total number of points in the trajectory, $\mathbf{f}$ is the acquisition frame rate, and $\mathbf{p}$ is the pixel

resolution. Here we set a window size, w= 40 frames. All analysis and visualizations from these

experiments where done using MATLAB R2015a (Mathworks).

Acknowledgements

This work was supported by grants from the HHMI (D.K.N.), NASA

(NNX12AD93G, D.K.N.), Jane Coffin Childs Memorial Fund (B.J.B.), NIH (K99GM126141,

B.J.B.), and Ford Foundation Predoctoral Fellowship (J.d.A.), and Army Research Office (W911NF-18-1-

0254, GW). We thank Dr. Eric Giraud for his generous gift of *A. afraspera* seeds and training on

Aeschynomene symbioses and Drs. Hans-Martin Fischer and Raphael Ledermann for plasmids and

technical advice for the genetic transformation of *B. diazoefficiens*. Dr. Nathan Dalleska of the

Environmental Analysis Center at Caltech was instrumental in providing training and support for GC-MS

analysis of acetylene reduction. We are grateful to Dr. Gargi Kulkarni and other members of the Newman

lab, as well as Drs. Elliot Meyerowitz and Rob Phillips, for their collegiality and thoughtful discussions

about this work. We are indebted to Ms. Shannon Park and Ms. Kristy Nguyen for providing the

administrative assistance that allows us to focus on our research.

References

1.   Belin BJ, Busset N, Giraud E, Molinaro A, Silipo A, Newman DK. Hopanoid lipids: from

membranes to plant-bacteria interactions. Nat Rev Microbiol. England; 2018;16: 304–315.

doi:10.1038/nrmicro.2017.173

2.   Saenz JP, Grosser D, Bradley AS, Lagny TJ, Lavrynenko O, Broda M, et al. Hopanoids as

functional analogues of cholesterol in bacterial membranes. Proc Natl Acad Sci U S A. United

States; 2015;112: 11971–11976. doi:10.1073/pnas.1515607112

3.   Wu C-H, Bialecka-Fornal M, Newman DK. Methylation at the C-2 position of hopanoids increases

rigidity in native bacterial membranes. Elife. England; 2015;4. doi:10.7554/eLife.05663

4.   Fantini J, Barrantes FJ. How cholesterol interacts with membrane proteins: an exploration of

cholesterol-binding sites including CRAC, CARC, and tilted domains. Front Physiol. Frontiers;

2013;4: 31. doi:10.3389/fphys.2013.00031

5.   Ochs D, Kaletta C, Entian K, Beck-sickinger A, Porallal K, Chemie O, et al. Cloning, Expression,

and Sequencing of Squalene-Hopene Cyclase,. 1992;174: 298–302.

6.   Syren P-O, Henche S, Eichler A, Nestl BM, Hauer B. Squalene-hopene cyclases-evolution,

dynamics and catalytic scope. Curr Opin Struct Biol. England; 2016;41: 73–82.

doi:10.1016/j.sbi.2016.05.019

7.   Welander P V, Coleman ML, Sessions AL, Summons RE, Newman DK. Identification of a

methylase required for 2-methylhopanoid production and implications for the interpretation of

sedimentary hopanes. Proc Natl Acad Sci U S A. United States; 2010;107: 8537–8542.

doi:10.1073/pnas.0912949107

8.   Welander P V, Doughty DM, Wu C, Mehay S, Roger E. Identification and characterization of

Rhodopseudomonas palustris TIE-1 hopanoid biosynthesis mutants. 2012;10: 163–177.

9.   Schmerk CL, Welander P V, Hamad MA, Bain KL, Bernards MA, Summons RE, et al. Elucidation

of the Burkholderia cenocepacia hopanoid biosynthesis pathway uncovers functions for conserved

proteins in hopanoid-producing bacteria. Environ Microbiol. England; 2015;17: 735–750.

doi:10.1111/1462-2920.12509

10.   Silipo A, Vitiello G, Gully D, Sturiale L, Chaintreuil C, Fardoux J, et al. Covalently linked hopanoid-

lipid A improves outer-membrane resistance of a Bradyrhizobium symbiont of legumes. Nat

Commun. England; 2014;5: 5106. doi:10.1038/ncomms6106

11.   Kulkarni G, Busset N, Molinaro A, Gargani D, Chaintreuil C, Silipo A, et al. Specific hopanoid

classes differentially affect free-living and symbiotic states of Bradyrhizobium diazoefficiens.

MBio. United States; 2015;6: e01251-15. doi:10.1128/mBio.01251-15

12.   Komaniecka I, Choma A, Mazur A, Duda KA, Lindner B, Schwudke D, et al. Occurrence of an

unusual hopanoid-containing lipid A among lipopolysaccharides from Bradyrhizobium species. J

Biol Chem. United States; 2014;289: 35644–35655. doi:10.1074/jbc.M114.614529

13.   Ricci JN, Coleman ML, Welander P V., Sessions AL, Summons RE, Spear JR, et al. Diverse

capacity for 2-methylhopanoid production correlates with a specific ecological niche. ISME J.

Nature Publishing Group; 2014;8: 675–684. doi:10.1038/ismej.2013.191

14.   Ourisson G, Albrecht P, Rohmer M. The Hopanoids: palaeochemistry and biochemistry of a group

of natural products. Pure Appl Chem. 1979;51: 709–729. doi:10.1351/pac197951040709

15.   Racolta S, Juhl PB, Sirim D, Pleiss J. The triterpene cyclase protein family: a systematic analysis.

Proteins. United States; 2012;80: 2009–2019. doi:10.1002/prot.24089

16.   Rodríguez-Navarro DN, Oliver IM, Contreras MA, Ruiz-Sainz JE. Soybean interactions with soil

microbes, agronomical and molecular aspects. doi:10.1051/agro/2010023

17.   Vilcheze C, Llopiz P, Neunlist S, Poralla K, Rohmer M. Prokaryotic triterpenoids: New hopanoids

from the nitrogen-fixing bacteria Azotobacter vinelandii, Beijerinckia indica and Beijerinckia

mobilis. Microbiology. 1994;140: 2749–2753. doi:10.1099/00221287-140-10-2749

18.   Parsons R, Silvester WB, Harris S, Gruijters WT, Bullivant S. Frankia vesicles provide inducible

and absolute oxygen protection for nitrogenase. Plant Physiol. United States; 1987;83: 728–731.

19.   Abeysekera RM, Newcomb W, Silvester WB, Torrey JG. A freeze-fracture electron microscopic

study of *Frankia* in root nodules of *Alnus incana* grown at three oxygen tensions. Can J Microbiol.

NRC Research Press Ottawa, Canada ; 1990;36: 97–108. doi:10.1139/m90-019

20.   Wheatley RM, Poole PS. Mechanisms of bacterial attachment to roots. FEMS Microbiol Rev.

England; 2018;42: 448–461. doi:10.1093/femsre/fuy014

21.   Kanbe M, Yagasaki J, Zehner S, Göttfert M, Aizawa S-I. Characterization of two sets of subpolar

flagella in Bradyrhizobium japonicum. J Bacteriol. American Society for Microbiology Journals;

2007;189: 1083–9. doi:10.1128/JB.01405-06

22.   Quelas JI, Althabegoiti MJ, Jimenez-Sanchez C, Melgarejo AA, Marconi VI, Mongiardini EJ, et al.

Swimming performance of Bradyrhizobium diazoefficiens is an emergent property of its two

flagellar systems OPEN. 2016; doi:10.1038/srep23841

23.   Althabegoiti MJ, Covelli JM, Pérez-Giménez J, Quelas JI, Mongiardini EJ, López MF, et al.

Analysis of the role of the two flagella of Bradyrhizobium japonicum in competition for nodulation of

soybean. FEMS Microbiol Lett. 2011;319: 133–139. doi:10.1111/j.1574-6968.2011.02280.x

24.   López-García SL, Perticari A, Piccinetti C, Ventimiglia L, Arias N, De Battista JJ, et al. In-Furrow

Inoculation and Selection for Higher Motility Enhances the Efficacy of Nodulation. Agron J.

American Society of Agronomy; 2009;101: 357. doi:10.2134/agronj2008.0155x

25.   Bonaldi K, Gargani D, Prin Y, Fardoux J, Gully D, Nouwen N, et al. Nodulation of Aeschynomene

afraspera and A. indica by photosynthetic Bradyrhizobium Sp. strain ORS285: the nod-dependent

versus the nod-independent symbiotic interaction. Mol Plant Microbe Interact. United States;

2011;24: 1359–1371. doi:10.1094/MPMI-04-11-0093

26.   Patriarca EJ, Tatè R, Ferraioli S, Iaccarino M. Organogenesis of Legume Root Nodules. Int Rev

Cytol. Academic Press; 2004;234: 201–262. doi:10.1016/S0074-7696(04)34005-2

27.   Ledermann R, Bartsch I, Remus-Emsermann MN, Vorholt JA, Fischer H-M. Stable Fluorescent

and Enzymatic Tagging of Bradyrhizobium diazoefficiens to Analyze Host-Plant Infection and

Colonization. Mol Plant Microbe Interact. United States; 2015;28: 959–967. doi:10.1094/MPMI-03-

15-0054-TA

28.   Pierre O, Engler G, Hopkins J, Brau F, Boncompagni E, Hérouart D. Peribacteroid space

acidification: A marker of mature bacteroid functioning in Medicago truncatula nodules. Plant, Cell

Environ. 2013;36: 2059–2070. doi:10.1111/pce.12116

29.   Denison RF, Okano Y. Leghaemoglobin oxygenation gradients in alfalfa and yellow sweetclover

nodules. J Exp Bot. 2003;54: 1085–1091. doi:10.1093/jxb/erg097

30.   Kereszt A, Mergaert P, Montiel J, Endre G, Kondorosi E. Impact of Plant Peptides on Symbiotic

Nodule Development and Functioning. Front Plant Sci. Switzerland; 2018;9: 1026.

doi:10.3389/fpls.2018.01026

31.   Glazebrook J, Ichige A, Walker GC. A Rhizobium meliloti homolog of the Escherichia coli peptide-

antibiotic transport protein SbmA is essential for bacteroid development. Genes Dev. Cold Spring

Harbor Laboratory Press; 1993;7: 1485–97. doi:10.1101/GAD.7.8.1485

32.   Szparaga A, Kocira S. Generalized logistic functions in modelling emergence of Brassica napus L.

PLoS One. 2018;13: 1–14. doi:10.1371/journal.pone.0201980

33.   Richards FJ. NA Flexible Growth Function for Empirical Useo Title. J Exp Bot. 1959;10: 290–301.

doi:10.1093/jxb/10.2.290

34.   Lee CK, de Anda J, Baker AE, Bennett RR, Luo Y, Lee EY, et al. Multigenerational memory and

adaptive adhesion in early bacterial biofilm communities. Proc Natl Acad Sci U S A. National

Academy of Sciences; 2018;115: 4471–4476. doi:10.1073/pnas.1720071115

35.   Gupta R, Sharma M, Mittal A. Effects of membrane tension on nanopropeller driven bacterial

motion. J Nanosci Nanotechnol. 2006;6: 3854–3862. doi:10.1166/jnn.2006.630

36.   Petrova OE, Sauer K. Sticky situations: Key components that control bacterial surface attachment.

J Bacteriol. 2012;194: 2413–2425. doi:10.1128/JB.00003-12

37.   Persat A. Bacterial mechanotransduction. Curr Opin Microbiol. Elsevier Ltd; 2017;36: 1–6.

doi:10.1016/j.mib.2016.12.002

38.   Mä J, Driessen R, Galajda P, Keymer JE, Dekker C. Bacterial growth and motility in sub-micron

constrictions [Internet]. 2009. Available: www.pnas.org/cgi/content/full/

39.   Tuson HH, Auer GK, Renner LD, Hasebe M, Tropini C, Salick M, et al. Measuring the stiffness of

bacterial cells from growth rates in hydrogels of tunable elasticity. Mol Microbiol. Wiley/Blackwell

(10.1111); 2012;84: 874–891. doi:10.1111/j.1365-2958.2012.08063.x

40.   Busset N, Di Lorenzo F, Palmigiano A, Sturiale L, Gressent F, Fardoux J, et al. The Very Long

Chain Fatty Acid (C26:25OH) Linked to the Lipid A Is Important for the Fitness of the

Photosynthetic Bradyrhizobium Strain ORS278 and the Establishment of a Successful Symbiosis

with Aeschynomene Legumes. Front Microbiol. Switzerland; 2017;8: 1821.

doi:10.3389/fmicb.2017.01821

41.   Bradley AS, Swanson PK, Muller EEL, Bringel F, Caroll SM, Pearson A, et al. Hopanoid-free

Methylobacterium extorquens DM4 overproduces carotenoids and has widespread growth

impairment. PLoS One. United States; 2017;12: e0173323. doi:10.1371/journal.pone.0173323

42.   Neubauer C, Dalleska NF, Cowley ES, Shikuma NJ, Wu C-H, Sessions AL, et al. Lipid remodeling

in Rhodopseudomonas palustris TIE-1 upon loss of hopanoids and hopanoid methylation.

Geobiology. England; 2015;13: 443–453. doi:10.1111/gbi.12143

43.   Keymer A. Cross-kingdom lipid transfer in arbuscular mycorrhiza symbiosis and beyond. Curr

Opin Plant Biol. Elsevier Current Trends; 2018;44: 137–144. doi:10.1016/J.PBI.2018.04.005

44.   Ferguson BJ, Mens C, Hastwell AH, Zhang M, Su H, Jones CH, et al. Legume nodulation: The

host controls the party. Plant Cell Environ. Wiley/Blackwell (10.1111); 2018;

doi:10.1111/pce.13348

45.   Tsikou D, Yan Z, Holt DB, Abel NB, Reid DE, Madsen LH, et al. Systemic control of legume

susceptibility to rhizobial infection by a mobile microRNA. Science (80-). 2018;6907: eaat6907.

doi:10.1126/SCIENCE.AAT6907

46.   Boiero L, Perrig D, Masciarelli O, Penna C, Cassán F, Luna V. Phytohormone production by three

strains of Bradyrhizobium japonicum and possible physiological and technological implications.

Appl Microbiol Biotechnol. 2007;74: 874–880. doi:10.1007/s00253-006-0731-9

47.   Hammad Y, Nalin R, Marechal J, Fiasson K, Pepin R, Berry AM, et al. A possible role for phenyl

acetic acid (PAA) on Alnus glutinosa nodulation by Frankia. Plant Soil. Kluwer Academic

Publishers; 2003;254: 193–205. doi:10.1023/A:1024971417777

48.   Podlešáková K, Fardoux J, Patrel D, Bonaldi K, Novák O, Strnad M, et al. Rhizobial Synthesized

Cytokinins Contribute to But Are Not Essential for the Symbiotic Interaction Between

Photosynthetic Bradyrhizobia and *Aeschynomene* Legumes. Mol Plant-Microbe Interact. The

American Phytopathological Society ; 2013;26: 1232–1238. doi:10.1094/MPMI-03-13-0076-R

49.   Ledermann R, Bartsch I, Muller B, Wulser J, Fischer H-M. A Functional General Stress Response

926   of Bradyrhizobium diazoefficiens Is Required for Early Stages of Host Plant Infection. Mol Plant

927   Microbe Interact. United States; 2018;31: 537–547. doi:10.1094/MPMI-11-17-0284-R

928 50. Burghardt LT, Epstein B, Guhlin J, Nelson MS, Taylor MR, Young ND, et al. Select and

929   resequence reveals relative fitness of bacteria in symbiotic and free-living environments. Proc Natl

930   Acad Sci U S A. National Academy of Sciences; 2018;115: 2425–2430.

931   doi:10.1073/pnas.1714246115

932 51. Sugiyama A, Yazaki K. Root Exudates of Legume Plants and Their Involvement in Interactions

933   with Soil Microbes. doi:10.1007/978-3-642-23047-9_2

934 52. Pini F, East AK, Appia-Ayme C, Tomek J, Karunakaran R, Mendoza-Suarez M, et al. Bacterial

935   Biosensors for in Vivo Spatiotemporal Mapping of Root Secretion. Plant Physiol. United States;

936   2017;174: 1289–1306. doi:10.1104/pp.16.01302

937 53. FAO. Regional Overview of Food Security and Nutrition in Africa 2017. The food security and

938   nutrition–conflict nexus: building resilience for food security, nutrition and peace [Internet]. 2017.

939   Available: www.fao.org/publications

940 54. Delgado-Baquerizo M, Oliverio AM, Brewer TE, Benavent-Gonzalez A, Eldridge DJ, Bardgett RD,

941   et al. A global atlas of the dominant bacteria found in soil. Science. United States; 2018;359: 320–

942   325. doi:10.1126/science.aap9516

943 55. Regensburger B, Hennecke H. RNA polymerase from Rhizobium japonicum. Arch Microbiol.

944   Springer-Verlag; 1983;135: 103–109. doi:10.1007/BF00408017

945 56. Ehrhardt DW, Atkinson EM, Long SR. Depolarization of alfalfa root hair membrane potential by

946   Rhizobium meliloti Nod factors. Science. American Association for the Advancement of Science;

947   1992;256: 998–1000. doi:10.1126/SCIENCE.10744524

948 57. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: An open-

949   source platform for biological-image analysis [Internet]. Nature Methods. 2012. pp. 676–682.

950   doi:10.1038/nmeth.2019

951 58. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of Image Analysis HHS

952   Public Access [Internet]. Nat Methods. 2012. Available:

953   https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5554542/pdf/nihms876304.pdf

59.   Heckbert PS, Karel. Graphics gems IV [Internet]. Graphics gems IV. AP Professional; 1994.

Available: https://dl.acm.org/citation.cfm?id=180940

Figure Legends

Figure 1. (**a**) Chemical structure of the extended hopanoid 2-Methyl Bacteriohopanetetrol (2Me-BHT),

consisting of a central pentacyclic core synthesized by the *shc* gene product, a C2 methylation site added

by the product of *hpnP* (grey shading, left), and a tetrol group added by the *hpnH* product (grey shading,

right). (**b**) Average shoot heights and nodules per plant at 24 dpi for *A. afraspera* plants inoculated with

wild-type, Δ*hpnH* or Δ*hpnP B. diazoefficiens.* (**c**) Average acetylene reduction per plant and per nodule at

24 dpi for *A. asfrapera* plants inoculated with wild-type, Δ*hpnH* or Δ*hpnP*. (**d**) Representative confocal

images of cross-sections of wild type- and Δ*hpnH*-infected nodules at 24 dpi illustrating plant cell walls

(Calcofluor, cyan), live bacteria (SYTO9, yellow) and membrane-compromised bacteria and plant nuclei

(propidium iodide, magenta). (**e**) Average nodule dry mass and acetylene reduction per nodule dry mass

at 24 dpi for plants inoculated with wild-type, Δ*hpnH* or Δ*hpnP*. Data shown in (**b**), (**c**) and (**e**) was

collected from n = 8 plants, with error bars representing one standard deviation. Results of two-tailed t-

tests between wild type and Δ*hpnH* or Δ*hpnP* are denoted as follows: n.s., $p>0.01$; *****, $p<0.01$; ******, $p<0.001$;

***, $p<0.0001$.

Figure 2. (**a**) Average acetylene reduction per plant (n=4 plants per bar) and (**b**) average nodule dry

mass per plant (n=8 plants per bar) for *A. asfrapera* inoculated with wild-type or Δ*hpnH* over time. Error

bars representing one standard deviation. Results of two-tailed t-tests between wild type and Δ*hpnH* are

denoted as follows: n.s., $p>0.05$; *****, $p<0.05$; *******, $p<0.0001$. (**c-d**) *A. afraspera* inoculated with wild type or

Δ*hpnH* at (**c**) 20 dpi (left) and at (**d**) 40 dpi (right). (**e-f**) Distributions of nodule diameters at 40 dpi for *A.*

afraspera inoculated with (**e**) Δ*hpnH* (right; n=268 nodules pooled from 10 plants) or (**f**) wild type (left;

n=227 nodules pooled from 10 plants).

Figure 3. (**a**) Comparison of the development of selected wild type- and Δ*hpnH*-infected nodules over

time. (**b**) Nodule growth plots for 74 wild type-infected nodules tracked from 10 plants. (**c**) Nodule growth

plots for 84 Δ*hpnH*-infected nodules tracked from 16 plants. (**d**) Schematic of nodule development in *A.*

afraspera. From the left, bacteria (in blue) colonize and invade plant roots (green) and intracellularly infect

a root cell (pink); the time of this initial intracellular infection is considered t_i and the nodule volume can be

described as the volume of the single infected root cell, V_i. This infected cell proliferates to form a

spherical nodule that is visible to the naked eye, at time t_{min} and volume V_{min}. The infected plant cells

continue to proliferate at rate **dV/dt** until the nodule has fully matured at time t_{max} and volume V_{max}. (**e**)

Fitted growth curve for a sample wild-type nodule illustrating the positions of t_{min}, V_{min}, **dV/dt**, t_{max}, and

V_{max}. (**f-g**) Jitter and box plots of (**f**) **dV/dt** and (**g**) V_{max} values for all wild type- and Δ*hpnH*-infected

nodules. Results of KS-tests between wild-type and Δ*hpnH* nodules are denoted as follows: ***, $p<10^{-6}$.

(**h**) Scatter plots of **dV/dt** vs. V_{max} values for wild-type and Δ*hpnH* nodules. Values of **dV/dt** and V_{max}

below what is observed in the wild-type dataset are highlighted in green. (**i**) Distributions of t_{min} values (as

observed by eye) for nodules from wild type- (white bars) or Δ*hpnH*- (grey bars) infected plants. N=457

wild-type nodules across 20 plants and 479 Δ*hpnH* nodules across 20 plants.

Figure 4. (**a**) Confocal sections of small (<0.5 mm radius) Δ*hpnH*-infected nodules harvested at 40 dpi

and small (<0.5 mm radius) wild type-infected nodules harvested at 10 and 25 dpi. (**b**) Confocal sections

of larger (>0.5 mm radius) Δ*hpnH*- or wild type-infected nodules harvested at 40 dpi.

1000

Figure 5. (**a**) Confocal sections of nodules from plants co-inoculated with wild type-YFP and Δ*hpnH*-

mCherry harvested at 45-55 dpi. Sections were stained with Calcofluor (cyan) and are expressing YFP

(yellow) and mCherry (magenta). (**b**) Scatter plot of median YFP intensity per pixel normalized by

propidium iodide intensity per pixel (*e.g.* bacteroid density) within infection zones of nodules from plants

co-inoculated with wild type-YFP and wild type, as a function of the percentage of wild type-YFP in the

inoculum. (**c**) Scatter plot of median propidium iodide intensity per pixel (*e.g.* bacteroid density) within

infection zones of nodules from plants co-inoculated with YFP-tagged wild type and untagged wild type,

as a function of the percentage of WT-YFP in the inoculum. (**d**) Colony forming units/mL in wild type and

Δ*hpnH* cultures grown in BNM supplemented with varying concentrations of kanamycin and

spectinomycin at various times post-inoculation. (**e**) Average nodules per plant at 40 dpi for plants

inoculated with either wild type or Δ*hpnH* and treated with 50 µg/mL (Δ*hpnH*) or 100 µg/mL (wild type)

kanamycin and streptomycin at various time points post-inoculation. Nodule counts are normalized to

those observed in non-antibiotic treated plants. (**f**) Images of inoculated plants at 40 dpi after antibiotic

treatment at various time points. Untreated plants are shown on the left, with increasing time of antibiotic

addition. Error bars represent one standard deviation.

1016

Figure 6. (**a**) Sample time course of wild type and Δ*hpnH* colony expansion on low-agar PSY plates (dpi

= days post-inoculation). Scale bars represent 2 cm. (**b**) Average colony sizes of wild type and Δ*hpnH*

over time. N=4 plates per strain; error bars indicate one standard deviation. (**c**) Trajectories of individual

wild type (top) and Δ*hpnH* (bottom) cells over a 5 minute time course in PSY. (**d**) Distributions of mean-

speed s for motile wild type (N=359) and Δ*hpnH* (N=91) cells for trajectories in **d**. Dotted lines indicate the

means of the distributions. (**e**) Trajectories of individual wild type (top) and Δ*hpnH* (bottom) cells over a 5

minute time course in BNM. (**f**) Distributions of mean-speeds for motile WT (N=421) and Δ*hpnH* (N=141)

cells in BNM for trajectories in **e**. Dotted lines indicate the means of the distributions. (g) Jitter and box

plots of surface attachment (*e.g.* the percent of the field of view covered with cells) of WT and Δ*hpnH*

after 2 hours of incubation on glass in PSY or BNM. N=40 fields of view per condition. Results of two-

tailed t-tests between wild type and Δ*hpnH* are denoted as follows: n.s., p>0.05; *****, p<0.00001.

1028

Figure 7. Schematic representation of *A. afraspera* wild-type root nodule development (top row; white

background) and defects in development associated with extended hopanoid loss (bottom row; grey

background). Early in development, fewer Δ*hpnH* cells are motile (**a**) and competent to attach to root

surfaces (**b**), leading to a delay in establishment of stable root colonies. At later stages, slow growth of

Δ*hpnH* into the root interior, or poor uptake by and division within host cells (**c**) may generate "patchy", or

under-populated infection zone that is propagated as the nodule grows (**d**). Alternately, fully-infected

Δ*hpnH* nodules may lose symbionts to symbiont cell death (**e**) via poor bacteroid survival or plant-directed

symbiosome degradation.

    Supplemental Figures

    Figure S1. Confocal images of cross-sections of wild type-infected *A. afraspera* nodules at 24 dpi

    illustrating plant cell walls (Calcofluor, cyan), live bacteria (SYTO9, yellow) and membrane-compromised

    bacteria and plant nuclei (propidium iodide, magenta). Nodules were collected from 3 plants.

    Figure S2. Confocal images of cross-sections of Δ*hpnH*-infected *A. afraspera* nodules at 24 dpi

    illustrating plant cell walls (Calcofluor, cyan), live bacteria (SYTO9, yellow) and dead bacteria and plant

    nuclei (propidium iodide, magenta). Nodules were collected from 3 plants. White boxes highlight small

    nodules. White arrow indicates a likely plant defense reaction.

    Figure S3. Average (**a**) shoot height, (**b**) nodules per plant, (**c**) nodule dry weight per plant, (**d**) acetylene

    reduction per plant, (**e**) acetylene reduction per nodule, and (**f**) acetylene reduction per nodule dry weight

    for *A. asfrapera* inoculated with wild-type or Δ*hpnH* at 40 dpi. N=4 plants per bar; error bars represent one

    standard deviation. Results of two-tailed t-tests between wild type and Δ*hpnH* are denoted as follows:

    n.s., p>0.05.

    Figure S4. Reconstructed images of the root system of a wild type-infected *A. afraspera* plant. Nodules

    fully visible in at least five time points are indicated with black arrowheads.

    Figure S5. Reconstructed images of the root system of a Δ*hpnH*-infected *A. afraspera* plant. Nodules

    fully visible in at least five time points are indicated with black arrowheads.

    Figure S6. Nodule growth plots for all 74 wild type-infected nodules fit with quadratic (orange; long

    dashed lines), exponential (yellow; short dashed lines), or sigmoidal (blue; solid lines) models. Standard

    errors (SE) for each model are shown.

1065   **Figure S7.** Nodule growth plots for all 84 Δ*hpnH*-infected nodules fit with quadratic (orange; long dashed

1066   lines), exponential (yellow; short dashed lines), or sigmoidal (blue; solid lines) models. Standard errors

1067   (SE) for each model are shown.

1068

Figure S8. (**a**) Jitter and box plots of t_{max} values for all wild type- and Δ*hpnH*-infected nodules. (**b**) Jitter

and box plots of maximum growth windows for all wild type- and Δ*hpnH*-infected nodules. (**c**) Jitter and

box plots of t_{min} values (as determined by extrapolation using sigmoidal fits of nodule growth curves) for

all wild type- and Δ*hpnH*-infected nodules, in which V_{min} is defined as 0.05 mm^3, 0.1 mm^3, 0.2 mm^3.

Green shading highlights negative t_{min} values. Results of KS-tests between wild-type and Δ*hpnH* nodules

are denoted as follows: *, $p<0.05$; n.s., $p>0.05$.

1075

1076   **Figure S9.** (**a-b**) Scatter plots of t_{max} vs. (**a**) **dV/dt** and (**b**) V_{max} for all wild type- (open circles) and

1077   Δ*hpnH*- (grey circles) infected nodules. Green regions highlight values below what is observed for wild

1078   type. (**c-d**) Scatter plots of maximum growth windows vs. (**c**) **dV/dt** and (**d**) V_{max}. (**e-f**) Scatter plots of t_{min}

1079   vs. (**c**) **dV/dt** and (**d**) V_{max}. (**g-h**) Scatter plots of t_{min} vs. (**a**) t_{max} and (**b**) maximum growth windows.

1080

1081   **Figure S10**. Confocal sections of small (<0.5 mm radius) Δ*hpnH*-infected nodules harvested at 40 dpi.

1082   Sections were stained with Calcofluor (cyan), SYTO9 (yellow), and propidium iodide (magenta). N=74

1083   nodules harvested from 5 plants. White boxes highlight under-infected nodules. Magenta boxes indicate

1084   nodules primarily containing membrane-compromised cells.

1085

1086   **Figure S11**. Confocal sections of large (>0.5 mm radius) Δ*hpnH*-infected nodules harvested at 40 dpi.

1087   Sections were stained with Calcofluor (cyan), SYTO9 (yellow), and propidium iodide (magenta). N=87

1088   nodules harvested from 5 plants. White boxes highlight under-infected nodules.

1089

1090   **Figure S12**. Confocal sections of small (<0.5 mm radius) wild type-infected nodules harvested at 10 dpi.

1091   Sections were stained with Calcofluor (cyan), SYTO9 (yellow), and propidium iodide (magenta). N=80

1092   nodules harvested from 5 plants. White boxes highlight under-infected nodules.

1093

1094 **Figure S13**. Confocal sections of small (<0.5 mm radius) wild type-infected nodules harvested at 25 dpi.

1095 Sections were stained with Calcofluor (cyan), SYTO9 (yellow), and propidium iodide (magenta). N=82

1096 nodules harvested from 5 plants. Magenta boxes indicate nodules primarily containing membrane-

1097 compromised cells.

1098

1099 **Figure S14**. Confocal sections of wild type-infected nodules harvested at 40 dpi. Sections were stained

1100 with Calcofluor (cyan), SYTO9 (yellow), and propidium iodide (magenta). N=117 nodules harvested from

1101 5 plants. Magenta boxes indicate nodules primarily containing membrane-compromised cells.

1102

1103 **Figure S15**. Confocal sections of nodules from plants co-inoculated with 10% WT-YFP:90% Δ*hpnH*-

1104 mCherry. Nodules were harvested between 45 and 50 dpi. Sections were stained with Calcofluor (cyan)

1105 and express YFP (yellow) mCherry (magenta). N=132 nodules harvested from 8 plants.

1106

1107 **Figure S16**. Confocal sections of nodules from plants co-inoculated with 25% WT-YFP:75% Δ*hpnH*-

1108 mCherry. Nodules were harvested between 45 and 50 dpi. Sections were stained with Calcofluor (cyan)

1109 and express YFP (yellow) mCherry (magenta). N=125 nodules harvested from 7 plants.

1110

1111 **Figure S17**. Confocal sections of nodules from plants co-inoculated with 50% WT-YFP:50% Δ*hpnH*-

1112 mCherry. Nodules were harvested between 45 and 50 dpi. Sections were stained with Calcofluor (cyan)

1113 and express YFP (yellow) mCherry (magenta). N=143 nodules harvested from 7 plants.

1114

1115 **Figure S18**. Confocal sections of nodules from plants co-inoculated with 75% WT-YFP:25% Δ*hpnH*-

1116 mCherry. Nodules were harvested between 45 and 50 dpi. Sections were stained with Calcofluor (cyan)

1117 and express YFP (yellow) mCherry (magenta). N=143 nodules harvested from 8 plants.

1118

1119 **Figure S19**. Confocal sections of nodules from plants co-inoculated with 90% WT-YFP:10% Δ*hpnH*-

1120 mCherry. Nodules were harvested between 45 and 50 dpi. Sections were stained with Calcofluor (cyan)

1121 and express YFP (yellow) mCherry (magenta). N=110 nodules harvested from 7 plants.

1122

Figure S20. Average shoot height (**a**) and number of nodules (**b**) for plants co-inoculated with Δ*hpnH*-

mCherry and WT-YFP strains, recorded at 45 dpi. Average shoot height (**c**) and number of nodules (**d**) for

plants co-inoculated with WT and WT-YFP strains, recorded at 40 dpi. Average shoot height (**e**) and

number of nodules (**f**) for plants co-inoculated with Δ*hpnH* and Δ*hpnH*-mCherry strains, recorded at 50

dpi. N=7-8 plants per bar for all panels. Error bars represent one standard deviation. Results of two-tailed

t-tests are denoted as follows: n.s., $p>0.05$; ***, $p<0.0001$.

1129

Figure S21. (**a**-**d**) Intensity ratio of YFP to mCherry (**a**), mCherry intensity (**b**), and YFP intensity (**c**) per

pixel within infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and WT-YFP strains. (**d**)

Cross-sectional area of infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and WT-YFP

strains. For (**a**-**d**), N=132, 125, 143, 143 and 110 nodules for 10%, 25%, 50%, 75% and 90% WT-YFP

strain mixtures, respectively, which were sectioned and fixed between 45-50 dpi. (**e**) Nodule volume

distributions from plants co-inoculated with Δ*hpnH*-mCherry and WT-YFP strains at 45 dpi. Sample sizes

are N = 251, 200, 227, 204, and 149 nodules pooled from N = 8, 7, 7, 8, and 7 plants for the 10%, 25%,

50%, 75% and 90% WT-YFP strain mixtures, respectively. (**f**) Scatter plots of mCherry vs. YFP intensities

per pixel within infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and WT-YFP strains. (**g**-**h**)

Scatter plots of YFP/mCherry intensity ratios per pixel in infection zones vs. infection zone (**g**) and nodule

(**h**) cross-section areas for nodules co-inoculated with Δ*hpnH*-mCherry and WT-YFP strains. Scatter plots

contain data pooled from all ratios.

1142

Figure S22. (**a**-**d**) Intensity ratio of YFP to propidium iodide (PI) (**a**), PI intensity (**b**), and YFP intensity (**c**)

per pixel within infection zones of nodules co-inoculated with WT and WT-YFP strains. (**d**) Cross-

sectional area of infection zones of nodules co-inoculated with WT and WT-YFP strains. For (**a**-**d**), N =

141, 95, 134, 147, 133, and 167 nodules for 10%, 25%, 50%, 75%, 90% and 100% WT-YFP strain

mixtures, respectively, which were sectioned and fixed between 40-45 dpi. (**e**) Nodule volume

distributions from plants co-inoculated with WT and WT-YFP strains at 40 dpi. Sample sizes are N = 183,

116, 161, 172, 232, and 248 nodules pooled from N = 8, 7, 8, 8, 8, and 8 plants for the 10%, 25%, 50%,

75% and 90% WT-YFP strain mixtures, respectively. (**f**) Scatter plots of PI vs. YFP intensities per pixel

within infection zones of nodules co-inoculated with WT and WT-YFP strains. (**g**-**h**) Scatter plots of

YFP/PI intensity ratios per pixel in infection zones vs. infection zone (**g**) and nodule (**h**) cross-section

areas for nodules co-inoculated with WT and WT-YFP strains. Scatter plots contain data pooled from all

ratios.

1155

Figure S23. (**a**-**d**) Intensity ratio of mCherry to SYTO9 (**a**), SYTO9 intensity (**b**), and mCherry intensity (**c**)

per pixel within infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and Δ*hpnH* strains. (**d**)

Cross-sectional area of infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and Δ*hpnH* strains.

For (**a**-**d**), N = 117, 107, 128, 137, 103 and 50 nodules for 10%, 25%, 50%, 75%, 90% and 100% Δ*hpnH*-

mCherry strain mixtures, respectively, which were sectioned and fixed between 50-55 dpi. (**e**) Nodule

volume distributions from plants co-inoculated with Δ*hpnH*-mCherry and Δ*hpnH* strains at 45 dpi. Sample

sizes are N = 150, 222, 191, 254, 297, and 236 nodules pooled from N = 7, 7, 7, 8, 8, and 8 plants for the

10%, 25%, 50%, 75% and 90% WT-YFP strain mixtures, respectively. (**f**) Scatter plots of mCherry vs.

SYTO9 intensities per pixel within infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and

Δ*hpnH* strains. (**g**-**h**) Scatter plots of mCherry/SYTO9 intensity ratios per pixel in infection zones vs.

infection zone (**g**) and nodule (**h**) cross-section areas for nodules co-inoculated with Δ*hpnH*-mCherry and

Δ*hpnH* strains. Scatter plots contain data pooled from all strain ratios.

1168

1169   **Figure S24**. Average (**a**) shoot height, (**b**) shoot dry mass and (**c**) root dry mass for non-inoculated *A.*

1170   *afraspera* plants grown in BNM supplemented with kanamycin, streptomycin or tetracycline for 2 weeks

1171   under normal growth conditions. N=4 plants per condition; error bars represent one standard deviation.

1172   (**d**-**e**) Images of *A. afraspera* plants after 2 weeks of antibiotic treatment. Asterisks indicate plants grown

1173   in tetracycline-supplemented medium.

1174

1175   **Figure S25**. Surface attachment of wild type (**a,c**) and Δ*hpnH* (**b,d**) incubated on glass coverslips in

1176   various media. For each panel, raw phase images (top row), background-subtracted images (middle row),

1177   and binary images with cells shown in black (bottom row) are shown. Scale bars represent 20 μm.

a.

hpnP   *shc*   *hpnH*

b.

avg. shoot height (cm)

WT   ΔhpnH   ΔhpnP
       **   n.s.

avg. nodules per plant

WT   ΔhpnH   ΔhpnP
      n.s.   n.s.

c.

avg. acetylene reduced per plant (a.u./hr)

WT   ΔhpnH   ΔhpnP
       **   n.s.

avg. acetylene reduced per nodule (a.u./hr)

WT   ΔhpnH   ΔhpnP
       *   n.s.

d.

WT

ΔhpnH

e.

avg. nodule dry mass per plant (mg)

WT   ΔhpnH   ΔhpnP
      ***   n.s.

avg. acetylene reduced per mg nodule (a.u./hr)

WT   ΔhpnH   ΔhpnP
      n.s.   n.s.

Figure 1.

a. avg. acetylene reduced per plant (a.u./hr)

□ WT
■ Δ*hpnH*

*** 16dpi * 24dpi n.s. 32dpi n.s. 40dpi

b. avg. nodule dry mass per plant (mg)

□ WT
■ Δ*hpnH*

* 16dpi *** 24dpi n.s. 32dpi n.s. 40dpi

e. count vs nodule radius (mm)

Δ*hpnH* 40dpi   avg. 0.73 mm

0.0 0.5 1.0 1.5 2.0

f. count vs nodule radius (mm)

WT 40dpi   avg. 0.88 mm

0.0 0.5 1.0 1.5 2.0

c. 20 dpi

5 cm

WT   Δ*hpnH*

d. 40 dpi

5 cm

WT   Δ*hpnH*

Figure 2.

Figure 3.

a.

ΔhpnH 40 dpi

WT 10 dpi

WT 20 dpi

b.

ΔhpnH 40 dpi

WT 40 dpi

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure S1.

Figure S2.

Figure S3.

07 dpi

09 dpi

11 dpi

15 dpi

17 dpi

19 dpi

21 dpi

23dpi

26dpi

29dpi

33dpi

39 dpi

Figure S4.

dpi

dpi

dpi

dpi

dpi

dpi

dpi

23dpi

26dpi

29dpi

33dpi

39 dpi

Figure S5.

Figure S6.

Figure S6 cont.

Figure S7.

Figure S7 cont.

Figure S8.

Figure S9.

Figure S10.

Figure S11.

Figure S12.

Figure S13.

Figure S14.

Figure S15.

Figure S16.

Figure S17.

Figure S18.

Figure S19.

Figure S20.

Figure S21.

Figure S22.

Figure S23.

a.

b.

c.

d.

Figure S24.

a. WT - PSY

f. Δ*hpnH* - PSY

c. WT - BNM

d. Δ*hpnH* - BNM

Figure S25.