Extended hopanoid loss reduces bacterial motility and surface attachment and leads to

heterogeneity in root nodule growth kinetics in a *Bradyrhizobium-Aeschynomene* symbiosis

Belin, B.J.,[1] Tookmanian, E.T.,[2] de Anda, J.,[4] Wong, G. C. L.,[4] Newman, D.K.[1,3]

[1]Division of Biology & Bioengineering, [2]Division of Chemistry & Chemical Engineering, [3]Division of

Geological & Planetary Sciences, California Institute of Technology, Pasadena, CA.

[4]Department of Bioengineering, Department of Chemistry and Biochemistry, and California NanoSystems

Institute, University of California Los Angeles, Los Angeles, CA 90095-1600, USA University of California

Los Angeles, Los Angeles, CA.

Abstract

Hopanoids are steroid-like bacterial lipids that enhance membrane rigidity and promote bacterial growth

under diverse stresses. Hopanoid biosynthesis genes are conserved in nitrogen-fixing plant symbionts,

and we previously found that the extended (C_{35}) class of hopanoids in *Bradyrhizobium diazoefficiens* are

required for efficient symbiotic nitrogen fixation in the tropical legume host *Aeschynomene afraspera*.

Here we demonstrate that the nitrogen fixation defect conferred by extended loss can fully be explained

by a reduction in root nodule sizes rather than per-bacteroid nitrogen fixation levels. Using a single-

nodule tracking approach to track *A. afraspera* nodule development, we provide a quantitative model of

root nodule development in this host, uncovering both the baseline growth parameters for wild-type

nodules and a surprising heterogeneity of extended hopanoid mutant developmental phenotypes. These

phenotypes include a delay in root nodule initiation and presence of a subpopulation of nodules with slow

growth rates and low final volumes, which are correlated with reduced motility and surface attachment *in*

vitro and lower bacteroid densities *in planta*, respectively. This work provides a quantitative reference

point for understanding the phenotypic diversity of ineffective symbionts in *A. afraspera* and identifies

specific developmental stages affected by extended hopanoid loss for future mechanistic work.

Introduction

Hopanoids are steroid-like lipids that support bacterial survival under stress (reviewed in Belin et al. 2018). They are synthesized by the squalene-hopene cyclase (*shc*) family of enzymes (Ochs et al. 1992; Syren et al. 2016), which generate the pentacyclic, C_{30} hopanoid core from squalene. In many organisms, the C_{30} hopanoids can be further modified, including methylation at the C-2 position via the enzyme HpnP (Welander et al. 2010) and addition of a ribose-derived side chain by the enzyme HpnH (Fig. 1a) (Welander et al. 2012). Side chain-containing hopanoids are known collectively as the C_{35} or "extended" hopanoids and commonly include molecules with aminotriol-, polyol-, and adenosyl- side-chain moieties (Schmerk et al. 2015). Organism-specific side chains have also been observed, including a hopanoid-lipid A conjugate known as HoLA (Silipo et al. 2014; Kulkarni et al. 2015; Komaniecka et al. 2014) that so far has only been found in *Bradyrhizobiaceae*.

It is thought that hopanoids primarily promote bacterial survival by rigidifying and decreasing the permeability of membranes (Saenz et al. 2015; Wu et al. 2015), providing a better barrier against external stress. Structurally distinct hopanoids have different capacities to alter the biophysical properties of membranes and can also differ in the degrees of stress resistance they confer (reviewed in Belin et al. 2018). In the *Bradyrhizobia* genus of legume symbionts, hopanoids promote growth of free-living cultures under acid, salt, detergent, antibiotic, and redox stresses (Silipo et al. 2014; Kulkarni et al. 2015), and we previously showed that these stress resistance phenotypes are largely mediated by the extended hopanoid class (Kulkarni et al. 2015).

We also analyzed an extended hopanoid-deficient mutant of *Bradyrhizobium diazoefficiens* USDA110 in symbiosis with two legumes: the native soybean host for this strain and *Aeschynomene afraspera,* the native host of the closely related photosynthetic *Bradyrhizobia*. *A. afraspera* is a flood-tolerant legume from tropical West Africa, where it has been used in rice intercropping systems (Somado et al. 2003) and to accelerate wound healing in traditional medicine (Swapna et al. 2011; Chifundera 2001; Caamal-Fuentes et al. 2015; Lei et al. 2018). We found that extended hopanoid-deficient mutants of *B. diazoefficiens* fixed less nitrogen per nodule in *A. afraspera* than wild type, while this strain did not appear to have a defect in its native soybean host. Microscopy analyses of a small sample of extended hopanoid mutant-infected *A. afraspera* nodules revealed several aberrant cytological phenotypes,

including both nodules containing necrotic signatures, disorganized infection zones, and visible starch

granule accumulation (Kulkarni et al. 2015).

These phenotypes are common signatures of poor symbiont performance, yet the lack of genetic

tools for *A. afraspera*, the limited literature on this host's response to non-cooperators compared to model

plants, and low number of nodules examined made it difficult to determine the underlying cause. While it

has been proposed that hopanoids may enable high rates of symbiotic nitrogen fixation in some hosts by

limiting oxygen diffusion across cell membranes (Vilcheze et al. 1994; Parsons et al. 1987; Abeysekera et

al. 1990), from our previous assays, we could not determine whether the poor symbiotic performance of

extended hopanoid mutants reflects ineffective nitrogen fixation *per se*, or is simply a consequence of

lower general stress resistance. Because we did not observe an extended hopanoid mutant phenotype in

soybean, we instead suggested that the extended hopanoid mutant may not survive exposure to nodule

cysteine-rich (NCR) peptides, which are synthesized by *A. afraspera* (Czernic et al. 2015) but absent in

soybean.

Here, we sought to dissect further the symbiotic phenotypes of *B. diazoefficiens* extended

hopanoid mutants in association with *A. afraspera*. We found that the lower nitrogen fixation of extended

hopanoid mutants can be fully explained by a reduction in root nodule sizes and rhizobial occupancy,

indicating that the underlying defect is unrelated to per-bacteroid nitrogen fixation levels. Using a novel

single-nodule tracking approach to quantify *A. afraspera* nodule development, we uncovered both the

baseline growth parameters for wild-type nodules and a surprising heterogeneity of extended hopanoid

mutant developmental phenotypes. These results challenge the conclusions of our prior study (Kulkarni et

al. 2015) and identify new, potentially hopanoid-dependent stages in the *B. diazoefficiens-A. afraspera*

symbiosis for future mechanistic work. This work also provides a quantitative reference point for

understanding the impact of symbiotically ineffective strains on *A. afraspera* nodule development.

Results

Loss of extended hopanoids results in reduced nodule size

Previously, we observed a symbiotic defect for an extended hopanoid-deficient (Δ*hpnH*) strain of

B. diazoefficiens in association with *A. afraspera* (Kulkarni et al. 2015). To further validate this defect, we

inoculated *A. afraspera* plants with Δ*hpnH* (lacking extended hopanoids)*,* Δ*hpnP* (lacking 2-Me

hopanoids), or wild-type *B. diazoefficiens*. At 24 days post-inoculation (dpi), plants inoculated with Δ*hpnH*

were shorter than wild type-inoculated plants, although both strains produced equivalent numbers of

nodules (Fig. 1b). Δ*hpnH*-inoculated plants also exhibited a roughly 50% decrease in the rate of

acetylene gas reduction compared to wild type-inoculated plants at this time point (Fig. 1c). In contrast,

the Δ*hpnP* mutant was similar to wild type (Fig. 1b-c). These results are consistent with our previous

findings (Kulkarni et al. 2015).

To assess Δ*hpnH* viability within *A. afraspera* nodules, we performed morphological analyses of

nodules using confocal fluorescent microscopy. Fifty-seven wild-type and 67 Δ*hpnH* nodule cross-

sections were stained with a bacterial Live:Dead kit, consisting of the cell-permeable SYTO9 dye (staining

all cells) and propidium iodide (PI) (staining only cells with a compromised membrane). We did not

observe an increase in predominantly PI-stained nodules for Δ*hpnH* compared to wild type (Fig. 1d; Fig.

S1,S2). Signatures of plant necrosis, which we previously associated with Δ*hpnH* when we observed a

smaller number of nodules (Kulkarni et al. 2015), occurred prominently in only 1/67 Δ*hpnH* nodules

examined (Fig. S2).

The most apparent phenotype of Δ*hpnH* nodules was their relatively small size (Fig. S1,S2). We

repeated acetylene reduction assays for wild type- and Δ*hpnH*-inoculated plants and then calculated the

total nodule dry mass for each plant at 24 dpi. We found a decrease in the nodule dry mass per plant for

Δ*hpnH*-inoculated plants that is sufficient to explain the decrease in acetylene reduction rates (Fig. 1e).

This result rules out the possibility that nitrogenase functions ineffectively in the absence of extended

hopanoids due to inactivation by oxygen, as has been suggested in *Frankia* (Vilcheze et al. 1994;

Parsons et al. 1987; Abeysekera et al. 1990), as the per-mg nitrogen fixation rates are not affected by

extended hopanoid loss.

Δ*hpnH nodules are more variable in size than wild-type nodules*

We next measured acetylene reduction per plant across an extended 40 dpi period, and we

observed that the differences in both acetylene reduction rates and nodule dry masses between wild type

and Δ*hpnH* steadily decreased with time (Fig. 2a-b). By 40 dpi the overall symbiotic efficiencies of wild

type and Δ*hpnH* per plant were indistinguishable, in terms of the plants' qualitative appearance (Fig. 2c-d)

as well as their average shoot heights and acetylene reduction rates (Fig. S3). Total nodule counts per

plant also did not differ between wild type and Δ*hpnH* at 40 dpi, indicating that the increase in total nodule

mass reflects growing nodules rather than more frequent nodulation (Fig. S3).

We also measured the radii of individual nodules on ten plants for each strain at 40 dpi (Fig. 2e-f).

Interestingly, although *average* nodule sizes did become similar between strains by this time point (0.73

vs. 0.88 mm average radii), their underlying distributions were markedly distinct. Wild-type nodule radii

appear to form a roughly normal distribution, whereas the Δ*hpnH* nodule radius distribution is bimodal,

consisting of a subpopulation of small nodules with small radii (<0.5 mm) that are rarely observed in wild

type, as well as a second, larger subpopulation that has a similar median radius as wild type but is

skewed towards larger radii (>1.5mm). These data demonstrate that the small-nodule phenotype of

Δ*hpnH* persists throughout a 40 dpi time course, but is compensated by greater size heterogeneity, in

which a handful of "mega" nodules offset smaller nodules over time.

*Δ*hpnH *nodule size heterogeneity reflects variable nodule growth rates*

To better evaluate the possible origins of the Δ*hpnH* nodule size defect, we studied the kinetics of

single nodule development. Beginning one week after inoculation, we collected images of entire plant

roots every 3-5 days up to ~40 days post-inoculation (Fig. S4,S5). From these images, we identified

nodules that were clearly visible (*e.g.* not obscured by lateral roots or more recently emerged nodules) in

at least five time points (Fig. 3a) and measured their radii. We then calculated nodule volumes by

approximating nodules as spheres and plotted the volume of the tracked nodules over time. While we

again observed that many Δ*hpnH* nodules were smaller at 40 dpi than any of the wild type nodules, we

also found that nodule growth was highly variable both within and between strains (Fig. 3b-c).

We then developed a simple framework for quantifying nodule development, in which nodule

growth is defined by the following variables: the time (t_i) of the initial intracellular infection event and the

volume of the nascent nodule (V_i), equivalent to the volume of one infected *A. afraspera* cortical cell; the

time (t_{min}) and volume (V_{min}) at which a clearly visible, spherical nodule has developed; the rate of growth

of a nodule once it has become visible (dV/dt); and the time (t_{max}) and volume (V_{max}) of a nodule when its

growth has stopped (Fig. 3d). To calculate these variables, we fit each nodule's growth over time to three

different growth models: exponential, quadratic, and a generalized logistic (*e.g.* sigmoidal) equation

commonly used for plant growth (Szparaga and Kocira 2018; Richards 1959) (see Methods for complete

details). Sigmoidal models generally provided the best fit to the experimental data, so these models were

used for growth parameter calculation (Fig. 3e; Fig. S6, S7).

The growth rates of Δ*hpnH* nodules were lower on average than wild-type nodules (Fig. 3f), with

roughly a third of tracked nodules exhibiting growth rates lower than observed for wild type (<0.1

mm^3/dpi). A similar fraction of nodules had smaller final volumes than wild type (Fig. 3g). We further found

that the growth rate of a nodule and its maximum size are positively linearly correlated for both strains,

with Pearson coefficients of ~0.64 ($p<10^{-9}$) for wild type and ~0.75 ($p<10^{-15}$) for Δ*hpnH,* and that the

subpopulation of nodules with lower-than-wild-type growth rates and small nodule sizes are the same

(Fig. 3h). We interpret these data to suggest that host proliferation is slower in a subset of nodules

infected with Δ*hpnH,* and that this largely accounts for the low final volume of these nodules.

We also noted that Δ*hpnH* nodule sizes at 40 dpi differed between these single-nodule volume

measurements (Fig. 3g) and our previous 40 dpi end-point measurements of nodule radii (Fig. 2e), in that

we did not observe larger-than-wild-type "mega" nodules in the single-nodule dataset. This discrepancy

likely reflects the smaller sample size in our single-nodule tracking experiments (84 compared to 268 end-

point nodules), and the low frequency of "mega" nodule formation. To verify this, we selected 10,000

random subsets of 84 nodules from the 268 Δ*hpnH* nodules shown in Figure 2e, converted the nodule

radii to volumes, and found that there is no statistically significant difference (p<0.05) between a random

subset of Fig. 2e and the Δ*hpnH* single-nodule tracking data in ~92% (9184/10000) of cases. Thus the

differences in nodule size distributions in Figure 3g and Figure 2e are consistent with sampling error.

We also calculated each nodule's window of maximum growth, defined as the time required for a

nodule to increase from 10% to 90% of its final volume. Neither the time at which a nodule reaches 90%

of its maximum volume, t_{max}, nor the window of maximum growth differs significantly between Δ*hpnH* and

wild type (Fig. S8a-b). The window of maximum growth for each nodule is also uncorrelated with their

final volume or growth rate, indicating that small nodules are not prematurely aborted; rather, their growth

periods are similar to larger nodules (Fig. S9a-d).

To better understand the subpopulation of small, slow-growing Δ*hpnH* nodules, we isolated

nodules with <0.5 mm radius, sectioned and stained them with SYTO9, PI and Calcofluor, and imaged

them with confocal microscopy. We found that while most small Δ*hpnH* nodules contained a single,

continuous infection zone, a large fraction were un- or under-infected with bacteria, often exhibiting

disorganized central infection zones (~37%; 28/75) (Fig. 4a; Fig. S10). Of the fully infected small Δ*hpnH*

nodules, a subset contained primarily PI-stained, likely dead bacterial cells (~25%; 12/47) (Fig. 4a; Fig.

S10). Similar proportions of under-infected nodules or nodules primarily occupied with membrane-

compromised bacteria did not occur in larger Δ*hpnH* nodules harvested at the same time point, although

fragmented infection zones were still common (Fig. 4b; Fig. S11). We also compared the subpopulation of

small Δ*hpnH* nodules at 40 dpi to two wild-type nodule populations: similarly small nodules harvested at

10 and 25 dpi (Fig. 4a; Fig S12; Fig. S13), and nodules harvested at the same 40 dpi time point (Fig. 4b;

Fig. S14). Again, we found that high proportions of under-infected nodules and membrane-compromised

bacteria were unique to the Δ*hpnH* small-nodule subset.

*Δ*hpnH *nodule emergence is delayed*

The "true" beginning of nodule formation is the time when the first *A. afraspera* cortical cell is

infected, t_i (Fig. 3d). However, this initial infection event is not visible at the root surface, and it is difficult

to extrapolate from sigmoidal models in which the growth curves approach the initial volume $V_i \sim 0$ mm^3

asymptotically. As a proxy for t_i, we defined three alternate t_{min} as the times at which nodules reached

three arbitrarily small volumes: $V = 0.05$ mm^3, $V = 0.1$ mm^3, and $V = 0.2$ mm^3. When t_{min} is defined by $V =$

0.05 mm^3 or 0.1 mm^3, t_{min} could not be accurately calculated for all nodules, as the sigmoidal models

sometimes predicted an impossible $t_{min} < 0$ (Fig. S8c). These nodule volumes are also too small to be

seen on the root surface, and we had no experimental means to determine the accuracy of the

calculations in this low-volume regime. When t_{min} is defined by $V = 0.2$ mm^3 (the smallest nodule volume

that we could identify in our single-nodule tracking assays), there is a small but statistically significant

increase for Δ*hpnH* relative to wild type (Fig. S8c).

To independently verify this delay in nodule emergence, we inspected the roots of 20 wild type-

and 20 Δ*hpnH*-inoculated plants over 40 dpi and recorded the number of visible nodules per plant each

day. We found a more even distribution of observed t_{min} for $\Delta hpnH$ relative to wild type, with a 1-3 day

shift in the most frequent dpi. Surprisingly, we also found that the formation of new nodules is periodic,

with a new "burst" of nodules emerging roughly every 18 days (Fig. 3i). This periodicity of nodule

emergence appears to be similar between strains.

While the slight t_{min} delay for $\Delta hpnH$ is consistent with longer times required to initiate the

symbiosis (*e.g.* root surface colonization, invasion of the root epidermis and cortex, and intracellular

uptake), it is also possible that a delay in t_{min} simply reflects a lower rate of nodule growth immediately

after the first intracellular infection. To address this, we compared the calculated value of t_{min} (defined by

$V = 0.2$ mm^3) to the maximum growth rates and volumes for each nodule (Fig. S9e-f). We did not find that

nodules with lower growth rates and final volumes than wild type were more likely to have a later t_{min},

supporting the interpretation that the delay in t_{min} of $\Delta hpnH$ could be due to a separate initiation defect.

Interestingly, t_{min} is also not correlated with the period in which maximum nodule growth occurs, such that

later-emerging nodules have similar growth period to nodules formed within a few dpi (Fig. S9g-h). This

indicates that although nodule emergence is restricted to narrow, periodic windows (Fig. 3i), once a

nodule has entered its maximum growth phase, its continued growth is comparatively unconstrained.

*Δ*hpnH *is delayed in a pre-endosymbiont stage*

We next performed competition assays using a standard fluorescence labeling approach. We first

generated $\Delta hpnH$ and wild-type strains expressing chromosomally-integrated fluorescent proteins, and

then we co-inoculated *A. afraspera* with different ratios of these two strains. As control experiments, we

also co-inoculated each tagged strain with its untagged counterpart, in order to determine the effect of

fluorescent protein overexpression on each strain's competitiveness. After 40 dpi we measured the size

of nodules on plants inoculated with each strain combination and ratio, then sectioned and fixed nodules

for imaging. Although we expected each nodule to contain a clonal population of symbionts based on

previous work (Bonaldi et al. 2011; Ledermann et al. 2015), the majority of nodules instead contained a

mixture of both strains (Fig. 5a).

We quantified the relative abundance of each strain in each nodule by fluorescence imaging; in

our control experiments, in which only one fluorophore-expressing strain was present, a DNA dye was

used to label all bacteria. Both WT-YFP and Δ*hpnH*-mCherry were significantly out-competed by their

corresponding untagged strains, with higher proportions of tagged strains correlating with lower bacterial

DNA abundance and smaller nodule and/or infection zone sizes (Fig. 5b-c; Fig. S16-S18). Additionally,

plants co-inoculated with untagged-Δ*hpnH* and Δ*hpnH*-mCherry were significantly shorter than plants

inoculated with untagged-Δ*hpnH* only, suggesting Δ*hpnH*-mCherry is symbiotically defective (Fig. S15).

These effects of fluorophore overexpression made it difficult to interpret our WT-YFP and Δ*hpnH*-

mCherry competition data, so we developed an alternative, antibiotics-based method to study the timing

of early symbiotic initiation. First, we identified antibiotics that were effective against *B. diazoefficiens* but

would minimally affect *A. afraspera* growth. We tested three antibiotics (100 µg/ml streptomycin, 100

µg/ml kanamycin, and 20 µg/ml tetracycline) and treated non-inoculated plants with these antibiotics for

two weeks, alone and in combination. After this treatment, we found that neither kanamycin nor

streptomycin, nor the combination of the two, significantly affected plant appearance, shoot height, or root

and shoot dry masses compared to untreated controls (Fig. S19). Plants treated with tetracycline were

noticeably more yellow in color, indicating chlorosis, and the roots and plant medium became brown;

these plants also had lower shoot and root dry masses than untreated plants (Fig. S19).

Next, because the Δ*hpnH* strain is more sensitive to antibiotics than wild type (Kulkarni et al.

2015), we tested various concentrations of the non-plant-perturbing antibiotics streptomycin and

kanamycin to identify concentrations that would result in the same rates of cell death for both strains. We

inoculated plant growth media with wild type or Δ*hpnH* to the same cell densities and under the same

environmental conditions as in plant inoculation experiments. The wild-type culture was supplemented

with 100 µg/ml streptomycin plus 100 µg/ml kanamycin, and Δ*hpnH* cultures were supplemented with

decreasing concentrations of these antibiotics: 75, 50 and 25 µg/mL each. Samples of the cultures were

then collected, serially diluted and added to PSY plates to estimate colony-forming units (cfus) per mL

over time. At 50 µg/mL kanamycin plus 50 µg/mL streptomycin, the rate of decrease in cfus/mL for Δ*hpnH*

was equivalent to that of wild type treated with 100 µg/ml kanamycin plus streptomycin (Fig. 5d).

Finally, we inoculated 40 plants each with wild type or Δ*hpnH* and added streptomycin or

kanamycin to 100 µg/mL each or 50 µg/mL each, respectively, at various points post-inoculation. After 40

days we counted the number of nodules per plant, and found that antibiotics were able to block nodule

formation over a ~50% longer window in Δ*hpnH* compared to wild type (Fig. 5e). The decrease in nodules

formed at different antibiotic treatment time points was also evident in the overall appearance of the

plants (Fig. 5f). These results suggest that Δ*hpnH* requires more time on average to reach the

intracellular stage of the symbiosis, at which point we presume that the bacteria are protected from

antibiotic by the host cells. These data are consistent with Δ*hpnH* requiring more time to colonize the root

surface, invade the root epidermis, and/or be internalized by host cells.

Extended hopanoids support surface attachment and motility in vitro

Because we found that expression of genetic tags in wild type and Δ*hpnH* perturbed their

symbiosis with *A. afraspera,* and because we found that the hopanoid mutant viability is reduced by

sonication, centrifugation, and mechanical or detergent-based tissue disruption techniques required to re-

isolate bacteria from plants, we could not confidently follow these strains *in planta*. Instead, we used an *in*

vitro approach to study two steps in the initiation of the symbiosis: (1) bacterial motility toward or along

the *A. afraspera* root, and (2) stable attachment of bacteria to the root surface (Wheatley and Poole

2018). To determine whether Δ*hpnH* is less motile than wild type, we inoculated low-agar, PSY plates

with Δ*hpnH* or wild type and measured the rate of zone of swimming over time. We observed that

diameter of motility was reduced in Δ*hpnH* compared to wild type (Fig. 6a-b), consistent with a swimming

motility defect; however, because we have previously shown that Δ*hpnH* grows more slowly in this

medium than wild type (Kulkarni et al. 2015), we could not rule out the possibility that slower zone

expansion simply reflects a longer doubling time.

To investigate the nature of the plate motility defect, we studied the motility of single *B.*

diazoefficiens cells. We inoculated cells into a glass-bottom, sterile PSY flow cells with 100μL of each

strain and recorded the movement of cells near the glass surface at 5 ms time resolution. Trajectories of

individual swimming cells, defined as having super-diffusive motion and a trajectory radius of gyration

>2.5 μm, were calculated and analyzed in MATLAB (Lee et al. 2018). In agreement with our low-agar

swimming plates, we found significantly fewer ($p < 0.0001$) motile cells for Δ*hpnH* (N = 65 ± 29) than wild

type (N= 368 ± 60) in PSY medium (Fig. 6c; Table S1). The average mean-speed among motile cells

were similar between strains: $<V>_{\Delta hpnH}$ = 24.83 ± 7.0 μm/sec and $<V>_{wt}$ = 22.75 ± 6.7 μm/sec (Fig. 6d;

Table S1). Because the composition of PSY differs greatly from that of the plant growth medium (BNM),

we repeated these assays in BNM supplemented with arabinose. Under this condition, we again

observed a lower fraction of motile Δ*hpnH* cells than wild type ($N_{\Delta hpnH}$ = 54 ± 59, N_{wt}= 450 ± 310) with

similar mean speeds between strains (Fig. 6e-f; Table S1).

We next tested the surface attachment capabilities of Δ*hpnH* and wild type by incubating dense

bacterial cultures on glass coverslips and quantifying the fraction of the surface covered with stably

adherent cells after two hours. In PSY medium, both strains adhered poorly, and there was no significant

difference in their attachment efficiencies (Fig 6g; Fig. S20). In BNM supplemented with arabinose, both

strains adhered to glass better than in PSY, and Δ*hpnH* attachment levels were significantly lower than

wild type (Fig. 6g; Fig. S20). The decreased adhesion and reduced motile cell population of Δ*hpnH*

suggest that stable root colonization by this strain may be less efficient.

Discussion

Hopanoids are well-established mediators of bacterial survival under stress, and previously we

showed that the capacity for hopanoid production is enriched in plant-associated environments (Ricci et

al. 2014) and required for optimal *Bradyrhizobia-Aeschynomene* spp. symbioses (Silipo et al. 2014;

Kulkarni et al. 2015). Here we performed a detailed, quantitative evaluation of the extended hopanoid

phenotypes in the *Bradyrhizobium diazoefficiens-Aeschynomene afraspera* symbiosis. We determined

that extended hopanoid mutants fix nitrogen at similar rates as wild type on a per-bacteroid level,

demonstrating that in this host, extended hopanoids are not required to protect nitrogenase from oxygen,

as often has been speculated (reviewed in Belin et al. 2018). Instead, we found that the extended

hopanoid mutants' lower *in planta* productivity can be fully attributed to changes in the kinetics of nodule

development. By tracking the development of individual root nodules, we observed later nodule

emergence times in Δ*hpnH*-inoculated plants. *In vitro*, Δ*hpnH* cells adhered poorly to glass and were less

motile than wild type, suggesting they may colonize roots less efficiently (Fig. 7a-b). A third of Δ*hpnH*

nodules also grew significantly slower than wild type and were smaller at maturity. Many of these small

nodules contained low symbiont densities; a subset of larger Δ*hpnH* nodules also had lower symbiont

loads, due to infection zone fragmentation. The origin of this under-infection is unclear. It is possible that

bacteria are inefficiently internalized or retained, and this phenotype is simply propagated as nodules

develop (Fig. 7c-d). Alternatively, low symbiont densities may reflect symbiont degradation in a previously

fully infected nodule (Fig. 7e).

These observations challenge two conclusions from our previous work, requiring a refinement of

our interpretation of the roles of extended hopanoids in the plant context (Kulkarni et al. 2015). First, we

reported that there was no symbiotic defect of the Δ*hpnH* strain in soybean, based on the observation

that nitrogen fixation per mg nodule dry weight was similar to wild type. Given that this study revealed that

a reduction in nodule dry weight explains the Δ*hpnH* defect in *A. afraspera*, it is possible that this strain is

also defective in its native host, but this defect was obscured by differences in normalization between the

soybean and *A. afraspera* datasets. Second, the majority of Δ*hpnH* nodules had wild type-like growth

kinetics and morphologies, with a few "mega" nodules displaying unusually fast growth. This finding

appears inconsistent with an inability to survive NCR peptides, unless NCR peptide expression levels are

extremely variable, or if the mechanisms that compensate for extended hopanoid loss are inconsistent.

What other mechanisms might underpin these extended hopanoid mutant phenotypes? Perhaps

they are simply consequences of less rigid *B. diazoefficiens* membranes. The fraction of motile cells in *E.*

coli populations has been suggested to be sensitive to changes to the mechanical properties of the outer

membrane (Gupta et al. 2006), and membrane-based mechanotransduction is required by diverse

bacteria to stimulate extracellular matrix production and cement their attachment to surfaces (Petrova and

Sauer 2012; Persat 2017). *B. diazoefficiens* mutants with weakened cell walls also have been shown to

be deficient in symbiosis with *A. afraspera* through an NCR peptide-independent mechanism (Barriere et

al. 2017), which may be elicited by Δ*hpnH*. Extended hopanoid loss may also have secondary effects on

Bradyrhizobium-Aeschynomene signaling. In the *Frankia*-actinorhizal symbiosis, bacterial extended

hopanoids can contain the auxinomimetic compound phenyl-acetic acid (PAA) (Hammad et al. 2003), and

though the effects of hopanoid loss on the bacterial metabolome have not been examined, changes in

hopanoid production may impact the synthesis and/or secretion of symbiotically active compounds.

Future work will be required to determine whether changes in signaling or membrane mechanics

dominate the hopanoid mutant phenotypes, and at which developmental stages.

Regardless of the underlying mechanism, it is curious that the absence of extended hopanoids is

not a death knell for the *B. diazoefficiens-A. afraspera* symbiosis at any stage. In our *in vitro* studies,

mean speeds among motile Δ*hpnH* cells were indistinguishable from wild type, and though we cannot

rule out more subtle defects in the direction of movement or chemotaxis, this suggests that motility

systems of Δ*hpnH* cells function properly once induced. Similarly *in planta*, Δ*hpnH* nodules developing at

wild-type rates and reaching average wild-type volumes did occur – and, in the case of "mega" nodules,

some exceeded their wild-type counterparts.

Why do Δ*hpnH* populations form two distinct populations (wild-type-like or defective) rather than

falling on a continuous distribution of behavior? Bimodality can reflect switch-like, or threshold-based,

regulation, and perhaps in the Δ*hpnH* strain, a fraction of cells cannot support levels of signaling above

the threshold required for proper function. Nodules may also differ in the extent to which extended

hopanoid loss is compensated. In *Methylobacterium extorquens* and *Rhodopseudomonas palustris*

(Bradley et al. 2017; Neubauer et al. 2015), hopanoid loss results in upregulation of other membrane-

rigidifying lipids including carotenoids and cardiolipins, and in other plant-microbe systems, lipid exchange

between hosts and microbes has been observed (Keymer 2018), suggesting that Δ*hpnH* nodule

phenotypes may relate to the local availability of structurally or functionally similar metabolites. Because

of these diverse possible explanations for Δ*hpnH* heterogeneity, a detailed comparison of wild type-like

and defective nodules, including the distributions of lipids and other metabolites, bacteroid morphology

and penetrance, and gene expression variability, will be required to determine why some Δ*hpnH* nodules

succeed and others do not.

Beyond hopanoids, our results provide insight into the developmental control of nodule formation

by *A. afraspera* hosts. We find that nodulation occurs in bursts separated by fixed 18-day intervals, and

that the timing of these bursts is unrelated to net fixed nitrogen production across the root, more likely

reflecting the inherent dynamics of the underlying signaling networks. The growth period of individual

nodules is similarly deterministic, suggesting that *A. afraspera* hosts do not respond to ineffective

symbionts by prematurely aborting nodule development. Rather, we find that *A. afraspera* nodules can be

primarily distinguished by their growth rates, *e.g.* the frequencies of infected host cell division. This finding

suggests that in *A. afraspera* host cell mitosis and symbiont performance may be coupled, enabling future

studies on the molecular signals through which this coupling occurs.

Finally, our results underscore the importance of identifying the most informative, least perturbing

tools for interrogating legume-microbe symbiosis. Employing quantitative, time-resolved, single-nodule

and single-cell approaches rather than bulk measurements were essential for uncovering the diverse

phenotypes of the *B. diazoefficiens* extended hopanoid mutants and yielded unexpected information on

regulation of nodule development by *A. afraspera*. We have also shown the limitations of introducing

overexpressed genetic tags into bacteria. While use of these tags has undoubtedly enhanced our

understanding of legume-microbe symbiosis (Ledermann et al. 2018), they may not fully capture the

behavior of native organisms. Additionally, our work is one of many to emphasize the importance of

appropriate culture models for mimicking the host environment, as the Δ*hpnH* surface attachment defect

was observed in plant growth medium but not in a standard richer medium. A more detailed analysis of

the host environment, including the full milieu of root exudates (Sugiyama and Yazaki 2012), available

carbon sources (Pini et al. 2017) and trace metals specific to each legume, will improve *in vitro* models of

legume-bacteria interactions and may allow selection of strains with improved performance in agriculture.

Methods

B. diazoefficiens *culture and strain generation*

B. diazoefficiens hopanoid biosynthesis mutants were generated previously (Kulkarni et al. 2015).

For construction of YFP- and mCherry-expressing strains, fluorophore expression vectors pRJPaph-YFP

and pRJPaph-mCherry (Ledermann et al. 2015) were provided as a gift from Prof. Dr. Hans-Martin

Fischer (ETH Zurich). These vectors were introduced into *B. diazoefficiens* by conjugation with the β2155

DAP auxotroph strain of *E.coli*, using the following protocol: *B. diazoefficiens* wild type and Δ*hpnH* were

grown in 5 mL PSY medium(Regensburger and Hennecke 1983) at 30°C and 250 rpm to an OD_{600} of

~1.0 (wild type) or of 0.5-0.8 (Δ*hpnH*). β2155 strains carrying pRJPaph vectors were grown to an OD_{600} of

0.5-0.8 in 5 mL LB supplemented with 10 µg/mL tetracycline and 300 µm DAP at 37°C and 250 rpm. Both

B. diazoefficiens and β2155 donor cultures were pelleted at 3250 x g for 30 minutes, washed three times

in 0.9% sterile saline, and resuspended in 0.9% sterile saline to a final OD_{600} of 1.0. *B. diazoefficiens*

strains and β2155 donor cells were combined at a 4:1 ratio, respectively, and mixed by repeated

pipetting. Aliquots (50 µl) of these 4:1 mixtures were dropped to PSY plates supplemented with 300 µm

DAP, dried in a biosafety cabinet, and incubated for 48 hours at 30°C. Conjugation pastes were then

removed from plates and resuspended in 5 mL sterile saline, pelleted at 3250 x g for 30 minutes and

washed twice, in order to remove residual DAP. Washed cells were pelleted a final time and resuspended

to 200 μl in 0.9% sterile saline and plated onto PSY plates supplemented with 20 μg/mL (wild type) or 10

μg/mL (Δ*hpnH*) tetracycline. Colonies appeared after 7-10 days (wild type) or 10-14 days (Δ*hpnH*) and

were streaked onto fresh PSY/tetracycline plates, then screened for fluorescence using a Lumascope 720

fluorescent microscope (Etaluma). Fluorescent colonies were then sequenced to verify insertion of the

pRJPaph vectors into the *scoI* locus.

A. afraspera *cultivation and inoculation with* B. diazoefficiens

A. afraspera seeds were obtained as a gift from the laboratory of Dr. Eric Giraud (LSTM/Cirad,

Montpelier, France). Seeds were sterilized and scarified by incubation in 95% sulfuric acid at RT for 45

minutes, followed by 5 washes in sterile-filtered nanopure water and a second incubation in 95% ethanol

for 5 minutes at RT. After ethanol treatment seeds were washed 5X and incubated overnight in sterile-

filtered nanopure water. Seeds were then transferred to freshly poured water/agar plates using sterile,

single-use forceps in a biosafety cabinet, and germinated for 24-72 hours in the dark at 28-32°C.

Seedlings were then placed in clear glass test tubes containing 100 mL of sterile, nitrogen-free

Buffered Nodulation Medium (BNM)(Ehrhardt et al. 1992) and grown for 7-10 days in plant growth

chambers (Percival) under the following settings: 28°C, 80-90% humidity, and 16 hour photoperiod under

photosynthetic light bulbs (General Electric) emitting ~4000 lumens/ft^2. In parallel, *B. diazoefficiens*

strains were grown in 5-10 mL PSY liquid culture at 30°C and 250 rpm to stationary phase (OD_{600} > 1.4).

Stationary phase cultures were diluted into PSY one day prior to plant inoculation to reach an OD_{600} of

~0.8 at the time of inoculation. OD_{600} ~ 0.8 cultures were pelleted at 3250 x g for 30 minutes at RT,

washed once in PSY, then resuspended in PSY to a final OD_{600} of 1.0. Resuspended *B. diazoefficiens*

cultures were directly inoculated into the plant medium in a sterile biosafety cabinet; 1 mL of OD_{600}=1.0

culture was added per plant. Inoculated plants were then returned to growth chambers and maintained for

the times indicated for each experiment. For longer experiments (lasting longer than ~30 days post-

inoculation), plant growth tubes were refilled with sterile-filered nanopure water as needed. To minimize

cross-contamination, inoculated plants and non-inoculated plants were cultivated in separate growth

chambers, and growth chambers were sterilized with 70% ethanol followed by UV irradiation for at least

24 hours between experiments.

Acetylene reduction experiments

Individual plants were transferred to clear glass 150 mL Balch-type anaerobic culture bottles containing

15 mL BNM medium and sealed under a gas-tight septum. After sealing, 15 mL of headspace gas (10%

of the culture bottle volume) was removed and replaced with 15 mL of acetylene gas (Airgas). Plants in

culture bottles were incubated in the light at 28°C in growth chambers for 3-6 hours. A 100 µl sample of

the headspace gas was removed using a gas-tight syringe (Hamilton), and this sample was injected and

analyzed for ethylene signal intensities using a Hewlett Packard 5890 Series II GC with Hewlett Packard

5972 Mass Spectrometer with a 30mx0.320mm GasPro Column (Agilent Technologies) and a 2 mm ID

splitless liner (Restek Corporation). Following acetylene reduction measurements, plants were removed

from jars and plant shoot heights and number of nodules per plant were recorded. When nodule dry mass

measurements were performed, nodules were harvested with a razor blade, transferred into pre-weighed

Eppendorf tubes, dried at 50°C for a minimum of 48 hours, then weighed again.

Live:Dead staining and imaging of nodule cross-sections

Nodules were hand-sectioned with razor blades and immediately transferred into a fresh solution

of 5 µM SYTO9 (diluted 1:100 from a 500 uM stock in DMSO at -20°C; Thermo Fisher) and 0.02 mg/mL

(30 µM) propidium iodide (diluted 1:50 from a 1 mg/mL stock stored in water at 4°C; Thermo Fisher) in

PBS. Nodule sections were incubated in this SYTO9/propidium iodide solution at room temperature for 30

minutes in the dark with gentle shaking, washed 5X in PBS, and fixed in 4% paraformaldehyde (Electron

Microscopy Sciences) in PBS overnight in the dark at 4°C. Fixed sections were washed 5X in PBS and

transferred to a freshly prepared solution of 0.1 mg/mL Calcofluor White (Fluorescence Brightener 28;

Sigma) in PBS. The sections were incubated in the Calcofluor solution in the dark for 1 hour at RT with

gentle shaking and washed 5X in PBS to remove excess dye.

Prior to imaging, sections were transferred to 30 mm imaging dishes with 20 mm, #0 coverglass

bottoms (MatTek) and overlaid with sterile 50% glycerol. Nodule images were collected on a Leica TCS

SPE laser-scanning confocal (model DMI4000B-CS) using a 10X/0.3 NA APO ACS objective and solid-

state laser lines for fluorophore excitation at the following settings for each dye: Calcofluor, 405 nm

excitation/410-500 nm emission; SYTO9, 488 nm excitation/510-570 nm emission; PI, 532 nm

excitation/600-650 nm emission. These images were then processed to enhance brightness and contrast

in FIJI(Schindelin et al. 2012; Schneider et al. 2012).

Nodule diameter and volume measurements

Inoculated *A. afraspera* root nodules were imaging using a high-definition Keyence VHX-600

digital microscope at 20X magnification. For end-point root nodule volume measurements at 40 days

post-inoculation, plants were removed from the growth chamber and imaged at RT on paper towels, then

discarded. Nodule diameters were measured using the line tool in FIJI and recorded using a custom FIJI

macro. For tracking nodule volumes over time, plants were serially removed from their growth chambers

and transferred to a plastic dish containing 150 mL of sterile BNM pre-warmed to 28°C. Images of

sections of the plant root were collected serially from the hypocotyl to the root tip. Following collection of

images, plants were immediately returned to their original growth tubes in the growth chamber. Plastic

dishes were sterilized for 10 minutes in 10% bleach, washed three times in sterile-filtered nanopure

water, sprayed with 70% ethanol/water, and air-dried before each new plant was imaged. A fresh aliquot

of sterile, pre-warmed BNM also was used for each plant. After the time course was completed, images of

entire plant root systems were reconstructed by eye for each plant at each time point. For nodules

appearing in at least five time points, nodule diameters were measured as described for the end-point

measurements and were converted to approximate volumes in R using the equation $V = 4/3\pi r^3$.

Nodule growth curve fitting and analysis

All analyses of nodule growth, and corresponding plots, were generated in R. For nodule growth

curve fitting, three model equations were used to identify the best fit, as follows:

(1) exponential function:

$$V = ae^{-bt} + c$$

476   (2) quadratic function:

$$V = at^2 + bt + c$$

478   (3) generalized logistic function (expressed as a Richard's function with a time shift):

$$V = \frac{a}{(1 + e^{-b(t-c)})^{(\frac{1}{d})}}$$

Calculation of the optimal parameter values for each equation (e.g. the values of **a**, **b**, **c**, and **d**) and the

standard error for each curve compared to the raw data were performed using the built-in function *nlm()* in

R. In some cases, *nlm()* could not produce a best-fit model without specifying initial values for the function

parameters. For exponential models, an equation of best fit could be successfully determined without

specification of initial values for parameters **a**, **b** and **c**. For quadratic models, initial parameter values

were required and were set to **a**=0, **b**=10 and **c**=0 for each nodule plot, after identifying these initial

parameter values as broadly optimal based on an initial parameter sweep of -50 to 50 for each plot. For

sigmoidal models, no broadly optimal initial values could be identified, so a parameter sweep was

performed for each plot with the initial value of **a** set to the maximum observed nodule volume (as **a**

describes the upper asymptote of the sigmoidal curve), **b** ranging from 0.1 to 1, **c** ranging from 0 to 10,

and **d** ranging from 0.01 to 1.0. In the sigmoidal plots, an initial point of (0,0) was added to the nodule

volume time series to improve fitting.

Because the sigmoidal model provided the best fits, extrapolation of nodule growth characteristics

was performed on sigmoidal models only. The maximum nodule volume, **V**$_{max}$, is defined as the upper

asymptote of the sigmoidal growth curve, *e.g.* **a**. The nodule initiation time, **t**$_{min}$, was defined in three

separate ways: the times at which the nodule volume is equal to 0.05, 0.1, or 0.2 mm^3 (*e.g.* through

solving 0.05, 0.1, or 0.2 = **a**/((1+e$^{(-b(t-c))}$)$^{(1/d)}$) for *t*). The maximum nodule growth rate, **dV/dt**, was defined

as the average rate of growth (*e.g.* slope) between the time at which the volume is 10% of V_{max} and the

time at which the volume is 90% of V_{max}. The time at which each nodule reaches its maximum size, t_{max},

was approximated as the time at which the volume is 90% of V_{max}, since the "true" maximum volume is

asymptotic to the growth curve and is therefore never fully reached in the model.

Competition assays

mCherry-tagged Δ*hpnH* and YFP-tagged wild type *B. diazoefficiens* were grown to stationary

phase (OD_{600} > 1.4) in 10 mL PSY cultures supplemented with 20 μg/mL (wild type) or 10 μg/mL (Δ*hpnH*)

tetracycline; untagged strains were grown in PSY. On the day prior to inoculation, all strains were diluted

into 50-150 mL tetracycline-free PSY to reach an OD_{600} of ~0.8 at the time of inoculation. *A. afraspera*

plants were cultivated pre-inoculation in test tubes as described above, with the addition of covering the

growth tubes in foil to minimize the production of chlorophyll in the plant roots, which spectrally overlaps

with mCherry. At the time of inoculation, all cultures were pelleted at 3250 x g for 30 minutes at RT,

washed three times, then resuspended in PSY to a final OD_{600} of 1.0. A 10 mL culture of each strain ratio

for inoculation was generated a sterile 15mL Falcon tube; for example, for a 50:50 mixture of mCherry-

tagged Δ*hpnH* and YFP-tagged wild type, 5 mL of each strain was combined. These cultures were mixed

thoroughly by gentle pipetting, and 1 mL of the mixtures was added to directly to the plant medium for 7-8

plants per strain mixture.

After 45-60 days, plants were harvested. First, plant heights and the number of nodules per plant

were recorded. Then, the roots were cut from the stem and images of all nodules for each plant were

collected on a high-definition Keyence VHX-600 digital microscope at 20X magnification. These nodules

were then cross-sectioned and immediately transferred to Eppendorfs containing 4% paraformaldehyde

(Electron Microscopy Sciences) in PBS. Fresh sections were fixed overnight in the dark at 4°C, washed

5X in PBS, and stored in PBS supplemented with 0.1% azide in the dark at 4°C until imaging.

Fixed sections were stained in Calcofluor (all strain combinations), SYTO9 (WT-YFP and WT co-

inoculation only) or propidium iodide (mCherry-Δ*hpnH* and Δ*hpnH* co-inoculation only) as described for

Live:Dead staining. Imaging was performed as described for Live:Dead staining using a 5X objective.

Given the high autofluorescence of these nodules and low mCherry and YFP signal intensities, the

following excitation/emission settings were used: Calcofluor, 405 nm excitation/410-460 nm emission;

YFP/SYTO9, 488 nm excitation/500-550 nm emisasion; mCherry, 532 nm excitation/600-650 nm

emission.

Quantification of nodule statistics (including nodule and infection zone areas, signal intensity of

YFP, mCherry, SYTO9 and propidium iodide) was performed on raw images using a custom FIJI macro.

Briefly, nodule images were opened at random, infection zones (IZs) and whole nodules were circled by

hand and saved as discrete regions of interest (ROIs), and the area and intensity in each channel were

measured automatically for all ROIs. These measurements were exported as a text table and various

parameters from these measurements were calculated using custom Python scripts, as indicated in the

Results. Plots of all parameters and statistical comparisons were generated using custom R scripts.

Antibiotic treatment of inoculated plants

A. afraspera plants were cultivated as described above and the following antibiotics were added

to non-inoculated plants 7 days after rooting in 100 mL BNM growth tubes: kanamycin to 100 µg/mL,

streptomycin to 100 µg/mL, tetracycline to 20 µg/mL, kanamycin plus tetracycline, kanamycin plus

streptomycin, streptomycin plus tetracycline. Plants were grown in antibiotics under normal plant growth

conditions for 14 days, after which plants were visually inspected. Plant heights were also recorded, and

the root and shoot systems were separated with a razor blade, transferred into pre-weighed 15 mL Falcon

tubes, dried at 50°C for a minimum of 48 hours, then weighed again.

Antibiotic treatments of ΔhpnH and wild-type *B. diazoefficiens* were performed by growing

antibiotic 5 mL PSY cultures of each strain to stationary phase (OD_{600} >1.4) and diluting strains in fresh

PSY to reach an OD_{600} of ~0.8 at the time of antibiotic treatment – *e.g.* as they would be grown prior to

plant inoculation. Cultures were pelleted at 3250 x g for 30 minutes at RT, washed three times, then

resuspended in PSY to a final OD_{600} of 1.0. Four 100 µl aliquots of these culture were diluted 1:00 into

separate 10 mL BNM cultures in clear glass tubes in plant growth chambers. Kanamycin (at 25, 50, 75,

and 100 µg/mL) and streptomycin (at 25, 50, 75, and 100 µg/mL) were added directly to the BNM

cultures, and 100 µl samples were taken immediately prior to antibiotic treatment and at 2, 4, 6, 8, and 10

hours post-antibiotic addition. These 100 µl samples were immediately diluted 1:10 in 900 µl and mixed

vigorously by repeated pipetting. Vortexing was avoided as we found that this method reduces Δ*hpnH*

viability. Ten serial 1:10 dilutions were performed, and three 10 µl samples of each dilution for each strain

were spotted and dripped across PSY plates. After 7 days (wild type) or 10 days (Δ*hpnH*), colonies were

counted manually and recorded for each dilution exhibiting discrete colonies. Log plots of colony counts

over time were generated in R.

Plants were then inoculated with Δ*hpnH* and wild-type *B. diazoefficiens* as described above, and

kanamycin and streptomycin were added to Δ*hpnH*-inoculated plants to 50 µg/mL each, and to wild type-

inoculated plants to 100 µg/mL at 12 hours and 36 hours and at 2, 2.5, 3, 3.5, 4, 4.5, 5, 6.5, 8.5, 9.5, 10.5,

and 12.5 days post-inoculation. Four plants were treated per time point per strain, with an additional four

plants each as an untreated control. At 40 dpi, the number of nodules per plant was recorded.

Bulk motility assays

Swimming motility assays were performed as previously described, with some modifications

(Althabegoiti et al. 2008). WT and Δ*hpnH* were grown to turbidity in 5 mL of PSY at 30°C and 250 rpm,

diluted to an OD_{600} of 0.02 in 5 mL of fresh PSY, and grown to exponential phase (OD_{600} = 0.3-0.5).

Exponential cultures then were diluted to an OD_{600} of 0.06 in fresh PSY and 2 µL of the adjusted cultures

into the center of swimming plate containing 0.3% agar/PSY. After inoculation, the plates were wrapped

with parafilm to prevent dehydration and incubated in a humidity-controlled environmental chamber

(Percival) at 30°C for 10 days total, with daily scans after 5 days. The resulting images were analyzed in

FIJI to measure the area of the swimming colony.

Surface attachment assays

Δ*hpnH* and wild-type *B. diazoefficiens* were grown in 5 mL PSY cultures to stationary phase

(OD_{600} >1.4) then diluted in fresh PSY to reach an OD_{600} of ~0.8 at the time of surface attachment

assays. Cultures were pelleted at 3250 x g for 30 minutes at RT, washed twice in the indicated

attachment medium, then resuspended in attachment medium to an OD_{600} of 1.0. These cultures were

mixed thoroughly by repeated pipetting, and 2 mL samples were added to sterile imaging dishes (30 mm

dishes with 20 mm, #1.5 coverglass bottoms; MatTek). Cultures were incubated on imaging dishes

without shaking at 30°C for two hours. To remove non-adhered cells, imaging dishes were immersed in

50 mL of attachment media in a 100 mL glass beaker on an orbital shaker and shaken gently at RT for 5

minutes; direct application of washing medium to the coverglass surface was avoided, as we found that

this creates a shear force sufficient to wash away adhered cells. Imaging dishes were then gently lifted

out of the washing medium and imaged with a 100X objective on a Lumascope 720 fluorescence

microscope (Etaluma). Forty fields of view were recorded for each strain and media combination. These

images were processed in FIJI using the Enhanced Local Contrast (CLAHE) plugin (Heckbert and Karel

1994) and converted into a binary image to determine the area of the imaging window covered with

adhered cells. Calculation of the fraction of the surface was performed in Excel and statistical analyses

were conducted in R. Areas of the surface containing groups of cells larger than 10 μm^2 in area were

ignored in the calculations, as these likely do not represent true attachment events rather than

sedimentation of larger cell clumps. BNM used for attachment assays was prepared as described above,

with the addition of 1.0 g/mL arabinose. Because BNM contains salt crystals that can sediment onto

coverglass and occlude or obscure adhered cells, this medium was passed through a 2 μm filter

(Millipore) prior to the attachment experiments.

Single-cell motility assays and analysis

B. diazoefficiens wild-type and ΔhpnH were grown in 12.5 ml PSY medium at 30°C and 200 rpm

to an OD_{600} = 0.6-0.8 from an AG medium plate culture. Then, a 1:10 dilution of cell culture was

subcultured in PSY medium to a final volume of 12.5 ml and regrown to an OD_{600} of ~0.6. Two aliquots of

750 μL were sampled from the regrowth culture and pelleted at 3500 x g for 20 min (wild-type) or for 30

min (ΔhpnH) at RT. The supernatant was removed, and one pellet was resuspended in 500 μL PSY and

the other in 500 μL BNM medium. Because BNM contains salt crystals that can sediment onto coverglass

and occlude or obscure adhered cells, this medium was passed through a 2 μm filter (Millipore) prior to

usage for these experiments. The two medium conditions were then incubated for 2.5 hrs (wild-type) or

for 3.5 hrs (ΔhpnH) at 30°C; given the difference in growth time ΔhpnH incubated for longer. Right before

imaging, each culture was diluted at a 1:10 ratio with its respective medium. The bacteria were then

injected into a sterile flow cell (ibidi sticky-Slide VI0.4 with a glass coverslip). The flow cell was attached to

a heating stage set to 30°C.

The imaging protocol involved high-speed bright-field imaging for 5 min at a single XYZ location

per experimental repeat. High speed bright-field recordings used a Phantom V12.1 high speed camera

(Vision Research); images were taken with a 5 ms exposure at 200 fps and a resolution of 512×512

pixels (0.1 µm/pixel). This protocol was performed on an Olympus IX83 microscope equipped with a 100×

oil objective, a 2× multiplier lens, and a Zero Drift Correction autofocus system. The recorded movies

were extracted into single frames from the .cine files using PCC 2.8 (Phantom Software). Image

processing and cell tracking algorithms are adapted from previous work (Lee et al. 2018) and written in

MATLAB R2015a (Mathworks).

We identified cells swimming near the surface as cells with a trajectory radius of gyration greater

than 2.5 µm and a mean-squared displacement (MSD) slope greater than 1.5. Setting a minimum radius

of gyration selects for cells with a minimum net translation on the across the surface, while a minimum

MSD slope threshold ensured the cells are moving super-diffusively (MSD slope $\cong$ 1, diffusive motion;

MSD slope $\cong$ 2, super-diffusive motion). For each tracked cell, the mean-speed, **v**, was calculated by

averaging a moving window, **w**, of the displacement over the cell's full trajectory, using the following

equation:

$$< v > = \ Avg\left(\sum_{=1}^{N-w} \frac{\sqrt{(x_{t+w} - x_t)^2 + (y_{t+w} - y_t)^2}}{w} * f * p \right)$$

where **N** is the total number of points in the trajectory, **f** is the acquisition frame rate, and **p** is the pixel

resolution. Here we set a window size, w= 40 frames. All analysis and visualizations from these

experiments where done using MATLAB R2015a (Mathworks).

Acknowledgements

This work was supported by grants from the HHMI (D.K.N.), NASA

(NNX12AD93G, D.K.N.), Jane Coffin Childs Memorial Fund (B.J.B.), NIH (K99GM126141,

B.J.B.), and Army Research Office (W911NF-18-1-0254, GW) and predoctoral fellowships from NSF

(E.T.) and the Ford Foundation (J.d.A.). We thank Dr. Eric Giraud for his generous gift of *A. afraspera*

seeds and training on *Aeschynomene* symbioses and Drs. Hans-Martin Fischer and Raphael Ledermann

for plasmids and technical advice for the genetic transformation of *B. diazoefficiens*. Dr. Nathan Dalleska

of the Environmental Analysis Center at Caltech was instrumental in providing training and support for

GC-MS analysis of acetylene reduction. We are grateful to Dr. Gargi Kulkarni and other members of the

Newman lab, as well as Drs. Elliot Meyerowitz and Rob Phillips, for their collegiality and thoughtful

discussions about this work. We are indebted to Ms. Shannon Park and Ms. Kristy Nguyen for providing

the administrative assistance that allows us to focus on our research.

References

Abeysekera, R. M., Newcomb, W., Silvester, W. B., and Torrey, J. G. 1990. A freeze-fracture electron

microscopic study of *Frankia* in root nodules of *Alnus incana* grown at three oxygen tensions. Can.

J. Microbiol. 36:97–108

Barriere, Q., Guefrachi, I., Gully, D., Lamouche, F., Pierre, O., Fardoux, J., Chaintreuil, C., Alunni, B.,

Timchenko, T., Giraud, E., and Mergaert, P. 2017. Integrated roles of BclA and DD-

carboxypeptidase 1 in *Bradyrhizobium* differentiation within NCR-producing and NCR-lacking root

nodules. Sci. Rep. 7:9063

Belin, B. J., Busset, N., Giraud, E., Molinaro, A., Silipo, A., and Newman, D. K. 2018. Hopanoid lipids:

from membranes to plant-bacteria interactions. Nat. Rev. Microbiol. 16:304–315

Bonaldi, K., Gargani, D., Prin, Y., Fardoux, J., Gully, D., Nouwen, N., Goormachtig, S., and Giraud, E.

2011. Nodulation of *Aeschynomene afraspera* and *A. indica* by photosynthetic *Bradyrhizobium* Sp.

strain ORS285: the nod-dependent versus the nod-independent symbiotic interaction. Mol. Plant.

Microbe. Interact. 24:1359–1371

Bradley, A. S., Swanson, P. K., Muller, E. E. L., Bringel, F., Caroll, S. M., Pearson, A., Vuilleumier, S., and

Marx, C. J. 2017. Hopanoid-free *Methylobacterium extorquens* DM4 overproduces carotenoids and

has widespread growth impairment. PLoS One. 12:e0173323

Caamal-Fuentes, E., Peraza-Sánchez, S., Torres-Tapia, L., and Moo-Puc, R. 2015. Isolation and

identification of cytotoxic compounds from *Aeschynomene fascicularis*, a Mayan medicinal plant.

Molecules. 20:13563–13574

Chifundera, K. 2001. Contribution to the inventory of medicinal plants from the Bushi area, South Kivu

Province, Democratic Republic of Congo. Fitoterapia. 72:351–368

Czernic, P., Gully, D., Cartieaux, F., Moulin, L., Guefrachi, I., Patrel, D., Pierre, O., Fardoux, J.,

Chaintreuil, C., Nguyen, P., Gressent, F., Da Silva, C., Poulain, J., Wincker, P., Rofidal, V., Hem, S.,

Barrière, Q., Arrighi, J.-F., Mergaert, P., and Giraud, E. 2015. Convergent evolution of endosymbiont

differentiation in Dalbergioid and Inverted Repeat-Lacking Clade legumes mediated by nodule-

specific cysteine-rich peptides. Plant Physiol. 169:1254–1265

Ehrhardt, D. W., Atkinson, E. M., and Long, S. R. 1992. Depolarization of alfalfa root hair membrane

potential by *Rhizobium meliloti* Nod factors. Science. 256:998–1000

Gupta, R., Sharma, M., and Mittal, A. 2006. Effects of membrane tension on nanopropeller driven

bacterial motion. J Nanosci Nanotechnol. 6:3854–3862

Hammad, Y., Nalin, R., Marechal, J., Fiasson, K., Pepin, R., Berry, A. M., Normand, P., and Domenach,

A.-M. 2003. A possible role for phenyl acetic acid (PAA) on *Alnus glutinosa* nodulation by *Frankia*.

Plant Soil. 254:193–205

Heckbert, P. S., and Karel. 1994. *Graphics gems IV*. AP Professional.

Keymer, A. 2018. Cross-kingdom lipid transfer in arbuscular mycorrhiza symbiosis and beyond. Curr.

Opin. Plant Biol. 44:137–144

Komaniecka, I., Choma, A., Mazur, A., Duda, K. A., Lindner, B., Schwudke, D., and Holst, O. 2014.

Occurrence of an unusual hopanoid-containing lipid A among lipopolysaccharides from

Bradyrhizobium species. J. Biol. Chem. 289:35644–35655

Kulkarni, G., Busset, N., Molinaro, A., Gargani, D., Chaintreuil, C., Silipo, A., Giraud, E., and Newman, D.

K. 2015. Specific hopanoid classes differentially affect free-living and symbiotic states of

Bradyrhizobium diazoefficiens. MBio. 6:e01251-15

Ledermann, R., Bartsch, I., Muller, B., Wulser, J., and Fischer, H.-M. 2018. A functional general stress

response of *Bradyrhizobium diazoefficiens* is required for early stages of host plant infection. Mol.

Plant. Microbe. Interact. 31:537–547

Ledermann, R., Bartsch, I., Remus-Emsermann, M. N., Vorholt, J. A., and Fischer, H.-M. 2015. Stable

fluorescent and enzymatic tagging of *Bradyrhizobium diazoefficiens* to analyze host-plant infection

and colonization. Mol. Plant. Microbe. Interact. 28:959–967

Lee, C. K., de Anda, J., Baker, A. E., Bennett, R. R., Luo, Y., Lee, E. Y., Keefe, J. A., Helali, J. S., Ma, J.,

Zhao, K., Golestanian, R., O'Toole, G. A., and Wong, G. C. L. 2018. Multigenerational memory and

adaptive adhesion in early bacterial biofilm communities. Proc. Natl. Acad. Sci. U. S. A. 115:4471–

4476

Lei, Z.-Y., Chen, J.-J., Cao, Z.-J., Ao, M.-Z., and Yu, L.-J. 2018. Efficacy of *Aeschynomene indica* Linn

leaves for wound healing and isolation of active constituent. J. Ethnopharmacol. 228:156-163

Neubauer, C., Dalleska, N. F., Cowley, E. S., Shikuma, N. J., Wu, C.-H., Sessions, A. L., and Newman,

D. K. 2015. Lipid remodeling in *Rhodopseudomonas palustris* TIE-1 upon loss of hopanoids and

hopanoid methylation. Geobiology. 13:443–453

Ochs, D., Kaletta, C., Entian, K., Beck-sickinger, A., Porallal, K., Chemie, O., Tubingen, U., and Institut, B.

1992. Cloning, expression, and sequencing of squalene-hopene cyclase,. 174:298–302

Parsons, R., Silvester, W. B., Harris, S., Gruijters, W. T., and Bullivant, S. 1987. *Frankia* vesicles provide

inducible and absolute oxygen protection for nitrogenase. Plant Physiol. 83:728–731

Persat, A. 2017. Bacterial mechanotransduction. Curr. Opin. Microbiol. 36:1–6

Petrova, O. E., and Sauer, K. 2012. Sticky situations: Key components that control bacterial surface

attachment. J. Bacteriol. 194:2413–2425

Pini, F., East, A. K., Appia-Ayme, C., Tomek, J., Karunakaran, R., Mendoza-Suarez, M., Edwards, A.,

Terpolilli, J. J., Roworth, J., Downie, J. A., and Poole, P. S. 2017. Bacterial biosensors for *in vivo*

spatiotemporal mapping of root secretion. Plant Physiol. 174:1289–1306

Regensburger, B., and Hennecke, H. 1983. RNA polymerase from *Rhizobium japonicum*. Arch. Microbiol.

135:103–109

Ricci, J. N., Coleman, M. L., Welander, P. V., Sessions, A. L., Summons, R. E., Spear, J. R., and

Newman, D. K. 2014. Diverse capacity for 2-methylhopanoid production correlates with a specific

ecological niche. ISME J. 8:675–684

Richards, F. J. 1959. NA Flexible growth function for empirical use. J. Exp. Bot. 10:290–301

Saenz, J. P., Grosser, D., Bradley, A. S., Lagny, T. J., Lavrynenko, O., Broda, M., and Simons, K. 2015.

Hopanoids as functional analogues of cholesterol in bacterial membranes. Proc. Natl. Acad. Sci. U.

S. A. 112:11971–11976

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,

Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K.,

Tomancak, P., and Cardona, A. 2012. Fiji: An open-source platform for biological-image analysis.

Nat. Methods. 9:676–682

Schmerk, C. L., Welander, P. V, Hamad, M. A., Bain, K. L., Bernards, M. A., Summons, R. E., and

Valvano, M. A. 2015. Elucidation of the *Burkholderia cenocepacia* hopanoid biosynthesis pathway

uncovers functions for conserved proteins in hopanoid-producing bacteria. Environ. Microbiol.

17:735–750

Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. 2012. NIH Image to ImageJ: 25 years of image

analysis. Nat. Methods. 9: 671-675

Silipo, A., Vitiello, G., Gully, D., Sturiale, L., Chaintreuil, C., Fardoux, J., Gargani, D., Lee, H.-I., Kulkarni,

G., Busset, N., Marchetti, R., Palmigiano, A., Moll, H., Engel, R., Lanzetta, R., Paduano, L., Parrilli,

M., Chang, W.-S., Holst, O., Newman, D. K., Garozzo, D., D'Errico, G., Giraud, E., and Molinaro, A.

2014. Covalently linked hopanoid-lipid A improves outer-membrane resistance of a *Bradyrhizobium*

symbiont of legumes. Nat. Commun. 5:5106

Somado, E. A., Becker, M., Kuehne, R. F., Sahrawat, K. L., and Vlek, P. L. G. 2003. Combined effects of

legumes with rock phosphorus on rice in West Africa. Agron. J. 95:1172

Sugiyama, A., and Yazaki, K. 2012. Root exudates of legume plants and their involvement in interactions

with soil microbes. In: Vivanco J., Baluška F. (eds) Secretions and exudates in biological systems.

signaling and communication in plants, vol 12. Springer, Berlin, Heidelberg

Swapna M.M, S., R., P., K.P., A., C.N, M., and N.P, R. 2011. A review on the medicinal and edible

aspects of aquatic and wetland plants of India. J. Med. Plants Res. 5:7163–7176

Syren, P.-O., Henche, S., Eichler, A., Nestl, B. M., and Hauer, B. 2016. Squalene-hopene cyclases-

evolution, dynamics and catalytic scope. Curr. Opin. Struct. Biol. 41:73–82

Szparaga, A., and Kocira, S. 2018. Generalized logistic functions in modelling emergence of *Brassica*

napus L. PLoS One. 13:1–14

Vilcheze, C., Llopiz, P., Neunlist, S., Poralla, K., and Rohmer, M. 1994. Prokaryotic triterpenoids: New

hopanoids from the nitrogen-fixing bacteria *Azotobacter vinelandii*, *Beijerinckia indica* and

Beijerinckia mobilis. Microbiology. 140:2749–2753

Welander, P. V, Coleman, M. L., Sessions, A. L., Summons, R. E., and Newman, D. K. 2010.

Identification of a methylase required for 2-methylhopanoid production and implications for the

interpretation of sedimentary hopanes. Proc. Natl. Acad. Sci. U. S. A. 107:8537–8542

Welander, P. V, Doughty, D. M., Wu, C., Mehay, S., and Roger, E. 2012. Identification and

characterization of *Rhodopseudomonas palustris* TIE-1 hopanoid biosynthesis mutants. 10:163–177

Wheatley, R. M., and Poole, P. S. 2018. Mechanisms of bacterial attachment to roots. FEMS Microbiol.

Rev. 42:448–461

Wu, C.-H., Bialecka-Fornal, M., and Newman, D. K. 2015. Methylation at the C-2 position of hopanoids

increases rigidity in native bacterial membranes. Elife. 4

Figure Legends

Figure 1. The nitrogen fixation defect of Δ*hpnH* results from a reduction in nodule sizes. (a)

Chemical structure of the extended hopanoid 2-Methyl Bacteriohopanetetrol (2Me-BHT), consisting of a

central pentacyclic core synthesized by the *shc* gene product, a C2 methylation site added by the product

of *hpnP* (grey shading, left), and a tetrol group added by the *hpnH* product (grey shading, right). (**b**)

Average shoot heights and nodules per plant at 24 dpi for *A. afraspera* plants inoculated with wild-type,

Δ*hpnH* or Δ*hpnP B. diazoefficiens*. (**c**) Average acetylene reduction per plant and per nodule at 24 dpi for

A. asfrapera plants inoculated with wild-type, Δ*hpnH* or Δ*hpnP*. (**d**) Representative confocal images of

cross-sections of wild type- and Δ*hpnH*-infected nodules at 24 dpi illustrating plant cell walls (Calcofluor,

cyan), live bacteria (SYTO9, yellow) and membrane-compromised bacteria and plant nuclei (propidium

iodide, magenta). (**e**) Average nodule dry mass and acetylene reduction per nodule dry mass at 24 dpi for

plants inoculated with wild-type, Δ*hpnH* or Δ*hpnP*. Data shown in (**b**), (**c**) and (**e**) was collected from n = 8

plants, with error bars representing one standard deviation. Results of two-tailed t-tests between wild type

and Δ*hpnH* or Δ*hpnP* are denoted as follows: n.s., p>0.01; *****, p<0.01; ******, p<0.001; *******, p<0.0001.

Figure 2. Smaller Δ*hpnH* nodules are offset by increased nodule size heterogeneity over time. (**a**)

Average acetylene reduction per plant (n=4 plants per bar) and (**b**) average nodule dry mass per plant

(n=8 plants per bar) for *A. asfrapera* inoculated with wild-type or Δ*hpnH* over time. Error bars representing

one standard deviation. Results of two-tailed t-tests between wild type and Δ*hpnH* are denoted as follows:

n.s., p>0.05; *****, p<0.05; *******, p<0.0001. (**c-d**) *A. afraspera* inoculated with wild type or Δ*hpnH* at (**c**) 20 dpi

(left) and at (**d**) 40 dpi (right). (**e-f**) Distributions of nodule diameters at 40 dpi for *A. afraspera* inoculated

with (**e**) Δ*hpnH* (right; n=268 nodules pooled from 10 plants) or (**f**) wild type (left; n=227 nodules pooled

from 10 plants).

Figure 3. Nodules containing Δ*hpnH* emerge later and have more heterogeneous growth rates and

final volumes than wild type. (**a**) Comparison of the development of selected wild type- and Δ*hpnH*-

infected nodules over time. (**b**) Nodule growth plots for 74 wild type-infected nodules tracked from 10

plants. (**c**) Nodule growth plots for 84 Δ*hpnH*-infected nodules tracked from 16 plants. (**d**) Schematic of

nodule development in *A. afraspera*. From the left, bacteria (in blue) colonize and invade plant roots

(green) and intracellularly infect a root cell (pink); the time of this initial intracellular infection is considered

t_i and the nodule volume can be described as the volume of the single infected root cell, V_i. This infected

cell proliferates to form a spherical nodule that is visible to the naked eye, at time t_{min} and volume V_{min}.

The infected plant cells continue to proliferate at rate dV/dt until the nodule has fully matured at time t_{max}

and volume V_{max}. (**e**) Fitted growth curve for a sample wild-type nodule illustrating the positions of t_{min},

V_{min}, dV/dt, t_{max}, and V_{max}. (**f-g**) Jitter and box plots of (**f**) dV/dt and (**g**) V_{max} values for all wild type- and

Δ*hpnH*-infected nodules. Results of KS-tests between wild-type and Δ*hpnH* nodules are denoted as

follows: *******, $p<10^{-6}$. (**h**) Scatter plots of dV/dt vs. V_{max} values for wild-type and Δ*hpnH* nodules. Values of

dV/dt and V_{max} below what is observed in the wild-type dataset are highlighted in green. (**i**) Distributions

of t_{min} values (as observed by eye) for nodules from wild type- (white bars) or Δ*hpnH*- (grey bars) infected

plants. N=457 wild-type nodules across 20 plants and 479 Δ*hpnH* nodules across 20 plants.

Figure 4. Small Δ*hpnH* nodules are under-infected compared to wild type. (**a**) Confocal sections of small (<0.5 mm radius) Δ*hpnH*-infected nodules harvested at 40 dpi and small (<0.5 mm radius) wild type-infected nodules harvested at 10 and 25 dpi. (**b**) Confocal sections of larger (>0.5 mm radius) Δ*hpnH*- or wild type-infected nodules harvested at 40 dpi.

Figure 5. Extended hopanoid mutants are delayed at pre-intracellular stage(s) in symbiosis development. (**a**) Confocal sections of nodules from plants co-inoculated with wild type-YFP and Δ*hpnH*-mCherry harvested at 45-55 dpi. Sections were stained with Calcofluor (cyan) and are expressing YFP (yellow) and mCherry (magenta). (**b**) Scatter plot of median YFP intensity per pixel normalized by propidium iodide intensity per pixel (*e.g.* bacteroid density) within infection zones of nodules from plants co-inoculated with wild type-YFP and wild type, as a function of the percentage of wild type-YFP in the inoculum. (**c**) Scatter plot of median propidium iodide intensity per pixel (*e.g.* bacteroid density) within infection zones of nodules from plants co-inoculated with YFP-tagged wild type and untagged wild type, as a function of the percentage of WT-YFP in the inoculum. (**d**) Colony forming units/mL in wild type and Δ*hpnH* cultures grown in BNM supplemented with varying concentrations of kanamycin and spectinomycin at various times post-inoculation. (**e**) Average nodules per plant at 40 dpi for plants inoculated with either wild type or Δ*hpnH* and treated with 50 µg/mL (Δ*hpnH*) or 100 µg/mL (wild type) kanamycin and streptomycin at various time points post-inoculation. Nodule counts are normalized to those observed in non-antibiotic treated plants. (**f**) Images of inoculated plants at 40 dpi after antibiotic treatment at various time points. Untreated plants are shown on the left, with increasing time of antibiotic addition. Error bars represent one standard deviation.

Figure 6. Extended hopanoid mutants are less motile than wild type and attach poorly to surfaces *in vitro*. (**a**) Sample time course of wild type and Δ*hpnH* colony expansion on low-agar PSY plates (dpi = days post-inoculation). Scale bars represent 2 cm. (**b**) Average colony sizes of wild type and Δ*hpnH* over time. N=4 plates per strain; error bars indicate one standard deviation. (**c**) Trajectories of individual wild type (top) and Δ*hpnH* (bottom) cells over a 5 minute time course in PSY. (**d**) Distributions of mean-speed

s for motile wild type (N=359) and Δ*hpnH* (N=91) cells for trajectories in **d**. Dotted lines indicate the

means of the distributions. (**e**) Trajectories of individual wild type (top) and Δ*hpnH* (bottom) cells over a 5

minute time course in BNM. (**f**) Distributions of mean-speeds for motile WT (N=421) and Δ*hpnH* (N=141)

cells in BNM for trajectories in **e**. Dotted lines indicate the means of the distributions. (g) Jitter and box

plots of surface attachment (*e.g.* the percent of the field of view covered with cells) of WT and Δ*hpnH*

after 2 hours of incubation on glass in PSY or BNM. N=40 fields of view per condition. Results of two-

tailed t-tests between wild type and Δ*hpnH* are denoted as follows: n.s., p>0.05; ***, p<0.00001.

Figure 7. Consequences of extended hopanoid loss in *A. afraspera* nodule development.

Schematic representation of *A. afraspera* wild-type root nodule development (top row; white background)

and defects in development associated with extended hopanoid loss (bottom row; grey background).

Early in development, fewer Δ*hpnH* cells are motile (**a**) and competent to attach to root surfaces (**b**),

leading to a delay in establishment of stable root colonies. At later stages, slow growth of Δ*hpnH* into the

root interior, or poor uptake by and division within host cells (**c**) may generate "patchy", or under-

populated infection zone that is propagated as the nodule grows (**d**). Alternately, fully-infected Δ*hpnH*

nodules may lose symbionts to symbiont cell death (**e**) via poor bacteroid survival or plant-directed

symbiosome degradation.

Supplemental Data

Table S1. Motile cell counts and mean swimming speeds for wild-type and Δ*hpnH B. diazoefficiens*.

a.

hpnP　*shc*　*hpnH*

b.

avg. shoot height (cm)

WT　*ΔhpnH*　*ΔhpnP*

** n.s.

avg. nodules per plant

WT　*ΔhpnH*　*ΔhpnP*

n.s. n.s.

c.

avg. acetylene reduced per plant (a.u./hr)

WT　*ΔhpnH*　*ΔhpnP*

** n.s.

avg. acetylene reduced per nodule (a.u./hr)

WT　*ΔhpnH*　*ΔhpnP*

* n.s.

d.

WT

ΔhpnH

e.

avg. nodule dry mass per plant (mg)

WT　*ΔhpnH*　*ΔhpnP*

*** n.s.

avg. acetylene reduced per mg nodule (a.u./hr)

WT　*ΔhpnH*　*ΔhpnP*

n.s. n.s.

Figure 1.

a. avg. acetylene reduced per plant (a.u./hr)

WT □ Δ*hpnH* ▨

***   *   n.s.   n.s.

16dpi 24dpi 32dpi 40dpi

b. avg. nodule dry mass per plant (mg)

WT □ Δ*hpnH* ▨

*   ***   n.s.   n.s.

16dpi 24dpi 32dpi 40dpi

e. count

Δ*hpnH* 40dpi   avg. 0.73 mm

0.0   0.5   1.0   1.5   2.0
nodule radius (mm)

f. count

WT 40dpi   avg. 0.88 mm

0.0   0.5   1.0   1.5   2.0
nodule radius (mm)

c. 20 dpi

5 cm

WT   Δ*hpnH*

d. 40 dpi

5 cm

WT   Δ*hpnH*

Figure 2.

Figure 3.

Figure 4.

a.

b. WT-YFP + WT co-inoculation

median YFP intensity per bacteroid

$R^2 = 0.87$

3.5 X decrease

%WT-YFP in inoculum

c. WT-YFP + WT co-inoculation

median bacteroid density per infection zone (a.u./pixel)

$R^2 = 0.761$

%WT-YFP in inoculum

d.

log(cfu/mL)

time in Kan/Strep (hours)

WT: 100 μg/mL
Δ*hpnH*: 75 μg/mL
Δ*hpnH*: 50 μg/mL
Δ*hpnH*: 25 μg/mL

e.

nodules at 40 dpi (relative to untreated plants)

WT

Δ*hpnH*

time of Kan/Strep addition (dpi)

f.

WT

5 cm

ctrl 0.5 1.5 2.5 3.5 4.5 6.5 8.5 10.5 12.5

time of Kan/Strep addition (dpi)

Δ*hpnH*

ctrl 0.5 1.5 2.5 3.5 4.5 6.5 8.5 10.5 12.5

time of Kan/Strep addition (dpi)

Figure 5.

Figure 6.

Figure 7.

Figure S1. Confocal images of cross-sections of wild type-infected *A. afraspera* nodules at 24 dpi illustrating plant cell walls (Calcofluor, cyan), live bacteria (SYTO9, yellow) and membrane-compromised bacteria and plant nuclei (propidium iodide, magenta). Nodules were collected from 3 plants.

Figure S2. Confocal images of cross-sections of Δ*hpnH*-infected *A. afraspera* nodules at 24 dpi illustrating plant cell walls (Calcofluor, cyan), live bacteria (SYTO9, yellow) and dead bacteria and plant nuclei (propidium iodide, magenta). Nodules were collected from 3 plants. White boxes highlight small nodules. White arrow indicates a likely plant defense reaction.

Figure S3. Average (**a**) shoot height, (**b**) nodules per plant, (**c**) nodule dry weight per plant, (**d**) acetylene reduction per plant, (**e**) acetylene reduction per nodule, and (**f**) acetylene reduction per nodule dry weight for *A. asfrapera* inoculated with wild-type or Δ*hpnH* at 40 dpi. N=4 plants per bar; error bars represent one standard deviation. Results of two-tailed t-tests between wild type and Δ*hpnH* are denoted as follows: n.s., p>0.05.

Figure S4. Reconstructed images of the root system of a wild type-infected *A. afraspera* plant. Nodules fully visible in at least five time points are indicated with black arrowheads.

Figure S5. Reconstructed images of the root system of a Δ*hpnH*-infected *A. afraspera* plant. Nodules fully visible in at least five time points are indicated with black arrowheads.

Figure S6. Nodule growth plots for all 74 wild type-infected nodules fit with quadratic (orange; long dashed lines), exponential (yellow; short dashed lines), or sigmoidal (blue; solid lines) models. Standard errors (SE) for each model are shown.

Figure S7. Nodule growth plots for all 84 Δ*hpnH*-infected nodules fit with quadratic (orange; long dashed lines), exponential (yellow; short dashed lines), or sigmoidal (blue; solid lines) models. Standard errors (SE) for each model are shown.

Figure S8. (**a**) Jitter and box plots of t_{max} values for all wild type- and Δ*hpnH*-infected nodules. (**b**) Jitter and box plots of maximum growth windows for all wild type- and Δ*hpnH*-infected nodules. (**c**) Jitter and box plots of t_{min} values (as determined by extrapolation using sigmoidal fits of nodule growth curves) for all wild type- and Δ*hpnH*-infected nodules, in which V_{min} is defined as 0.05 mm³, 0.1 mm³, 0.2 mm³. Green shading highlights negative t_{min} values. Results of KS-tests between wild-type and Δ*hpnH* nodules are denoted as follows: *, $p<0.05$; n.s., $p>0.05$.

Figure S9. (**a-b**) Scatter plots of t_{max} vs. (**a**) **dV/dt** and (**b**) V_{max} for all wild type- (open circles) and Δ*hpnH*- (grey circles) infected nodules. Green regions highlight values below what is observed for wild type. (**c-d**) Scatter plots of maximum growth windows vs. (**c**) **dV/dt** and (**d**) V_{max}. (**e-f**) Scatter plots of t_{min} vs. (**c**) **dV/dt** and (**d**) V_{max}. (**g-h**) Scatter plots of t_{min} vs. (**a**) t_{max} and (**b**) maximum growth windows.

Figure S10. Confocal sections of small (<0.5 mm radius) Δ*hpnH*-infected nodules harvested at 40 dpi. Sections were stained with Calcofluor (cyan), SYTO9 (yellow), and propidium iodide (magenta). N=74 nodules harvested from 5 plants. White boxes highlight under-infected nodules. Magenta boxes indicate nodules primarily containing membrane-compromised cells.

Figure S11. Confocal sections of large (>0.5 mm radius) Δ*hpnH*-infected nodules harvested at 40 dpi. Sections were stained with Calcofluor (cyan), SYTO9 (yellow), and propidium iodide (magenta). N=87 nodules harvested from 5 plants. White boxes highlight under-infected nodules.

Figure S12. Confocal sections of small (<0.5 mm radius) wild type-infected nodules harvested at 10 dpi. Sections were stained with Calcofluor (cyan), SYTO9 (yellow), and propidium iodide (magenta). N=80 nodules harvested from 5 plants. White boxes highlight under-infected nodules.

Figure S13. Confocal sections of small (<0.5 mm radius) wild type-infected nodules harvested at 25 dpi. Sections were stained with Calcofluor (cyan), SYTO9 (yellow), and propidium iodide (magenta). N=82 nodules harvested from 5 plants. Magenta boxes indicate nodules primarily containing membrane-compromised cells.

Figure S14. Confocal sections of wild type-infected nodules harvested at 40 dpi. Sections were stained with Calcofluor (cyan), SYTO9 (yellow), and propidium iodide (magenta). N=117 nodules harvested from 5 plants. Magenta boxes indicate nodules primarily containing membrane-compromised cells.

Figure S15. Average shoot height (**a**) and number of nodules (**b**) for plants co-inoculated with Δ*hpnH*-mCherry and WT-YFP strains, recorded at 45 dpi. Average shoot height (**c**) and number of nodules (**d**) for plants co-inoculated with WT and WT-YFP strains, recorded at 40 dpi. Average shoot height (**e**) and number of nodules (**f**) for plants co-inoculated with Δ*hpnH* and Δ*hpnH*-mCherry strains, recorded at 50 dpi. N=7-8 plants per bar for all panels. Error bars represent one standard deviation. Results of two-tailed t-tests are denoted as follows: n.s., p>0.05; ***, p<0.0001.

Figure S16. (**a**-**d**) Intensity ratio of YFP to mCherry (**a**), mCherry intensity (**b**), and YFP intensity (**c**) per pixel within infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and WT-YFP strains. (**d**) Cross-sectional area of infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and WT-YFP strains. For (**a**-**d**), N=132, 125, 143, 143 and 110 nodules for 10%, 25%, 50%, 75% and 90% WT-YFP strain mixtures, respectively, which were sectioned and fixed between 45-50 dpi. (**e**) Nodule volume distributions from plants co-inoculated with Δ*hpnH*-mCherry and WT-YFP strains at 45 dpi. Sample sizes are N = 251, 200, 227, 204, and 149 nodules pooled from N = 8, 7, 7, 8, and 7 plants for the 10%, 25%, 50%, 75% and 90% WT-YFP strain mixtures, respectively. (**f**) Scatter plots of mCherry vs. YFP intensities per pixel within infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and WT-YFP strains. (**g**-**h**) Scatter plots of YFP/mCherry intensity ratios per pixel in infection zones vs. infection zone (**g**) and nodule (**h**) cross-section areas for nodules co-inoculated with Δ*hpnH*-mCherry and WT-YFP. Scatter plots contain data pooled from all ratios.

Figure S17. (**a**-**d**) Intensity ratio of YFP to propidium iodide (PI) (**a**), PI intensity (**b**), and YFP intensity (**c**) per pixel within infection zones of nodules co-inoculated with WT and WT-YFP strains. (**d**) Cross-sectional area of infection zones of nodules co-inoculated with WT and WT-YFP strains. For (**a**-**d**), N = 141, 95, 134, 147, 133, and 167 nodules for 10%, 25%, 50%, 75%, 90% and 100% WT-YFP strain mixtures, respectively, which were sectioned and fixed between 40-45 dpi. (**e**) Nodule volume distributions from plants co-inoculated with WT and WT-YFP strains at 40 dpi. Sample sizes are N = 183, 116, 161, 172, 232, and 248 nodules pooled from N = 8, 7, 8, 8, 8, and 8 plants for the 10%, 25%, 50%, 75% and 90% WT-YFP strain mixtures, respectively. (**f**) Scatter plots of PI vs. YFP intensities per pixel within infection zones of nodules co-inoculated with WT and WT-YFP strains. (**g**-**h**) Scatter plots of YFP/PI intensity ratios per pixel in infection zones vs. infection zone (**g**) and nodule (**h**) cross-section areas for nodules co-inoculated with WT and WT-YFP strains. Scatter plots contain data pooled from all ratios.

Figure S18. (**a**-**d**) Intensity ratio of mCherry to SYTO9 (**a**), SYTO9 intensity (**b**), and mCherry intensity (**c**) per pixel within infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and Δ*hpnH* strains. (**d**) Cross-sectional area of infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and Δ*hpnH* strains. For (**a**-**d**), N = 117, 107, 128, 137, 103 and 50 nodules for 10%, 25%, 50%, 75%, 90% and 100% Δ*hpnH*-mCherry strain mixtures, respectively, which were sectioned and fixed between 50-55 dpi. (**e**) Nodule volume distributions from plants co-inoculated with Δ*hpnH*-mCherry and Δ*hpnH* strains at 45 dpi. Sample sizes are N = 150, 222, 191, 254, 297, and 236 nodules pooled from N = 7, 7, 7, 8, 8, and 8 plants for the 10%, 25%, 50%, 75% and 90% WT-YFP strain mixtures, respectively. (**f**) Scatter plots of mCherry vs. SYTO9 intensities per pixel within infection zones of nodules co-inoculated with Δ*hpnH*-mCherry and Δ*hpnH* strains. (**g**-**h**) Scatter plots of mCherry/SYTO9 intensity ratios per pixel in infection zones vs. infection zone (**g**) and nodule (**h**) cross-section areas for nodules co-inoculated with Δ*hpnH*-mCherry and Δ*hpnH* strains. Scatter plots contain data pooled from all strain ratios.

Figure S19. Average (**a**) shoot height, (**b**) shoot dry mass and (**c**) root dry mass for non-inoculated *A. afraspera* plants grown in BNM supplemented with kanamycin, streptomycin or tetracycline for 2 weeks under normal growth conditions. N=4 plants per condition; error bars represent one standard deviation. (**d-e**) Images of *A. afraspera* plants after 2 weeks of antibiotic treatment. Asterisks indicate plants grown in tetracycline-supplemented medium.

Figure S20. Surface attachment of wild type (**a,c**) and Δ*hpnH* (**b,d**) incubated on glass coverslips in various media. For each panel, raw phase images (top row), background-subtracted images (middle row), and binary images with cells shown in black (bottom row) are shown. Scale bars represent 20 μm.