**Effects of two centuries of global environmental variation on phenology and physiology of**

*Arabidopsis thaliana*

200 years of *A. thaliana* phenotypic variation

Victoria L. DeLeo[1], Duncan N. L. Menge[2], Ephraim M. Hanks[3], Thomas E. Juenger[4], Jesse R.

Lasky[5,6,7]

[1]Intercollege Graduate Degree Program in Plant Biology

Pennsylvania State University

University Park, PA 16802

[2]Department of Ecology, Evolution, and Environmental Biology

Columbia University

New York, NY 10027

dm2972@columbia.edu

[3]Department of Statistics

Pennsylvania State University

University Park, PA 16802

hanks@psu.edu

[4]Department of Integrative Biology

University of Texas at Austin

Austin, TX 78712

tjuenger@austin.utexas.edu

[5]Department of Biology

Pennsylvania State University

University Park, PA 16802

lasky@psu.edu

Past affiliations:

[6]Department of Ecology, Evolution, and Environmental Biology

Columbia University

New York, NY 10027

[7]Earth Institute

Columbia University

New York, NY 10025

Corresponding information:

Victoria DeLeo

814-863-8854

vldeleo@psu.edu

**Abstract**

Intraspecific trait variation is caused by genetic and plastic responses to environment. This intraspecific diversity is captured in immense natural history collections, giving us a window into trait variation across continents and through centuries of environmental shifts. Here we tested if hypotheses based on life history and the leaf economics spectrum explain intraspecific trait changes across global spatiotemporal environmental gradients. We measured phenotypes on a 216-year time series of *Arabidopsis thaliana* accessions from across the native range and applied spatially varying coefficient models to quantify region-specific trends in trait coordination and trait responses to climate gradients. All traits exhibited significant change across space and/or through time. For example, $\delta^{15}N$ decreased over time across much of the range and leaf C:N increased, consistent with predictions based on anthropogenic changes in land use and atmosphere. Plants were collected later in the growing season in more recent years in many regions, possibly because populations shifted toward more spring germination and summer flowering as opposed to fall germination and spring flowering. When climate variables were considered, collection dates were earlier in warmer years, while summer rainfall had opposing associations with collection date depending on regions. There was only a modest correlation among traits, indicating a lack of a single life history/physiology axis. Nevertheless, leaf C:N was low for summer- versus spring-collected plants, consistent with a life history-physiology axis from slow-growing winter annuals to fast-growing spring/summer annuals. Regional heterogeneity in phenotype trends indicates complex responses to spatiotemporal environmental gradients potentially due to geographic genetic variation and climate interactions with other aspects of environment. Our study demonstrates how natural history collections can be used to

67    broadly characterize trait responses to environment, revealing heterogeneity in response to

68    anthropogenic change.

69    **Keywords**: Generalized Additive Model, abiotic stress, climate change, $\Delta^{13}C$, phenology

## Introduction

An organism's fitness is determined by the interaction between its traits and its environment. Organisms respond to environmental gradients in diverse ways, including genetic and plastic shifts in life history, phenology, and physiology (Burghardt, Metcalf, Wilczek, Schmitt, & Donohue, 2015; Reich, 2014; Wright et al., 2004). Spatial and temporal environmental gradients can promote phenotypic plasticity or generate varying selection across which populations adapt to local conditions (Bradshaw, 1965; Henn et al., 2018; Joshi et al., 2001; Leimu & Fischer, 2008; Linhart & Grant, 2002; Matesanz, Gianoli, & Valladares, 2010; Turesson, 1922). By studying how phenology and physiology change across environments through space and time we can learn about mechanisms of adaptive environmental response and biological constraints.

Anthropogenic global change has led to dramatic phenotypic changes in many organisms. For example, many species are shifting their ranges poleward and temperate spring phenology is advancing (Parmesan & Yohe, 2003). However, anthropogenic global change is multi-faceted, involving climate, nitrogen deposition, atmospheric $CO_2$, and land use. Our understanding of the specific environmental drivers of phenotypic change has been hampered both by insufficient long-term datasets and by the complexities of interacting and correlated environmental variables. Furthermore, many populations and species do not exhibit the stereotypic advancing temperate phenology and poleward range shifts (Both et al., 2004; CaraDonna, Iler, & Inouye, 2014; Park et al., 2018). These diverse responses can be caused by geographic variation in the rate of environmental change or by intraspecific genetic variation, clouding our understanding of anthropogenic impacts. To address these challenges, we collected physiology and phenology data for *Arabidopsis thaliana* (hereafter, *Arabidopsis*) specimens over the last 200 years and across its native range, and we tested relationships between climate and traits using spatial

generalized additive models (GAMs) to account for geographic structure in environmental

response.

*Arabidopsis* is a powerful system for studying phenotypic change across space and climate

gradients. Past studies have found that *Arabidopsis* populations exhibit genetic differences

among populations likely due to isolation by distance (Alonso-Blanco et al., 2016; Horton et al.,

2012; Ostrowski et al., 2006; Platt et al., 2010) and local adaptation (Fournier-Level et al. 2011,

Hancock et al. 2011, Lasky et al. 2012) . Genetic differences in flowering time among

populations may be due to local adaptation (Atwell et al., 2010; Burghardt et al., 2015; Tabas-

Madrid et al., 2018), with northern genotypes having later flowering times (Atwell et al., 2010;

Stinchcombe et al., 2004). *Arabidopsis* physiology also shows evidence of a role in local

adaptation. Genotypes from regions of greater precipitation have faster growth and lower leaf

vein density, and genotypes from colder temperatures have increased leaf thickness and wider

leaf minor vein cross section (Adams, Stewart, Cohu, Muller, & Demmig-Adams, 2016; Sack et

al., 2012; Sartori et al., 2018). These findings provide a lens through which to interpret

phenotypic variation among plants in nature. In turn, museum collections offer broadly

distributed sampling in space and time to test the relationships between phenotypes and

environment in nature (Lang, Willems, Scheepens, Burbano, & Bossdorf, 2018; Willis et al.,

2017).

Multiple frameworks of plant life history and physiology variation correspond to a continuum of

fast to slow life histories (e.g. Grime, 1977; Westoby, 1998; I. J. Wright et al., 2004). Individuals

that use a fast strategy are characterized by fast relative growth, early reproduction, and intensive

use of nutrients or water, while individuals with a slow strategy are characterized by slow

growth, late reproduction, and more measured use of nutrients and water. The Leaf Economics

Spectrum (LES) hypothesizes that leaves vary along single life history-physiology axis from fast

to slow in association with large-scale climate gradients (Reich, 2014; Reich et al., 2003; Wright

et al., 2004). The LES predicts that lower nitrogen concentration leaves (high C:N, low

proportion N) are found in drier and in hotter areas, possibly in part because of investment in

non-photosynthetic leaf features, e.g. veins (Blonder, Violle, Bentley, & Enquist, 2011; Easlon et

al., 2014; Sack et al., 2012) (see Table 1 for phenotype/environment hypotheses and how they

relate to the fast/slow framework). Low N leaves are thicker (high mass to area) and provide

protection against stress (drought) at the expense of a N investment in photosynthesis, resulting

in a slower life cycle (Evans, 1989; Stocking & Ongun, 1962). Although community-wide

turnover in mean traits across natural environments often follows LES predictions, *within* species

trait variation often defies LES predictions (Anderegg et al., 2018; Hu et al., 2015; J. P. Wright

& Sutton-Grier, 2012). Nevertheless, *Arabidopsis* exhibits genetic variation in traits that

corresponds to LES predictions (Easlon et al., 2014; Sartori et al., 2018; Vasseur, Violle,

Enquist, Granier, & Vile, 2012); individuals with rapid life histories have physiological traits tied

to fast growth and resource acquisition (e.g. high stomatal conductance, high leaf area relative

mass (Specific Leaf Area, or SLA)) (Lovell et al., 2013; McKay, Richards, & Mitchell-Olds,

2003; Sartori et al., 2018; Wolfe & Tonsor, 2014).

Isotopic signatures provide clues to organismal life histories. In plants, $\Delta^{13}C$ measures

discrimination against $^{13}C$ in photosynthesis and is an indicator of $pCO_2$ within leaves ($C_i$)

relative to atmospheric $pCO_2$ ($C_a$) (Farquhar, O'Leary, & Berry, 1982). $C_i$ declines when stomata

are closed, which may be a conservative life history response to soil drying, while $C_a$ declines

with elevation. Thus, we expect $\Delta^{13}C$ to increase in moist growing environments and decrease

with elevation (Diefendorf, Mueller, Wing, Koch, & Freeman, 2010; Farquhar et al., 1982; Zhu,

Siegwolf, Durka, & Körner, 2010) (Table 1). $\delta^{15}N$ (the ratio of $^{15}N$ to $^{14}N$) can be affected by

nitrogen allocation and so might reflect variation in C:N (Stock & Evans, 2006). However,

variation in leaf N and $\delta^{15}N$ may also directly reflect changing environments (N deposition,

biogeochemical cycling, e.g. Pardo *et al.* 2007) rather than plant traits, with both increasing with

temperature and leaf N increasing and $\delta^{15}N$ decreasing with rainfall (Table 1).

Fast/slow strategies also correspond to phenological variation, especially in annual plants. In

seasonal environments, phenology is constrained by seasonality and simultaneously determines

the environment encountered during vulnerable stages. *Arabidopsis* development can be highly

sensitive to moisture, temperature, and photoperiod (Burghardt et al., 2015; Wilczek et al.,

2009). For example, although warmth can increase growth rates, many *Arabidopsis* genotypes

require winter cold cues (known as vernalization) to transition to spring flowering. Fast life

histories can allow spring or summer annual life cycles, where a plant germinates and flowers

within a single season, while slow life histories and vernalization requirements result in a winter

annual cycle, where a plant germinates in the fall and flowers the following spring. Rapid

development and reproduction can allow *Arabidopsis* plants to escape drought (McKay et al.,

2003), while slower flowering plants can exhibit drought avoidance strategies of minimizing

water loss (e.g. through stomatal closure) or maximizing water uptake (Kenney, Mckay,

Richards, & Juenger, 2014; Ludlow, 1989). Because herbarium specimens are typically

reproductive, it is challenging to infer germination times based on collection dates. However,

information on physiology and climate preceding collection may provide information on life

history variation.

Standardized metrics facilitate the comparison of ecologically relevant phenological variation

among sites that differ in climate and seasonal timing.  Photothermal units (PTUs) integrate

developmental time under favorable temperatures and light and account for much of the

environmental influence on flowering dates in *Arabidopsis* (Brachi et al., 2010; Wilczek et al.,

2009). In essence, PTUs estimate how far along in a growing season an event occurs. Measures

of developmental time standardized to environmental conditions can better capture genetic

variation in development compared to raw flowering dates in *Arabidopsis*, the latter of which are

strongly driven by environment (Brachi et al., 2010). Without knowing an exact germination

date, it is impossible to perfectly approximate the climate experienced in the wild, yet even a

rough estimate using an arbitrary date allows us to compare changes in the climate experienced

at flowering. Variation in herbarium collection dates is a reliable proxy for variation in

phenology of flowering date (Davis, Willis, Connolly, Kelly, & Ellison, 2015; MacGillivray,

Hudson, & Lowe, 2010; Miller-Rushing, Primack, Primack, & Mukunda, 2006). In addition, low

PTUs at collection hints at a winter annual growing pattern (because growth is occurring before

PTU calculations begin), so we can use this measurement to search for regional variation in life

history. Combined with existing knowledge the phenology of ecotypes from different sites, PTUs

may help reveal phenological adaptation along spatial and temporal environmental gradients.

Here, we leverage the immense fieldwork underlying natural history collections to investigate

how intraspecific diversity is structured through time and along spatiotemporal climate gradients.

Specifically, we use thousands of *Arabidopsis* specimens that span over 200 years of sampling

across *Arabidopsis*' range in Eurasia and Northern Africa. We quantify the spatial patterns of

*Arabidopsis*' phenotypic variation along environmental gradients, which allows us to put

temporal trends in context. We hypothesized that for natural *Arabidopsis* populations,

phenotype-environment correlations would follow fast-slow predictions of LES and phenology

traits (Table 1). We combine these records with global gridded climate data to ask three

questions about *Arabidopsis* in nature:

1.  To what degree does intraspecific trait variation among wild individuals fall along a

single coordinated life history-physiology axis?

2.  Do life history and physiology vary across spatial environmental gradients in long-term

average conditions, suggesting adaptive responses consistent with the LES?

3.  Have life history and physiology changed over the last two centuries? In particular, have

changes tracked climate fluctuations, suggesting adaptive responses consistent with the

LES?

 **Materials & Methods**

 *Samples*

Our set of samples (N= 3443) included *Arabidopsis thaliana* herbarium and germplasm

accessions with known collection dates between 1794 and 2010 from the native range of

*Arabidopsis* in Europe, the Middle East, Central Asia, and North Africa (Hoffmann, 2002).

Wild-collected germplasm accessions with known collection date and location (N = 447) were

included only in models of phenology. Information on germplasm accessions came from the

*Arabidopsis* Biological Resource Center (https://abrc.osu.edu/). For each herbarium specimen

(N=2663) we visually verified species identification and reproductive status as simultaneously

flowering and fruiting. Samples that were only fruited/senesced, only flowering, or had neither

open flowers nor fruits were excluded to focus on a relatively uniform developmental stage (see

Supplementary Table 1). This consistency is important for assessing C:N, since progression of

plant development involves reallocation of nutrients, and for a meaningful characterization of

phenology with collection date (Himelblau & Amasino, 2001). Furthermore, too few samples

(178/2663) were in other phenological stages to allow for a rigorous comparison. We excluded

dozens of misidentified specimens, highlighting the importance of verification of information in

natural history collections (cf. unverified data in some online databases). Samples with too low

precision in collection date (less precise than a single month) were excluded from phenological

analysis.

*Leaf traits*

To test LES hypotheses for response to environment, we measured $\Delta^{13}$C, $\delta^{15}$N, and C:N from

leaf tissue of herbarium samples. We removed and pulverized leaf samples (mean weight = 2.75

mg) of a subset of our quality-checked herbarium specimens and sent them to the UC-Davis

Stable Isotope Facility. In total, we obtained values for $\delta^{15}$N, $\delta^{13}$C, C:N, and proportion N in 459

accessions, although 5 samples failed for $\delta^{13}$C and 1 sample was missing failed for both C:N and

proportion N values.

We measured leaf $\delta^{13}$C (isotope ratio), but atmospheric $\delta^{13}$C has changed dramatically over the

time period of this study due to fossil fuel emissions. Thus we converted leaf isotope ratio ($\delta^{13}$C)

to discrimination ($\Delta^{13}$C) using an estimate of the atmospheric $\delta^{13}$C time series (McCarroll &

Loader, 2004) from 1850 to 2000, continuing linear extrapolation beyond 2000, using the 1850

value for earlier specimens, and the equation of Farquhar et al. (1989), $\Delta = \frac{\delta_a - \delta_p}{1 + \delta_p}$, where $\delta_a$ is the

isotope ratio in the atmosphere and $\delta_p$ is the isotope ratio in plant tissue (ratios relative to a

standard).

*Phenology*

To estimate accumulated photothermal units (PTU) at date of collection, we used the equation of

Burghardt et al. (2015) to model the hourly temperature values for the accumulation of sunlight

degree hours between January 1 and dusk on the date of collection at each accession's

coordinate. Daylength was approximated with the R package geosphere (Hijmans, 2017).

Monthly temperature values for the period 1900-2010 came from the Climate Research Unit time

series dataset v4.01 (Harris, Jones, Osborn, & Lister, 2014). PTUs were only calculated for

specimens collected after 1901 (N = 2488) due to the historical limit of the monthly temperature

data. Daily temperatures were interpolated from monthly temperatures using the function

splinefun in R on the "periodic" setting. PTUs calculated from January 1[st] will not completely

account for the climate experienced by plants that germinate in the fall. However, for the same

developmental time (PTUs), determined by weather conditions, winter annuals are expected to

flower earlier in a growing season compared to spring annuals. Comparing changes in PTU at

collection to changes in date of collection might provide clues as to where climate is driving

flowering time shifts and where flowering time is responding to pressures other than

temperature.

*Statistical analysis*

*Arabidopsis* displays substantial genetic diversity in environmental response between genotypes

from different regions (e.g. Exposito-Alonso et al., 2018; Lasky, Forester, & Reimherr, 2018).

Thus, we employed a regression model with spatially varying coefficients (generalized additive

models, GAMs) to account for regional differences in responses to environment, much of which

may have a genetic component (Wheeler & Waller, 2009; Wood, 2006). GAMs allow fitting of

parameters that vary smoothly in space (i.e. parameter surfaces) and can thus capture spatially

varying relationships between predictors and the response of interest, such as we see in

ecological processes (Yee & Mackenzie, 2002; Yee & Mitchell, 2006). The spatially varying

coefficients fit by GAM allow us to infer from the data where relationships between variables

change, as opposed to binning data into a set of fixed (and possibly artificially defined) regions.

In a standard linear statistical model, the effect of a covariate x (i.e., the effect of x = January

Minimum Temperature) at site i is a linear function of the covariate: $x_i * \beta$. Note that differences

across sites in $x_i * \beta$ are completely controlled by differences in the covariate x at different sites.

In the spatially-varying GAMs we consider in this study, we allow the effect of covariates to

vary across space, with the effect of a covariate x at location i being $x_i*\beta_i$ where $\beta_i$ is the linear

effect of x at the $i^{th}$ spatial location. That is, we model allow the effect of the covariate x to vary

across space. Thus, the effect of (for example) January Minimum Temperature might be different

in Europe than it would be in Southeast Asia.

It would be impossible to uniquely identify a completely different effect $\beta_i$ at every location, as

we only have one replicate at each site, and there would be no replication. Spatially-varying

GAMs address this identifiability by smoothing $\beta_i$ across space and requiring that effects $\beta_i$ and

$\beta_j$ for sites i and j that are close in space be very similar to each other. The degree of spatial

smoothness in the GAM is chosen by cross-validation to ensure that the GAM does not overfit

the data. Thus, spatially-varying coefficient models provide an approach for modeling variability

in the effect of covariates across space, with the effect constrained to vary slowly across space to

ensure identifiability and combat overfitting.

Each cell in our 140x200 grid model rasters corresponded to 53.1 km East/West at the lowest

latitude (28.16°, versus 20.3 km at 68.18° N) and 31.8 km North/South (calculated using

Vincenty ellipsoid distances in the geosphere package). Smaller grid cells allow for more finely

smoothed slope values. Model predictions farther than 200km from a sampled accession were

discarded when visualizing results.

We selected climate variables based on knowledge of critical *Arabidopsis* developmental times

and likely environmental stressors: average temperature in April, when warmth is expected to

accelerate development (AprilMean in the models), minimum temperature in January, when

vernalization cues are likely accumulating or when Mediterranean plants are in early growth

(JanMinimum), and July aridity index (AI), when summer drought may be most likely (Fournier-

Level et al., 2013; Hoffmann, 2002; Lasky et al., 2012; Wilczek, Cooper, Korves, & Schmitt,

2014). Our analyses should not be highly sensitive to the exact calendar month chosen, given the

high correlation in conditions between consecutive months (e.g. warm Aprils tend to be followed

by warm Mays). Aridity index was calculated from July precipitation divided by July potential

evapotranspiration (PET) (United Nations Environment Program, 1997). These climate gradients

were generally not strongly correlated (July Aridity to April Mean Temperature r = -0.33; July

Aridity to January Minimum Temperature r = -0.12; January Minimum Temperature to April

Mean Temperature r = 0.71 by Pearson's product-moment correlation). We took temperature,

precipitation, and PET values from the Climate Research Unit time series dataset, using values

for the year of collection (New, Hulme, & Jones, 2000).

First, to study trait correlations that might indicate a fast-slow life history (Question 1), we

performed a Principal Components Analysis of flowering time, $\Delta^{13}C$, $\delta^{15}N$, and C:N ratio and

tested pairwise associations between traits.  We considered how traits co-vary by calculating the

Pearson's correlation coefficients between traits and by Principal Components Analysis. We also

fit GAMs (described in detail below) with spatially varying intercepts allowing measured

phenotypes as both response and predictor variables to observe how the correspondence of traits

changes through space.

Second, to study phenotypic responses to spatial gradients in long-term average climates

(Question 2), we fit models with spatially varying coefficients for long-term, 50-year climate

averages at each location ("spatial climate models", Equation 1). We scaled these climate

covariates and year of collection to unit standard deviation (Hijmans, Cameron, Parra, Jones, &

Jarvis, 2005). In these models of responses to long-term average conditions, year of collection

can be considered a nuisance variable, accounting for temporal variation at a location that may

be important but is not the focus of this specific model. In spatial climate models, we used a

single global intercept. Spatial climate models included specimens from all years with phenotype

data.

$$\text{Phenotype}_{ij} = \beta_1 \text{year}_j + \beta_{2j} \text{AI}_j + \beta_{3j} \text{JanMinimum}_j + \beta_{4j} \text{AprilMean}_j + \mu + \text{error}_{ij} \qquad \text{(eqn 1)}$$

In all models, the subscript j denotes location and i denotes year of collection. For the temporal

climate anomaly model, the spatially varying intercept is denoted by $\mu_j$, where the "j" subscript

indicates that the intercept varies with location.  The errors are assumed to be independent, be

normally distributed, and have constant variance.

Next, to assess how phenotypes have changed across the last two centuries (Question 3), we

tested a model with spatially-varying coefficients for the effect of year, allowing for geographic

variation in temporal trends (hereafter, "year models"). The model also included spatially

varying intercepts to account for regional differences in long-term mean phenotypes. Year

models included all specimens with data for a particular phenotype.

Finally, to assess how temporal fluctuations in climate drive phenotypic change (Question 3), we

fit models with the three climate covariates for the year of collection (Equation 2). We converted

climate covariates to local anomalies by standardizing them relative to the entire time-series for a

given grid cell to unit standard deviation and mean zero ("temporal climate anomaly models").

In standardizing climate fluctuations to the climate record of a location, we assume that the effect

of an anomaly on the response variable is best captured by the relative strength (and direction) of

an anomaly (relative to an average anomaly) rather than how extreme an anomaly is in general.

The model also included spatially varying intercepts to account for regional differences in long-

term mean phenotypes. Temporal climate models only included specimens from after 1900,

when we had data on monthly climate from CRU. These models had the following structure

$\text{Phenotype}_{ij} = \beta_1 \text{year}_{dev, ij} + \beta_2 \text{AI}_{dev, ij} + \beta_3 \text{JanMinimum}_{dev, ij} + \beta_4 \text{AprilMean}_{dev, ij} + \mu_j + \text{error}_{ij}$

(eqn 2)

Models were fit in R (version 3.5.0, R Core Team 2011) using the 'gam' function in package

mgcv (version 1.8-17, Wood 2011). We allowed the model fitting to penalize covariates to 0 so

that covariates weakly associated with phenotypes could be completely removed from the model;

thus, using the mgcv package we can achieve model selection through joint penalization of

multiple model terms. Coefficients in spatially varying coefficient models represent the

relationship between each term and phenotype at each geographic point (indexed by *j* in our

models).

We considered two other spatially-varying environmental variables of interest: elevation and N

deposition. We left elevation and nitrogen deposition covariates out of the final models because

inclusion resulted in instability in the numerical routines the GAM software (mgcv) used to

estimate parameters and approximate Hessian matrices needed for confidence intervals. See

supplemental material for more information on these covariates. Including only the variables of

the three climate covariates and year resulted in numerically stable estimates. In addition, scaling

of year and climate variables tended to reduce the concurvity of variables and increase stability.

Code for all the models and plots will be included as a supplement and will be available on

github.

**Results**

*Distribution of samples through time and space*

Samples were broadly distributed, with dense collections in Norway/Sweden, the Netherlands, and Spain (reflecting major herbaria used in the study), and sparser collections to the east (Figure 1). The subset of samples with tissue analysis spanned the extent of the geographic distribution of all samples. Tissue sampling was most dense in Norway, Spain, and the UK and sparser elsewhere in the range. The earliest collection date we used was 1794, but a greater number of samples were available from the 1900s onwards.

Figure 1: (A) Locations of collections used in our analysis. Color of circle corresponds to year of

collection. Accessions that were sampled for tissue are outlined in black. (B) Distribution of

years of collection.  (C) Sample herbarium record from Nepal on April 5, 1952.

*Correlations among phenotypes (question 1)*

We found generally weak correlations among phenotypes of *Arabidopsis* individuals (Question

1). The first two principal components explained only 36.2% and 24.1%, respectively, of the

variance in the five phenotypes of $\Delta^{13}C$, $\delta^{15}N$, date of collection, C:N, and PTU (N = 397). The

first principal component corresponded to a negative correlation between C:N versus day of

collection (bivariate r = -0.189) and PTU (bivariate r = -0.101). Inspecting the relationship

between collection date and C:N further revealed a triangle shape (Figure 2B). That is, there

were no late-collected individuals with high C:N, potentially indicating that the late-collected

plants that we hypothesize are spring/summer annuals also exhibit fast growth strategies. Plants

with the lowest C:N have a less negative relationship to day of collection than plants with a

higher C:N, and ANOVA showed there to be a significant difference between the slopes of the

regression of the 25$^{th}$ and 75$^{th}$ percentiles of C:N (p = 0.0002, Figure S2) (Koenker & Koenker,

2011). When allowing the relationship between C:N and phenology to vary spatially (GAM with

spatially varying coefficients), we found both date of collection and PTU were negatively

correlated with C:N across the *Arabidopsis* native range, but this correlation was not

significantly different from 0 (the 95% confidence interval included 0) (Figure S3). The second

PC corresponded to a negative correlation between $\Delta^{13}C$ and $\delta^{15}N$ (bivariate r = -0.218). C:N and

leaf proportion N are highly correlated (bivariate r = -0.816, Figure S1), so we focus on C:N. See

supplementary material for leaf N results (Figure S16 and S17).

Figure 2: (A) Variation in phenotypes across the native range of *Arabidopsis* for $\Delta^{13}$C, $\delta^{15}$N,

C:N, collection date, and photothermal units (PTU) at collection. Color indicates the fitted mean

value (spatially-varying intercept) of the phenotype from the year-only model. For example,

collection date is earlier in the Mediterranean in comparison to other regions; however, PTUs are

lowest in the central part of the range. (B) Correlations between phenotypes in this study.

Histograms of the measured values of each phenotype are plotted along the diagonal. (C) PCA of

phenotypes. Correlations of phenotypes with principal components are plotted as arrows, with

length multiplied by 3 for ease of viewing. Latitude (correlation with first PC r = -0.182) and

longitude (correlation with first PC r = -0.115) are plotted for geographic context, though they

were not included in PCA. Arrows for latitude and longitude are scaled equally to the arrows for

phenotype correlations.

*Spatial variation in long-term average phenotypes (questions 2 and 3)*

We visualized spatial diversity in phenotypes by plotting the spatial intercept surfaces in the year

only models (Figure 2A). All phenotypes showed significant spatial variation (all GAM smooth

terms significantly different from zero). $\Delta^{13}$C was lower in the Iberian Peninsula and higher in

Russia (GAM smooth term, p = 0.0002). $\delta^{15}$N varied across the range, but with less pronounced

spatial gradients (GAM smooth term, p = 0.01). C:N was higher in the Iberian Peninsula and

central Asia and lower in Russia (GAM smooth term, p = 9e-05). Collection day was earlier

along the Atlantic coast and Mediterranean (GAM smooth term, p = <2e-16). Despite this, PTU

at collection still was higher in the Mediterranean region as well as at far northern, continental

sites (GAM smooth term, p = <2e-16).

## A.   Day of Collection

## B.   Photothermal Units

## C.   δ ¹⁵N

## D.   C:N

Figure 3: Change in phenotypes across years for collection date (A), photothermal units (B), $\delta$

nitrogen (C), and C:N (D). Color indicates the value of the coefficient for year in the model

excluding climate variables, gray shading indicates regions where estimated coefficient is not

significantly different from 0. For example, day of collection and photothermal units have

significantly increased over time in most of the range, but with some exceptions for day of

collection in the south. Inset scatterplot in A shows the significant increase in collection date

with year for samples in the boxed Mediterranean region. Plots to the left of A show the density

of collection dates through the year remains stable through time for Scandinavian collections

within the boxed region (top) but shift toward more collections late in the year in the boxed

Mediterranean collections (bottom).

Several phenotypes have changed significantly across large regions over the study period (1794-

2010, Figure 3 above). For example, C:N ratio increased in later years in much of southwestern

Europe. $\delta^{15}N$ decreased significantly throughout most of the range. Collection date and PTUs

became significantly later in many regions from the Mediterranean to Central Asia, although

collection date became significantly earlier in the extreme south (Morocco and Himalayas).

There was no significant temporal trend in $\Delta^{13}C$ (not shown).

The year trends in phenotypes across the study period were likely partly related to underlying

climate variation. However, collection date, C:N, and $\delta^{15}N$ were still significantly associated

with year of collection even when accounting for temporal anomalies in climate from 1901-2010

(Figures S6, S12, S14).  PTUs were even more negatively related to year of collection after

controlling for temporal anomalies (Figure S8). A notable discrepancy is that Iberian collections

were collected significantly earlier in later years when yearly climate anomalies were accounted

for (Figure S6). There was still no significant temporal trend in $\Delta^{13}$C.

*Phenotype associations with spatiotemporal climate gradients (questions 2 and 3)*

**Date of collection –** In years (temporal climate anomaly models) with a relatively warm April

plants were collected significantly earlier (Figure 4A). Similarly, in locations (spatial climate

models) with warmer temperatures plants were on average collected earlier in the year, though in

many regions these coefficients were non-significant (Figure S7A, B). We also tested

associations with July aridity index (precipitation/PET) and found that plants were collected

significantly earlier in years (temporal climate anomaly models) with dry summers in

Central/Eastern Europe (Figure 4B).

A. Mean April Temperature

B. July Aridity Index

Figure 4: Association between collection day of *Arabidopsis* temporal mean April temperature

anomalies (A) and July aridity index anomalies (B) (compared to 50-year average). Color

indicates the value of the coefficient of the April mean temperature anomaly or July aridity index

anomaly term. In years where April was warmer (positive anomalies), plants were collected

earlier (a negative relationship). In wetter years, plants were collected later in Eastern Europe.

Shading indicates regions where estimated coefficient is not significantly different from 0.

Scatterplots of phenotype measures for individuals within the boxed areas show a decreasing

collection date with mean April temperature anomaly and an increasing collection date with July

aridity index anomaly in Eastern Europe and a decreasing collection date in Central Asia.

**Photothermal units –** To standardize spatiotemporal variation in developmental periods, we also modeled climate associations with PTUs. As expected, there were few areas where temperature anomalies were significantly associated with PTUs, likely due to the ability of PTUs to account for plastic responses (Figure S8). However, in some areas, accumulated PTUs at collection changed significantly in association with spatial temperature gradients, perhaps indicating spatial genetic differences in phenology (Figure S9). Locations with warmer Aprils had plants collected at more PTUs in East-Central Europe and fewer PTUs around the Aegean and Northern Asia. In the spatial climate model, plants from wetter areas in the Mediterranean were collected at lower PTUs (Figure S8, S9).

**Δ $^{13}$Carbon –** Wet summers were not significantly related to $\Delta^{13}$C in any region in either the temporal climate anomaly or spatial climate models (Figure S10, S11). Spatial variation in mean April temperatures was not significantly related to $\Delta^{13}$C, but plants from locations of colder Januaries did have lower $\Delta^{13}$C in Northern Asia and the Iberian Peninsula. (Figure S11B). Although elevation was not included in final models for reasons discussed above, replacing year with elevation in the temporal climate anomaly model showed a significant negative association between elevation and $\Delta^{13}$C (Figure S18D). Including elevation reduced the significance of the relationship between spatial variation in January temperature and $\Delta^{13}$C.

**δ** **¹⁵Nitrogen** – $\delta^{15}$N was significantly higher in wetter years in Iberia, Asia, and Central Europe,

(Figure S12C), but lower in the North of France. Spatial variation in minimum January

temperatures was significantly positively related to $\delta^{15}$N around the North Sea (Figure S13B).

**Leaf C:N** – For the temporal climate anomaly model, plants collected in years with warmer

winters in Iberia had significantly lower C:N ratios (Figure S14B. Leaf C:N differed in response

to April mean temperature and January minimum temperature among locations (spatial climate

models, Figure S15), although the patterns were mostly insignificant.

**Discussion**

Widely distributed species often exhibit considerable phenotypic diversity, a large portion of

which may be driven by adaptive plastic and evolutionary responses to environmental gradients.

Previous studies of intraspecific trait variation in response to environment have tended to focus

on genetic variation of environmental responses in common gardens (*e.g.* Wilczek *et al.* 2009;

Kenney *et al.* 2014), temporal trends in phenology from well-monitored sites (*e.g.* CaraDonna et

al., 2014), or field sampling of individuals from a small number of sites (*e.g.* Jung, Violle,

Mondy, Hoffmann, & Muller, 2010). Here, we complement this literature by studying change in

traits across an entire species range over two centuries, giving us a window into drivers of

intraspecific diversity and regional differences in global change biology. From the accumulated

effort contained in natural history collections, we tested hypotheses about variation in life history

and physiology in response to environment. We observed modest evidence of coordinated

phenological-physiological axes of variation for *Arabidopsis* in nature. We found later flowering

times and higher accumulated photothermal units over the study period across most of the range

and lower $\delta^{15}$N and higher C:N in more recent collections. Additionally, we observed distinct

regional differences in phenology, $\Delta^{13}$C, and C:N in response to rainfall and temperature,

potentially due to genetic differences among populations.

*Intraspecific variation in life history and physiology shows little coordination along a single*

*major axis (question 1)*

We found little evidence for tight coordination among studied phenotypes, fitting with some past

studies that found weak to no support for a single major axis in intraspecific trait variation in

response to environment across diverse plant growth forms (e.g. Albert *et al.* 2010; Wright &

Sutton☐Grier 2012). Common garden experiments often find substantial genetic covariation

between the traits we studied possibly due to pleiotropy or selection maintaining correlated

variation (Des Marais et al., 2012; Kenney et al., 2014; McKay et al., 2003). By contrast, the

massively complex environmental variation organisms experience in the wild may combine with

genotype-by-environment interactions to generate high dimensional trait variation among

individuals in nature.

Nevertheless, we found modest evidence of a life history-physiology axis: plants collected later

in the year had low leaf C:N, indicative of a fast-growing resource acquisitive strategy with low

investment in C for structure and high investment in N for photosynthesis. The C:N/collection

date axis is probably not due to later collections being at later developmental stages, since we

would expect plants collected later in development to have allocated nitrogen away from leaves,

lowering C:N. Instead, a strategy of higher leaf C:N may be adaptive for rapid-cycling plants

germinating and flowering within a season (spring/summer annuals), which we expect to be

collected later in the year due to later germination, contrasted with slower-growing genotypes

known to require vernalization for early spring flowering over a winter annual habit. Indeed, Des

Marais et al. (2012) found that vernalization-requiring (winter annual) *Arabidopsis* genotypes

had lower leaf N than genotypes not requiring vernalization for flowering, the latter of which

could also behave as spring or summer annuals.

The negative correlation we observed between $\Delta^{13}C$ and $\delta^{15}N$ has been reported by other authors

and suggested to be a result of independent responses to multiple correlated environmental

variables rather than a biological constraint. Environmental variables with opposing effects on

$\Delta^{13}C$ and $\delta^{15}N$ include soil, temperature, and rainfall patterns (Hartman & Danin, 2010; Liu et

al., 2007; Peri et al., 2012), due to depletion of soil N and changes in stomatal opening, and

atmospheric carbon (Bloom, Burger, Asensio, & Cousins, 2010), which increases carbon uptake

while suppressing nitrate assimilation.

Arabidopsis *life history and physiology vary across spatial environmental gradients, suggesting*

*adaptive responses to long-term environmental conditions (question 2)*

Geographic clines in traits in nature may be due to adaptive responses to environment. However,

the spatial differences in traits or trait changes through time we observed are difficult to ascribe

to genetic or plastic causes because of unknown genotype-environment interactions in the field

and the confounding of environmental gradients and population genetic structure. The 1001

Genomes Project identified genetic clusters of *Arabidopsis* that were somewhat geographically

structured but noted that these clusters overlapped and were distributed across a wide range of

environments (Alonso-Blanco et al., 2016) (see Figure S5 for a map of the clusters). The patterns

of significant phenotype-environment relationships we observed spanned multiple genetic

clusters, making it unclear how much of a role these broad genetic groupings play in determining

environmental response. The phenotype-environment relationships we observed followed our

expectations for how phenology could affect fitness through earlier flowering times in response

to warmth and both earlier and later flowering in response to drier environments. Physiological

traits were less well aligned with our predictions for adaptive response; we did not find low $\Delta^{13}C$

and high C:N associated with environments or years of water stress.

*Physiology, lack of correspondence to the Leaf Economic Spectrum*

The Leaf Economic Spectrum and fast/slow life history predictions were not well supported by

our results for how C:N, $\Delta^{13}$C, and $\delta^{15}$N respond to climate, since we saw both positive and

negative trends with temperature and aridity depending on geographical region. This may be due

to the intraspecific nature of our study, as opposed to the interspecific data often used to support

the LES (Albert et al., 2010; Elmore, Craine, Nelson, & Guinn, 2017). In addition, our study may

have overlooked the effects of edaphic conditions on C:N and $\delta^{15}$N. C:N over most of the native

range was insignificantly related to spatial and temporal gradients of temperature and aridity

index (July precipitation/PET) but increased with year as seen in grassland communities due to

rising $C_a$ (Gill et al., 2002). Likewise, $\delta^{15}$N over most of the range neither decreased with aridity

index nor responded to temperature as expected, but did decrease with year as previously

reported (McLauchlan, Ferguson, Wilson, Ocheltree, & Craine, 2010), possibly due to $CO_2$

enrichment.

Similarly, we did not see strong relationships between aridity index and $\Delta^{13}$C. $\Delta^{13}$C was

expected to be related to rainfall and temperature due to $\Delta^{13}$C being a proxy for stomatal gas

exchange (Diefendorf et al., 2010; Farquhar et al., 1989). There are at least three potential

explanations for weak $\Delta^{13}$C relationships with climate. First, we observed both positive and

negative trends for aridity and date of collection, consistent with the hypothesis that *Arabidopsis*

exhibits both drought escaping and drought avoiding genotypes. The phenological response to

moisture of rapid flowering (drought escape strategy) could confine growth to periods of high

moisture, obviating any stomatal closure in response to soil drying (and hence no effect on

$\Delta^{13}$C). Stated simply, phenology and physiology cannot be treated as completely independent

traits. Second, variation in plant traits we did not directly consider may affect $\Delta^{13}$C. Gas

exchange and carbon assimilation depend in part on leaf architecture and physiology traits like

venation, root allocation, and mesophyll conductance (Brodribb, Feild, & Jordan, 2007; Easlon

et al., 2014; Schulze, Turner, Nicolle, & Schumacher, 2006), which could limit responses in

$\Delta^{13}$C. For example, given the role of roots in sensing drought and triggering stomatal response

(Christmann, Hoffmann, Teplova, Grill, & Muller, 2004), greater investment in roots could allow

plants in relatively drier conditions to maintain open stomata, preventing decreases in $C_i$ and

leading to no observed climate effect on $\Delta^{13}$C. Third, elevated atmospheric partial $CO_2$ could

mitigate climate effects on $\Delta^{13}$C by increasing the efficiency of stomatal gas exchange (Drake,

Hanson, Lowrey, & Sharp, 2017). Local investigations of the patterns we found could

complement our results by characterizing the underlying ecophysiological and life history

mechanisms driving intraspecific variation.

*Phenology, high variation across space*

We found strong spatial gradients in two measures of phenology, suggesting that adaptive

responses to climate drive long-term trait differences among regions. Locations that were

warmer than average in either April or January corresponded to significantly earlier collection

dates, consistent with temperature's positive effect on growth rate (Wilczek et al., 2009). In

addition, our models provided support that some phenological variation did reflect seasonality of

moisture availability. We found that *Arabidopsis* was collected significantly earlier in years with

dry summers in central Europe and at significantly lower PTU in regions of wet summers around

the Mediterranean, suggesting drought escape or avoidance strategies, respectively, could be

important in those regions. Alternatively, later collections in wetter years could be the result of

multiple successful generations due to the extra rainfall. This ambiguity illustrates an important

caveat in using collection date and PTU to study *Arabidopsis* in the wild. Because we cannot

differentiate changes in total life span from changes in germination start date, shifts in phenology

(the timing of an organism's life to seasonal conditions) does not necessarily indicate shifts in

life history. Nevertheless, the observed regional variation in the response of collection date and

PTU to climate variables provides valuable insight into how different populations experience

their environment and suggest areas for more direct study.

*Changes in* Arabidopsis *life history and physiology over the last two centuries track climate,*

*suggesting adaptive responses (question 3)*

Increasing global temperatures were expected to increase relative growth rate and hasten

germination, decreasing flowering time as measured by collection date. In addition, atmospheric

$CO_2$ enrichment was expected to increase $\Delta^{13}C$ (Drake et al., 2017) and C:N and decrease $\delta^{15}N$

(Bloom et al., 2010). With increasing environmental nitrogen due to human activity (Galloway et

al., 2004), we expected $\delta^{15}N$ to decrease.

Our findings were largely consistent with these hypotheses in the year models for leaf

physiology. $\Delta^{13}C$ did not significantly change through time across the native range, which could

be due to life history shifts as mentioned above or differential response to aridity gradients across

the locations sampled masking the effect of elevated $CO_2$ (Drake et al., 2017). Geographic

variation in the strength of the relationships for other traits could be due to underlying genetic

variation or interaction with environmental factors we did not account for. Nevertheless, C:N

increased and $\delta^{15}N$ decreased as expected across large portions of the native range.

For collection date and PTU, however, our models returned the surprising result of later

collection rather than earlier, despite earlier collections in warmer years. The fact that the

relationship between warmth and collection date was spatially variable, and insignificant in some

regions, may indicate areas of contrasting phenological response, perhaps due to lost

vernalization signal or variable effects on germination (Burghardt, Edwards, & Donohue, 2016).

Variation across space in phenological response to climate change has been shown before and

may be due to genetic differences among populations or due to interactions with other

environmental variables (Park et al., 2018). *Arabidopsis* is known to complete a generation

within a single season, climate permitting, and warmer climates may even allow for fall

flowering (Fournier-Level et al., 2013; Wilczek et al., 2009). If warmer temperatures enable a

greater number of spring or summer germinants to flower before winter in regions such as

Central Europe, we would expect to see later collection dates in more recent years (Burghardt et

al., 2015). Regions that did not show later collection dates through time might be limited in

generational cycles due to summer drought or very short growing seasons. For instance, early

flowering in the spring has been implicated as an important strategy for *Arabidopsis* in the

Iberian Peninsula to escape seasonal heat and water limitation that curtail growth in later months

(Wolfe & Tonsor, 2014).

Our findings of later collection dates through the study period (1798-2010) may surprise some

readers due to previously observed acceleration of temperate spring phenology (Parmesan &

Yohe, 2003). However, we modeled changes in mean phenological response to environment,

which can be weakly related to either tail of phenology trait distributions (CaraDonna et al.,

2014). Individuals on the extremes, such as first-flowering individuals, are often the primary

focus of studies showing accelerated spring phenology in recent years. Why might *Arabidopsis*

flower later even as global temperatures rise? First, anthropogenic land use change may drive

phenology by favoring spring germinants, *e.g.* if disturbances favor faster life cycles. Second,

warming climate or increasing atmospheric $pCO_2$ may favor alternate life histories by increasing

relative growth rate, thus allowing spring or fall germinants to complete their life cycle before

conditions degrade at the end of a growing season. Later collections in more recent years might

represent an increasing proportion of fast-growing spring or summer annuals as opposed to

winter annuals. Whatever the cause of *Arabidopsis* flowering later, these phenological changes

may have important ecological effects, such as altered biotic interactions.

We found evidence that trait correlations may be changing through time. For example, in some

regions both C:N and date of collection have significantly increased over the past 200 years

(around the Eastern Mediterranean), while in other regions date of collection decreased while

C:N increased over time (Morocco and the Iberian Peninsula) (Figure 1). If the negative

relationship between leaf C:N and flowering time that we observed is truly an axis of adaptive

tradeoff between fast and slow life histories, this tradeoff may be changing at different rates

among regions with time. Changing environments might reshape biological constraints on

adaptive plant responses (Sgrò & Hoffmann, 2004).

*Our approach, technical limitations in herbaria data to surmount in future studies*

Understanding how environmental variation drives the intraspecific diversity in broadly

distributed species has been challenging due to logistics of large spatiotemporal scales. However,

advances in digitization of museum specimens and the generation of global gridded

spatiotemporal environmental data are opening a new window into large scale patterns of

biodiversity. One challenge of herbarium specimens is that they typically present a single

observation of a mature, reproductive individuals. Thus, these specimens contain limited

information on phenology and physiology at earlier life stages (*e.g.* seedling plants), which can

have subsequently strong impacts on later observed stages. Use of developmental models

(Burghardt et al., 2015) might allow one to backcast potential developmental trajectories using

herbarium specimens and climate data, to make predictions about phenology of germination and

transition to flowering. In addition, herbaria collections are often biased by factors such as

geography, species, and climate (Daru et al., 2018; Loiselle et al., 2008). Hierarchical sampling

through repeated collections in the same region could improve the confidence of our model in

representing phenotypic change through time.

Generalized additive models are a flexible approach to model phenotype responses to

environment that might differ spatially among populations (MacGillivray et al., 2010). These

models allow the data to inform on spatial variation in the trends studied, unlike approaches that

bin individuals into discrete and arbitrarily bounded regions. Herbarium records represent

imperfect and biased samples of natural populations (Daru et al., 2018), and future efforts may

benefit from additional information that might allow us to account for these biases. Here, we

sampled a very large number of specimens across continents and centuries, perhaps reducing the

effect of biases associated with specific collectors. Nevertheless, as museum informatics advance

it may become possible to explicitly model potential sources of bias, for example those arising

from collecting behavior of specific researchers.

*Conclusion*

Widely distributed species often harbor extensive intraspecific trait diversity. Natural history

collections offer a window into this diversity and in particular allow investigation of long-term

responses to anthropogenic change across species ranges. Here we show that spatiotemporal

climate gradients explain much of this diversity but nevertheless much of the phenotypic

diversity in nature for the model plant remains to be explained.

**Acknowledgements**

Hundreds of professional and amateur botanists collected the specimens studied here. The staff at

herbaria at Oslo Natural History Museum, Kew Gardens, Real Jardin Botanico, Komarov

Botanical Garden, the New York Botanical Garden, and the British Museum of Natural History

gave permission for tissue sampling. Michelle Brown provided essential assistance in collecting

herbarium tissue. Jason Bonnette helped coordinate sample preparation. Major assistance in

digitizing of specimens was provided by Patrick Herné. Eugene Shakirov aided in translating

Russian specimen labels. Data from the MNHN in Paris were obtained thanks to the

participatory science program Les Herbonautes" (MNHN/Tela Botanica) which is part of

Infrastructure Nationale e-RECOLNAT: ANR-11-INBS-0004. Additional volunteers from the

Atlas of Living Australia helped with digitization and georeferencing. Funding was provided by

an Earth Institute fellowship to JRL.

The authors report no commercial or other relationships relevant to the content of this article that

would represent a conflict of interest.

**Tables**

Table 1: Hypothesized responses of phenotypes to increases in temperature, rainfall, or year, or

how traits would change along a faster life history strategy.

Year trends are predicted due to elevated $CO_2$, nitrogen deposition, or elevated temperatures.

Citations for hypotheses: (Amundson et al., 2003[1]; BassiriRad et al., 2003[2]; Burghardt et al.,

2015[3]; Craine et al., 2009[4]; Diefendorf et al., 2010[5]; Drake, Hanson, Lowrey, & Sharp, 2017[6];

Gill et al., 2002[7]; McLauchlan, Ferguson, Wilson, Ocheltree, & Craine, 2010[8]; Menzel et al.,

2006[9]; Ordoñez et al., 2009[10]; Peñuelas et al., 2004[11]; Reich, Hungate, & Luo, 2006[12]; Seibt,

Rajabi, Griffiths, & Berry, 2008[13]; Sparks & Carey, 2006[14]; Stock & Evans, 2006[15]; I. J. Wright

et al., 2004[16])

|  | Temperature | Rainfall | Year | Fast Life History |
|---|---|---|---|---|
| $\Delta^{13}C$ | +[13] | +[5] | +[6] | + |
| $\delta^{15}N$ | +[1,4] | -[1,4] | -[2,8,15] | No change |
| C:N | -[10] in cold regions +[16] in warm regions | -[16] | +[7,8,12] | - |
| Photothermal Units | + or no change | + | No change | - |
| Collection Date | -[3,9] | +[11] | -[9,14] | - |

739 *Comparison of results to hypotheses*

740 Table 2: Expected phenotype responses to increases in temperature, rainfall, or year and

741 observed model output.

742 For some phenotypes a single trend over time was observed; however, most phenotypes showed

743 variation in responses to temperature and rainfall across the range.

| | Temperature | | Rainfall | | Year | |
|---|---|---|---|---|---|---|
| | Expected | Observed | Expected | Observed | Expected | Observed |
| $\Delta^{13}C$ | - | -, mainly insignificant | + | -, mainly insignificant | + | insignificant |
| $\delta^{15}N$ | + | +, -, mainly insignificant | - | + | - | - |
| C:N | -<br>+ | +, -, mainly insignificant | - | insignificant | + | + |
| Photothermal Units | +<br>or no change | +, - | + | +, - | No change | + |
| Collection Date | - | -, + | + | +, - | - | +, - |

## References

Adams, W. W., Stewart, J. J., Cohu, C. M., Muller, O., & Demmig-Adams, B. (2016). Habitat

Temperature and Precipitation of *Arabidopsis thaliana* Ecotypes Determine the Response of

Foliar Vasculature, Photosynthesis, and Transpiration to Growth Temperature. *Frontiers in*

*Plant Science*, *7*. https://doi.org/10.3389/fpls.2016.01026

Albert, C. H., Thuiller, W., Yoccoz, N. G., Soudant, A., Boucher, F., Saccone, P., & Lavorel, S.

(2010). Intraspecific functional variability: extent, structure and sources of variation.

*Journal of Ecology*, *98*(3), 604–613. https://doi.org/10.1111/j.1365-2745.2010.01651.x

Alonso-Blanco, C., Andrade, J., Becker, C., Bemm, F., Bergelson, J., Borgwardt, K. M. M., …

Zhou, X. (2016). 1,135 genomes reveal the global pattern of polymorphism in *Arabidopsis*

*thaliana*. *Cell*, *166*(2), 481–491. https://doi.org/10.1016/j.cell.2016.05.063

Amundson, R., Austin, A. T., Schuur, E. A. G., Yoo, K., Matzek, V., Kendall, C., … Baisden,

W. T. (2003). Global patterns of the isotopic composition of soil and plant nitrogen. *Global*

*Biogeochemical Cycles*, *17*(1).

Anderegg, L. D. L., Berner, L. T., Badgley, G., Sethi, M. L., Law, B. E., & HilleRisLambers, J.

(2018). Within-species patterns challenge our understanding of the leaf economics

spectrum. *Ecology Letters*. https://doi.org/10.1111/ele.12945

Atwell, S., Huang, Y. S., Vilhjálmsson, B. J., Willems, G., Horton, M., Li, Y., … Nordborg, M.

(2010). Genome-wide association study of 107 phenotypes in *Arabidopsis thaliana* inbred

lines. *Nature*. https://doi.org/10.1038/nature08800

BassiriRad, H., Constable, J. V. H., Lussenhop, J., Kimball, B. A., Norbys, R. J., Oechel, W. C.,

… Silim, S. (2003). Widespread foliage $\delta^{15}N$ depletion under elevated $CO_2$: Inferences for

the nitrogen cycle. *Global Change Biology*. https://doi.org/10.1046/j.1365-

2486.2003.00679.x

Blonder, B., Violle, C., Bentley, L. P., & Enquist, B. J. (2011). Venation networks and the origin

of the leaf economics spectrum. *Ecology Letters*, *14*(2), 91–100.

https://doi.org/10.1111/j.1461-0248.2010.01554.x

Bloom, A. J., Burger, M., Asensio, J. S. R., & Cousins, A. B. (2010). Carbon dioxide enrichment

inhibits nitrate assimilation in wheat and *arabidopsis*. *Science*.

https://doi.org/10.1126/science.1186440

Both, C., Artemyev, A. V., Blaauw, B., Cowie, R. J., Dekhuijzen, A. J., Eeva, T., … Visser, M.

E. (2004). Large-scale geographical variation confirms that climate change causes birds to

lay earlier. *Proceedings of the Royal Society B: Biological Sciences*, *271*(1549), 1657–1662.

https://doi.org/10.1098/rspb.2004.2770

Brachi, B., Faure, N., Horton, M., Flahauw, E., Vazquez, A., Nordborg, M., … Roux, F. (2010).

Linkage and association mapping of *Arabidopsis thaliana* flowering time in nature. *PLoS*

*Genetics*, *6*(5), 40. https://doi.org/10.1371/journal.pgen.1000940

Bradshaw, A. D. (1965). Evolutionary significance of phenotypic plasticity in plants. *Advances*

*in Genetics*. https://doi.org/10.1016/S0065-2660(08)60048-6

Brodribb, T. J., Feild, T. S., & Jordan, G. J. (2007). Leaf maximum photosynthetic rate and

venation are linked by hydraulics. *Plant Physiology*, *144*(4), 1890–1898.

https://doi.org/10.1104/pp.107.101352

Burghardt, L. T., Edwards, B. R., & Donohue, K. (2016). Multiple paths to similar germination

behavior in *Arabidopsis thaliana*. *New Phytologist*, *209*(3), 1301–1312.

https://doi.org/10.1111/nph.13685

Burghardt, L. T., Metcalf, C. J. E., Wilczek, A. M., Schmitt, J., & Donohue, K. (2015). Modeling

the influence of genetic and environmental variation on the expression of plant life cycles

across landscapes. *The American Naturalist*, *185*(2), 212–227.

CaraDonna, P. J., Iler, A. M., & Inouye, D. W. (2014). Shifts in flowering phenology reshape a

subalpine plant community. *Proceedings of the National Academy of Sciences*, *111*(13),

4916–4921. https://doi.org/10.1073/pnas.1323073111

Christmann, A., Hoffmann, T., Teplova, I., Grill, E., & Muller, A. (2004). Generation of active

pools of abscisic acid revealed by *in vivo* imaging of water-stressed *Arabidopsis*. *Plant

Physiology*, *137*(1), 209–219. https://doi.org/10.1104/pp.104.053082

Craine, J. M., Elmore, A. J., Aidar, M. P. M., Bustamante, M., Dawson, T. E., Hobbie, E. A., …

Wright, I. J. (2009). Global patterns of foliar nitrogen isotopes and their relationships with

climate, mycorrhizal fungi, foliar nutrient concentrations, and nitrogen availability. *New

Phytologist*, *183*(4), 980–992. https://doi.org/10.1111/j.1469-8137.2009.02917.x

Daru, B. H., Park, D. S., Primack, R. B., Willis, C. G., Barrington, D. S., Whitfeld, T. J. S., …

Davis, C. C. (2018). Widespread sampling biases in herbaria revealed from large-scale

digitization. *New Phytologist*. https://doi.org/10.1111/nph.14855

Davis, C. C., Willis, C. G., Connolly, B., Kelly, C., & Ellison, A. M. (2015). Herbarium records

are reliable sources of phenological change driven by climate and provide novel insights

into species' phenological cueing mechanisms. *American Journal of Botany*, *102*(10),

1599–1609. https://doi.org/10.3732/ajb.1500237

Des Marais, D. L., McKay, J. K., Richards, J. H., Sen, S., Wayne, T., & Juenger, T. E. (2012).

Physiological genomics of response to soil drying in diverse *Arabidopsis* accessions. *The*

*Plant Cell*. https://doi.org/10.1105/tpc.112.096180

Diefendorf, A. F., Mueller, K. E., Wing, S. L., Koch, P. L., & Freeman, K. H. (2010). Global

patterns in leaf $^{13}$C discrimination and implications for studies of past and future climate.

*Proceedings of the National Academy of Sciences*, *107*(13), 5738–5743.

https://doi.org/10.1073/pnas.0910513107

Drake, B. L., Hanson, D. T., Lowrey, T. K., & Sharp, Z. D. (2017). The carbon fertilization

effect over a century of anthropogenic $CO_2$ emissions: higher intracellular $CO_2$ and more

drought resistance among invasive and native grass species contrasts with increased water

use efficiency for woody plant. *Global Change Biology*, *23*(2), 782–792.

https://doi.org/10.1111/gcb.13449

Easlon, H. M., Nemali, K. S., Richards, J. H., Hanson, D. T., Juenger, T. E., & McKay, J. K.

(2014). The physiological basis for genetic variation in water use efficiency and carbon

isotope composition in *Arabidopsis thaliana*. *Photosynthesis Research*, *119*(1–2), 119–129.

Elmore, A. J., Craine, J. M., Nelson, D. M., & Guinn, S. M. (2017). Continental scale variability

of foliar nitrogen and carbon isotopes in *Populus balsamifera* and their relationships with

climate. *Scientific Reports*, *7*(1). https://doi.org/10.1038/s41598-017-08156-x

Evans, J. R. (1989). Photosynthesis and nitrogen relationships in leaves of C3 plants. *Oecologia*,

*78*(1), 9–19.

Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H. A., & Weigel, D. (2018).

Genomic basis and evolutionary potential for extreme drought adaptation in *Arabidopsis*

*thaliana*. *Nature Ecology and Evolution*. https://doi.org/10.1038/s41559-017-0423-0

Farquhar, G., Ehleringer, J., & Hubick, K. (1989). Carbon isotope discrimination and

photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology*, *40*(1),

503–537. https://doi.org/10.1146/annurev.pp.40.060189.002443

Farquhar, G., O'Leary, M., & Berry, J. (1982). On the relationship between carbon isotope

discrimination and the intercellular carbon dioxide concentration in leaves. *Australian*

*Journal of Plant Physiology*, *9*(2), 121–137. https://doi.org/10.1071/PP9820121

Fournier-Level, A., Wilczek, A. M., Cooper, M. D., Roe, J. L., Anderson, J., Eaton, D., …

Schmitt, J. (2013). Paths to selection on life history loci in different natural environments

across the native range of *Arabidopsis thaliana*. *Molecular Ecology*, *22*(13), 3552–3566.

https://doi.org/10.1111/mec.12285

Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P.,

… Holland, E. A. (2004). Nitrogen cycles: past, present, and future. *Biogeochemistry*,

*70*(2), 153–226.

Gill, R. A., Polley, H. W., Johnson, H. B., Anderson, L. J., Maherali, H., & Jackson, R. B.

(2002). Nonlinear grassland responses to past and future atmospheric $CO_2$. *Nature*.

https://doi.org/10.1038/417279a

Grime, J. P. (1977). Evidence for the existence of three primary strategies in plants and its

relevance to ecological and evolutionary theory. *The American Naturalist*.

https://doi.org/10.1086/283244

Harris, I., Jones, P. D., Osborn, T. J., & Lister, D. H. (2014). Updated high☐resolution grids of

monthly climatic observations–the CRU TS3. 10 Dataset. *International Journal of*

*Climatology*, *34*(3), 623–642.

Hartman, G., & Danin, A. (2010). Isotopic values of plants in relation to water availability in the

Eastern Mediterranean region. *Oecologia*. https://doi.org/10.1007/s00442-009-1514-7

Henn, J. J., Buzzard, V., Enquist, B. J., Halbritter, A. H., Klanderud, K., Maitner, B. S., …

Vandvik, V. (2018). Intraspecific trait variation and phenotypic plasticity mediate alpine

plant species response to climate change. *Frontiers in Plant Science*.

https://doi.org/10.3389/fpls.2018.01548

Hijmans, R. J. (2017). geosphere: spherical trigonometry. Retrieved from https://cran.r-

project.org/package=geosphere

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high

resolution interpolated climate surfaces for global land areas. *International Journal of*

*Climatology*, *25*(15), 1965–1978.

Himelblau, E., & Amasino, R. M. (2001). Nutrients mobilized from leaves of *Arabidopsis*

*thaliana* during leaf senescence. *Journal of Plant Physiology*. https://doi.org/10.1078/0176-

1617-00608

Hoffmann, M. H. (2002). Biogeography of *Arabidopsis thaliana* (L.) Heynh. (Brassicaceae).

*Journal of Biogeography*, *29*(1), 125–134. https://doi.org/10.1046/j.1365-

2699.2002.00647.x

Horton, M. W., Hancock, A. M., Huang, Y. S., Toomajian, C., Atwell, S., Auton, A., …

Bergelson, J. (2012). Genome-wide patterns of genetic variation in worldwide *Arabidopsis*

*thaliana* accessions from the RegMap panel. *Nature Genetics*.

https://doi.org/10.1038/ng.1042

Hu, Y.-K., Pan, X., Liu, G.-F., Li, W.-B., Dai, W.-H., Tang, S.-L., … Dong, M. (2015). Novel

evidence for within-species leaf economics spectrum at multiple spatial scales. *Frontiers in*

*Plant Science*. https://doi.org/10.3389/fpls.2015.00901

Joshi, J., Schmid, B., Caldeira, M. C., Dimitrakopoulos, P. G., Good, J., Harris, R., … Lawton, J.

H. (2001). Local adaptation enhances performance of common plant species. *Ecology*

*Letters*. https://doi.org/10.1046/j.1461-0248.2001.00262.x

Jung, V., Violle, C., Mondy, C., Hoffmann, L., & Muller, S. (2010). Intraspecific variability and

trait-based community assembly. *Journal of Ecology*, *98*(5), 1134–1140.

https://doi.org/10.1111/j.1365-2745.2010.01687.x

Kenney, A. M., Mckay, J. K., Richards, J. H., & Juenger, T. E. (2014). Direct and indirect

selection on flowering time, water-use efficiency (WUE, $\delta^{13}C$), and WUE plasticity to

drought in *Arabidopsis thaliana*. *Ecology and Evolution*, *4*(23), 4505–4521.

https://doi.org/10.1002/ece3.1270

Koenker, A. R., & Koenker, M. R. (2011). Package ' quantreg .' *Annals of Statistics*.

Lang, P. L. M., Willems, F. M., Scheepens, J. F., Burbano, H. A., & Bossdorf, O. (2018). Using

herbaria to study global environmental change. *New Phytologist*.

https://doi.org/10.1111/nph.15401

Lasky, J. R., Des Marais, D. L., McKay, J. K., Richards, J. H., Juenger, T. E., & Keitt, T. H.

(2012). Characterizing genomic variation of *Arabidopsis thaliana*: the roles of geography

and climate. *Molecular Ecology*, *21*(22), 5512–5529. https://doi.org/10.1111/j.1365-

294X.2012.05709.x

Lasky, J. R., Forester, B. R., & Reimherr, M. (2018). Coherent synthesis of genomic associations

with phenotypes and home environments. *Molecular Ecology Resources*, *18*(1), 91–106.

Leimu, R., & Fischer, M. (2008). A meta-analysis of local adaptation in plants. *PLoS ONE*.

https://doi.org/10.1371/journal.pone.0004010

Linhart, Y. B., & Grant, M. C. (2002). Evolutionary significance of local genetic differentiation

in plants. *Annual Review of Ecology and Systematics*.

https://doi.org/10.1146/annurev.ecolsys.27.1.237

Liu, X., Zhao, L., Gasaw, M., Gao, D., Qin, D., & Ren, J. (2007). Foliar $\delta^{13}$C and $\delta^{15}$N values of

C3 plants in the Ethiopia Rift Valley and their environmental controls. *Chinese Science*

*Bulletin*. https://doi.org/10.1007/s11434-007-0165-5

Loiselle, B. A., Jørgensen, P. M., Consiglio, T., Jiménez, I., Blake, J. G., Lohmann, L. G., &

Montiel, O. M. (2008). Predicting species distributions from herbarium collections: Does

climate bias in collection sampling influence model outcomes? *Journal of Biogeography*.

https://doi.org/10.1111/j.1365-2699.2007.01779.x

Lovell, J. T., Juenger, T. E., Michaels, S. D., Lasky, J. R., Platt, A., Richards, J. H., … Mckay, J.

K. (2013). Pleiotropy of FRIGIDA enhances the potential for multivariate adaptation.

*Proceedings of the Royal Society B: Biological Sciences*.

https://doi.org/10.1098/rspb.2013.1043

Ludlow, M. M. (1989). Strategies of response to water stress. In K. H. Kreeb, H. Richter, & T.

M. Hinckley (Eds.), *Structural and functional responses to environmental stresses: water*

*shortage* (pp. 269–281). Berlin: SPB Academic Publishers.

MacGillivray, F., Hudson, I. L., & Lowe, A. J. (2010). Herbarium collections and photographic

images: Alternative data sources for phenological research. In *Phenological Research:*

*Methods for Environmental and Climate Change Analysis* (pp. 425–461). Dordrecht:

Springer Netherlands. https://doi.org/10.1007/978-90-481-3335-2_19

Matesanz, S., Gianoli, E., & Valladares, F. (2010). Global change and the evolution of

phenotypic plasticity in plants. *Annals of the New York Academy of Sciences*.

https://doi.org/10.1111/j.1749-6632.2010.05704.x

McCarroll, D., & Loader, N. J. (2004). Stable isotopes in tree rings. *Quaternary Science*

*Reviews*, *23*(7–8), 771–801.

McKay, J. K., Richards, J. H., & Mitchell-Olds, T. (2003). Genetics of drought adaptation in

*Arabidopsis thaliana*: I. Pleiotropy contributes to genetic correlations among ecological

traits. *Molecular Ecology*. https://doi.org/10.1046/j.1365-294X.2003.01833.x

McLauchlan, K. K., Ferguson, C. J., Wilson, I. E., Ocheltree, T. W., & Craine, J. M. (2010).

Thirteen decades of foliar isotopes indicate declining nitrogen availability in central North

American grasslands. *New Phytologist*, *187*(4), 1135–1145. https://doi.org/10.1111/j.1469-

8137.2010.03322.x

Menzel, A., Sparks, T. H., Estrella, N., Koch, E., Aaasa, A., Ahas, R., … Zust, A. (2006).

European phenological response to climate change matches the warming pattern. *Global*

*Change Biology*. https://doi.org/10.1111/j.1365-2486.2006.01193.x

Miller-Rushing, A. J., Primack, R. B., Primack, D., & Mukunda, S. (2006). Photographs and

herbarium specimens as tools to document phenological changes in response to global

warming. *American Journal of Botany*, *93*(11), 1667–1674.

https://doi.org/10.3732/ajb.93.11.1667

New, M., Hulme, M., & Jones, P. (2000). Representing twentieth-century space-time climate

variability. Part II: Development of 1901-96 monthly grids of terrestrial surface climate.

*Journal of Climate*, *13*(13), 2217–2238. https://doi.org/10.1175/1520-

0442(2000)013<2217:RTCSTC>2.0.CO;2

Ordoñez, J. C., Van Bodegom, P. M., Witte, J. P. M., Wright, I. J., Reich, P. B., & Aerts, R.

(2009). A global study of relationships between leaf traits, climate and soil measures of

nutrient fertility. *Global Ecology and Biogeography*, *18*(2), 137–149.

https://doi.org/10.1111/j.1466-8238.2008.00441.x

Ostrowski, M. F., David, J., Santoni, S., McKhann, H., Reboud, X., Le Corre, V., … Bataillon,

T. (2006). Evidence for a large-scale population structure among accessions of *Arabidopsis*

*thaliana*: Possible causes and consequences for the distribution of linkage disequilibrium.

*Molecular Ecology*. https://doi.org/10.1111/j.1365-294X.2006.02865.x

Pardo, L. H., McNulty, S. G., Boggs, J. L., & Duke, S. (2007). Regional patterns in foliar [15]N

across a gradient of nitrogen deposition in the northeastern US. *Environmental Pollution*,

*149*(3), 293–302. https://doi.org/10.1016/j.envpol.2007.05.030

Park, D. S., Breckheimer, I., Williams, A. C., Law, E., Ellison, A. M., & Davis, C. C. (2018).

Herbarium specimens reveal substantial and unexpected variation in phenological

sensitivity across the eastern United States. *Philosophical Transactions of the Royal Society*

*of London. Series B, Biological Sciences*. https://doi.org/10.1098/rstb.2017.0394

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts

across natural systems. *Nature*, *421*(6918), 37–42. https://doi.org/10.1038/nature01286

Peñuelas, J., Filella, I., Zhang, X., Llorens, L., Ogaya, R., Lloret, F., … Terradas, J. (2004).

Complex spatiotemporal phenological shifts as a response to rainfall changes. *New*

*Phytologist*. https://doi.org/10.1111/j.1469-8137.2004.01003.x

Peri, P. L., Ladd, B., Pepper, D. A., Bonser, S. P., Laffan, S. W., & Amelung, W. (2012). Carbon

($\delta^{13}$C) and nitrogen ($\delta^{15}$N) stable isotope composition in plant and soil in Southern

Patagonia's native forests. *Global Change Biology*. https://doi.org/10.1111/j.1365-

2486.2011.02494.x

Platt, A., Horton, M., Huang, Y. S., Li, Y., Anastasio, A. E., Mulyati, N. W., … Borevitz, J. O.

(2010). The scale of population structure in *Arabidopsis thaliana*. *PLoS Genetics*.

https://doi.org/10.1371/journal.pgen.1000843

R Core Team. (2018). R: a language and environment for statistical computing. Vienna, Austria.

https://doi.org/10.1007/978-3-540-74686-7

Reich, P. B. (2014). The world-wide "fast-slow" plant economics spectrum: a traits manifesto.

*Journal of Ecology*, *102*(2), 275–301. https://doi.org/10.1111/1365-2745.12211

Reich, P. B., Hungate, B. A., & Luo, Y. (2006). Carbon-nitrogen interactions in terrestrial

ecosystems in response to rising atmospheric carbon dioxide. *Annual Review of Ecology,*

*Evolution, and Systematics*, *37*(1), 611–636.

https://doi.org/10.1146/annurev.ecolsys.37.091305.110039

Reich, P. B., Wright, I. J., Cavender‑Bares, J., Craine, J. M., Oleksyn, J., Westoby, M., &

Walters, M. B. (2003). The evolution of plant functional variation: traits, spectra, and

strategies. *International Journal of Plant Sciences*, *164*(S3), S143–S164.

https://doi.org/10.1086/374368

Sack, L., Scoffoni, C., McKown, A. D., Frole, K., Rawls, M., Havran, J. C., … Tran, T. (2012).

Developmentally based scaling of leaf venation architecture explains global ecological

patterns. *Nature Communications*, *3*, 837. https://doi.org/10.1038/ncomms1835

Sartori, K. F. R., Vasseur, F., Violle, C., Baron, E., Gerard, M., Rowe, N., … Masclef, D. (2018).

Leaf economics guides slow-fast adaptation across the geographic range of *A. thaliana*.

*BioRxiv*, 487066.

Schulze, E. D., Turner, N. C., Nicolle, D., & Schumacher, J. (2006). Leaf and wood carbon

isotope ratios, specific leaf areas and wood growth of *Eucalyptus* species across a rainfall

gradient in Australia. *Tree Physiology*, *26*(4), 479–492.

https://doi.org/10.1093/treephys/26.4.479

Seibt, U., Rajabi, A., Griffiths, H., & Berry, J. A. (2008). Carbon isotopes and water use

efficiency: Sense and sensitivity. *Oecologia*. https://doi.org/10.1007/s00442-007-0932-7

Sgrò, C. M., & Hoffmann, A. A. (2004). Genetic correlations, tradeoffs and environmental

variation. *Heredity*. https://doi.org/10.1038/sj.hdy.6800532

Sparks, T. H., & Carey, P. D. (2006). The responses of species to climate over two centuries: an

analysis of the Marsham phenological record, 1736-1947. *The Journal of Ecology*.

https://doi.org/10.2307/2261570

Stinchcombe, J. R., Weinig, C., Ungerer, M., Olsen, K. M., Mays, C., Halldorsdottir, S. S., …

Schmitt, J. (2004). A latitudinal cline in flowering time in *Arabidopsis thaliana* modulated

by the flowering time gene FRIGIDA. *Proceedings of the National Academy of Sciences*.

https://doi.org/10.1073/pnas.0306401101

Stock, W. D., & Evans, J. R. (2006). Effects of water availability, nitrogen supply and

atmospheric $CO_2$ concentrations on plant nitrogen natural abundance values. *Functional

Plant Biology*, *33*(3), 219–227. https://doi.org/10.1071/FP05188

Stocking, C. R., & Ongun, A. (1962). The intracellular distribution of some metallic elements in

leaves. *American Journal of Botany*, *49*(3), 284–289.

Tabas-Madrid, D., Méndez-Vigo, B., Arteaga, N., Marcer, A., Pascual-Montano, A., Weigel, D.,

… Alonso-Blanco, C. (2018). Genome-wide signatures of flowering adaptation to climate

temperature: Regional analyses in a highly diverse native range of *Arabidopsis thaliana*.

*Plant Cell and Environment*, *41*(8), 1806–1820. https://doi.org/10.1111/pce.13189

Turesson, G. (1922). The genotypical response of the plant species to the habitat. *Hereditas*.

https://doi.org/10.1111/j.1601-5223.1922.tb02734.x

United Nations Environment Program. (1997). *World atlas of desertification*. (N. Middleton &

D. Thomas, Eds.) (2nd ed.).

Vasseur, F., Violle, C., Enquist, B. J., Granier, C., & Vile, D. (2012). A common genetic basis to

the origin of the leaf economics spectrum and metabolic scaling allometry. *Ecology Letters*,

*15*(10), 1149–1157. https://doi.org/10.1111/j.1461-0248.2012.01839.x

Westoby, M. (1998). A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil*.

https://doi.org/10.1023/A:1004327224729

Wheeler, D. C., & Waller, L. A. (2009). Comparing spatially varying coefficient models: a case

study examining violent crime rates and their relationships to alcohol outlets and illegal

drug arrests. *Journal of Geographical Systems*, *11*(1), 1–22. https://doi.org/10.1007/s10109-

008-0073-5

Wilczek, A. M., Cooper, M. D., Korves, T. M., & Schmitt, J. (2014). Lagging adaptation to

warming climate in *Arabidopsis thaliana*. *Proceedings of the National Academy of*

*Sciences*, *111*(22), 7906–7913. https://doi.org/10.1073/pnas.1406314111

Wilczek, A. M., Roe, J. L., Knapp, M. C., Cooper, M. D., Lopez-Gallego, C., Martin, L. J., …

Schmitt, J. (2009). Effects of genetic perturbation on seasonal life history plasticity.

*Science*, *323*(5916), 930–934. https://doi.org/10.1126/science.1165826

Willis, C. G., Ellwood, E. R., Primack, R. B., Davis, C. C., Pearson, K. D., Gallinat, A. S., …

Soltis, P. S. (2017). Old plants, new tricks: phenological research using herbarium

specimens. *Trends in Ecology and Evolution*, *32*(7), 531–546.

https://doi.org/10.1016/j.tree.2017.03.015

Wolfe, M. D., & Tonsor, S. J. (2014). Adaptation to spring heat and drought in northeastern

Spanish *Arabidopsis thaliana*. *New Phytologist*. https://doi.org/10.1111/nph.12485

Wood, S. N. (2006). *Generalized additive models: an introduction with R. Texts in Statistical*

*Science* (1st ed.). Chapman and Hall/CRC. https://doi.org/10.1111/j.1541-

0420.2007.00905_3.x

Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood

estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society. Series B: Statistical Methodology*, *73*(1), 3–36. https://doi.org/10.1111/j.1467-9868.2010.00749.x

Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., … Villar, R. (2004). The worldwide leaf economics spectrum. *Nature*, *428*(6985), 821–827. https://doi.org/10.1038/nature02403

Wright, J. P., & Sutton-Grier, A. (2012). Does the leaf economic spectrum hold within local species pools across varying environmental conditions? *Functional Ecology*, *26*(6), 1390–1398. https://doi.org/10.1111/1365-2435.12001

Yee, T. W., & Mackenzie, M. (2002). Vector generalized additive models in plant ecology. *Ecological Modelling*. https://doi.org/10.1016/S0304-3800(02)00192-8

Yee, T. W., & Mitchell, N. D. (2006). Generalized additive models in plant ecology. *Journal of Vegetation Science*. https://doi.org/10.2307/3236170

Zhu, Y., Siegwolf, R. T. W., Durka, W., & Körner, C. (2010). Phylogenetically balanced evidence for structural and carbon isotope responses in plants along elevational gradients. *Oecologia*, *162*(4), 853–863. https://doi.org/10.1007/s00442-009-1515-6