Atypically larger variability of resource allocation accounts for

visual working memory deficits in schizophrenia

Yi-Jie Zhao[1,2,3,4†], Tianye Ma[2†], Xuemei Ran[2], Li Zhang[2], Ru-Yuan
Zhang[5,6*], Yixuan Ku[1,2,7*]

[1]Guangdong Provincial Key Laboratory of Social Cognitive Neuroscience and
Mental Health, Department of Psychology, Sun Yat-Sen University, Guangzhou,
China.
[2]The Shanghai Key Lab of Brain Functional Genomics, Shanghai Changning-
ECNU Mental Health Center, School of Psychology and Cognitive Science, East
China Normal University, Shanghai, China.
[3]Institute of Science and Technology for Brain-Inspired Intelligence, Fudan
University, Shanghai, China.
[4]Key Laboratory of Computational Neuroscience and Brain-Inspired Intelligence,
Fudan University, Ministry of Education, China.
[5]Shanghai Key Laboratory of Psychotic Disorders, Shanghai Mental Health Center,
Shanghai Jiao Tong University School of Medicine, Shanghai, China.
[6]Institute of Psychology and Behavioral Science, Shanghai Jiao Tong University,
Shanghai, China.
[7]Peng Cheng Laboratory, Shenzhen, China.
† co-first authors
* co-corresponding authors

Correspondence should be addressed to:
Yixuan Ku, Department of Psychology, Sun Yat-Sen University, Guangzhou,
510006, China. E-mail: kuyixuan@gmail.com
and
Ru-Yuan Zhang, Shanghai Mental Health Center, School of Medicine, Shanghai
Jiao Tong University, Shanghai, China. E-mail: ruyuanzhang@gmail.com

Abstract

Schizophrenia patients are known to have profound deficits in visual working memory (VWM), and almost all previous studies attribute the deficits to decreased memory capacity. This account, however, ignores the potential contributions of other VWM components (e.g., memory precision). Here, we measure the VWM performance of schizophrenia patients and healthy control subjects on two classical delay-estimation tasks. Moreover, we thoroughly evaluate several established computational models of VWM to compare the performance of the two groups. We find that the model assuming variable precision across items and trials is the best model to explain the performance of both groups. According to the variable-precision model, schizophrenia subjects exhibit abnormally larger variability of allocating memory resources rather than resources per se. These results invite a rethink of the widely accepted decreased-capacity theory and propose a new perspective on the diagnosis and rehabilitation of schizophrenia.

Keywords: Schizophrenia, Visual working memory, Memory precision, Memory capacity, Bayesian inference, Perceptual variability

Schizophrenia is a severe mental disorder accompanied by a range of dysfunctions

in perceptual and cognitive behavior, among which working memory deficit is

considered as a core feature [1–4]. Working memory refers to the ability to temporally

store and manipulate information in order to guide appropriate behavior, and it has

been shown to link with a broad range of other brain functions, including perception,

attention, problem-solving and executive control [5–8]. Dysfunctions in working

memory therefore might cascade into multiple mental processes, causing a wide

spectrum of negative consequences [2,3,9].

A well-established finding in lab-based experiments is that people with

schizophrenia (SZ) exhibit worse performance than healthy control (HC) in visual

working memory (VWM) tasks [2]. This phenomenon has long been attributed to

decreased VWM capacity in SZ [2,10,11]. The theory of decreased capacity was

supported by previous studies using various VWM or other WM tasks, including

the 'span' tasks (e.g., digit span, spatial span, verbal span) [12,13], the N-back task [14–

16], the delayed-response task [17–19], the change detection task [20–24], and the delay-

estimation task [25–27]. Despite the considerable differences across behavioral tasks,

almost all previous studies converged to the same conclusion that decreased

capacity is the major cause of the VWM deficits in SZ.

Besides capacity, people have increasingly recognized memory *precision* as

another pivotal determinant of VWM performance in the basic research of VWM [28].

Precision reflects the amount of memory resources assigned to individual items—a

larger amount of resources leads to higher memory precision. At the neural level,

low perceptual precision might arise from either the intrinsic noise in neural

processing [29–31] or the fluctuations of cognitive factors (e.g., arousal, attention) [31,32].

Atypically increased variability in both behavioral and neural responses has been

discovered in patients with mental diseases such as autism spectrum disorder [33,34],

dyslexia [35], and attention-deficit/hyperactivity disorder [36]. These theoretical and

empirical studies raise the possibility that SZ and HC might differ in memory

precision rather than capacity—that is, these two groups might be able to remember

an equal number of items (i.e., comparable capacity) but SZ generally process and

maintain items in a less precise manner. Only a few studies have attempted to

simultaneously quantify memory capacity and precision in schizophrenic or
schizotypy subjects, and the results did not reach a consensus [25,26].

Despite the confound of different VWM components as the possible causes, it
is unclear whether SZ and HC employ the same computational strategies (i.e.,
observer model) in VWM. Most prior studies only used one model and implicitly
assumed that model is the best one for both SZ and HC. But without systematic
model comparisons model optimality cannot be firmly warranted, and endowed
results might be biased by the choice of a particular model. Given that several
influential models have been proposed to explain the VWM behavior in normal
subjects [28], it remains unclear which one is the best for SZ. If the best model for SZ
differs from the one for HC, it indicates that the two groups use qualitatively
different computational strategies to complete behavioral tasks. Otherwise, if SZ
and HC share the same best model, it indicates that they use the same memory
structure but quantitatively different parameters. These possibilities should be
thoroughly tested using transparent model comparisons.

In the present study, we aim to systematically disentangle the impact of
memory capacity and precision, as well as other factors (i.e., variability in
allocating resources and variability in choice) in SZ. In this study, the performance
of SZ and demographically matched HC was measured in two standard VWM
delayed-estimation tasks (Fig. 1). To obtain generalizable results, we tested separate
groups of subjects on the delay-estimation paradigm using two independent visual
features—color and orientation. Using standard tasks allows us to compare our
results with those from previous studies [25,37–40]. Most importantly, in contrast to
most prior studies, we evaluated and compared several mainstream computational
models in visual working memory research. This approach allows us to take an
unbiased perspective and search a large space of both models and parameters. We
believe that the well-controlled tasks and thorough computational modeling will
shed new light on the mechanisms of VWM deficits associated with schizophrenia.

Figure 1. Schematics of the color (***panel A***) and the orientation (***panel B***)
delay-estimation experiments. In the color experiment, subjects are instructed
to first memorize the colors of all squares (i.e., set size = 3 in this example trial)
on the screen, and after a 900ms delay choose the color of the probed square
(the one in the left lower visual field in this example) on a color wheel. The
orientation experiment is similar to the color experiment. Subjects are
instructed to remember the orientations of a set of bars (i.e., set size = 4 in this
example trial) and then adjust the orientation of the probed bar using a mouse.
Both color and orientation stimuli can be described using a circular feature
space of (0, 180]. Response error is the difference between the reported color
or orientation and the real color or orientation of the probe in this standard
feature space.

Results

Worse VWM performance in SZ in both color and orientation delay-estimation experiments

We tested subjects on two delay-estimation experiments. In the color delay-

estimation experiment, subjects viewed a set of squares on the screen, and were

instructed to memorize their colors. After a short blank, subjects were asked to

report the color of one of the squares (i.e., probe) by clicking the color on a color

wheel. In order to test a large cohort of subjects, we reduced the difficulty of the

task and tested them on two set size levels (i.e., the number of items to memorize).

Similarly, in the orientation delay-estimation experiment, subjects were instructed

to remember the orientations of a set of bars and then manually adjusted the orientation of the probe bar to the memorized orientation. To improve the efficacy of computational modeling, we increased the task difficulty and tested subjects on set size levels 1, 2, 4, 6 (see experimental details in Methods). These two tasks have been frequently used in prior VWM research [37,41,42]. Note that we recruited distinct groups of subjects for the two tasks in order to avoid task- or subject-specific confounding factors.

The precision of memory in each trial can be quantified as the difference between the reported color (or orientation) and the true color (respectively, orientation) of the probe. For the color experiment, a repeated-measure ANOVA was performed with circular standard deviation (CSD) as the dependent variable, set size (1/3) as the within-subject variable, group as the between-subject variable (Fig. 2A). Consistent with findings in previous literature, VWM performance was worse when set size was higher ($F(1, 119) = 641,703$, $p < 0.001$, partial $\eta^2 = 0.844$), and HC performed significantly better than SZ ($F(1,119) = 13.651$, $p < 0.001$, partial $\eta^2 = 0.103$). The interaction between set size and group was not significant ($F(1,119) = 0.229$, $p = 0.633$, partial $\eta^2 = 0.002$).

In the orientation experiment, similarly, we also found that HC achieved significant lower CSD (($F(3, 31) = 62.967$, $p < 0.001$, partial $\eta^2 = 0.656$). Unsurprisingly, both groups performed worse as set size increased ($F(3,31) = 87.682$, $p < 0.001$, partial $\eta^2 = 0.895$). The group by set size interaction was significant ($F(3,31) = 3.043$, $p = 0.043$, partial $\eta^2 = 0.228$), with smaller group differences on set size 1 and 6 and slightly larger group differences on set size 2 and 4 ($p < 0.001$ for all pairwise comparisons, Bonferroni corrected).

Taken together, we replicated the widely documented VWM deficits in SZ in both color and orientation delay-estimation tasks.

Figure 2. Behavioral performance of SZ and HC in the two tasks. Circular standard deviations of response errors in color (panel A) and orientation (panel B) delay-estimation tasks. SZ show higher CSDs (i.e., worse performance) than HC. All error bars represent SEM across subjects.

Variable-precision model accounts for VWM behavior in both HC and SZ

To systematically compare the VWM performance between SZ and HZ, we evaluated several mainstream computational models of VWM. We provide some brief introductions here, and readers may consider to skip the following paragraph to directly reach the result part or go to Supplementary Materials for detailed mathematical and intuitive explanations of the models, depending on reading preference.

The first one is the item-limit (IL) model. The IL model assumes no uncertainty in the sensory encoding stage, and that each subject has a fixed memory capacity and a fixed response variability across set size levels [43]. The second one is the mixture (MIX) model, similar to the IL model but assuming response variability is set-size dependent [25,26]. Compared with the MIX model, the slots-plus-averaging (SA) model [37] further elaborates the idea that memory resources manifest as discrete chunks, and these chunks can be flexibly assigned to multiple items. We also explored a modified version of the SA model, dubbed cosSA model, which inherits the idea of discrete memory resources and further assumes that response bias is stimulus-dependent and can be described as empirically derived periodic functions. The fifth one is the equal-precision (EP) model, which is similar to the

variable-precision (VP) model (Fig. 4) below but assumes that the memory
resources are evenly distributed across items and trials [44,45]. The VP model proposes
that memory resources are continuous, and the amount of resource assigned to
individual items varies across items and trials. Note that the VP model does not
include the capacity component thus we also constructed a variable-precision-with-
capacity (VPcap) model that not only acknowledges the variable precision
mechanisms but also explicitly estimates capacity of individual subjects. Note that
the IL, MIX, SA and cosSA, and VPcap models have the parameter of capacity, and
the EP and VP models do not. Here, capacity is operationally defined as the
maximum number of items that can be stored in memory. Some items are out of
memory if set size exceeds capacity, and the subject has to randomly guess the
color if one of the out-of-memory items is probed.

Figure 3. Model comparison results. A total of seven computational models
are compared in each subject. The pie charts illustrate the proportion (i.e., the
percent number shown in each slice) of subjects for whom each model is their
best-fitting model. Under both AIC and BIC model comparison metrics, the
VP model is the best-fitting model for the majority of subjects in both groups
and in both experiments. This result indicates both groups share a qualitatively
similar internal process of VWM.

We compared all seven models using the Akaike information criterion (AIC) and
the Bayesian information criterion (BIC) [46,47] in each subject. In the color
experiment, we found the VP model as the best-fitting model for over 84% of

subjects in the HC group under both metrics (Fig. 3), replicating previous results in
normal subjects [48,49]. Most importantly, the VP model was also the best-fitting
model for over 90% of subjects in the SZ group (Fig. 3). This result indicates that
both groups use the same observer model to perform the task.

The results were replicated in the orientation delay-estimation task. We
found that among 9 SZ subjects, the VP model was the best in 7 subjects using AIC
and in all 9 subjects using BIC. Similarly, in 26 HC subjects, the VP model
outperformed than other models in 21 and 24 subjects using AIC and BIC,
respectively.

Figure 4. Variable-precision model of VWM. *A*. Resource decay function.
The VP model assumes that the mean resource ($\bar{J}$) for processing a single item
declines as a power function of set size *N*, a trend characterized by two free
parameters—initial resources ($\bar{J}_1$) and decaying exponent (*a*). *B*. The resources
across items or trials follow a gamma distribution with the mean resource ($\bar{J}_1$)
determined by the resource decay function (panel A) and the resource
allocation variability (*τ*). Larger amounts of resources (*J*) indicate higher
precision and therefore generate narrower von Mises distributions (three small
axes indicating the precision equals to 5, 10 and 15 respectively) of stimulus
measurement (*m*). The widths of the von Mises distributions indicate the
degree of trial-by-trial sensory uncertainty. *C*. The eventual behavioral choice
given the internal stimulus measurement (*m*) is also uncertain, following a von
Mises distribution with the choice variability (κ_r) [50]. In the VP model, initial
resources ($\bar{J}$), decaying exponent (*a*), resource allocation variability (*τ*) and
choice variability (κ_r) are four free parameters to estimate (see details in SI and
van den Berg *et al*. [48]). All numbers here are only for illustration purposes and
not quantitatively related to the model fitting results in this paper.

The superior performance of the VP model indicates the important role of
variable precision in VWM processing. It is worth noting that the VP model, as the

best-fitting model, does not include a capacity term. As such, we cannot conclude
that the two groups must have the same capacity. But this result highlights the
importance of systematic model comparisons before comparing group differences
on model parameters.

Larger resource allocation variability in SZ

Analyses above have established that HC and SZ employ the qualitatively
same observer model to complete the VWM task. Their behavioral differences thus
should arise from the differences on some parameters in the observer model. We
next compared the fitted parameters of the VP model in the two groups. In the color
experiment, we found that the two groups had comparable resource decay functions
(Fig. 5A, initial resources, $t(119) = 0.689$, $p = 0.492$, $d = 0.125$; decaying exponent,
$t(119) = 1.065$, $p = 0.289$, $d = 0.194$), indicating a similar trend of diminished
memory resources as set size increases. SZ, however, had larger variability in
allocating resources (Fig. 5B, resource allocation variability, $t(119) = 4.03$, $p = 9.87$
$\times 10^{-5}$, $d = 0.733$). Furthermore, the VP model explicitly distinguishes the
variability in processing items and the variability in exerting a behavioral choice
(e.g., motor or decision noise). We found no significant group difference in the
choice variability (Fig. 5C, $t(119) = 1.7034$, $p = 0.091$, $d = 0.31$), excluding the
possibility that the atypical performance of SZ arises from larger variability at the
choice stage.

Highly consistent results were again discovered in the orientation
experiment: resource allocation variability was statistically higher in the SZ
subjects ($t(33) = 5.833$, $p = 1.576^{-6}$, $d = 2.256$), and no significant group differences
were detected in other parameters (initial resources, $t(33) = 0.437$, $p = 0.665$, $d = 0.169$; decaying exponent, $t(33) = 0.145$, $p = 0.886$, $d = 0.056$; choice variability,
$t(33) = 1.651$, $p = 0.108$, $d = 0.638$).

These results suggest that, although the two groups have on average the
same amount of memory resources across different set size levels, SZ allocate the
resources across items or trials in a more heterogeneous manner, with some items in
some trials receiving considerably larger amounts and vice versa in other cases.

This is unnatural since the probe was randomly chosen among all presented items

with an equal probability. The more common strategy therefore should be to assign

an equal amount of resources to every item and in every trial to tackle the

unpredictable target. Note that in the condition of set size equal to 1 (i.e., only one

item is presented), resource allocation cannot vary across items but may still vary

across trials, leading to worse performance in SZ. We further quantitatively

confirmed that increased resource allocation variability indeed leads to larger

behavioral response errors (see Supplementary Fig. S4).

Figure 5. Fitted parameters of the VP model in color (*panels A-C*) and
orientation (*panels D-F*) experiments. In the color experiment, no significant
group differences are noted between two groups in resource decay functions
(*panel A*), and choice variability (*panel C*). SZ have larger resource allocation
variability than HC (*panel B*). The results are replicated in the orientation
delay-estimation experiment (*panels D-F*). In panels A&D, the individual
resource decay functions are computed by $\bar{J} = \bar{J}_1 * N^{-a}$, where N is the set
size, and a are the estimated initial resources and the decaying exponent. The
solid lines represent the averaged resource decay functions across subjects.
The shaded areas in panels A and D, and all error bars in the other panels
represent ±SEM across subjects. Significance symbol conventions are ***: p <
0.001.

Resource allocation variability predicts the severity of schizophrenic symptoms

We next turned to investigate whether the results from the VP model can

predict clinical symptoms. In both tasks, a set of correlational analyses was carried

out to link the estimated resource allocation variability to the schizophrenia

symptomatology in each subject. BPRS, SANS, and SAPS questionnaires were

administered for each subject in the color experiment (Table 1 in Methods). The

Positive and Negative Syndrome Scale (PANSS)[51] was administered for each

subject in the orientation experiment. PANSS includes positive symptomatology,

negative symptomatology, as well as general psychopathology symptoms (Table 2

in Methods).

In the color experiment, we noticed that the estimated resource allocation

variability values of the SZ subjects correlated with their BPRS scores (Fig. 6A, r =

0.259, p = 0.045) and SANS scores (Fig. 6B, r = 0.302, p = 0.019). No significant

correlation was noted on the SAPS scores (Fig. 6C, r = -0.121, p = 0.358). In the

orientation experiment, we also found that resource allocation variability

significantly correlated with the severity of general psychopathology symptoms

(Fig. 6D, r = 0.717, p = 0.03) and negative symptoms (Fig. 6E, r = 0.882, p = 0.002),

but not positive symptoms (Fig. 6F, r = 0.551, p = 0.124).

Using different behavioral tasks and different measurements of

schizophrenia symptomatology, we found consistent results that the resource

allocation variability is linked to negative but not positive symptoms. These results

suggest that resource allocation variability not only is the key factor explaining

VWM behavior in SZ but also can quantitatively predict the severity of clinically

measured symptoms, highlighting its promising functionality in future diagnosis

and rehabilitations of schizophrenia.

Figure 6. Individual differences in resource allocation variability predict the scores in symptom assessments. In the color experiment, estimated resource allocation variability values in the SZ group significantly correlates with their scores on BPRS (*panel A*) and SANS (negative symptoms, *panel B*) but not on SAPS (positive symptoms, *panel C*). In the orientation experiment, resource allocation variability significantly correlates with the PANSS general scores (*panel D*), PANSS negative scores (*panel E*). This correlation is not significant for PANSS positive scores (*panel F*).

Control experiments

To further exclude the potential confounding factors, we run two additional control experiments.

In the first control experiment, we tested the color perception of the subjects in a color estimation task. This task is similar to the color delay-estimation task except that there was no delay period and the probe was always shown on the screen such that subjects could directly see the probe when choosing colors on the color-wheel (see details in Supplementary Materials). We found that the individuals' color perception thresholds cannot explain the group difference in resource allocation variability in Fig. 5B, indicating that the group differences are indeed memory related, not due to general worse perception in SZ.

In the second experiment, we aimed to exclude the possibility of imprecise model fitting due to low set size levels used in the color experiment. In order to recruit a large sample of subjects, we reduced the task difficulty and only tested two

relatively low set size levels (i.e., 1/3) in the color experiment. One caveat of this
approach is the possible imprecise model fitting because we did not challenge
subjects' ceiling performance. Although we obtained highly consistent results in the
orientation experiment using more and higher set size levels, there still exist the
possibilities that 1) the VP model might not be the best model, and more
importantly, 2) the estimation of the model parameters (e.g., capacity parameter K
in the VPcap model) might be inaccurate.

We further tested another large cohort of HC subjects (N=62) on the
identical color delay-estimation task but used higher set size levels (i.e., 2, 4, and
6). Memorizing the colors of 6 objects is extremely challenging, and previous
studies usually regarded 6 as the ceiling condition in a delay-estimation task [37,52,53].
We performed the same analyses and found that the VP model was still the best-
fitting model in over 70% of the subjects (Supplementary Fig. S3).

In sum, the three datasets (the color experiment, the orientation experiment,
and the control color experiment) including a total of 149 HC and 69 SZ subjects
demonstrate the converging result that the VP model is the best model for both HC
and SZ subjects. This finding is unlikely due to some idiosyncratic experimental
settings.

Discussion

The mechanisms of VWM deficits in schizophrenia have been a matter of debate
over the past few years. Decreased capacity has been widely proposed as the major
cause of the deficits in SZ. In the present study, we re-examine this conclusion by
comparing the performance of SZ and HC using several mainstream computational
models of VWM proposed so far. We first establish that the VP model is the best
model to characterize performance of both groups, indicating a qualitative similar
internal process in both groups. We then further evaluate different components in
the VP model and find that SZ have larger variability in allocating memory
resources. These findings are highly consistent in two independent samples of
subjects and in two independent behavioral tasks. Furthermore, individual
differences in resource allocation variability predict variation of patients' symptom

severity, highlighting its clinical functionality. Taken together, our results propose

for the first time that resource allocation variability is the key factor that limits

VWM performance in schizophrenia.

A large body of literature has documented that SZ perform poorly in various

forms of working memory tasks [2,3,54,55]. The majority of the studies made the same

conclusion: memory capacity is decreased in schizophrenia. However, through a

careful examination of the literature, we find that the definition of capacity varies

substantially across studies. Many studies directly equated worse performance with

decreased capacity without quantitatively demonstrating how capacity modulates

performance. For example, memory capacity was defined as the number of digits

that can be recalled in the longest strand in digit span tasks [12]. In N-back tasks,

capacity was defined as the number of backs corresponding to a certain accuracy

level [14–16]. Moreover, the calculation of capacity resembled the d-prime metric in

change detection tasks [22–24,43,56]. The majority of these metrics are actually

behavioral thresholds or accuracy related to capacity rather than direct

quantifications of capacity. This is partly because we lack appropriate

computational models for the majority of those tasks. The VP model is

advantageous as it describes the generative process of the delay-estimation task and

the change-detection task [48]. As such, it allows to disassociate the effect of capacity

from other VWM components. Note that here we only consider resource, capacity,

resource allocation, and choice as the dimensions of modeling. Some studies

suggest the existence of "binding error" in VWM [42,57]. Van den Berg et al 2014 [49]

explored a large space of model parameters and found that "binding error" only

account for small fraction of VWM limitation. As such we did not consider

"binding error" in this paper. Also, all computational models we used in this study

are mostly specific to the delay-estimation task. It is possible that the VWM deficits

in schizophrenia manifest differently in different tasks. Future studies are needed to

test whether larger resource allocation variability can also account for VWM

deficits of SZ in other WM tasks.

Only a few studies have quantitatively estimated capacity and precision in

schizophrenia. Gold et al [25] employed the same color delay-estimation task to that

in our study and estimated individual's capacity and precision using the MIX model. Results in that study echoed the decreased-capacity theory. The MIX model, however, does not consider two important factors. First, the model assumes an equal precision across items in memory. Second, the model does not separate the variability for processing stimuli (i.e., sensory uncertainty, κ in Supplementary Eq. S5) and the variability in exertion of a choice (i.e., choice uncertainty, κ_r in Supplementary Eq. S6). This distinction is important since it highlights different types of uncertainty in encoding and decoding stages of VWM. Mathematically, these two types of uncertainty can be distinguished by manipulating set size since the encoding variability depends on set size but the choice variability does not. The issues of the MIX model have been symmetrically addressed in recent work [58].

Compared with capacity and precision—the two diagnostic features of VWM—resource allocation variability emerges as a new concept in VWM. It refers to the heterogeneity of allocating resources across multiple items and trials. We speculate that resource allocation variability might reflect the stability of attentional control when the brain processes multiple objects. First, it has been shown that attention and WM are two core components of executive control and tightly linked with each other [59,60]. Second, schizophrenia is known to have deficits in top-down attentional modulation [55,59]. Particularly, recent studies discovered the phenomenon of spatial hyperfocusing in schizophrenia patients[19,61–63]. If schizophrenia patients overly attend to one item and ignore others in the memory encoding stage, unbalanced resource allocation will likely occur. Note that SZ also exhibit worse performance when only one target is presented. In this condition, resources do not vary across items (i.e., only one time) but could vary across trials. Also, we want to emphasize that such variability is not equivalent to attentional lapse. A higher attentional lapse rate may lead to overall fewer resources, a phenomenon we did not observe in this study.

What are the neural mechanisms of this resource allocation variability? Recent neurophysiological studies proposed that the neural representation of a stimulus may follow a doubly stochastic process [64,65], which suggests that the variability in encoding precision is a consequence of trial-to-trial and item-to-item

fluctuations in attentional gain [32,48,66]. A recent study combined functional magnetic

resonance imaging and the VP model, showing that the superior intraparietal sulcus

(IPS) is the cortical locus that controls the resource allocation [67]. Interestingly,

schizophrenia patients have been known to have IPS deficits [68]. Note that besides

top-down factors, we cannot rule out the contribution of bottom-up neural noise in

perceptual and cognitive processing [64,65], as found in several other mental diseases

[33–36]. Also, a recent study found that variable precision is even more likely to be

driven by stimulus-specific effects [69].

The current results also reveal links between resource allocation variability

and patients' negative symptoms, but not positive symptoms (Fig. 6). These

findings are consistent with several experimental and meta-analysis studies

claiming dissociable mechanisms underlying the cluster of negative symptoms

versus that of positive symptoms [70–73]. More broadly, a growing collection of

evidence suggests that visual perceptual deficits in schizophrenic patients are more

likely to link to negative rather than positive symptom severity [74–78]. Negative

symptoms in turn might produce improvised social functioning. Humans depend

heavily on VWM to interact with multiple agents and complete social tasks.

Deficits in distributing processing resources over multiple agents therefore might

cause disadvantages in social cognition.

In conclusion, our study proposes a new explanation that the resource

allocation variability accounts for the atypical VWM performance in schizophrenia.

This view differs from the decreased-capacity theory and provides a new direction

for future studies that attempt to promote diagnosis and rehabilitation for

schizophrenic patients.

Methods

467 **Ethics Statement.** All experimental protocols were approved by the institutional

468 review board at the East China Normal University. All research was performed in

469 accordance with relevant guidelines and regulations. Informed written consent was

470 obtained from all participants.

Color delay-estimation experiment

Subjects. 61 HC and 60 SZ participated in the experiment. SZ were clinically (symptom and medication) stable inpatients (N = 33) and outpatients (N = 27) who met DSM-IV criteria [79] for schizophrenia. Patients having the history of any other mental or neurological disorders were excluded. All patients were receiving antipsychotic medication (2 first-generation, 43 second-generation, 15 both). Symptom severity was evaluated by the Brief Psychiatric Rating Scale (BPRS) [80], the Scale for the Assessment of Negative (SANS) and Positive Symptoms (SAPS) [81,82]. HC were recruited by advertisement. All HC had no current diagnosis of axis 1 or 2 disorders as well as no family history of psychosis nor substance abuse or dependence. All subjects are right-handed with normal sight and color perception.

The two groups were matched in age ($t(119) = 1.58$, $p = 0.118$, $d = 0.284$), gender (31 females and 29 males) and education level of parents ($t(119) = 0.257$, $p = 0.798$, $d = 0.047$). Inevitably, the SZ had fewer years of education than the HC ($t(119) = 5.51$, $p = 2.09 \times 10^{-7}$, $d = 1.00$). The detailed demographic information is summarized in the Table 1.

Table 1. Demographics and clinical information of people with schizophrenia (SZ) and healthy control subjects (HC)

	SZ (N = 60)		HC (N = 61)	
	Mean	SD	Mean	SD
Age	35.67	6.58	33.82	9.90
range	23-48	n/a	21-57	n/a
Female/male	31/29	n/a	29/32	n/a
Inpatient/outpatient	33/27	n/a	n/a	n/a
Subject's education (in years)	12.03	2.24	15.13	3.70
Paternal education (in years) [a]	9.89	2.53	9.76	2.95
Maternal education (in years)	9.62	2.91	9.29	3.63
BPRS	27.25	6.27	n/a	n/a
SAPS	5.77	7.02	n/a	n/a
SANS	24.43	11.45	n/a	n/a

[a] Average of mother's and father's years of education

BPRS: Brief Psychiatric Rating Scale [80]; SAPS: Scale for the Assessment of

Positive Symptoms [82]; SANS: Scale for the Assessment of Negative Symptoms [81].

Stimuli and Task. The subjects sat 50 cm away from an LCD monitor. All

stimuli were generated by Matlab 8.1 and Psychtoolbox 3 [83,84], and then presented

on a LCD monitor.

In the color delay-estimation task (Fig. 1A), each trial began with a fixation

cross presented at center-of-gaze for a duration randomly chosen from a sequence

of 300, 350, 400, 450 and 500 ms. Subjects shall keep their fixation on the cross

throughout the whole experiment. A set of colored squares (set size = 1 or 3) was

shown on an invisible circle with 4^o radius. Our pilot experiment showed that SZ

patients exhibit a high dropout rate if the task is longer than 30 mins or too hard

(i.e., set size > 4). We therefore reduced the difficulty of the color task to set size

level 1 and 3. The sample array lasted 500 ms. Each square was $1.5^o \times 1.5^o$ of visual

angle. Their colors were randomly selected from the 180 colors that are equally

distributed along the wheel representing the CIE L*a*b color space. The color

wheel was centered at (L = 70, a = 20, b = 38) with a radius of 60 in the color space

[37]. The sample array then disappeared and was followed by a 900 ms blank period

for memory retention. After the delay, an equal number of outlined squares were

shown at the same location of each sample array item, with one of them bolded as

the probe. In the meantime, a randomly rotated color wheel was shown. The color

wheel was 2.1^o thick and centered on the monitor with the inner and the outer radius

as 7.8^o and 9.8^o respectively. Subjects were asked to choose the remembered color

of the probe by clicking a color on the color wheel using a computer mouse.

Subjects shall choose the color as precisely as possible and response time was not

constrained. Every subject completed 2 blocks for the set size 1 and 3, respectively.

The order of the two blocks was counterbalanced across subjects. Each block had

80 trials. The difference between the reported color and the true color of the target

is considered as the response error.

Orientation delay-estimation experiment

Subjects. Data from 26 HC and 9 SZ were obtained in the orientation delay-

estimation task (see Table 2 for subject information). The data from the HC subjects

has been previously presented in ref. [41]. SZ were all clinically stable inpatients who

met the DSM-IV criteria for schizophrenia[79]. Patients having a history of any other

mental or neurological disorders were excluded. All nine patients were receiving

second-generation antipsychotic medication. The Positive and Negative Syndrome

Scale (PANSS)[51] was used to evaluate the psychotic symptoms of the patients. This

scale includes positive symptomatology, negative symptomatology, as well as

general psychopathology symptoms. HC subjects were recruited by advertisement.

All HC had no current diagnosis of axis 1 or 2 disorders as well as no family history

of psychosis nor substance abuse or dependence. All subjects are right-handed with

normal sight.

Table 2. Demographics and clinical information of the subjects in the

orientation delay-estimation task

	SZ (N = 9)		HC (N = 26)	
	Mean	SD	Mean	SD
Age	39.33	9.35	21.31	1.94
range	27-56	n/a	19-26	n/a
Female/male	0/9	n/a	16/26	n/a
Inpatient/outpatient	9/9	n/a	n/a	n/a
Subjects' education (years)	9	2.12	14.12	1.63
PANSS - positive	11.33	4.33	n/a	n/a
PANSS - negative	22	8.14	n/a	n/a
PANSS - general	31.78	14.10	n/a	n/a

537   PANSS: The Positive and Negative Syndrome Scale[51]

539   _Stimuli and Task._ Stimuli were presented on a 60 Hz LCD monitor through

540   Matlab psychophysics Toolbox (Version 3). In the sample array, all items were

541   shown on an invisible circle with 10^o radius. The length and width of each item in

542   the sample array were 3.7^o and 0.6^o respectively. Then orientations of the bars were

randomly chosen from 1^o to 180^o. The procedure of this task was similar to the color delay-estimation task, except that the sample array was presented for 200 ms (Fig. 1B). In the probe array, one of the sample bars appeared as the probe, and subjects were required to adjust the orientation of the probed bar using a computer mouse.

The HC subjects completed 5 blocks with 100 trials in each block. The set size of each trial was randomly chosen from 1, 2, 3, 4, 6. The SZ patients complete the experiment in three visit sessions. In each visit, they were asked to finish 10 blocks with 40 trials in each block. The set sizes were 1, 2, 4, 6 and counter-balanced across blocks. We analyzed the behavioral performance using only the data of set size levels 1, 2, 4, 6 because they were the conditions shared by both groups. For model fitting, we included all set size levels (1/2/3/4/6) of the HC group and compared the likelihood, AIC, and BIC per trial to compensate the trial difference between groups and subjects. One patient only finished 9 blocks in the second visit and another patient only finished 8 blocks in the first visit. In total, seven SZ subjects completed 300 trials for each set size, one SZ completed 290 trials and one SZ completed 280 trials.

Data availability statement

The data are available from the corresponding author upon reasonable request.

Acknowledgments

We thank Zheng Ma, Ting Qian and Haojiang Ying for their invaluable comments on the manuscript. This work was supported by the National Social Science Foundation of China (17ZDA323), the Shanghai Committee of Science and Technology (19ZR1416700, 17JC1404101, 17JC1404105) to Y.K.

Author contributions

571   Y. K., Y. Z. and X. R. designed the experiments. X. R. and L. Z. performed the

572   experiments. Y. Z., T.M. and R-Y. Z. analyzed the data, Y. Z and R-Y. Z. wrote the

573   manuscript in consultation with Y. K. and L. Z.

References

1. Gold, J. M., Randolph, C. & Carpenter, C. Auditory working memory and Wisconsin Card Sorting Test Performance in Schizophrenia. *Arch. Gen. Psychiatry* **54**, 159–165 (1997).

2. Lee, J. & Park, S. Working Memory Impairments in Schizophrenia: A Meta-Analysis. *J. Abnorm. Psychol.* **114**, 599–611 (2005).

3. Forbes, N. F., Carrick, L. A., McIntosh, A. M. & Lawrie, S. M. Working memory in schizophrenia: a meta-analysis. *Psychol. Med.* **39**, 889–905 (2009).

4. Goldman-Rakic, P. S. Working memory dysfunction in schizophrenia. *J. Neuropsychiatry Clin. Neurosci.* **6**, 348–357 (1994).

5. Mayer, R. E. & Moreno, R. A split-attention effect in multimedia learning: Evidence for dual processing systems in working memory. *J. Educ. Psychol.* **90**, 312–320 (1998).

6. Postle, B. R. Working memory as an emergent property of the mind and brain. *Neuroscience* **139**, 23–38 (2006).

7. Nee, D. E. *et al.* A meta-Analysis of executive components of working memory. *Cereb. Cortex* **23**, 264–282 (2013).

8. Luck, S. J. & Vogel, E. K. Visual working memory capacity: From psychophysics and neurobiology to individual differences. *Trends Cogn. Sci.* **17**, 391–400 (2013).

9. Coleman, M. J., Krastoshevsky, O., Tu, X., Mendell, N. R. & Levy, D. L. The effects of perceptual encoding on the magnitude of object working memory impairment in schizophrenia. *Schizophr. Res.* **139**, 60–65 (2012).

10. Leonard, C. J. *et al.* Testing sensory and cognitive explanations of the antisaccade deficit in schizophrenia. *J. Abnorm. Psychol.* **122**, 1111–1120 (2013).

11. Johnson, M. K. *et al.* The relationship between working memory capacity and broad measures of cognitive ability in healthy adults and people with schizophrenia. *Neuropsychology* **27**, 220–229 (2013).

12. Conklin, H. M. Verbal Working Memory Impairment in Schizophrenia

Patients and Their First-Degree Relatives: Evidence From the Digit Span
Task. *Am. J. Psychiatry* **157**, 275–277 (2000).

13.   Chey, J., Lee, J., Kim, Y.-S., Kwon, S.-M. & Shin, Y.-M. Spatial working
memory span, delayed response and executive function in schizophrenia.
Psychiatry Res. **110**, 259–271 (2002).

14.   Callicott, J. Functional Magnetic Resonance Imaging Brain Mapping in
Psychiatry: Methodological Issues Illustrated in a Study of Working Memory
in Schizophrenia. *Neuropsychopharmacology* **18**, 186–196 (1998).

15.   Barch, D. M., Csernansky, J. G., Conturo, T. & Snyder, A. Z. Working and
long-term memory deficits in schizophrenia: Is there a common prefrontal
mechanism? *J. Abnorm. Psychol.* **111**, 478–494 (2002).

16.   Jansma, J. M., Ramsey, N. F., Van Der Wee, N. J. A. & Kahn, R. S. Working
memory capacity in schizophrenia: A parametric fMRI study. *Schizophr.*
Res. **68**, 159–171 (2004).

17.   Park, S. Schizophrenics Show Spatial Working Memory Deficits. *Arch. Gen.*
Psychiatry **49**, 975 (1992).

18.   Keedy, S. K., Ebens, C. L., Keshavan, M. S. & Sweeney, J. A. Functional
magnetic resonance imaging studies of eye movements in first episode
schizophrenia: Smooth pursuit, visually guided saccades and the oculomotor
delayed response task. *Psychiatry Res.* **146**, 199–211 (2006).

19.   Sawaki, R. *et al.* Hyperfocusing of attention on goal-related information in
schizophrenia: Evidence from electrophysiology. *J. Abnorm. Psychol.* **126**,
106–116 (2017).

20.   Gold, J. M., Wilk, C. M., McMahon, R. P., Buchanan, R. W. & Luck, S. J.
Working memory for visual features and conjunctions in schizophrenia. *J.*
Abnorm. Psychol. **112**, 61–71 (2003).

21.   Lencz, T. *et al.* Impairments in perceptual competency and maintenance on a
visual delayed match-to-sample test in first-episode schizophrenia. *Arch.*
Gen. Psychiatry **60**, 238–243 (2003).

22.   Erickson, M. A. *et al.* Impaired Working Memory Capacity Is Not Caused by
Failures of Selective Attention in Schizophrenia. *Schizophr. Bull.* **41**, 366–

373 (2015).

23. Erickson, M. A. *et al.* Enhanced vulnerability to distraction does not account for working memory capacity reduction in people with schizophrenia. *Schizophr. Res. Cogn.* **1**, 149–154 (2014).

24. Leonard, C. J. *et al.* Toward the Neural Mechanisms of Reduced Working Memory Capacity in Schizophrenia. *Cereb. Cortex* **23**, 1582–1592 (2013).

25. Gold, J. M. *et al.* Reduced Capacity but Spared Precision and Maintenance of Working Memory Representations in Schizophrenia. *Arch. Gen. Psychiatry* **67**, 570–577 (2010).

26. Xie, W. *et al.* Schizotypy is associated with reduced mnemonic precision in visual working memory. *Schizophr. Res.* **193**, 91–97 (2018).

27. Starc, M. *et al.* Schizophrenia is associated with a pattern of spatial working memory deficits consistent with cortical disinhibition. *Schizophr. Res.* **181**, 107–116 (2017).

28. Ma, W. J., Husain, M. & Bays, P. M. Changing concepts of working memory. *Nat. Neurosci.* **17**, 347–356 (2014).

29. Bialek, W. Physical Limits to Sensation and Perception. *Annu. Rev. Biophys. Biophys. Chem.* **16**, 455–478 (1987).

30. Rolls, E. T. & Deco, G. *The Noisy Brain: Stochastic Dynamics as a Principle of Brain Function*. (Oxford University Press, 2010).

31. Faisal, A. A., Selen, L. P. J. & Wolpert, D. M. Noise in the nervous system. *Nature Reviews Neuroscience* **9**, 292–303 (2008).

32. Goris, R. L. T., Movshon, J. A. & Simoncelli, E. P. Partitioning neuronal variability. *Nat. Neurosci.* **17**, 858–865 (2014).

33. Dinstein, I. *et al.* Unreliable Evoked Responses in Autism. *Neuron* **75**, 981–991 (2012).

34. Park, W. J., Schauder, K. B., Zhang, R., Bennetto, L. & Tadin, D. High internal noise and poor external noise filtering characterize perception in autism spectrum disorder. *Sci. Rep.* **7**, 17584 (2017).

35. Sperling, A. J., Lu, Z.-L., Manis, F. R. & Seidenberg, M. S. Deficits in perceptual noise exclusion in developmental dyslexia. *Nat. Neurosci.* **8**, 862–

863 (2005).

36. Bubl, E. *et al.* Elevated Background Noise in Adult Attention Deficit

Hyperactivity Disorder Is Associated with Inattention. *PLoS One* **10**,

e0118271 (2015).

37. Zhang, W. & Luck, S. J. Discrete fixed-resolution representations in visual

working memory. *Nature* **453**, 233–235 (2008).

38. Zhang, W. & Luck, S. J. Sudden Death and Gradual Decay in Visual

Working Memory: Research Report. *Psychol. Sci.* **20**, 423–428 (2009).

39. Zhang, W. & Luck, S. J. The number and quality of representations in

working memory. *Psychol. Sci.* **22**, 1434–1441 (2011).

40. Foster, J. J., Bsales, E. M., Jaffe, R. J. & Awh, E. Alpha-Band Activity

Reveals Spontaneous Representations of Spatial Position in Visual Working

Memory. *Curr. Biol.* **27**, 3216-3223.e6 (2017).

41. Zhao, Y., Kuai, S., Zanto, T. P. & Ku, Y. Neural Correlates Underlying the

Precision of Visual Working Memory. *Neuroscience* **425**, 301–311 (2020).

42. Bays, P. M., Catalao, R. F. G. & Husain, M. The precision of visual working

memory is set by allocation of a shared resource. *J. Vis.* **9**, 7.1-11 (2009).

43. Pashler, H. Familiarity and visual change detection. *Percept. Psychophys.* **44**,

369–378 (1988).

44. Palmer, J. Attentional limits on the perception and memory of visual

information. *J. Exp. Psychol. Hum. Percept. Perform.* **16**, 332–350 (1990).

45. Shaw, M. Identifying attentional and decision-making components in

information processing. in *Attention and Performance* (ed. Nickerson, R.)

227–296 (Erlbaum, 1980).

46. Wit, E., van den Heuvel, E. & Romeijn, J. W. 'All models are wrong. ': An

introduction to model uncertainty. *Stat. Neerl.* **66**, 217–236 (2012).

47. Burnham, K. P. & Anderson, D. R. *Model Selection and Multimodel*

Inference: A Practical Information-Theoretic Approach. *Ecological*

Modelling **172**, (Springer-Verlag, 2002).

48. van den Berg, R., Shin, H., Chou, W.-C., George, R. & Ma, W. J. Variability

in encoding precision accounts for visual short-term memory limitations.

Proc. Natl. Acad. Sci. **109**, 8780–8785 (2012).

49.   van den Berg, R., Awh, E. & Ma, W. J. Factorial comparison of working

memory models. *Psychol. Rev.* **121**, 124–149 (2014).

50.   Osborne, L. C., Lisberger, S. G. & Bialek, W. A sensory source for motor

variation. *Nature* **437**, 412–416 (2005).

51.   Kay, S. R., Opler, L. A. & Lindenmayer, J.-P. The Positive and Negative

Syndrome Scale (PANSS): Rationale and Standardisation. *Br. J. Psychiatry*

155, 59–65 (1989).

52.   Luck, S. J. & Vogel, E. K. The capacity of visual working memory for

features and conjunctions. *Nature* **390**, 279–281 (1997).

53.   Xu, Y. & Chun, M. M. Dissociable neural mechanisms supporting visual

short-term memory for objects. *Nature* **440**, 91–95 (2006).

54.   Shakow, D. The Worcester State Hospital research on schizophrenia (1927-

1946). *J. Abnorm. Psychol.* **80**, 67–110 (1972).

55.   Gold, J. M., Hahn, B., Strauss, G. P. & Waltz, J. A. Turning it upside down:

Areas of preserved cognitive function in schizophrenia. *Neuropsychol. Rev.*

19, 294–311 (2009).

56.   Cowan, N. The magical number 4 in short-term memory: A reconsideration

of mental storage capacity. *Behav. Brain Sci.* **24**, 87–114 (2001).

57.   Wheeler, M. E. & Treisman, A. M. Binding in short-term visual memory. *J.*

Exp. Psychol. Gen. **131**, 48–64 (2002).

58.   Ma, W. J. Problematic usage of the Zhang and Luck mixture model. *bioRxiv*

(2018). doi:10.1101/268961

59.   Gold, J. M. *et al.* Selective Attention, Working Memory, and Executive

Function as Potential Independent Sources of Cognitive Dysfunction in

Schizophrenia. *Schizophr. Bull.* **44**, 1227–1234 (2018).

60.   Christophel, T. B., Iamshchinina, P., Yan, C., Allefeld, C. & Haynes, J. D.

Cortical specialization for attended versus unattended working memory. *Nat.*

Neurosci. **21**, 494–496 (2018).

61.   Fuller, R. L. *et al.* Impaired control of visual attention in schizophrenia. *J.*

Abnorm. Psychol. **115**, 266–275 (2006).

62. Kreither, J. *et al.* Electrophysiological Evidence for Hyperfocusing of Spatial
Attention in Schizophrenia. *J. Neurosci.* **37**, 3813–3823 (2017).

63. Luck, S. J. *et al.* Hyperfocusing in schizophrenia: Evidence from interactions
between working memory and eye movements. *J. Abnorm. Psychol.* **123**,
783–795 (2014).

64. Churchland, A. K. *et al.* Variance as a Signature of Neural Computations
during Decision Making. *Neuron* **69**, 818–831 (2011).

65. Churchland, M. M. *et al.* Stimulus onset quenches neural variability: a
widespread cortical phenomenon. *Nat. Neurosci.* **13**, 369–378 (2010).

66. Nienborg, H. & Cumming, B. G. Decision-related activity in sensory neurons
reflects more than a neuron's causal effect. *Nature* **459**, 89–92 (2009).

67. Galeano Weber, E. M., Peters, B., Hahn, T., Bledowski, C. & Fiebach, C. J.
Superior Intraparietal Sulcus Controls the Variability of Visual Working
Memory Precision. *J. Neurosci.* **36**, 5623–5635 (2016).

68. Zhou, S.-Y. *et al.* Parietal lobe volume deficits in schizophrenia spectrum
disorders. *Schizophr. Res.* **89**, 35–48 (2007).

69. Shen, S. & Ma, W. J. Variable precision in visual perception. *Psychol. Rev.*
126, 89–132 (2019).

70. de Gracia Dominguez, M., Viechtbauer, W., Simons, C. J. P., van Os, J. &
Krabbendam, L. Are Psychotic Psychopathology and Neurocognition
Orthogonal? A Systematic Review of Their Associations. *Psychol. Bull.* **135**,
157–171 (2009).

71. Cameron, A. M. *et al.* Working memory correlates of three symptom clusters
in schizophrenia. *Psychiatry Res.* **110**, 49–61 (2002).

72. Carter, C. *et al.* Spatial working memory deficits and their relationship to
negative symptoms in unmedicated schizophrenia patients. *Biol. Psychiatry*
40, 930–932 (1996).

73. Park, S., Püschel, J., Sauter, B. H., Rentsch, M. & Hell, D. Visual object
working memory function and clinical symptoms in schizophrenia.
Schizophr. Res. **59**, 261–268 (2003).

74. Cadenhead, K. S. *et al.* Information processing deficits of schizophrenia

patients: Relationship to clinical ratings, gender and medication status. *Schizophr. Res.* **28**, 51–62 (1997).

75. Butler, P. D. & Javitt, D. C. Early-stage visual processing deficits in schizophrenia. *Current Opinion in Psychiatry* **18**, 151–157 (2005).

76. Kéri, S., Kiss, I., Kelemen, O., Benedek, G. & Janka, Z. Anomalous visual experiences, negative symptoms, perceptual organization and the magnocellular pathway in schizophrenia: A shared construct? *Psychol. Med.* **35**, 1445–1455 (2005).

77. Slaghuis, W. L. Spatio-temporal luminance contrast sensitivity and visual backward masking in schizophrenia. *Exp. brain Res.* **156**, 196–211 (2004).

78. Slaghuis, W. L. & Bishop, A. M. Luminance flicker sensitivity in positive- and negative-symptom schizophrenia. *Exp. Brain Res.* **138**, 88–99 (2001).

79. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders (4th ed.)*. *American Psychiatric Publishing* (1994).

80. Overall, J. E. & Gorham, D. R. The Brief Psychiatric Rating Scale. *Psychol. Rep.* **10**, 799–812 (1962).

81. Andreasen, N. C. *The Scale for the Assessment of Negative Symptoms (SANS)*. *University of Iowa* (1983).

82. Andreasen, N. C. *The Scale for the Assessment of Positive Symptoms (SAPS)*. *University of Iowa* (1984).

83. Brainard, D. H. The Psychophysics Toolbox. *Spat. Vis.* **10**, 433–436 (1997).

84. Pelli, D. G. The VideoToolbox software for visual psychophysics: Transforming numbers into movies. *Spat. Vis.* **10**, 437–442 (1997).