Atypically larger variability of resource allocation accounts for visual working memory deficits in schizophrenia

Yijie Zhao[1], Xuemei Ran[1], Li Zhang[1], Ruyuan Zhang[2*], Yixuan Ku[1,3*]

[1]The Shanghai Key Lab of Brain Functional Genomics, Shanghai Changning-
ECNU Mental Health Center, School of Psychology and Cognitive Science, East
China Normal University, Shanghai, 200062 China

[2]Center for Magnetic Resonance Research, Department of Neuroscience, University
of Minnesota, Minneapolis, MN 55455 USA

[3]NYU-ECNU Institute of Brain and Cognitive Science, NYU Shanghai and
Collaborative Innovation Center for Brain Science, Shanghai, 200062 China

* co-senior author

Correspondence:
Ruyuan Zhang ruyuanzhang@gmail.com
Yixuan Ku yk1616@nyu.edu
20

Authors' contributions. RZ, YK &YZ developed research idea and study concepts;
RZ, YK & YZ design the research; XR and LZ collected the data; RZ performed
the data analyses and modeling; RZ & YZ wrote the manuscript.

Acknowledgement. We thank Zheng Ma, Ting Qian and Haojiang Ying for their
invaluable comments on the manuscript. This work was supported by the National
Social Science Foundation of China (17ZDA323), the National Key Fundamental
Research Program of China (2013CB329501), the Major Program of Science and

Technology Commission Shanghai Municipal (17JC1404100), the Fundamental

Research Funds for the Central Universities (2018ECNU-QKT015), and the NYU-

ECNU Institute of Brain and Cognitive Science at NYU (YK).

ABSTRACT

Atypical performance of people with schizophrenia (PSZ) in visual working memory (VWM) has long been attributed to decreased capacity compared with healthy control subjects (HCS). This notion, however, largely ignores the effects of other VWM components, such as precision, on behavioral performance. Here, we measured the performance of 60 PSZ and 61 HCS in a classical delay-estimation task and disentangled the contribution of various VWM components by thoroughly comparing several influential computational models of VWM. Surprisingly, none of the models suggest group differences in memory capacity and in memory resources across set size levels–two diagnostic features of VWM. Notably, we find that the model assuming variable precision (VP) across items and trials is the best model to explain the performance of both groups, indicating the two groups employ the qualitatively same internal process in VWM. According to the VP model, PSZ only exhibit abnormally larger variability of allocating memory resources rather than resources *per se*. These results directly challenge the widely accepted decreased-capacity theory and establish the first time to our knowledge that the elevated resource allocation variability is the major determinant of the atypical VWM behavior in schizophrenia. Finally, we show that individual differences in the resource allocation variability predict variation of symptom severity in PSZ, highlighting its functional relevance to schizophrenic pathology. Taken together, our findings provide a novel account for the VWM deficits in schizophrenia and shed a new light on the utility of generative computational models to characterize mechanisms of mental deficits in clinical neuroscience.

Keywords:

Schizophrenia, Visual working memory, Memory precision, Generative model

SIGNIFICANCE STATEMENT

Working memory deficits in schizophrenia have long been thought to arise from abnormally decreased capacity. We revisit this topic by leveraging the most recent advances in research on visual working memory. Combining a classical visual working memory paradigm and generative computational modeling, we challenge the conventional decreased-capacity theory and demonstrate that the atypically larger variability of distributing memory resources across items and trials accounts for behavioral deficits observed in schizophrenia. The current study offers a new perspective for future studies aiming for characterizing the diagnostic pathology of schizophrenia.

INTRODUCTION

Schizophrenia is a severe mental disorder accompanied by a range of dysfunctions in perceptual and cognitive behavior, among which working memory deficit is considered as a core feature (1–4). Working memory refers to the ability to temporally store and manipulate information in order to guide appropriate behavior, and it has been shown to link with a broad range of other brain functions, including perception, attention, problem-solving and executive control (5–8). Dysfunctions in working memory therefore might cascade into multiple mental processes, causing a wide spectrum of negative consequences (2, 3, 9).

A well-established finding in lab-based experiments is that people with schizophrenia (PSZ) exhibit worse performance than healthy control subjects (HCS) in visual working memory (VWM) tasks (2). This phenomenon has long been attributed to decreased VWM *capacity* in PSZ (2, 10, 11), a theory supported by previous studies using various VWM tasks, including the 'span' tasks (e.g., digit span, spatial span, verbal span) (12, 13), the N-back task (14–16), the delayed-response task (17–19), the change detection task (20–24) and the delay-estimation task (25–27). Despite the considerable differences across tasks, almost all previous studies converged to the same conclusion that decreased-capacity is the major cause for the VWM deficits in PSZ.

In another line of research, people have increasingly recognized memory *precision* as a pivotal determinant of VWM performance (28). Precision indicates the amount of memory resources assigned to individual items–a larger amount of resources produce higher memory precision. At the neural level, low perceptual precision might arise from either the intrinsic noise in neural responses (29–31) or the fluctuations of internal cognitive factors (e.g., arousal, attention) (31, 32). Atypically increased variability in both behavioral and neural responses has been discovered in patients with mental diseases such as autism spectrum disorder (33, 34), dyslexia (35), ADHD (36). These theoretical and empirical studies raise the possibility that PSZ and HCS might differ in memory precision rather than capacity. That is, these two groups might be able to remember an equal number of items (i.e., comparable capacity) but PSZ generally process and maintain the items in a less

precise manner. Yet, few studies have attempted to directly estimate memory

capacity and precision at the same time in schizophrenic patients (25, 26). Using the

same delayed-estimation task and the same computational model, Gold et al (25)

found decreased capacity and intact precision in PSZ but Xie&Zhang (26) obtained

the opposite conclusion. Those studies, however, employed a phenomenological

model that categorizes response errors into a mixture of two distributions, and did

not consider the trial-by-trial sensory uncertainty and encoding-decoding

mechanisms underlying VWM, thereby offering limited insight into the mechanistic

differences between PSZ and HCS.

Disentangling the effects of different VWM components (e.g., capacity,

precision) on performance requires generative computational models that explicitly

describe how these components give rise to a behavioral response. van den Berg *et*

al. (37) proposed a variable precision (VP) model (Fig. 1), suggesting that memory

resources assigned to individual items are not only continuous but also variable

across items and trials. Variable resources cause variable processing precision and

thus affect the trial-by-trial response error. Furthermore, the VP model explicitly

distinguishes the precision of encoding items from the precision of exerting a

behavior choice (e.g., motor or decision noise), an aspect that most previous models

ignored. Tested on a large-scale benchmark database of VWM, the VP model has

been shown to outperform other conventional models (38).

Figure 1. Variable precision model of VWM. *A*. Resource decay

function. The VP model assumes that the mean resource ($\bar{J}$) for

processing a single item declines as a power function of set size *N*, a

trend characterized by two free parameters–initial resources ($\bar{J}_1$)

and decaying exponent (*a*). *B*. The resources across items or trials

follow a gamma distribution with the mean resource ($\bar{J}$) determined by the resource decay function (panel A) and the resource allocation variability (τ). Larger amounts of resources (J) indicate higher precision and therefore generate narrower von Mises distributions (three small axes indicating the precision equals to 5, 10 and 15 respectively) of stimulus measurement (m). The widths of the von Mises distributions indicate the degree of trial-by-trial sensory uncertainty. *C*. The eventual behavioral choice given the internal stimulus measurement (m) is also uncertain, following a von Mises distribution with the choice variability (κ_r) (39). In the VP model, initial resources ($\bar{J}_1$), decaying exponent (a), resource allocation variability (τ) and choice variability (κ_r) are four free parameters to estimate (see details in SI and van den Berg *et al*. (37)). All numbers here are only for illustration purposes and not quantitatively related to the model fitting results in this paper.

Leveraging recent advances in basic science, we aimed to use the VP model to understand the computational underpinnings of VWM deficits in schizophrenia. We sought to systematically disentangle the impact of memory capacity and resources, as well as other factors (i.e., variability in allocating resources and variability in choice) in PSZ. In this study, the performance of PSZ and demographically matched HCS was measured in a standard VWM delayed-estimation task (Fig. 2). Using a standard task allows us to compare our results to previous ones (25, 40–43). We believe that a well-controlled task and thorough computational modeling will shed a new light on the mechanisms of perceptual deficits associated with schizophrenia.

Fixation	Sample array	Delay	Probe array
300 ms - 500 ms	500 ms	900 ms	Until response

Time

Figure 2. The color delay-estimation task. This figure depicts an
example trial (set size = 3) of the color delay-estimation task.
Subjects are instructed to first memorize the colors of all squares on
the screen and after a 900 ms-delay choose the color of the probed
square (the one in the lower visual field in this example) on a color
wheel. Set size (i.e., number of squares in the sample array) varies
trial by trial. Response error is the difference between the reported
color and the real color of the probe in the standard color space.
167
168

RESULTS

We measured the performance of 60 PSZ and 61 HCS in a standard color delay-estimation task (Fig. 2). In the task, subjects were instructed to first memorize a set of colored squares. After a short delay, they chose the color of the probed square on a color wheel that represents the standard color space. The difference between the reported color and the true color of the target is considered as the response error.

Variable precision model accounts for VWM behavior in both HCS and PSZ

The VP model proposes that memory resources are continuous and the amount of resource assigned to individual items varies across items and trials. Previous studies have suggested the VP model, compared with other conventional models, as the best account of VWM so far in normal adults (37, 38). However, PSZ might use a qualitatively different observer model other than the VP model to perform the task. We therefore first compared the VP model against the other five computational models that summarize major existing theories of VWM (see model details in Supplementary Appendix note 1). The first one is the item-limit (IL) model. The IL model assumes no uncertainty in the sensory encoding stage and that each subject has a fixed memory capacity and a fixed response variability across set size levels (44). The second one is the mixture (MIX) model, similar to the IL model but assuming response variability is set-size dependent. Note that the MIX model has been used to study schizophrenia (25, 26), thus it enables us to directly compare our results with previous studies. Also, the MIX model is originally called "slots-plus-averaging model" in (45) but we denote it as the MIX model here to differentiate it from our third model described as follows. Compared with the MIX model, our slots-plus-averaging (SA) model (45) further elaborates the idea that memory resources manifest as discrete chunks and these chunks can be flexibly assigned to multiple items. The fourth one is the equal-precision (EP) model, which is similar to the VP model but assumes that the memory resources are equally distributed across items (46, 47). Note that the VP model does not include the capacity component thus one might argue that we cannot draw any direct conclusion about memory capacity by merely analyzing the VP model. To circumvent this, we also

constructed a variable-precision-with-capacity (VPcap) model that not only acknowledges the variable precision mechanisms and but also explicitly estimates the capacity of individual subjects.

Figure 3. Model comparison results. The VP model is tested against other five computational models. The pie charts illustrate the proportion of subjects for whom each model is their best-fitting model. The VP model is the best-fitting model for over 85% of subjects in both groups and under both AIC and BIC, indicating both groups share a qualitatively similar internal process of VWM.

We compared all six models using the Akaike information criterion (AIC) and the Bayesian information criterion (BIC) (48, 49). It turned out that, among all models, the VP model was the best-fitting model for over 85% of subjects in the HCS group under both metrics, replicating previous results in normal subjects (37, 38). Most importantly, the VP model was also the best-fitting model for over 87% of subjects in the PSZ group, indicating that both groups used a qualitatively same observer model to perform the task.

It is worth highlighting two comparisons here. First, the VP model outperformed the MIX model in both groups. This result directly questions the reliability of the previous study based on the MIX model (25). Second, we found that the VP model was better than the VPcap model, even though the later one

incorporates the additional capacity parameter. This result suggests that adding the
capacity parameter seems unnecessary from the modeling perspective. This result is
also in line with the literature showing that a fixed capacity might not exist in
VWM (50, 51). Although systematically examining the existence of a fixed
capacity is beyond the scope of this paper, this result invites a rethink of whether
memory capacity should be considered as a key factor that limits VWM behavior in
PSZ.

Larger resource allocation variability in PSZ

Analyses above have established that HCS and PSZ employ the qualitatively
same observer model to complete the VWM task. Their behavioral differences thus
can only be attributed to the parameter differences on some components of VWM.
We next compared the two groups' fitted parameters of the VP model. Results
showed that the two groups had comparable resource decay functions (Fig. 4A,
initial resources, $t(119) = 0.689$, $p = 0.492$, $d = 0.125$; decaying exponent, $t(119) =$
1.065, $p = 0.289$, $d = 0.194$), indicating a similar trend of diminished memory
resources as set size increases. PSZ, however, had larger variability in allocating
resources (Fig. 4B, resource allocation variability, $t(119) = 4.03$, $p = 9.87 \times 10^{-5}$, d
$= 0.733$). This suggests that, although the two groups have on average the same
amount of memory resources across different set size levels, PSZ allocate the
resources across items or trials in a more heterogeneous manner, with some items in
some trials receiving considerably larger amounts and vice versa in other cases.
This is theoretically suboptimal with respect to completing the task since the probe
was randomly chosen among all presented items with an equal probability. The
optimal strategy therefore should be to assign an equal amount of resources to every
item and in every trial to tackle the unpredictable target. Furthermore, our VP
model explicitly distinguishes the variability of processing items and the variability
in exerting a behavior choice (e.g., motor or decision noise). We found no
significant group difference in the choice variability (Fig. 4C, $t(119) = 1.7034$, $p =$
0.091, $d = 0.31$), diminishing the possibility that the atypical performance of PSZ
arises from larger variability at the choice stage.

Figure 4. Fitted parameters of the VP model. No significant group differences are noted between two groups in resource decay functions (panel A), and choice variability (panel C). PSZ have larger resource allocation variability than HCS (panel B). The individual resource decay functions are computed by $\bar{J_1} = \bar{J} * N^{-a}$, where N is the set size, $\bar{J_1}$ and a are the estimated initial resources and the decaying exponent of one subject. The solid lines represent the averaged resource decay functions across subjects. The shaded areas in panel A, and all error bars in panel B and C represent ±SEM across subjects. Significance symbol conventions are ***: p < 0.001; n.s.: non-significant.

Suboptimal models reveal no capacity difference between HCS and PSZ

Although the VP model was the most appropriate model for both groups, we further examined other suboptimal models. We believe this is valuable for several reasons. First, the VP model does not have the concept of capacity thus we cannot completely rule out the influence of capacity. One might argue that resource allocation variability and limited capacity might jointly manifest in PSZ thus a hybrid model that aggregates two factors might yield a better explanation. Second, the fitted parameters of all models might be derived from specific model settings or possible idiosyncratic model fitting processes.

We argue that the VPcap model is such a hybrid model that accommodates both the variable precision mechanism and a fixed capacity. However, this model is worse than the VP model. We also evaluated the fitted parameters in the VPcap model since it inherits the variable precision mechanism. The results from the VPcap model largely replicated the results of the VP model. Again, we found a

significantly larger resource allocation variability in PSZ (t(119) = 3.891, p =1.65 x
10^{-4}, d = 0.707, see full statistical results in Supplementary Appendix note 2), This
result suggests that the effect of resource allocation variability is quite robust even
though we altered the model structure.

We further examined the estimated capacity of all subjects in the IL, the
MIX, the SA and the VPcap model. Most importantly, none of the four models
suggest decreased capacity in PSZ (see full stats in Supplementary Appendix note
3). Particularly, no significant group difference in capacity was noted in the MIX
model, which contradicts the results of the previous study that used the same task
and the same MIX model (25).

In sum, we found robustly larger resource allocation variability in PSZ in
both the VP and the VPcap models, strongly supporting its key role in limiting
performance in PSZ. Also, we found no evidence for decreased capacity in PSZ in
all models that include the capacity parameter. These results directly challenge the
widely accepted decreased-capacity account, and highlight the role of resource
allocation variability in VWM deficits of PSZ.

Resource allocation variability predicts the severity of schizophrenic symptoms

We next turned to investigate whether the results from the VP model are
predictive of clinically measured symptoms. A set of correlational analyses was
carried out to link the estimated resource allocation variability from the VP model
to the schizophrenia symptomatology (BPRS, SANS, and SAPS).

We noticed that the estimated resource allocation variability of individual
subjects correlates with their BPRS scores (Fig. 5A, r = 0.259, p = 0.045) and the
SANS scores (Fig. 5B, r = 0.302, p = 0.019) in PSZ. No significant correlation was
noted on the SAPS scores (Fig. 5C, r = -0.121, p = 0.358). These results suggest
that resource allocation variability not only is the key factor from a computational
perspective but also can quantitatively predict the severity of clinically measured
symptoms.

Figure 5. Individual differences in resource allocation variability predict the
scores in symptom assessments. Estimated resource allocation variability
values in the PSZ group significantly correlates with their scores on BPRS
(panel A) and SANS (negative symptoms, panel B) but not on SAPS
(positive symptoms, panel C).

315
316

DISCUSSION

The mechanisms of VWM deficits in schizophrenia have been a matter of debate in the past few years. One widely accepted view proposes decreased capacity as the major cause of the deficits in PSZ. This conclusion, however, is merely based on estimated behavioral performance or suboptimal VWM models. In the present study, we re-examined this conclusion by comparing the performance of PSZ and HCS using the computational models that summarize several major theories of VWM. We first established that the VP model was the best model to characterize performance in both groups, indicating the presence of a qualitative similar internal process in both groups. We then further evaluated different components in the VP model as well as other suboptimal models, with special focuses on memory capacity and the declining trend of mean precision as a function of set size. Surprisingly, we found that PSZ and HCS differ in none of these two diagnostic features of VWM. Most importantly, we showed that PSZ had larger variability in allocating memory resources. Furthermore, individual differences in resource allocation variability predict variation of patients' symptom severity, highlighting the clinical functionality of this factor. Taken together, our results challenge the long-standing decreased-capacity explanation for the VWM deficits in schizophrenia and propose for the first time that resource allocation variability might be the key determinant that limits their performance.

A large body of literature has documented that PSZ perform poorly in various forms of working memory tasks (2, 3, 52, 53). Most studies reached the same conclusion: memory capacity is decreased in schizophrenia. However, a careful examination of the literature reveals that the definition of capacity varies across studies. Many studies directly equated worse performance with decreased capacity without quantitatively demonstrating how capacity modulates performance. For example, memory capacity is usually defined as the number of digits that can be recalled in the longest strand in digit span tasks (12). Also, in N-back tasks capacity is defined as the number of backs corresponding to a certain accuracy level (14–16). Moreover, the calculation of capacity resembles the d-prime metric in change detection tasks (22–24, 44, 54). The majority of these metrics are more like

behavioral thresholds related to capacity rather than the quantitative estimates of
capacity. Although these metrics indeed suggest worse performance in PSZ, we
cannot easily equate the worse performance with decreased capacity given the
presence of other components such as memory resource or choice variability. This
is partly because we lack appropriate generative models for the majority of the tasks
thus it is unclear how these components jointly determine performance. The VP
model can describe the generative process of the delay-estimation task and the
change-detection task (37), and allow us to disassociate the effect of capacity from
other VWM components.

Only a few studies have quantitatively estimated capacity and precision in
schizophrenia but they demonstrated inconsistent findings. Gold et al (25)
employed the same delay-estimation task and estimated individual's capacity and
precision using the MIX model (see introduction above and details in
Supplementary Appendix note 1). The model assumes that response errors arise
from a mixture distribution combining a von Mises distribution whose variance
reflects memory precision, and a uniform distribution accounting for the random
guessing if the set size exceeds capacity. Results obtained from the MIX model
echoed the decreased-capacity theory. The MIX model, however, does not take into
account two important factors. First, the model assumes an equal precision across
items held in memory. Second, the model does not separate the variability for
processing stimuli (i.e., sensory uncertainty) and the variability in exertion of a
choice (i.e., choice uncertainty), an issue most previous VWM models also ignored.
Such distinction is important since it highlights different types of uncertainty in
encoding and decoding stages of VWM. Mathematically, these two types of
uncertainty can be distinguished by manipulating set size since the encoding
variability depends on set size but the choice variability does not. The issues of the
MIX model have been symmetrically addressed in a recent work (55). Most
importantly, we showed in this study that the MIX model is not only conceptually
inappropriate but also quantitatively less accurate than the VP model. In addition,
we did not find any group difference in capacity even examining the fitted
parameters.

Compared with capacity and precision–two diagnostic features of VWM,

resource allocation variability is a new concept. It describes the heterogeneity of

allocating resources across multiple items and trials. It has been recently shown that

such variability might not only manifest in VWM and but also act as a ubiquitous

mechanism when processing multiple objects (56). We speculate that resource

allocation variability reflects the extent of attentional control when the brain

processes multiple objects. Two aspects of available evidence support this argument.

First, it has been shown that attention and WM are two core components of

executive control and tightly linked with each other (57, 58). Second, schizophrenia

is known to have deficits in top-down attentional modulation (53, 57). Particularly,

recent studies discovered the phenomenon of spatial hyperfocusing in schizophrenia

patients (19, 59–61). If schizophrenia patients overly attend one item and ignore

others in the memory encoding stage, unbalanced resource allocation will likely

occur.

What are the neural mechanisms of this resource allocation variability?

Recent neurophysiological studies proposed that the neural representation of a

stimulus may follow a doubly stochastic process (62, 63), which suggests that the

variability in encoding precision is a consequence of trial-to-trial and item-to-item

fluctuations in attentional gain (32, 37, 64). A recent study combined functional

magnetic resonance imaging and the VP model, showing that the superior

intraparietal sulcus (IPS) is the cortical locus that controls the resource allocation

(65). Interestingly, schizophrenic patients have been known to have IPS deficits

(66). Note that besides top-down factors, we cannot rule out the contribution of

bottom-up neural noise in perceptual and cognitive processing (62, 63), as found in

several other mental diseases (33–36).

The current results also reveal links between resource allocation variability

and patients' negative symptoms, but not positive symptoms (Fig. 5). These

findings are consistent with several experimental and meta-analysis studies

claiming dissociable mechanisms underlying the cluster of negative symptoms

versus that of positive symptoms (67–70). More broadly, a growing collection of

evidence suggests that visual perceptual deficits in schizophrenic patients are more

likely to link to negative rather than positive symptom severity (71–75). Negative
symptoms in turn might produce improvised social functioning. Humans depend
heavily on VWM to interact with multiple agents and complete social tasks.
Deficits in distributing processing resources over multiple agents therefore might
cause disadvantages in social cognition.

In conclusion, our study proposes a new explanation that the resource
allocation variability accounts for the atypical VWM performance in schizophrenia.
This view differs from the decreased-capacity theory and provides a new direction
for future studies that attempt to promote diagnosis and rehabilitation for
schizophrenic patients.

MATERIALS AND METHODS

Ethics Statement. All experimental protocols were approved by the institutional review board at the East China Normal University. All research was performed in accordance with relevant guidelines and regulations. Informed written consent was obtained from all participants.

Subjects. 61 HCS and 60 PSZ participated in the study. PSZ were clinically stable inpatients (N = 33) and outpatients (N = 27) who met DSM-IV criteria (76) for schizophrenia. All patients were receiving antipsychotic medication (2 first-generation, 43 second-generation, 15 both). Symptom severity was evaluated by the Brief Psychiatric Rating Scale (BPRS) (77), the Scale for the Assessment of Negative (SANS) and Positive Symptoms (SAPS) (78, 79). HCS were recruited by advertisement. All HCS had no current diagnosis of axis 1 or 2 disorders as well as no family history of psychosis nor substance abuse or dependence. All subjects are right-handed with normal sight and color perception.

The two groups were matched in age (t(119) = 1.58, p = 0.118, d = 0.284), gender (t(119) = 1.20, p = 0.234, d = 0.218) and education level of parents (t(119) = 0.257, p = 0.798, d = 0.047). Inevitably, the PSZ had fewer years of education than the HCS (t(119) = 5.51, p = 2.09×10^{-7}, d = 1.00). The detailed demographic information is summarized in the Table 1.

Table 1. Demographics and clinical information of people with schizophrenia (PSZ) and healthy control subjects (HCS)

	PSZ (N = 60)		HCS (N = 61)	
	Mean	SD	Mean	SD
age	35.67	6.58	33.82	9.90
range	23-48	n/a	21-57	n/a
Female/male	31/29	n/a	29/32	n/a
Inpatient/outpatient	33/27	n/a	n/a	n/a
Subject's education (in years)	12.03	2.24	15.13	3.70
Paternal education (in years) [a]	9.89	2.53	9.76	2.95
Maternal education (in years)	9.62	2.91	9.29	3.63
BPRS	27.25	6.27	n/a	n/a
SAPS	5.77	7.02	n/a	n/a
SANS	24.43	11.45	n/a	n/a

[a] Average of mother's and father's years of education

BPRS: Brief Psychiatric Rating Scale (77); SAPS: Scale for the Assessment of Positive Symptoms (79); SANA: Scale for the Assessment of Negative Symptoms (78).

Stimuli and Task. Subjects sat 50 cm away from an LCD monitor. All stimuli were generated by Matlab 8.1 and Psychtoolbox 3 (80, 81), and then presented on the LCD monitor.

In the color delay-estimation VWM task, each trial began with a fixation cross presented at center-of-gaze for a duration randomly chosen from a sequence of 300, 350, 400, 450 and 500 ms. Subjects shall keep their fixation on the cross throughout the whole experiment. A set of colored squares (set size = 1 or 3) was shown within an invisible area with 4^o radius. The sample array lasted 500 ms. Each square was $1.5^o \times 1.5^o$. Their colors were randomly selected from the 180 colors that are equally distributed along the wheel representing the CIE L*a*b color space. The color wheel was centered at (L = 70, a = 20, b = 38) with a radius of 60 in the color space (40). The sample array then disappeared and was followed by a 900 ms blank

period for memory retention. After the delay, an equal number of outlined squares

were shown at the same location of each sample array item, with one of them

bolded as the probe. At the meantime, a randomly rotated color wheel was shown.

The color wheel was 2.1^{o} thick and centered on the monitor with the inner and the

outer radius as 7.8^{o} and 9.8^{o} respectively. Subjects were asked to choose the

remembered color of the probe by clicking a color on the color wheel using a

computer mouse. Subjects shall choose the color as precisely as possible and

response time was not constrained. Every subject completed 2 blocks for the set

size 1 and 3, respectively. The order of the two blocks was counterbalanced across

subjects. Each block had 80 trials.

REFERENCES

1. Gold JM, Randolph C, Carpenter C (1997) Auditory working memory and Wisconsin Card Sorting Test Performance in Schizophrenia. *Arch Gen Psychiatry* 54:159–165.

2. Lee J, Park S (2005) Working Memory Impairments in Schizophrenia: A Meta-Analysis. *J Abnorm Psychol* 114:599–611.

3. Forbes NF, Carrick LA, McIntosh AM, Lawrie SM (2009) Working memory in schizophrenia: a meta-analysis. *Psychol Med* 39:889–905.

4. Goldman-Rakic PS (1994) Working memory dysfunction in schizophrenia. *J Neuropsychiatry Clin Neurosci* 6:348–357.

5. Mayer RE, Moreno R (1998) A split-attention effect in multimedia learning: Evidence for dual processing systems in working memory. *J Educ Psychol* 90:312–320.

6. Postle BR (2006) Working memory as an emergent property of the mind and brain. *Neuroscience* 139:23–38.

7. Nee DE, et al. (2013) A meta-Analysis of executive components of working memory. *Cereb Cortex* 23:264–282.

8. Luck SJ, Vogel EK (2013) Visual working memory capacity: From psychophysics and neurobiology to individual differences. *Trends Cogn Sci* 17:391–400.

9. Coleman MJ, Krastoshevsky O, Tu X, Mendell NR, Levy DL (2012) The effects of perceptual encoding on the magnitude of object working memory impairment in schizophrenia. *Schizophr Res* 139:60–65.

10. Leonard CJ, et al. (2013) Testing sensory and cognitive explanations of the antisaccade deficit in schizophrenia. *J Abnorm Psychol* 122:1111–1120.

11. Johnson MK, et al. (2013) The relationship between working memory capacity and broad measures of cognitive ability in healthy adults and people with schizophrenia. *Neuropsychology* 27:220–229.

12. Conklin HM (2000) Verbal Working Memory Impairment in Schizophrenia Patients and Their First-Degree Relatives: Evidence From the Digit Span Task. *Am J Psychiatry* 157:275–277.

13. Chey J, Lee J, Kim Y-S, Kwon S-M, Shin Y-M (2002) Spatial working memory span, delayed response and executive function in schizophrenia. *Psychiatry Res* 110:259–271.

14. Callicott J (1998) Functional Magnetic Resonance Imaging Brain Mapping in Psychiatry: Methodological Issues Illustrated in a Study of Working Memory in Schizophrenia. *Neuropsychopharmacology* 18:186–196.

15. Barch DM, Csernansky JG, Conturo T, Snyder AZ (2002) Working and long-term memory deficits in schizophrenia: Is there a common prefrontal mechanism? *J Abnorm Psychol* 111:478–494.

16. Jansma JM, Ramsey NF, Van Der Wee NJA, Kahn RS (2004) Working memory capacity in schizophrenia: A parametric fMRI study. *Schizophr Res* 68:159–171.

17. Park S, Holzman PS (1992) Schizophrenics Show Spatial Working Memory Deficits. *Arch Gen Psychiatry* 49:975–982.

18. Keedy SK, Ebens CL, Keshavan MS, Sweeney JA (2006) Functional
magnetic resonance imaging studies of eye movements in first episode
schizophrenia: Smooth pursuit, visually guided saccades and the oculomotor
delayed response task. *Psychiatry Res* 146:199–211.
19. Sawaki R, et al. (2017) Hyperfocusing of attention on goal-related
information in schizophrenia: Evidence from electrophysiology. *J Abnorm*
Psychol 126:106–116.
20. Gold JM, Wilk CM, McMahon RP, Buchanan RW, Luck SJ (2003) Working
memory for visual features and conjunctions in schizophrenia. *J Abnorm*
Psychol 112:61–71.
21. Lencz T, et al. (2003) Impairments in perceptual competency and
maintenance on a visual delayed match-to-sample test in first-episode
schizophrenia. *Arch Gen Psychiatry* 60:238–243.
22. Erickson MA, et al. (2015) Impaired Working Memory Capacity Is Not
Caused by Failures of Selective Attention in Schizophrenia. *Schizophr Bull*
41:366–373.
23. Erickson MA, et al. (2014) Enhanced vulnerability to distraction does not
account for working memory capacity reduction in people with schizophrenia.
Schizophr Res Cogn 1:149–154.
24. Leonard CJ, et al. (2013) Toward the Neural Mechanisms of Reduced
Working Memory Capacity in Schizophrenia. *Cereb Cortex* 23:1582–1592.
25. Gold JM, et al. (2010) Reduced Capacity but Spared Precision and
Maintenance of Working Memory Representations in Schizophrenia. *Arch*
Gen Psychiatry 67:570–577.
26. Xie W, et al. (2018) Schizotypy is associated with reduced mnemonic
precision in visual working memory. *Schizophr Res* 193:91–97.
27. Starc M, et al. (2017) Schizophrenia is associated with a pattern of spatial
working memory deficits consistent with cortical disinhibition. *Schizophr*
Res 181:107–116.
28. Ma WJ, Husain M, Bays PM (2014) Changing concepts of working memory.
Nat Neurosci 17:347–356.
29. Bialek W (1987) Physical Limits to Sensation and Perception. *Annu Rev*
Biophys Biophys Chem 16:455–478.
30. Rolls ET, Deco G (2010) *The Noisy BrainStochastic Dynamics as a*
Principle of Brain Function (Oxford University Press).
31. Faisal AA, Selen LPJ, Wolpert DM (2008) Noise in the nervous system. *Nat*
Rev Neurosci 9:292–303.
32. Goris RLT, Movshon JA, Simoncelli EP (2014) Partitioning neuronal
variability. *Nat Neurosci* 17:858–865.
33. Dinstein I, et al. (2012) Unreliable Evoked Responses in Autism. *Neuron*
75:981–991.
34. Park WJ, Schauder KB, Zhang R, Bennetto L, Tadin D (2017) High internal
noise and poor external noise filtering characterize perception in autism
spectrum disorder. *Sci Rep* 7:17584.
35. Sperling AJ, Lu Z-L, Manis FR, Seidenberg MS (2005) Deficits in
perceptual noise exclusion in developmental dyslexia. *Nat Neurosci* 8:862–

863.

36. Bubl E, et al. (2015) Elevated Background Noise in Adult Attention Deficit
Hyperactivity Disorder Is Associated with Inattention. *PLoS One*
10:e0118271.

37. van den Berg R, Shin H, Chou W-C, George R, Ma WJ (2012) Variability in
encoding precision accounts for visual short-term memory limitations. *Proc*
Natl Acad Sci 109:8780–8785.

38. van den Berg R, Awh E, Ma WJ (2014) Factorial comparison of working
memory models. *Psychol Rev* 121:124–149.

39. Osborne LC, Lisberger SG, Bialek W (2005) A sensory source for motor
variation. *Nature* 437:412–416.

40. Zhang WW, Luck SJ (2008) Discrete fixed-resolution representations in
visual working memory. *Nature* 453:233–235.

41. Zhang WW, Luck SJ (2009) Sudden Death and Gradual Decay in Visual
Working Memory: Research Report. *Psychol Sci* 20:423–428.

42. Zhang WW, Luck SJ (2011) The number and quality of representations in
working memory. *Psychol Sci* 22:1434–1441.

43. Foster JJ, Bsales EM, Jaffe RJ, Awh E (2017) Alpha-Band Activity Reveals
Spontaneous Representations of Spatial Position in Visual Working Memory.
Curr Biol 27:3216–3223.

44. Pashler H (1988) Familiarity and visual change detection. *Percept*
Psychophys 44:369–378.

45. Zhang WW, Luck SJ (2008) Discrete Fixed-Resolution Representations in
Visual Working Memory Weiwei. *Nature* 453:233–235.

46. Palmer J (1990) Attentional limits on the perception and memory of visual
information. *J Exp Psychol Hum Percept Perform* 16:332–350.

47. Shaw M (1980) Identifying attentional and decision-making components in
information processing. *Attention and Performance*, ed Nickerson R
(Erlbaum, Hillsdale, NJ), pp 227–296.

48. Wit E, van den Heuvel E, Romeijn JW (2012) 'All models are wrong. ': An
introduction to model uncertainty. *Stat Neerl* 66:217–236.

49. Burnham KP, Anderson DR (2002) *Model Selection and Multimodel*
Inference: A Practical Information-Theoretic Approach (Springer-Verlag,
New York, NY). 2nd Ed.

50. Bays PM, Husain M (2008) Dynamic Shifts of Limited Working Memory
Resources in Human Vision. *Science* 321:851–854.

51. Bays PM, Catalao RFG, Husain M (2009) The precision of visual working
memory is set by allocation of a shared resource. *J Vis* 9:7.1-11.

52. Shakow D (1972) The Worcester State Hospital research on schizophrenia
(1927-1946). *J Abnorm Psychol* 80:67–110.

53. Gold JM, Hahn B, Strauss GP, Waltz JA (2009) Turning it upside down:
Areas of preserved cognitive function in schizophrenia. *Neuropsychol Rev*
19:294–311.

54. Cowan N (2001) The magical number 4 in short-term memory: A
reconsideration of mental storage capacity. *Behav Brain Sci* 24:87–114.

55. Ma WJ (2018) Problematic usage of the Zhang and Luck mixture model.

bioRxiv. doi:10.1101/268961.

56. Shen S, Ma WJ (2018) Variable precision in visual perception. *bioRxiv*.
doi:10.1101/153650.

57. Gold JM, et al. (2017) Selective Attention, Working Memory, and Executive
Function as Potential Independent Sources of Cognitive Dysfunction in
Schizophrenia. *Schizophr Bull* 12:1–8.

58. Christophel TB, Iamshchinina P, Yan C, Allefeld C, Haynes JD (2018)
Cortical specialization for attended versus unattended working memory. *Nat*
Neurosci 21:494–496.

59. Fuller RL, et al. (2006) Impaired control of visual attention in schizophrenia.
J Abnorm Psychol 115:266–275.

60. Kreither J, et al. (2017) Electrophysiological Evidence for Hyperfocusing of
Spatial Attention in Schizophrenia. *J Neurosci* 37:3813–3823.

61. Luck SJ, et al. (2014) Hyperfocusing in schizophrenia: Evidence from
interactions between working memory and eye movements. *J Abnorm*
Psychol 123:783–795.

62. Churchland AK, et al. (2011) Variance as a Signature of Neural
Computations during Decision Making. *Neuron* 69:818–831.

63. Churchland MM, et al. (2010) Stimulus onset quenches neural variability: a
widespread cortical phenomenon. *Nat Neurosci* 13:369–378.

64. Nienborg H, Cumming BG (2009) Decision-related activity in sensory
neurons reflects more than a neuron's causal effect. *Nature* 459:89–92.

65. Galeano Weber EM, Peters B, Hahn T, Bledowski C, Fiebach CJ (2016)
Superior Intraparietal Sulcus Controls the Variability of Visual Working
Memory Precision. *J Neurosci* 36:5623–5635.

66. Zhou S-Y, et al. (2007) Parietal lobe volume deficits in schizophrenia
spectrum disorders. *Schizophr Res* 89:35–48.

67. de Gracia Dominguez M, Viechtbauer W, Simons CJP, van Os J,
Krabbendam L (2009) Are Psychotic Psychopathology and Neurocognition
Orthogonal? A Systematic Review of Their Associations. *Psychol Bull*
135:157–171.

68. Cameron AM, et al. (2002) Working memory correlates of three symptom
clusters in schizophrenia. *Psychiatry Res* 110:49–61.

69. Carter C, et al. (1996) Spatial working memory deficits and their relationship
to negative symptoms in unmedicated schizophrenia patients. *Biol Psychiatry*
40:930–932.

70. Park S, Sauter BH, Rentsch M, Hell D (2002) Visual object working memory
function and clinical symptoms in schizophrenia. *Schizophr Res* 59:261–268.

71. Cadenhead KS, et al. (1997) Information processing deficits of schizophrenia
patients: Relationship to clinical ratings, gender and medication status.
Schizophr Res 28:51–62.

72. Butler PD, Javitt DC (2005) Early-stage visual processing deficits in
schizophrenia. *Curr Opin Psychiatry* 18:151–157.

73. Kéri S, Kiss I, Kelemen O, Benedek G, Janka Z (2005) Anomalous visual
experiences, negative symptoms, perceptual organization and the
magnocellular pathway in schizophrenia: A shared construct? *Psychol Med*

35:1445–1455.

74.   Slaghuis WL (2004) Spatio-temporal luminance contrast sensitivity and
visual backward masking in schizophrenia. *Exp brain Res* 156:196–211.

75.   Slaghuis WL, Bishop AM (2001) Luminance flicker sensitivity in positive-
and negative-symptom schizophrenia. *Exp Brain Res* 138:88–99.

76.   American Psychiatric Association (1994) *Diagnostic and statistical manual*
of mental disorders (4th ed.) (Washington, DC).

77.   Overall JE, Gorham DR (1962) The Brief Psychiatric Rating Scale. *Psychol*
Rep 10:799–812.

78.   Andreasen NC (1983) *The Scale for the Assessment of Negative Symptoms*
(SANS) (Iowa City, IA).

79.   Andreasen NC (1984) *The Scale for the Assessment of Positive Symptoms*
(SAPS) (Iowa City, IA).

80.   Brainard DH (1997) The Psychophysics Toolbox. *Spat Vis* 10:433–436.

81.   Pelli DG (1997) The VideoToolbox software for visual psychophysics:
Transforming numbers into movies. *Spat Vis* 10:437–442.

Supplementary Information for

Atypically larger variability of resource allocation accounts for visual working memory deficits in schizophrenia

Yijie Zhao, Xuemei Ran, Li Zhang, Ruyuan Zhang, Yixuan Ku

Ruyuan Zhang
Email: ruyuanzhang@gmail.com

Yixuan Ku
Email: yk1616@nyu.edu

This PDF file includes:

Supplementary Note 1: Computational modeling of VWM

VP model. The variable precision (VP) model has been shown as the state-of-the-art computational model of VWM. Details of the VP model have been documented in several previous studies (1, 2) and the model codes are publicly available (http://www.cns.nyu.edu/malab/resources.html).

The VP model assumes a resource decaying function describing the decreasing trend of mean memory resource ($\overline{J}$) assigned to individual items as the set size (N) increases (3, 4):

$$\overline{J} = \overline{J_1} * N^{-a} , \tag{S1}$$

where $\overline{J_1}$ is the initial resources when only 1 item ($N = 1$) should be memorized and a is the decaying exponent. The key component of the VP model is that the memory resources J across items and trials follow a Gamma distribution with the mean $\overline{J}$ and the scale parameter τ:

$$J \sim Gamma(\overline{J}, \tau) , \tag{S2}$$

Intuitively, a larger τ indicates a more uneven distribution of memory resources across items or trials, with some items in some trials receiving a larger amount of resource while others receive comparative fewer. Note that a larger amount of memory resource produces a higher precision. Thus, we do not explicitly distinguish resource and precision and denote them as J. Defining precision as Fisher information (5), precision J can be linked to the variance of the von Mises distribution of sensory measurement:

$$J = k \frac{I_1(\kappa)}{I_0(\kappa)}, \tag{S3}$$

where I_0 and I_1 are modified Bessel functions of the first kind of order 0 and 1 respectively, with the concentration parameter κ. Eq. S3 specifies a one-on-one mapping between precision J and variance κ. We can rewrite their relationship as:

$$\kappa = \Phi(J) , \tag{S4}$$

where Φ is the mapping function. The distribution of sensory measurement (m) given the input stimulus (s) can be written as:

$$p(m \mid s) = \frac{1}{2\pi I_0(\kappa)} e^{\kappa \cos(m-s)} \equiv VM(m; s, \kappa),$$ (S5)

We further assume that the reported color ($\hat{s}$) by participants also follows a von Mises distribution:

$$p(\hat{s} \mid m) = \frac{1}{2\pi I_0(\kappa_r)} e^{\kappa_r \cos(\hat{s}-m)} \equiv VM(\hat{s}; m, \kappa_r),$$ (S6)

where k_r represents the variability at the choice stage.

Taken together, this VP model has four free parameters: $\overline{J}_1$, a, τ and k_r.

Given the four free parameters and stimulus color s in a trial, we can derive the probability of the observed response in a trial by marginalizing over sensory measurement m and variable precision J:

$$
\begin{aligned}
p(\hat{s} \mid s; \overline{J}, \tau) &= \int p(\hat{s} \mid s; J) p(J \mid \overline{J}; \tau) dJ \\
&= \int VM(\hat{s}; s, \Phi(J)) Gamma(J; \overline{J}, \tau) dJ \\
&= \iint VM(\hat{s}; m, k_r) VM(m; s, \Phi(J)) Gamma(J; \overline{J}, \tau) dJ dm \\
&= \int \frac{I_0(\sqrt{\Phi(J)^2 + k_r^2 + 2\Phi(J)k_r \cos(\hat{s}-s)})}{2\pi I_0(k_r) I_0(\Phi(J))} Gamma(J; \overline{J}, \tau) dJ
\end{aligned}
$$ (S7)

Note that in Eq. S7, sensory measurement (m) can be analytically eliminated. Since precision J is a random variable across items and trials, we sampled it 10000 times from the Gamma distribution with mean $\overline{J}$ and scale parameter τ. Note that (1) confirmed that 500 samples are enough in the model fitting. We then used all the samples to calculate response probability in each trial.

Variable-precision-with-capacity model. The variable-precision-with-capacity (VPcap) model inherits all parameters and the structure of the VP model above, except that an additional capacity parameter (K) is introduced to estimate memory capacity of individuals. If the set size N is smaller than capacity K, the VPcap model is identical to the VP model. If the set size N exceeds the capacity K, the model assumes that the probe is stored in the VWM with the probability K/N, and out of memory with the probability 1- K/N. In the latter case, a participant randomly guesses a color. The response probability therefore can be written as:

$$p(\hat{s}\,|\,s) = \begin{cases} \dfrac{K}{N}p(\hat{s}\,|\,s;\overline{J},\tau) + (1-\dfrac{K}{N})\dfrac{1}{2\pi}, & K <= N \\[2ex] p(\hat{s}\,|\,s;\overline{J},\tau), & K > N \end{cases}, \qquad (S8)$$

Where $p(\hat{s}\,|\,s;\overline{J},\tau)$ is defined in Eq. S7. In essence, the VPcap model is a mixture model of the VP model and a random guessing process when the set size exceeds the participant's capacity. The VPcap model has five parameters, four as the same in the VP model and the additional capacity parameter (K).

Item-limit model. The item-limit (IL) model assumes no uncertainty in the sensory encoding stage such that the internal sensory measurement m is equal to the input stimulus s. But there exists choice variability from measurement m to the reported color ($\hat{s}$). Such choice variability does not vary across set size levels. The IL model also assumes a fixed capacity K. The response probability is:

$$p(\hat{s}\,|\,s) \equiv p(\hat{s}\,|\,m) = \begin{cases} \dfrac{K}{N}VM(\hat{s}\,|\,s,\kappa_r) + (1-\dfrac{K}{N})\dfrac{1}{2\pi}, & K <= N \\[2ex] VM(\hat{s}\,|\,s,\kappa_r), & K > N \end{cases}, \qquad (S9)$$

The IL model has two free parameters: choice variability (κ_r) and capacity (K).

Slots-plus-averaging model. The slots-plus-averaging (SA) model was originally proposed in ref. (6) and further elaborated in ref. (1). Unlike the IL model, the SA model acknowledges the presence of noise in sensory encoding stage, however the memory resource is discrete chunks and a single or multiple chunks can be assigned to one item. For one item, the SA model assumes the Eq. S4 still holds as the relationship between the resource assigned to that item and the width of the Von Mises distribution:

$$\kappa = \Phi(SJ_s), \qquad (S10)$$

where S is the number of chunks and J_s is the resource of one chunk. The SA model also assumes a capacity K. When $N > K$, an item should receive either 0 or 1 chunk. Then the allocation should be similar to the IL model. When $N \leq K$, some items receive either one

or more chunks. Assuming that the resource chunks should be assigned as equally as possible across items, the S can be calculated as:

$$S = \begin{cases} \left\lfloor \dfrac{K}{N} \right\rfloor, & \text{with probability } 1 - \dfrac{K \bmod N}{N} \\[4mm] \left\lfloor \dfrac{K}{N} \right\rfloor + 1, & \text{with probability } \dfrac{K \bmod N}{N} \end{cases}, \qquad \text{(S11)}$$

where $\lfloor x \rfloor$ represents the *floor* function in Matlab. The response probability in the SA model can be written as:

$$p(\hat{s}\,|\,s) = \begin{cases} \dfrac{K}{N} VM(\hat{s}\,|\,s, \Phi(J_s)) + (1 - \dfrac{K}{N}) \dfrac{1}{2\pi}, & K < N \\[4mm] (1 - \dfrac{K \bmod N}{N}) VM(\hat{s}\,|\,s, \Phi(\left\lfloor \dfrac{K}{N} \right\rfloor J_s)) + \dfrac{K \bmod N}{N} VM(\hat{s}\,|\,s, \Phi(\left\lfloor \dfrac{K}{N} \right\rfloor + 1\, J_s)), & K >= N \end{cases}, \text{(S12)}$$

The SA model has two free parameters: unit resource J_s and capacity K.

Mixture model. The mixture model (MIX) has been used in previous clinical research (7). Similar to the IL model, the MIX model only assumes the uncertainty from stimulus s to the reported color ($\hat{s}$) and a fixed capacity K. The difference is that the uncertainty (κ) reflects both sensory noise and choice variability, and thus the uncertainty is set-size dependent (each set size has one κ). The response probability can be written as:

$$p(\hat{s}\,|\,s) = \begin{cases} \dfrac{K}{N} VM(\hat{s}\,|\,s, \kappa_{1/3}) + (1 - \dfrac{K}{N}) \dfrac{1}{2\pi}, & K <= N \\[4mm] VM(\hat{s}\,|\,s, \kappa_{1/3}), & K > N \end{cases}, \qquad \text{(S13)}$$

where κ_1 and κ_3 denote the uncertainty for set size 1 and 3, respectively. The MIX model has three parameters: uncertainty levels κ_1, κ_3 and capacity K.

Equal-precision model. The equal precision (EP) model is very similar to the VP model, except that an equal amount of resources is assigned to every item and in any trial. Namely, the Eq. S2 does not apply to the EP model. In the EP model, the resource assigned to one item declines as a power function (as Eq. S1). Then the resource at each

set size level can be converted to the width of the Von Mises distribution using (Eq. S4). The response probability is given by:

$$p(\hat{s} \mid s; J_1, a, \kappa_r) = \frac{I_0\left(\sqrt{\Phi(J_1 N^{-a})^2 + k_r^2 + 2\Phi(J_1 N^{-a})k_r \cos(\hat{s} - s)}\right)}{2\pi I_0(\kappa_r) I_0(\Phi(J_1 N^{-a}))} \ , \tag{S14}$$

where J_1 is the resource when set size is 1 (initial resources). The EP model has three free parameters: initial resources (J_1), decaying exponent (a), and choice variability (κ_r).

Model fitting. The BADS optimization toolbox in MATLAB (8) was used to search the best-fit parameters that maximize the likelihood of responses. BADS has been shown to outperform other default nonlinear optimization algorithms in MATLAB, especially in the problems where gradients on loss function are not available or hard to compute (8). We fit all models separately to each participant. To avoid local minima, we repeated the optimization process with 20 different initial seeds that are equally spaced within a lower and an upper bound. Parameters bounds were set to be very broad to avoid bias. The parameters with the maximum likelihood value were used as the best-fit parameters for one subject.

Supplementary Note 2: Model comparisons

We compared the performance of all models fitted in this study, including the VP model, the VPcap model, the EP model, the SA model, the MIX model and the IL model. Model comparisons were performed for both groups using both Akaike information criterion (AIC) and Bayesian information criterion (BIC) (9, 10) metrics (Fig. S1). All models were compared for each subject and we derived the best model for each subject. Results showed that the VP model outperformed all other models over 85% of subjects in both groups and with respect to both AIC and BIC (Fig. S2). Particularly, the VP model is the best-fitting model in 52 out of 61 (85%) HCS and in 55 out of 60 PSZ (92%) under the AIC. Using the BIC, the VP model is the best-fitting model in 52 out of 61 HCS (85%) and 52 out of 60 (87%) PSZ. These results strongly support the idea that the VP model assuming no fixed capacity better explains the VWM behavior. This result also questions

the conventional theory whether capacity really acts as a key determinant of limiting VWM performance in PSZ.

Supplementary Note 3: Results of other suboptimal models

Fitted parameters of the VPcap model. The VPcap model is a variant of the VP model and explicitly incorporates the capacity parameter. Estimated parameters in the VPcap model largely replicated the results of the VP model (Fig. S2). Again, PSZ have larger resource allocation variability than HCS (Fig. S2B, t(119) = 3.891, p =1.65 x 10^{-4}, d = 0.707) and the two groups did not significantly differ in the resource decay function (Fig. S2A, initial resources, t(119) = 0.012, p = 0.990, d = 0.002; decaying exponent, t(119) = 1.142, p = 0.256, d = 0.208). We observed a significant larger choice variability in HCS (Fig. S2C, choice variability, t(119) = 2.365, p = 0.02, d = 0.43). Most importantly, the estimated capacity values of two groups were statistically comparable (Fig. S2D, t(119) = 0.459, p = 0.647, d = 0.083).

Comparing capacity of the two groups in suboptimal models. We further investigated the estimated capacity of all subjects in the IL, the SA, the MIX and the VPcap model, the four models having the capacity parameter. We found no significant group difference in capacity measured by all four models (Fig. S3, IL model, t(119) = 1.554, p = 0.123, d = 0.283; SA model, t(119) = 1.03, p = 0.306, d = 0.187; MIX model, t(119) = 0.273, p = 0.786, d = 0.050; VPcap model, t(119) = 0.459, p = 0.647, d = 0.083). These results directly reject the decreased-capacity account of PSZ. Notably, one previous study employed the same task and fit the same MIX model and found smaller capacity of PSZ compared with HCS (7). But we could not replicate this finding here.

Figure S1. Positive log-likelihood (panels A, D), AIC (panels B, E) and BIC (panels C, F) values for all models. Note that here we display the positive log-likelihood values to help visually compare models since maximum negative log-likelihood values are equivalent to minimum positive log-likelihood values. Thus, in all panels a lower y-axis value indicates a better model. The upper (panels A-C) and lower (panels D-F) rows depict the model comparison results for HCS and PSZ respectively. The best-fitting model is the VP model for both groups and using both AIC and BIC metrics (also see the Fig. 3 in the main text).

Figure S2. Fitted parameters (panel A: resource decay functions; panel B: resource allocation variability; panel C: choice variability; panel D: capacity) of the VPcap model. The results replicate the results in the Fig. 4. Furthermore, this model estimates capacity in individual subjects and the result show that the two groups have comparable capacity (panel D). All error bars are ± SEM across subjects. Other figure captions are the same as in the Fig. 4 in the main text. Significance symbol conventions are *:p < 0.05; ***: p < 0.001; n.s.: non-significant.

Figure S3. Capacity of the two groups measured by four suboptimal models. None of the four models reveal the significant group differences in capacity. These results directly challenge the conventional decreased-capacity account of PSZ. All error bars are ± SEM across subjects.

References

1. van den Berg R, Shin H, Chou W-C, George R, Ma WJ (2012) Variability in encoding precision accounts for visual short-term memory limitations. *Proc Natl Acad Sci* 109:8780–8785.

2. van den Berg R, Awh E, Ma WJ (2014) Factorial comparison of working memory models. *Psychol Rev* 121:124–149.

3. Bays PM, Husain M (2008) Dynamic Shifts of Limited Working Memory Resources in Human Vision. *Science* 321:851–854.

4. Bays PM, Catalao RFG, Husain M (2009) The precision of visual working memory is set by allocation of a shared resource. *J Vis* 9:7.1-11.

5. Ma WJ, Beck JM, Latham PE, Pouget A (2006) Bayesian inference with probabilistic population codes. *Nat Neurosci* 9:1432–1438.

6. Zhang WW, Luck SJ (2008) Discrete fixed-resolution representations in visual working memory. *Nature* 453:233–235.

7. Gold JM, et al. (2010) Reduced Capacity but Spared Precision and Maintenance of Working Memory Representations in Schizophrenia. *Arch Gen Psychiatry* 67:570–577.

8. Acerbi L, Ma WJ (2017) Practical Bayesian Optimization for Model Fitting with Bayesian Adaptive Direct Search. *Advances in Neural Information Processing Systems 30*, pp 1836–1846.

9. Wit E, van den Heuvel E, Romeijn JW (2012) 'All models are wrong. ': An introduction to model uncertainty. *Stat Neerl* 66:217–236.

10. Burnham KP, Anderson DR (2002) *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach* (Springer-Verlag, New York, NY). 2nd Ed.