When do shifts in trait dynamics precede population declines?

Gaurav Baruah[1], Christopher F. Clements[1,2], Fréderic Guillaume[1], Arpat Ozgul[1]

[1] Department of Evolutionary Biology and Environmental studies, University of Zurich

[2] School of Biosciences, University of Melbourne

*Corresponding Author: Gaurav Baruah

Email: gaurav.baruah@ieu.uzh.ch

Keywords*: population decline, quantitative trait, early warning signals, demography*

Manuscript type*: Article*

Number of words in main-text*: 5572*

Number of words in Abstract*: 200*

Abstract

Predicting population responses to environmental change is an on-going challenge in ecology. Studies investigating the links between fitness-related phenotypic traits and demography have shown that trait dynamic responses to environmental change can sometimes precede population dynamic responses, and thus, can be used as an early warning signal. However, it is still unknown under which ecological and evolutionary circumstances, shifts in fitness-related traits can precede population responses to environmental perturbation. Here, we take a trait-based demographic approach and investigate both trait and population dynamics in a density-regulated population in response to a gradual change in the environment. We explore the ecological and evolutionary constraints under which shifts in a fitness-related trait precedes a decline in population size. We show both analytically and with experimental data that under medium-to-slow rate of environmental change, shifts in trait value can precede population decline. We further show the positive influence of environmental predictability, average reproductive rate, plasticity, and genetic variation on shifts in trait dynamics preceding potential population declines. These results still hold under non-constant genetic variation and environmental stochasticity. Our study highlights ecological and evolutionary circumstances under which a fitness-related trait can be used as an early warning signal of an impending population decline.

1. **Introduction**

Exogenous pressure can force complex systems with alternative stable states

towards so-called tipping points, the point at which the state system can rapidly and

substantially change in response to a small perturbation (May 1977; Hutchings and

Reynolds 2004; Frank et al. 2011) . Examples of such transitions are documented

from rapid shifts in shrub cover in grasslands (Kéfi et al. 2007) to the collapse of

fisheries (Jackson et al. 2001; Frank et al. 2011), and have been shown to be

experimentally inducible in laboratory systems (Dai et al. 2012; Lei Dai, Korolev,

and Gore 2013). The non-linear nature of such transitions makes them difficult to

predict, but may be possible through the identification of statistical signals

embedded in time series data, typically termed early warning signals (EWS) (Wissel

1984; Dakos et al. 2008; Marten Scheffer et al. 2009). Such statistical signals arise

from a systems behavior prior to a critical transition (van Nes and Scheffer 2007; M.

Scheffer et al. 2012) – whereby it takes longer to return to it's original equilibrium

state after every perturbation. This behavior of the system is known as critical

slowing down (CSD).

CSD behavior is predicted to lead to increasing variance and autocorrelation in the

abundance time-series in the region of a bifurcation. However, shifts in variance or

autocorrelation in the abundance time series might not be the only indicators of

whether a population is nearing a tipping point (Clements and Ozgul 2016). External

environmental change has been shown to also substantially affect phenotypic trait

distributions along with changes in demography (Traill, Schindler, and Coulson

2014; Pigeon et al. 2017). Phenotypic traits, for example body size, are intimately

linked with an individual survival and reproductive success (McNamara & Houston,

2007), vulnerability of a population to extinction and fluctuations in population size

(Olden, Hogan, and Zanden 2007; van Benthem et al. 2017). Changes in body size

have also been shown to influence resilience of food webs to disturbance

(Woodward et al. 2005). Furthermore, recent work has also suggested the inclusion

of body size-based measures of stability, as shifts in body size of diatoms were

detected before a regime shift in a lake ecosystem (Spanbauer et al. 2016). Thus,

tracking shifts in such fitness-related phenotypic trait values might allow us to infer

how stable the population is, which can inform us about an impending population

decline.

Empirical observations from experiments and natural populations have shown that

including information from shifts in phenotypic traits such as body size into

abundance-based stability indicators improves the accuracy of predictions of

population collapse (Clements and Ozgul 2016; Clements et al. 2017), but the

circumstances under which doing so is informative is still unknown. The response of

a population to a change in the environment can be a combination of ecological and

evolutionary responses governed by factors such as genetic variation or adaptive

plasticity (Charmantier et al. 2008; Ozgul et al. 2012). Plastic responses to a change

in the environment can be fast, and such responses have been shown to stabilize

population dynamics (Reed et al. 2010; Charmantier et al. 2008). However, if the

environment keeps on changing, populations might deplete its plastic capacity and

standing genetic variation, causing it to eventually decline. This decline in population

size will also be dependent on the reproductive rate i.e., whether a population is

growing faster or slower (Hutchings et al. 2012; Juan-Jordá et al. 2015). In addition

to this, due to directional change in the environment, selection on the trait will act on

the standing genetic variation of that trait, and higher the genetic variation faster is

the evolutionary response (assuming the phenotype heritable). Such theoretical

expectations raise important practical questions: can a shift in phenotypic trait

dynamics occur before an eventual decline in population size? Under what

circumstances is this possible? If such a shift in trait dynamics occurs, what are the

factors that govern the earlier occurrence of this shift?

To answer these questions, we used a combination of theoretical and experimental

approaches to understand the circumstances under which information from trait

values can be useful to predict a potential population decline. First, we integrated

quantitative genetics and population dynamics in a theoretical approach, and

showed both analytically and numerically whether, and under what circumstances,

shifts in trait dynamics can precede population declines. We then experimentally

test our predictions using microcosm data where replicate protist populations were

forced to collapse under different environmental perturbations (Clements and Ozgul

2016). Finally, we evaluate, through numerical simulations, how genetic variation,

adaptive plasticity, and reproductive rate affect when shifts in trait dynamics

precede decline in population size.

2. **Methods and modeling framework**

2.1 The model

In our model, we consider a closed population that has non-overlapping generations,

and is subjected to density-dependent population regulation. We assume in our

population that all individuals experience the same environment; i.e. there is no

spatial heterogeneity. Fitness of individuals in the population is determined by a

quantitative trait z that is under stabilizing selection. Under these assumptions the

dynamics of the population and the mean value of the trait can be written as (Chevin

and Lande 2010; Gomulkiewicz and Holt 1995)

$$N_{t+1} = N_t \overline{R_t} \qquad\qquad (1)$$

$$\overline{z}_{t+1} = \overline{a}_t + \sigma_a^2 \frac{\partial ln\overline{W}}{\partial \overline{a}} + bu_t \qquad\qquad (2)$$

$\overline{R_t}$ is the average fitness of the population at generation t. $\overline{z}_{t+1}$ is the mean value of

the trait at generation $t+1$, $\overline{a}_t$ is the mean breeding value of the population at

generation t, σ_a^2 is the additive genetic variance, $\frac{\partial ln\overline{W}}{\partial \overline{a}}$ is the gradient of selection on

the mean trait $\overline{z}$, $\overline{W}$ is the mean fitness due to the trait z and bu_t quantifies the

average plastic response of the trait (Table 1).

$\overline{R}$ in equation (1) can be expanded as :

$$\overline{R} = \int R(z)p(z)dz \qquad\qquad (3)$$

where $R(z)$ is the absolute multiplicative fitness of an individual with trait z and $p(z)$

is the distribution of the quantitative trait z which is normal. $R(z)$ can further be

decomposed into:

$$R(z) = R_0^{(1-\frac{f(N_t)}{f(K)})} W(z) \qquad\qquad (4)$$

where, R_0 is the net reproductive rate, $\frac{f(N_t)}{f(K)}$ is the density dependence function

and $W(z)$ is the Gaussian stabilizing fitness function given as $W(z, \theta) = e^{-(z-\theta)^2/2w_z^2}$

with width w_z^2 and optimum phenotype of θ. Hence an individual's reproductive

fitness will thus be determined not only by how far its trait z is from the optimum

phenotype θ, but will also concurrently be dependent on the current density of the

population given by the density-dependence function.

Finally, the response of the primary trait z to the environment is modeled using

linear reaction norms (Gavrilets and Scheiner 1993). The phenotype of an individual

at any generation t in the population is given by

$z_i = a_i + bu + \epsilon$ (5)

where a is the breeding value of the individual i, which is normally distributed with

mean $\overline{a}$ and additive genetic variance of σ_a^2. ϵ is the residual component with mean 0

and variance σ_ϵ^2. The slope b in our model determines how plastic the trait is and is

modeled as a constant value meaning that plasticity in the trait cannot evolve.

Hence, variance of the phenotype z is $\sigma_z^2 = \sigma_a^2 + \sigma_\epsilon^2$ (we assume σ_ϵ^2 to be zero in our

model). The environment in our model determines the optimal phenotypic

value θ for the primary phenotype z and also cues the plastic response. θ is assumed

to be linearly dependent on the environment E that selects for a particular

phenotypic value such that at any time t, $\theta_t = BE_t$. The environmental cue

u quantifies how an individual on average perceives the environment. For example,

snow cover could be one of the environmental cues for ground squirrels to come out

of their hibernation that correlates with resource availability, the environmental

factor (Lane et al. 2012). This means that the cue u_t and the selecting environment

E_t are related (Charmantier et al. 2008; Reed et al. 2010). We model this relation by

making the environmental cue a function of the selecting environment such

that $u_t = f(E_t)$. In the case when u_t is a linear function of the selecting environment

E_t, changes in the selecting environment triggers a change in the cue response. If

u_t is modeled as a constant value and independent of the selecting environment E_t,

then the cue response will be different to the change in the selecting environment. In

this case, individuals in the population will not be able to perceive changes in the

selecting environment.

Finally we can arrive at equation (2), which describes the dynamical changes in the

mean trait due to changes in the mean fitness $\overline{W}$ caused by changes in the selecting

environment E_t. The mean fitness $\overline{W}$ can be calculated by: $\overline{W} = \int W(z)p(z)dz =$

$\sqrt{\dfrac{w_z^2}{\sigma_z^2 + w_z^2}}\; e^{-(\overline{z_t} - \theta_t)^2 / 2(w_z^2 + \sigma_z^2)}$

In our model we specifically address questions linked with the decline phase of the

population and not evolutionary rescue. Evolutionary rescue is a long-term process,

which occurs when genetic evolution rescues a population from extinction in

response to changes in the environment. Whether this rescue happens will depend

on the initial decline phase of the population, as the population might collapse before

it can adapt to changing environmental conditions (Gonzalez et al. 2013). Hence,

predicting this initial decline phase is of foremost importance if one has to mitigate

the demographic response of the population, before evolutionary rescue takes place

at a later phase.

2.1 Analytical framework:

Optimal phenotypic change in our analyses is directional and given by:

$$\theta_t = \begin{cases} B & , t < t_{500} \\ B\,E_t = \; B\,e^{-\left(\frac{t}{500}C+C\right)} & , \; t \geq t_{500} \end{cases} \qquad (6)$$

Where $B = 2$, $E_t = \; e^{-\left(\frac{t}{500}C+C\right)}$, t is the time in generations and C is a parameter that

controls how fast the optimal phenotype changes over time. We vary C over a range

of values to create a gradient of environmental change from fast to slow. Without

loss of generality, we consider that the average trait value is at its optimum, i.e.

$\bar{z} = \; \theta_0 = B$ and the environment shifts at t = 500.

Following an environmental change scenario, shifts in trait and population dynamics

are calculated by $\frac{(\bar{z_0} - \bar{z_t})}{\bar{z_0}}$ and $\frac{(N_0 - N_t)}{N_0}$ respectively, where $\bar{z_0}$ is the initial average trait

value and $\bar{z_t}$ is the average trait value after the optimum phenotype shifts. N_0 is the

abundance at carrying capacity and N_t is the abundance after the environment shifts.

We consider in the analytical cases below without the effect of plasticity, i.e. $b = 0$

(appendix A2 for plasticity affect). Hence, all the individuals in the population differ

only in terms of their breeding value. For analytical simplicity, we show two very

simplistic cases of one-generation change in trait and the population after the

optimum environment shifts. *Case 1*: the environmental change scenario that causes

$\frac{(\bar{z_0} - \bar{z_1})}{\bar{z_0}} > \frac{(N_0 - N_1)}{N_0}$ i.e., shift in trait value before population decline and, *Case 2*: the

environmental change scenario that causes $\frac{(\bar{z_0}-\bar{z_1})}{\bar{z_0}} < \frac{(N_0-N_1)}{N_0}$ i.e., shift in trait value

after a population decline. We consider in the above cases two extreme ends of

environmental change: abrupt, large shift in one generation and slow, small shift in

one generation. In the case of abrupt and large shift in the environment, the

optimum phenotype is allowed to shift in *one generation* by a large magnitude to a

new value of $\theta_1 = 0.5$ from a value of $\theta_0 = 2$ such that, $\delta = (\theta_0 - \theta_1) = (\bar{z} - \theta_1) =$

1.5. Specifically, δ can be termed as the initial phenotypic lag of the mean trait to

optimum phenotype in 1 generation. The level of this lag is dependent on how fast

the optimum phenotype shifts. A shift of $\delta \geq 1.5$, or a lag of 1.5 in just *one generation*

causes a substantial population decline (Fig. 1). Moreover, such a jump in the

optimum phenotype to a new value introduces a novel optimum that is beyond the

distribution of the adapted trait distribution (with $\sigma_a^2 = 0.5$). In the case of slow and

small shift in the environment, the optimum phenotype in *one generation* is allowed

to shift by a very small amount, $\delta \leq 0.2$. This value corresponds to a shift in the

optimum that is not novel and within the realms of the adapted trait distribution.

We discuss the analytical results of case 1 and case 2 in the *Result* section.

2.2 Numerical simulations

We performed deterministic numerical simulations of the model described above.

We also did stochastic numerical simulations of the model (for details see appendix

A3). Dynamics of the trait, population and optimum environment were iteratively

updated using equations (1-6). We used Gompertz density function as the form of

density-dependence in our model simulations (Chevin and Lande 2010; Gompertz

1825). Without loss of generality, we assumed that the mean trait value is at its

optimum $\bar{z} = \theta = B$ at the start of each simulation. Environmental change was

introduced only after the population has reached its carrying capacity which was at t

=500. By varying the parameter C (equation 6), we simulated a range of

environmental change scenarios: from very slow (C=10.5) to fast (C= 120.5). In all of

our simulations of different rates of environmental change, the optimum

environment switched to a new value at t = 500. The magnitude of this switch,

however, depended on C (equation 6). For example, if C=20.5, the optimum took

around ~ 30 generations to switch by a magnitude of 1.5 units and if C=120.5, the

optimum took just ~5 generations to switch. Next, to quantify whether the trait or

population abundance shifted first when the environmental conditions changed, we

calculated the area under the curves (AUC) of both the trait shift $\frac{(\bar{z_0} - \bar{z_t})}{z_0}$ and

population shift $\frac{(N_0 - N_t)}{N_0}$ over 20 generations after the optimum environment

changed. 1-generation change was used only in the two extreme cases to elucidate

the mathematical simplicity behind the environmental change scenario under which

trait could shift before a potential population decline. AUC, on the other hand, was

used to graphically extend the analytical results of 1-generation change to 20-

generation change. AUC quantified both the magnitude and the timing of

standardized shifts. With this, we then calculated the metric, $\Delta shift$ = (AUC of shift in

mean value of the trait – AUC of population abundance shift). We then compared the

size of $\Delta shift$ to the rate of change in the environment (for stochastic change in the

environment see appendix A3). If $\Delta shift$ was negative, for a particular rate of change

in the environment, it indicated that the population shift was larger and hence the

population declined/shifted earlier than a shift in the mean trait value; if *Δshift* was

positive it meant a shift in the trait value occurred before a decline/shift in

population size. This allowed us to quantify under what environmental change

scenarios (from slow to fast) we could expect shift in average trait value to occur

before a potential population decline. We then evaluated how the metric *Δshift,* for

the range of environmental change scenarios, was affected by factors like genetic

variation, strength in plasticity, reproductive rate, and environmental predictability.

In the appendix we show how relaxing the assumptions of constant genetic variation

(appendix A4) and stochastic environmental change (appendix A3) affect *Δshift*.

2.2.1 Rate of environmental change

We varied the parameter C from a value of 10.5 to 120.5 with a step size of 2 to

simulate a gradient of perturbations from slow to fast environmental change. We

then calculated from equation (6) the time in generations the optimum took to shift

by a magnitude of ~1.5 units. This ~1.5 unit of change in the optimum in 5

generations (when, C=120.5) or in more than ~30 generations (values of $C \leq 30$)

was enough to cause a significant population decline (Fig. 1). We then assessed how

Δshift was affected by the rate of environmental change.

2.2.2 Environmental predictability

We simulated two specific scenarios: 1) when the cue was a linear function of the

optimum environment, and hence individuals of the population could perfectly

predict changes in the optimum environment, 2) when the cue was a constant value

and hence individuals had zero predictability of changes in the optimum

environment. We then assessed how these two scenarios of environmental

predictability affected *Δshift*.

2.2.3 Genetic variation, strength of adaptive plasticity and average reproductive rate

We ran a range of numerical simulations with different levels of genetic variation

(low to high): $\sigma_a^2 = 0.05, 0.08, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6$; low to high strength of

adaptive plasticity b = 0.05, 0.1, 0.2,0.3,0.4, 0.5, 0.6; and low to high average

reproductive rate R_0 = 1.1, 1.15, 1.2, 1.25, 1.3, 1.35, 1.5. While varying levels of

genetic variation from low to high, we kept adaptive plasticity strength at b = 0.2 and

R_0 at 1.2. While varying levels of *b*, we kept σ_a^2 at 0.5 and R_0 at 1.2. Finally, while

varying R_0, we kept σ_a^2 and *b* at 0.5 and 0.2 respectively. For a range of

environmental change scenarios, we then evaluated the effect of each parameter

value on *Δshift*.

2.2.4 Stochasticity in optimum phenotypic change

All of the simulations and calculations of *Δshift* in the above sections were

deterministic. However, any recorded environmental parameter in nature, always

fluctuates around a mean expectation over time (García Molinos and Donohue

2011). We thus wanted to assess whether adding stochasticity to the changes in the

optimum affected our deterministic simulation results. We redid the numerical

simulations of the above theoretical model but added stochasticity in the optimum

phenotypic change (for stochasticity in the growth rate, see appendix A8). For details

see appendix A3. We then assessed how *Δshift* performed under stochasticity in

optimum phenotypic change and how the abovementioned factors affected *Δshift.*

2.2.5 Changing genetic variance

All of the above simulations were done assuming that genetic variation remained

constant. This is true when a quantitative genetic trait was assumed to be controlled

by infinite number of loci (Falconer and Mackay 1996). Here, we took into account

the decreases in genetic variation that might occur due to directional change of the

optimum phenotype. For an asexually reproducing population, the variance of the

distribution of the breeding values was given by the equation (Bürger 2000):

$\Delta \sigma_a^2 = -\left(\frac{\sigma_a^4}{\sigma_a^2 + w_z^2} \right)$ (7)

We used numerical iteration to solve the change in genetic variation over time using

equation (7). We repeated all the simulations above and varied the levels of

plasticity, reproductive rate, starting genetic variation, and environmental

predictability, and calculated *Δshift* as before (appendix A4).

2.5.6. *EWS*

We evaluated how shifts in EWS compared with shifts in mean trait in response to

changes in the environment. First, we showed that our model exhibits non-

catastrophic transcritical bifurcation (appendix A6). Next, we estimated standard

deviation and autocorrelation at first-lag from abundance data as two main EWS

indicators. Following this, shifts in EWSs and shifts in mean trait value were

compared by calculating a metric called *Δshift$_{ews}$* (appendix A6.4-A7). We evaluated

Δshift_{ews} for different levels of plasticity, genetic variation and reproductive rate.

Furthermore, we modified our model to allow for catastrophic fold bifurcation by

introducing an allee threshold at equation 2 (Dai et al. 2012; Hilker 2010). Similarly,

for this fold-bifurcation model we evaluated *Δshift_{ews}* (see appendix A6-A7 for

details).

2.3 Experimental Data

In addition, we analyzed an experimental data where microcosm populations were

forced to collapse by varying the rate of decline in food availability (Clements &

Ozgul, 2016). Clements & Ozgul (2016) used replicate populations of protozoan

ciliate *Didinium nasutum* that fed on *Paramecium caudatum.* In the experiment, four

different treatments of rates of decline of prey availability were chosen: 1) fast; 2)

medium; 3) slow, as well as a constant prey availability as the control treatment. A

total of 60 replicate populations, 15 per treatment were used for the experiment. In

our study, we used data only from the deteriorating environment treatments (i.e.,

fast, medium and slow decline in prey availability). We analyzed each population's

abundance and mean body size time series independently. We then calculated AUC

and Δ*shift* to qualitatively verify our theoretical simulation results (see appendix A5

for details).

3. **Results**

3.1 Analytical results

Before a shift in the optimum environment occurred, we assume that the population

is perfectly adapted to its optimum phenotype $(\theta - \bar{z}) = 0$. Considering the starting

population size to be at K with no plasticity (b =0), equilibrium population size at

any time t from equation (1), (3) and (4) is

$N = \dfrac{w_z}{\sqrt{w_z^2 + \sigma_z^2}} K$ (appendix A1)

Case #1: *Population declines before a trait shift:*

Let $N_0 = \dfrac{w_z}{\sqrt{w_z^2 + \sigma_z^2}} K$ be the population size when the population is at its equilibrium.

Next, when the environment changes by a large magnitude in one generation such

that $\delta \geq 1.5$, the standardized population shift in one generation is (appendix A1):

$$\frac{N_0 - N_1}{N_0} = \frac{N_0 \;-\; N_0 e^{-\frac{(\delta)^2}{2w_z^2 + 2\sigma_z^2}}}{N_0} = 1 - e^{-\frac{(\delta)^2}{2w_z^2 + 2\sigma_z^2}} > 0 \; ;$$

327   and, $1 - e^{-\frac{(\delta)^2}{2w_z^2 + 2\sigma_z^2}} > \dfrac{(\overline{z_0} - \overline{z_1})}{\overline{z_0}}$

328   Case #2: *Trait shifts before population decline:*

329   When the environment shifts by a small magnitude such that: $\delta \leq 0.2$ in one

330   generation. The standardized population shift becomes (appendix A1):

$$\frac{N_0 \;-\; N_0 e^{-\frac{(\delta)^2}{2w_z^2 + 2\sigma_z^2}}}{N_0} = \frac{N_0 \;-\; N_0}{N_0} \approx 0,$$

because, $e^{-\frac{(\delta)^2}{2w_z^2 + 2\sigma_z^2}} \approx 1$ as δ is very low conditional to the fact that $w_z^2 \gg \sigma_z^2$.

In this case, proportional shift in a trait is greater than zero and hence greater than

the population shift (see appendix A1).

$$\frac{(\overline{z_0}-\overline{z_1})}{\overline{z_0}} = \frac{(\theta_1-\overline{z_0})\,(\frac{w_z\sigma_a^2\gamma}{\sigma_z^2+w_z^2})}{\overline{z_0}} > \frac{N_0-N_t}{N_0} \approx 0$$

Our two extreme analytical cases showed that, if the optimum phenotype shifted by

a large magnitude in just one generation, the decline in population size preceded

shift in mean trait value immediately. However, this was not true in the scenario

when the optimum shifts by a very small magnitude over the course of a single

generation.

For the effect of adaptive plasticity see appendix A2.

3.2 Simulation results

3.2.1 Environmental predictability

Higher predictability of the optimum phenotypic change caused *Δshift* to be positive

when compared with lower predictability (Fig. 2). Low predictability of the optimum

phenotypic change caused *Δshift* to be negative even when the optimum phenotype

shifts by a magnitude of ~1.5 units in more than 15 generations (slow

environmental shift) (Fig. 2).

3.2.2 Genetic variation

Higher genetic variation σ_a^2 caused *Δshift* to be positive but this happened when the

optimum environment shifted by a magnitude of 1.5 units in more than 15

generations (slow environmental shift) (Fig. 3C). If the optimum environment

shifted to a new value by a magnitude of 1.5 units in less than 15 generations (fast

shift), population size always declined before any visible shift in the trait value.

3.2.3 Adaptive plasticity

Values of plasticity ($b > 0.2$) caused *Δshift* to be positive (i.e., trait shifted earlier than

population decline) even when the optimum shifted by a magnitude of 1.5 units in

less than 15 generations (fast shift) (Fig. 3B). However, for lower values of plasticity

(b <0.2), earlier shift in trait value would occur before the population declined only

when the optimum environment shifted slowly (>15 generations) (Fig. 3B).

3.2.4 Reproductive rate

For R_0 of 1.2, trait shift occurred earlier than decline in population size i.e., *Δshift* > 0,

given the optimum phenotype shifted slowly (by a magnitude of 1.5 units in ~15

generations or more). If the optimum phenotype shifted faster than that, average

reproductive rate should have to be higher than 1.2 for *Δshift* > 0 and hence shift in

the trait value would then be informative of a population decline (Fig. 3A).

3.2.5 Stochastic change in the optimum phenotype

Higher σ_a^2, higher *b*, higher environmental predictability, and higher R_0 substantially

caused *Δshift* to be positive for this particular scenario (see appendix A3).

3.2.6 Evolving genetic variation

In comparison to simulations with constant genetic variation, *Δshift* was smaller

(less positive) for simulations where genetic variation evolved over time. In other

words, $\Delta shift_{constant\ variation} > \Delta shift_{evolving\ variation}$ (see appendix A4).

3.2.7 EWS and shift in mean trait

We found that for both transcritical and fold-bifurcation models, shifts in mean trait

could occur significantly before shifts in the two EWS indicators (Appendix A6-A7).

This was particularly evident for medium to slow changes in the environment (Fig

A13-A16).

3.3 Experimental results

Experimental data supported our analytical and simulation results qualitatively.

Decline in prey availability from ~300 individuals of *P. caudatum* to zero happened

over a period of 10, 15 and 20 days indicating three different rates: fast, medium and

slow change in the environment respectively. Fast change in prey availability

resulted in population decline preceding shift in mean body size in 7 out of 9

replicates (Fig. 4A). However, during medium rate of prey decline, shift in average

body size preceded decline in population size in 7 out 9 replicate populations in and

4 out of 7 in slow decline in prey availability, qualitatively proving our analytical as

well as our simulation results correct (Fig. 4A).

4. Discussion

Predicting the fate of populations in response to environmental change is a key

challenge in ecology. Recently developed methods have suggested that tracking

fitness related traits might help achieve this goal (Clements and Ozgul 2016;

Clements et al. 2017). Our analytical and simulation results suggested that when the

optimum environment changes relatively slowly, shifts in average trait value may

occur earlier than shifts in the population abundance. However, this was affected by

the amount of genetic variation, strength in adaptive plasticity, environmental

predictability, and speed of life history.

In our simulations, due to higher adaptive plasticity (b>0.3), shifts in the average

trait value occurred before a decline in population size in response to a fast change

in the optimum environment (i.e., shift of 1.5 magnitude by the optimum

environment in less than 15 generations)(Fig. 3B, Fig. A2). Such a result was

dependent on how predictable the environment was, as plastic response of the trait

was mediated by how well the cue was related to a change in the optimum

environment. If the cue was a function of the selecting environment, very high

adaptive plasticity (b = 0.5) would then lead to a shift in average trait value before a

decline in population size was observed even during a significant fast shift in the

optimum phenotype (Fig. 3B, *b=0.5*).

While the positive influence of higher genetic variation on population persistence

(Willi and Hoffmann 2009) and evolutionary rescue (Hufbauer et al. 2015;

Gomulkiewicz, Krone, and Remien 2017) is relatively well studied, little is known

about its transient effect on shifts in average trait value in response to changes in

the optimum environment. Our result suggest that in response to a relatively slow

directional change in the environment, high and constant genetic variation in a

fitness-related trait will promote faster evolution in the average trait value and

consequently will cause a faster shift in the average trait value before a decline in

population size (Fig. 3C, Fig. A3). In addition, additive genetic variance is also

expected to decrease over time due to directional selection acting on the trait

(Barton and Keightley 2002). Here, in addition to the simulations with constant

genetic variation, we carried out a set of simulations where we relaxed the

assumption of constant genetic variation, and this slowed evolutionary change due

to the depletion of genetic variation caused by directional selection (Fig. A6). For this

reason, the magnitude of shift in average trait value was smaller and slower when

compared with another trait shift under the assumption of constant genetic variation

for the same scenarios of optimum environmental change (see appendix Fig. A6, A7).

Besides directional selection, a decline in population size can also lead to a decrease

in genetic variance (Ashander, Chevin, and Baskett 2016). Irrespective of the cause,

low genetic variance will impede both evolutionary rescue (Hufbauer et al. 2015;

Gomulkiewicz, Krone, and Remien 2017) as well as the predictability of population

decline with the help of trait information.

Deterministic and stochastic modeling of population persistence in response to a

changing environment had earlier revealed the positive affect of reproductive rate

on influencing adaptation (Yvonne and Hoffmann 2009). Earlier studies have also

reported that larger species tend to decline in population size more rapidly than

smaller species (Olden, Hogan, and Zanden 2007; Collette et al. 2011) due to

differences in life-history strategies and intrinsic growth rates between large and

small organisms (Hutchings et al. 2012). Populations of fish with slow life histories

(in the family Scombridae) declined faster than those which had comparatively

faster speed of life history (Juan-Jordá et al. 2015). Our modeling results also

reiterated that during a fast shift in the environment (shift of the optimum

environment by 1.5 units in <15 generations), slow growing populations ($R_0 < 1.2$)

would decline before a shift in the average trait value could be observed (Fig. 3A).

When we compared two populations with different reproductive rates, the

magnitude of *Δshift* was found to be substantially larger for populations with higher

R_0 for a same rate of environmental shift (Fig. 3A, Fig. A9), due to the rapid declines

of the populations with lower R_0 .

Our results also suggested that environmental predictability (the correlation

between the cue and the optimum environment) was a key factor in determining the

earlier occurrence of a shift in the average trait value in response to a change in the

optimum environment. Such environmental predictability acted as an interactive

factor, determining the speed and magnitude of shifts in trait value, which was

driven particularly by the strength in plasticity. Earlier studies had also indicated the

positive interactive effects of environmental predictability and adaptive plasticity on

population dynamics (Reed et al. 2010; Ashander, Chevin, and Baskett 2016) .

Introducing stochasticity in the change in the optimum phenotype or in the growth

rate (appendix A8) did not change our above results substantially (Fig. A4). In the

case of stochastic change in the optimum phenotype, environmental predictability

was particularly essential as the plastic response of the trait tracked the changes in

the optimum which led to an earlier trait shift before the population declined

(Fig.4B, Fig. A4).

EWS are shown to exist in models showing both non-catastrophic and catastrophic

transitions (Kéfi et al. 2013). In relation to this, our results suggest that regardless of

whether our model exhibited non-catastrophic transcritical or catastrophic fold

bifurcation, shifts in mean trait value could occur before shifts in EWS. This was

particularly evident for medium to slow change in the optimum environment

(FigA10-A16). Such a shift in mean trait value occurring before EWS, was however

slightly sensitive to variation in plasticity, genetic variation and net reproductive

rate. Nevertheless, shift in mean trait value in conjunction with shift in EWS could be

used as an indicator of imminent population declines (Clements and Ozgul 2016).

Our analytical and simulation results were qualitatively supported by experimental

data (Fig. 4A). In both medium and slow decline in prey availability treatments,

shifts in body size occurred before decline in population size. Shifts in body size in

response to decline in prey availability that were seen in the experimental data could

mostly be attributed to plasticity in body size, as the experimental population was

clonal. During fast decline in prey availability, the plastic response of body size over

time was not large enough to keep up with the pace of decline in prey availability

and hence could not stabilize the loss in fitness. This led to rapid decline in

population size before a significant body size shift. However, during medium decline

in prey availability, plastic shift in body size was able to track the decline in prey

availability in consequence of which a positive growth rate was maintained.

However, since the decline in prey availability continued, the plastic capacity of body

size in the population was depleted causing the population to eventually decline,

which was later than shift in body size was observed. In case of slow decline in prey

availability, the change was very slow that led to small plastic shifts in body size

before a decline in abundance was seen. These small shifts in body size were not

large enough in comparison to decline in abundance, which was reflected in some of

the replicates. Shifts in body size thus could be an obvious indicator of

environmental deterioration before a response in the population dynamics could be

observed. Whether a trait could be considered as an additional indicator of how

stressed a population is, would depend not only on the identity of the trait but also

on the kind of environmental forcing.

The results that we presented here were specific to the parameter space, but were

not restricted to any specific model system (see Chevin, Lande, and Mace 2010). In

our model, we made two main assumptions. Firstly, adaptive plasticity in our trait

remained constant and hence could not evolve. Studies have observed adaptive

phenotypic change without being able to attribute it to genetic change (Ghalambor et

al. 2007; Hendry, Farrugia, and Kinnison 2008). Our results, under the assumption of

constant adaptive plasticity, still held since we explicitly dealt with transient

dynamics after the optimum environment shifted. Secondly, we assumed linear

reaction norms. Nonlinear reaction norms are modeled for secondary traits that are

components of fitness, but such nonlinear reaction norms can evolve to linear ones

in the long term (Gavrilets and Scheiner 1993).

In conclusion, we show that shifts in average trait value could precede shifts in EWS

and population declines in response to a change in the optimum environment, and

higher levels of genetic variation, adaptive plasticity, environmental predictability,

and reproductive rate strengthened such an earlier shift in the average trait value.

Using an experimental data we also showed that shifts in average body size could

precede declines in population size, and hence could be indicative of a future

population decline. Such a shift in mean body size preceding a decline in population

size was possible if the change in the optimum environment was not fast relative to

the generation time of the organism. Thus, shifts in traits may be useful for

predicting population collapses in species where life histories are fast, the rate of

change of the environment is relatively slow, and the environmental predictability is

relatively high, giving hope that methods can be developed that for these signals in

real world populations.

514 **5. Acknowledgements**

515   This research was supported by the European Research Council Grant #337785 and

516   the Swiss National Science Foundation Grant #31003A_146445 to xx.

517 **6. References**

Ashander, Jaime, Luis-Miguel Chevin, and Marissa L. Baskett. 2016. 'Predicting
Evolutionary Rescue via Evolving Plasticity in Stochastic Environments'.
Proceedings of the Royal Society B: Biological Sciences.
doi:10.1098/rspb.2016.1690.

Barton, N H, and P D Keightley. 2002. 'Understanding Quantitative Genetic Variation.'
Nature Reviews. Genetics 3 (1): 11–21. doi:10.1038/nrg700.

Bürger, R. (Reinhard). 2000. *The Mathematical Theory of Selection, Recombination,*
and Mutation. Wiley.

Charmantier, Anne, Robin H. McCleery, Lionel R. Cole, Chris Perrins, Loeske E. B.
Kruuk, and Ben C. Sheldon. 2008. 'Adaptive Phenotypic Plasticity in Response to
Climate Change in a Wild Bird Population.' *Science (New York, N.Y.)* 320 (5877):
800–803. doi:10.1126/science.1157174.

Chevin, and Lande. 2010. 'When Do Adaptive Plasticity and Genetic Evolution
Prevent Extinction of a Density-Regulated Population?' *Evolution* 64 (4): 1143–
50. doi:10.1111/j.1558-5646.2009.00875.x.

Chevin, Luis Miguel, Russel Lande, and Georgina M. Mace. 2010. 'Adaptation,
Plasticity and Extintion in a Changing Enviroment: Towards a Predictive
Theory'. Edited by Joel G Kingsolver. *PLoS Biology* 8 (4). Public Library of
Science: 1–8.

Clements, Julia L. Blanchard, Kirsty L. Nash, Mark A. Hindell, and Arpat Ozgul. 2017.
'Body Size Shifts and Early Warning Signals Precede the Historic Collapse of
Whale Stocks'. *Nature Ecology & Evolution*. doi:10.1038/s41559-017-0188.

Clements, and Arpat Ozgul. 2016. 'Including Trait-Based Early Warning Signals Helps
Predict Population Collapse'. *Nature Communications* 7: 10984.
http://www.nature.com/doifinder/10.1038/ncomms10984.

Collette, B B, K E Carpenter, B A Polidoro, M J Juan-Jordá, A Boustany, D J Die, C Elfes,
et al. 2011. 'High Value and Long Life - Double Jeopardy for Tunas and
Billfishes'. *Science*. doi:10.1126/science.1208730.

Dai, L., D. Vorselen, K. S. Korolev, and J. Gore. 2012. 'Generic Indicators for Loss of
Resilience Before a Tipping Point Leading to Population Collapse'. *Science* 336
(6085): 1175–77. doi:10.1126/science.1219805.

Dai, Lei, Kirill S. Korolev, and Jeff Gore. 2013. 'Slower Recovery in Space before
Collapse of Connected Populations'. *Nature*. doi:10.1038/nature12071.

Dakos, Vasilis, Marten Scheffer, Egbert H van Nes, Victor Brovkin, Vladimir
Petoukhov, and Hermann Held. 2008. 'Slowing down as an Early Warning Signal
for Abrupt Climate Change.' *Proceedings of the National Academy of Sciences of
the United States of America* 105 (38): 14308–12.
doi:10.1073/pnas.0802430105.

Drake, John M, and Blaine D Griffen. 2010. 'Early Warning Signals of Extinction in
Deteriorating Environments.' *Nature* 467 (7314). Nature Publishing Group:
456–59. doi:10.1038/nature09389.

Falconer, Douglas S, and Trudy F C Mackay. 1996. *Introduction to Quantitative
Genetics (4th Edition)*. *Trends in Genetics*. Vol. 12.
http://www.amazon.com/Introduction-Quantitative-Genetics-Douglas-
Falconer/dp/0582243025.

Frank, Kenneth T, Brian Petrie, Jonathan a D Fisher, and William C Leggett. 2011.
'Transient Dynamics of an Altered Large Marine Ecosystem.' *Nature* 477 (7362):
86–89. http://www.ncbi.nlm.nih.gov/pubmed/21796120.

García Molinos, Jorge, and Ian Donohue. 2011. 'Temporal Variability within
Disturbance Events Regulates Their Effects on Natural Communities'. *Oecologia*.

doi:10.1007/s00442-011-1923-2.

Gavrilets, Sergey, and Samuel M Scheiner. 1993. 'The Genetics of Phenotypic of
Reaction Norm Shape V . Evolution'. *Journal of Evolutionary Biology* 6 (1): 31–
48. doi:10.1046/j.1420-9101.1993.6010031.x.

Ghalambor, C. K., J. K. McKay, S. P. Carroll, and D. N. Reznick. 2007. 'Adaptive versus
Non-Adaptive Phenotypic Plasticity and the Potential for Contemporary
Adaptation in New Environments'. *Functional Ecology*. doi:10.1111/j.1365-
2435.2007.01283.x.

Gompertz, B. 1825. 'On the Nature and Function Expressive of the Law of Human
Mortality, and on a New Mode of Determining the Value of Life Contingencies'.
Philosophical Transactions of the Royal Society of London .

Gomulkiewicz, Richard, and Robert D Holt. 1995. 'When Does Evolution by Natural
Selection Prevent Extinction?' *Evolution* 49 (1): 201–7.
http://www.jstor.org/stable/10.2307/2410305%5Cnpapers2://publication/u
uid/AF75FB13-CA45-471F-9595-604002DDAB20.

Gomulkiewicz, Richard, Stephen M. Krone, and Christopher H. Remien. 2017.
'Evolution and the Duration of a Doomed Population'. *Evolutionary Applications*.
doi:10.1111/eva.12467.

Gonzalez, Andrew, Ophélie Ronce, Regis Ferriere, and Michael E Hochberg. 2013.
'Evolutionary Rescue: An Emerging Focus at the Intersection between Ecology
and Evolution.' *Philosophical Transactions of the Royal Society of London. Series
B, Biological Sciences* 368 (1610). The Royal Society: 20120404.
doi:10.1098/rstb.2012.0404.

Hendry, Andrew P., Thomas J. Farrugia, and Michael T. Kinnison. 2008. 'Human
Influences on Rates of Phenotypic Change in Wild Animal Populations'.
Molecular Ecology. doi:10.1111/j.1365-294X.2007.03428.x.

Hilker, Frank M. 2010. 'Population Collapse to Extinction: The Catastrophic
Combination of Parasitism and Allee Effect'. *Journal of Biological Dynamics* 4 (1).
Taylor & Francis : 86–101. doi:10.1080/17513750903026429.

Hufbauer, Ruth A., Marianna Szűcs, Emily Kasyon, Courtney Youngberg, Michael J.
Koontz, Christopher Richards, Ty Tuff, and Brett A. Melbourne. 2015. 'Three
Types of Rescue Can Avert Extinction in a Changing Environment'. *Proceedings
of the National Academy of Sciences*. doi:10.1073/pnas.1504732112.

Hutchings, Jeffrey A., Ransom A. Myers, Veronica B. García, Luis O. Lucifora, and Anna
Kuparinen. 2012. 'Life-History Correlates of Extinction Risk and Recovery
Potential'. *Ecological Applications*. doi:10.1890/11-1313.1.

Hutchings, Jeffrey A., and John D. Reynolds. 2004. 'Marine Fish Population Collapses:
Consequences for Recovery and Extinction Risk'. *BioScience* 54 (4): 297–309.

http://bioscience.oxfordjournals.org/cgi/doi/10.1641/0006-
3568(2004)054[0297:MFPCCF]2.0.CO;2.

Jackson, J B, M X Kirby, W H Berger, K a Bjorndal, L W Botsford, B J Bourque, R H
Bradbury, et al. 2001. 'Historical Overfishing and the Recent Collapse of Coastal
Ecosystems.' *Science (New York, N.Y.)*. doi:10.1126/science.1059199.

Juan-Jordá, M. J., I. Mosqueira, J. Freire, and N. K. Dulvy. 2015. 'Population Declines of
Tuna and Relatives Depend on Their Speed of Life'. *Proceedings of the Royal*
Society B: Biological Sciences 282 (1811): 20150322.
doi:10.1098/rspb.2015.0322.

Kéfi, Sonia, Vasilis Dakos, Marten Scheffer, Egbert H. Van Nes, and Max Rietkerk.
2013. 'Early Warning Signals Also Precede Non-Catastrophic Transitions'. *Oikos*
122 (5). Wiley/Blackwell (10.1111): 641–48. doi:10.1111/j.1600-
0706.2012.20838.x.

Kéfi, Sonia, Max Rietkerk, Concepción L Alados, Yolanda Pueyo, Vasilios P
Papanastasis, Ahmed ElAich, Peter C de Ruiter, et al. 2007. 'Spatial Vegetation
Patterns and Imminent Desertification in Mediterranean Arid Ecosystems'.
Nature. doi:10.1038/nature06111.

Lane, Jeffrey E., Loeske E B Kruuk, Anne Charmantier, Jan O. Murie, and F. Stephen
Dobson. 2012. 'Delayed Phenology and Reduced Fitness Associated with Climate
Change in a Wild Hibernator'. *Nature*. doi:10.1038/nature11335.

May, Robert M. 1977. 'Thresholds and Breakpoints in Ecosystems with a Multiplicity
of Stable States'. *Nature*. doi:10.1038/269471a0.

Olden, Julian D., Zeb S. Hogan, and M. Jake Vander Zanden. 2007. 'Small Fish, Big Fish,
Red Fish, Blue Fish: Size-Biased Extinction Risk of the World's Freshwater and
Marine Fishes'. *Global Ecology and Biogeography*. doi:10.1111/j.1466-
8238.2007.00337.x.

Pigeon, Gabriel, Thomas H.G. Ezard, Marco Festa-Bianchet, David W. Coltman, and
Fanie Pelletier. 2017. 'Fluctuating Effects of Genetic and Plastic Changes in Body
Mass on Population Dynamics in a Large Herbivore'. *Ecology*.
doi:10.1002/ecy.1940.

Reed, Thomas E, Robin S Waples, Daniel E Schindler, Jeffrey J Hard, and Michael T
Kinnison. 2010. 'Phenotypic Plasticity and Population Viability: The Importance
of Environmental Predictability.' *Proceedings. Biological Sciences / The Royal*
Society 277 (1699): 3391–3400. doi:10.1098/rspb.2010.0771.

Scheffer, M., S. R. Carpenter, T. M. Lenton, J. Bascompte, W. Brock, V. Dakos, J. van de
Koppel, et al. 2012. 'Anticipating Critical Transitions'. *Science* 338 (6105): 344–
48. doi:10.1126/science.1225244.

Scheffer, Marten, Jordi Bascompte, William A. Brock, Victor Brovkin, Stephen R.

Carpenter, Vasilis Dakos, Hermann Held, Egbert H. van Nes, Max Rietkerk, and
George Sugihara. 2009. 'Early-Warning Signals for Critical Transitions.' *Nature*
461 (7260). Nature Publishing Group: 53–59. doi:10.1038/nature08227.

Sousa, Wayne P. 1984. 'The Role of Disturbance in Natural Communities'. *Annual*
Review of Ecology & Systematics.

Spanbauer, Trisha L, Craig R Allen, David G Angeler, Tarsha Eason, Sherilyn C Fritz,
Ahjond S Garmestani, Kirsty L Nash, Jeffery R Stone, Craig A Stow, and Shana M
Sundstrom. 2016. 'Body Size Distributions Signal a Regime Shift in a Lake
Ecosystem'. *Proceedings of the Royal Society B* 283: 20160249.
doi:http://dx.doi.org/10.1098/rspb.2016.0249.

Traill, L. W., S. Schindler, and T. Coulson. 2014. 'Demography, Not Inheritance, Drives
Phenotypic Change in Hunted Bighorn Sheep'. *Proceedings of the National*
Academy of Sciences. doi:10.1073/pnas.1407508111.

van Benthem, Koen J., Hannah Froy, Tim Coulson, Lowell L. Getz, Madan K. Oli, and
Arpat Ozgul. 2017. 'Trait–Demography Relationships Underlying Small Mammal
Population Fluctuations'. *Journal of Animal Ecology*. doi:10.1111/1365-
2656.12627.

van Nes, Egbert H, and Marten Scheffer. 2007. 'Slow Recovery from Perturbations as
a Generic Indicator of a Nearby Catastrophic Shift.' *The American Naturalist* 169
(6): 738–47. doi:10.1086/516845.

Wissel, C. 1984. 'A Universal Law of the Characteristic Return Time near Thresholds'.
Oecologia 65 (1): 101–7. doi:10.1007/BF00384470.

Woodward, Guy, Bo Ebenman, Mark Emmerson, Jose M. Montoya, Jens M. Olesen,
Alfredo Valido, and Philip H. Warren. 2005. 'Body Size in Ecological Networks'.
Trends in Ecology and Evolution 20 (7): 402–9. doi:10.1016/j.tree.2005.04.005.

Yvonne, Willi, and Ary A Hoffmann. 2009. 'Demographic Factors and Genetic
Variation Influence Population Persistence under Environmental Change'.
Journal of Evolutionary Biology.

List of tables and figures:

Parameters /variables	*Description*	*Value*

N_t	Abundance of the population at time t	....
$\bar{z}_t$; $\bar{a}_t$	Mean trait value at time t; mean breeding value at time t	....
σ_z^2; σ_a^2	Variance of the mean trait; additive genetic variance or variance of the mean breeding value	[0.05, 0.08, 0.1, 0.2, 0.3, 0.4,0.5, 0.6] ; [0.05, 0.08, 0.1, 0.2, 0.3, 0.4,0.5, 0.6]
b	Strength in phenotypic plasticity	0.05, 0.1, 0.2,0.3,0.4, 0.5, 0.6
R_0	Net reproductive rate	1.1, 1.15, 1.2, 1.25, 1.3,1.35, 1.5
u_t	Environmental cue at time t	Case 1: -0.0001 Case 2: $-\frac{B}{2} + \frac{B}{2} e^{-\left(\frac{t}{500}C+C\right)}$ where, C=4
θ_t	Optimum trait at time t	....
w_z^2	Width of the Gaussian stabilizing fitness function. A measure of the strength in selection	40
z_i, a_i	Trait value for an individual i ; breeding	....

	value for an individual i	
ϵ	Residual component or unexplained component of a trait z	0
σ_ϵ^2	Variance in that residual component ϵ	0
E_t	External environmental value at time t	....
B	Coefficient that captures the strength in the external environmental change	2
C	Parameter that controls the rate of change in the external environment E_t	Various: 10.5 to 120.5

675   Table 1: List of variables and parameter values used in the model. Note that

676   variables are not given any value, but parameters for which the model simulations

677   are tested are given values.

Figure 1. Population and trait dynamics under two different environmental scenarios. Left:

Population size declines by ~50%, when the optimum environment shifts fast (C=120.5;

blue line), whereas during slow change (C= 10.5; dashed line) population declines but to a

lesser extent. Right: Trait dynamics during fast (blue line; C = 120.5) and slow (dashed line;

C=10.5) environmental change. Data based on deterministic simulations of the theoretical

model. Parameters used: $\sigma_a^2 = 0.5$; $R_0 = 1.2$; $b = 0.2$; $w_z^2 = 40$.

Figure 2. Effect of environmental predictability on the magnitude of *Δshift*, data from

numerical simulations of the theoretical model. *Δshift* denotes how large and earlier the

shift occurs in trait dynamics before a decline in the population. If *Δshift* is +ve, shift in

average trait value precedes decline in population size and vice versa. The X-axis denotes

the time in generations it takes for the optimum to change by a magnitude of 1.5 units, and

hence 10 in the X-axis means the optimum shifts by a magnitude of 1.5 in 10 generations

indicating a fast change and 60 means the optimum takes 60 generations to shift by a

magnitude of 1.5 units indicating a very slow change. High environmental predictability

(blue line) facilitates earlier occurrence of trait shift (*Δshift* being +ve) compared to when

the environmental predictability is very low (dashed line). *Δshift* is always –ve (which

means population declines precede mean trait shifts) when the optimum environment shifts

in less than or equal to 15 generations by 1.5 units. Parameters used: $\sigma_a^2 = 0.5$; $R_0 = 1.2$; $b =$

0.2; $w_z^2 = 40$.

Figure 3. Contour plot showing the effect of net reproductive rate, plasticity and genetic

variation on *Δshift* (contour lines). Shown here are the results from numerical simulations

of the theoretical model. For a trait shift to be informative of a population decline, the *Δshift*

should be always greater than zero (contour line =0). The X-axis denotes the time in

generations it takes for the optimum to shift by a magnitude of 1.5 units.

Figure 4. Ability of shift in body size dynamics to precede population decline (i.e., *Δshift >0)*

under different environmental regimes, for the experimental data from Clements & Ozgul

(2016). Note that *Δshift* < 0 if population decline precedes shift in mean body size and *Δshift*

> 0 if shift in mean body size precedes population decline. (A) *Δshift* <0 for fast decline in

prey availability indicating that population decline preceded shift in mean body size. For

medium and slow decline in prey availability, *Δshift* was largely positive (*Δshift* >0). (B)

Replicate abundance and body size time-series are shown for *Δshift* < 0 (fast decline in prey

availability) and for *Δshift >0* (C and D) (Medium decline and slow decline in prey

availability respectively). The blue dashed line in B, C and D indicates the starting point of

environmental deterioration. The red line indicates *loess* smoothing across replicates.