Genome-Wide Associations with Resistance to Bipolaris Leaf Spot (*Bipolaris oryzae*

(Breda de Haan) Shoemaker) in a Northern Switchgrass Population (*Panicum virgatum*)

Kittikun Songsomboon[1][*], Ryan Crawford[*], Jamie Crawford[*], Julie Hansen[*], Jaime

Cummings[†], Neil Mattson[§], Gary Bergstrom[†], Donald Viands[*]

[*]Section of Plant Breeding and Genetics, School of Integrative Plant Science, Cornell

University, Ithaca, NY 14853, USA

[†]Section of Plant Pathology and Plant-Microbe Biology, School of Integrative Plant

Science, Cornell University, Ithaca, NY 14853, USA

[§]Section of Horticulture, School of Integrative Plant Science, Cornell University, Ith-

aca, NY 14853, USA

[1]Corresponding Author: K. Songsomboon, Email: ks899@cornell.edu Phone: +1 607

379 7332

RUNNING TITLE: Bipolaris leaf spot resistance in switchgrass

KEYWORDS:

Disease resistance, Bipolaris leaf spot, switchgrass, genome-wide association

[1]Corresponding Author: K. Songsomboon, Section of Plant Breeding and Genetics,

School of Integrative Plant Science, Cornell University, Ithaca, NY 14853, USA.

Email: ks899@cornell.edu Phone: +1 607 379 7332

ABSTRACT

Switchgrass (*Panicum virgatum* L.), a northern native perennial grass, suffers from yield reduction from Bipolaris leaf spot caused by *Bipolaris oryzae* (Breda de Haan) Shoemaker. To determine the candidate resistant populations via multiple phenotyping approaches and identify potential resistance genes, single- and multi-trait genome-wide association studies were conducted in the northern switchgrass population. The disease resistance was evaluated from both natural and artificial inoculations. The natural inoculation from the field was evaluated in two locations – NY and PA, and artificial inoculations by leaf detachment and leaf disk assay were conducted and evaluated via both vision and image analysis. The most resistant population based on a combination of leaf detachment via image analysis (DTIA)-leaf disk via image analysis (DSIA)-mean from two locations was 'SW805' population. There were only four reliable GWAS results from various trait combinations in various subsets of genotypes including 1) full set of 479 genotypes with DTIA-DSIA-highest score from NY (MNY), 2) tetraploid with MNY, 3) lowland with DTIA-DSIA-MNY and highest score from PA (MPA), and 4) upland with leaf detachment via vision (DTVI)-MNY showing non-overlapping regions. Overall, the resistance genes were associated with 18 markers on chromosomes 1b, 2a, 2b, 3a, 3b, 5a, 5b, 6a, 7a, 7b, 8a, 9a, and 9b and accumulatively explained phenotypic variance of DSIA by 26.72% at the most and DTIA by 6.32% at the least. Within linkage disequilibrium of 20 kb, the potential resistance genes included genes encoding Myb, cytochrome P450, isocitrate lyase, E3 ubiquitin-protein ligase, mitogen-activated kinases, glutathione S-transferase, Birefringence-like 32, and ABC transporter.

INTRODUCTION

Switchgrass (*Panicum virgatum* L.) is a perennial biomass crop native to North

America. Biomass and other agronomic traits have been the focus on improvement.

Although many diseases have been reported to cause deleterious effects on yield, re-

search on disease resistance, especially breeding for the resistance, is scarce. *Bipo-*

laris oryzae (Breda de Haan) Shoemaker (BO) is one of the major fungi causing Bipo-

laris leaf spot (BLS) in switchgrass that can reduce biomass by 70% (Fajolu, 2012).

The approach for the disease management was to breed for improvement of resistance

to BLS.

The natural distribution of switchgrass is broad latitudes across the USA east

of the Rocky Mountains. By simple morphological differentiation, switchgrass can be

separated into two groups of upland and lowland ecotypes. Upland ecotypes are well

adapted to higher latitudes and higher drought tolerance whereas lowland ecotypes

provide higher yield and require more water (Stroup et al., 2003). The more in-depth

genetic diversity was revealed by Network-Based single nucleotide polymorphism

(SNP) discovery protocol suggesting the differentiation of switchgrass group by isola-

tion-by-ploidy, the migration from south to north, and the incidence of tetraploid up-

land from octaploid upland (Lu et al., 2013). Such a large diversity within switchgrass

populations can provide a source of resistance.

Since BO is a necrotrophic pathogen, the interaction between switchgrass and

the fungus was modelled as an inverse gene-for-gene model (Friesen et al., 2007). In

general, the pathogen secretes necrotrophic effectors as host-selective toxins (HST)

that interact with host sensitivity genes, resulting in a compatible susceptible

interaction to trigger host cell death (Friesen et al., 2008). Such an interaction is

known as effector-triggered susceptibility (ETS). Interestingly, according to compara-

tive genome study (Condon et al., 2013), BO does not produce any HST. Regardless

of HST, BO produces ophiobolin A and B as a non-host-selective toxin (Xue et al.,

2015) triggering many groups of proteins and enzymes such as reactive oxygen spe-

cies (ROS) detoxification, protein phosphorylation, and ethylene production (Kim et

al., 2014; Bockhaven et al., 2015).

Such an ETS interaction naturally removes the individuals with the necrot-

rophic effectors from the population. Breeding for the improvement of resistance to

BLS can be done by eliminating susceptible genes from the population. However, the

recurrent phenotypic selection for the resistance in 'Shelter' and 'Cave-in-Rock' for

two cycles of selection did not provide any improvement (Songsomboon, 2018).

Breeding for the resistance by various approaches, stages, and populations was sug-

gested. Therefore, identification of resistant populations is a crucial step.

To accelerate breeding for disease resistance, genomics-assisted breeding is an

important approach. Basically, it begins with gene identification, isolation, cloning,

functional characterization, validation and utilization. There are two approaches to

identify resistance genes: linkage mapping and genome-wide association studies

(GWAS). Although GWAS cannot confirm the causal polymorphism, GWAS depends

on linkage disequilibrium (LD) between the markers and the causal polymorphisms.

With high allelic diversity and ancestral recombination events in the diversity panel,

GWAS yields a finer resolution than linkage mapping (Yu and Buckler, 2006). The

technique has been used to dissect flowering time in switchgrass (Grabowski et al.,

2017). Although GWAS has never been used to dissect disease resistance in

switchgrass, it has successfully dissected resistance to leaf rust caused by *Puccinia*

triticina Eriks., tan spot caused by *Pyrenophora tritici-repentis* (Die.) Shoemaker and

stripe rust caused by *Puccinia striiformis* West. in wheat (*Triticum aestivum* L.)

(Juliana et al., 2018). Despite no resistance genes identified in switchgrass, there were

13 SNP markers from quantitative trait loci (QTLs) linked with the resistance in rice

(Sato et al., 2008, 2015). This study will provide the first dissection of the resistance to

BLS in switchgrass.

The objectives were 1) to determine genotypes or populations from the

northern switchgrass association panel that can be candidates for resistance to BLS, 2)

to conduct a single- and multi-trait GWAS for resistance to BLS in the association

panel via both natural and artificial inoculation, and 3) to explore the genes linked to

the significant markers from the reliable GWAS model to determine potential

candidate genes for resistance to BLS.

111   **MATERIALS AND METHODS**

Switchgrass in the Northern Association Panel

The switchgrass population used in this study has been developed for acceler-

ating breeding progress especially for bioenergy traits at northern latitude (Lu et al.,

2013; Lipka et al., 2014). The population consists of 479 genotypes from 66 popula-

tions representing the mostly upland northern population and some southern lowland

population. It was initially planted in Ithaca, NY, in 2008 and then vegetatively cloned

and planted in a randomized, complete block design with three replicates in 0.9 m

spaced planting in Ithaca, NY, and Philipsburg, PA, in 2016. No fungicide was applied

to the field.

Disease evaluation and phenotype processing

Bipolaris leaf spot has been evaluated in both fields to determine the resistance

in natural incidence and laboratory to minimize environmental effects on the re-

sistance. Field evaluation was conducted in the same week in both locations in August

2017. Each plant was visually evaluated, the severity ranging from 0 to 5 as 0 = 0%

BLS, 1= 1-10% BLS, 2= 11-25% BLS, 3= 26-50% 134 BLS, 4= 51-75% BLS, and 5=

76-100% of the leaf area with BLS. With artificial inoculation in the laboratory, dis-

ease severity was evaluated by leaf detachment and leaf disk assay by taking leaf sam-

ples from one replicate of genotypes in Ithaca, NY. In the leaf detachment assay, three

healthy leaves were randomly selected from 30 centimeters below the top of the shoot

to control the same stage of leaf development. The leaves were cut into five centime-

ters, washed with deionized water, placed in a pre-wet petri dish bedded with filter pa-

per, kept cold in a cooler, and refrigerated overnight.

The inoculum was prepared from a subculture of a single conidium isolated

from a 'Carthage' switchgrass leaf in the warm season biofuels field experiment in Ith-

aca, NY, by Katie Waxman. The day after collecting the detached leaf, the three-

week-old plate was flooded, filtered via gauze, and the concentration adjusted to 10^5

conidia·ml^{-1} by hemocytometer with 2 drops of Tween-20 per 100 ml. An airbrush

pressuring at 10 psi was used to spray inoculum on each dry-surfaced adaxial side of

the leaf in each plate. After letting the droplets of inoculum dry and firmly attach to

the leaf surface, the plates were then sealed with a paraffin tape to maintain high mois-

ture condition. The plates were kept at room temperature with 24-hour light for seven

days (Figure 1). The disease evaluation was conducted by vision and image analysis

software. Both evaluations were based on percentage of leaf covered with lesions. The

evaluation by vision was from 0 to 100 % with 10% increment. Then each leaf was

dried and scanned with a flatbed scanner (Canon CanoScan LiDE 700F) at a resolution

of 1,200 dpi, and images were saved in .tiff format. Due to the limitation of laboratory

space for the plates and labor in leaf collection, 85 to 200 genotypes were sampled in

each set of the experiment, with some overlapping genotypes in one to three dupli-

cated plates and additional non-inoculated plates. The leaf samplings were done

weekly from May to June 2017. A total of seven sets of the experiment eventually

covered all of 482 genotypes. Such an unbalanced design was handled with best linear

unbiased prediction (BLUP) to extract random effect of the genotype by fitting the ex-

periment factor as a fixed effect.

In the leaf disk evaluation, all of 479 genotypes were collected on the same day

in August 2017 by cutting the same age leaves from each genotype and keeping them

refrigerated overnight. On the next day, the leaves were washed with deionized water,

surface dried, cut by keeping the midrib in the middle into an 8-mm disk, and placed

on a water agar plate. There were 28 leaf disks in each plate, including four check

disks (control and inoculated disks of resistant 'SW43_09' and susceptible

'SW122_02') and three replicated disks of eight genotypes (Figure 2). The inoculum

was prepared the same way described above. For each leaf, two µL of inoculum was

dropped in the middle of the right side of the leaf. After letting the droplet dry, the

plates were sealed with paraffin tape to keep high moisture and kept under 12-hour

light at room temperature. The disease evaluation was conducted at seven dpi both by

vision and image analysis software. The percent of leaf disk covered with lesions and

necrotic tissue was assessed by a vision from 0 to 100 with ten increments. Also, on

the same day, a photo was taken of each plate with a digital DSLR Cannon Rebel T6

DSLR camera with the resolution of 2644 x 3084 in .JPG.

Images from both leaf detachment and leaf disk assays were analyzed by using

ImageJ macro at the setting provided in the Supplement file. In brief, each detached

leaf or leaf disk was measured for total leaf area and total chlorosis lesion area (yellow

and black area). Percentage of area covered with lesions was computed by dividing the

total necrotic lesion area by total leaf area.

Package 'LME4' (Bates et al., 2015) in R was used to calculate Best Linear

Unbiased Prediction (BLUPs) from various disease evaluations. For the field evalua-

tion, BLUPs were computed based on each location separately and on the two loca-

tions combined using genotypes and replicates as random effects in each location

model. BLUPs for the two locations combined were fitted genotype, replicates, loca-

tions, the interaction between genotypes and location, as random effects. In addition

to generating BLUPs from field evaluation, the highest score, which is most severe

symptoms observed from each location and two locations over replicates, were used

due to successful QTLs of resistance to foliar symptoms caused by potato virus Y in

autotetraploid potato (*Solanum tuberosum* L.) (Silva et al., 2017). For disease evalua-

tions under laboratory condition, since all phenotypes were expressed as percentages

to leaf area, log transformation was performed before computing BLUPs. For the leaf

detachment assay, BLUPs were computed with experiment, genotype, interaction be-

tween genotypes and experiment, plates within the experiment, and leaf replicates of

genotypes as random effects. For the leaf disk assay, BLUPs were calculated with

plates as a fixed effect and genotype, replicates within plates, and genotype within

plates as random effects. Broad-sense heritability was estimated from variances in

each model. Moreover, phenotypic correlations were computed between resistance to

Bipolaris leaf spot to the other 20 morphological and biomass quality traits from Lipka

et al. (2014), such as plant height, anthesis date, acid detergent lignin, minerals, etha-

nol·g^{-1} dry weight, etc.

Therefore, the resistance to BLS was evaluated via three approaches – leaf

detachment, leaf disk, and field evaluation – providing phenotyped traits including

BLUPs of detached leaf percent lesion via vision and image analysis (DTVI and

DTIA), BLUPs of leaf disk percent lesion via vision and image analysis (DSVI and

DSIA), BLUPs of field evaluation from two locations, NY and PA (BTL, BNY, and

BPA), highest score from two locations, NY and PA (MTL, MNY, and MPA).

Genotyping, linkage disequilibrium analysis, and population structure

HAPMAP v.1 set from Evans et al. (2015) provided 1,377,841 single-nucleo-

tide polymorphisms covering 38,654 genic regions. In short, DNA of each genotype

was processed via exome-capture using the Roch-Nimblegen switchgrass exome-cap-

ture probe set (Evans et al., 2014) and DNA sequencing. The sequences were aligned

to the *P. virgatum* genome assembly v.1.1 (*P. virgatum* v.1.1, DOE-JGI, http://phyto-

zome.jgi.doe.gov/) for SNP discovery. In this calling, SNPs were required to be bi-al-

lelic, sequenced in all samples, be monomorphic in at least two samples, and have

more than 5X coverage in more than 95% of the samples. At each SNP, the genotype

dosages ranged from zero to two copies of minor alleles and can be nonintegers by us-

ing EM algorithm (Martin et al., 2010). Naturally, the switchgrass association panel

includes allopolyploid switchgrass: tetraploids (4x), octoploids (8x) and hexaploids

(6x). It is challenging to perform GWAS with polyploid models. In this study, we

modelled them under the assumption of disomic inheritance similar to the study of

GWAS of flowering time in Grabowski et al. (2017). This was because tetraploid

switchgrass was confirmed to show disomic inheritance (Okada et al., 2010). Alt-

hough octaploid switchgrass had four copies of each homologous chromosome, which

was difficult to precisely determine heterozygous genotypes, the disomic segregation

was used for the model with the caution of the increasing standard error of estimates in

GWAS. Moreover, linkage disequilibrium (LD) was computed via '-r2' tag in PLINK

as r^2 between all SNPs within 1 MB and recorded all values (Purcell et al., 2007).

Principle component analysis was used to evaluate the population structure via TAS-

SEL v.5 (Trait Analysis by aSSociation Evolution and Linkage) (Bradbury et al.,

2007)

Genome-wide association studies

Genome-wide Efficient Mixed Model Association (GEMMA) (Zhou and

Stephens, 2012) was used to implement a multivariate linear mixed model for GWAS

of single- and multiple-phenotyping approaches for resistance to BLS. Kinship was

included as a random effect. Also, the first three principal components (PCs) were

included as fixed covariates. Although the results of Bayesian information criterion re-

sulted in no improvement of the GWAS model with PC inclusion (Lipka et al., 2013),

the inclusion of PCs helped to improve the QQ-plot closer to the theoretical QQ-plot

for this resistance to BLS. The GWAS was conducted in five groups including all of

479, lowland, upland, tetraploid (4X), and octaploid (8X) genotypes to determine if

there were any SNPs linked to specific switchgrass population. Minor allele frequency

(MAF) at 0.05 was used to filter the minor alleles. To correct multiple testing, false

discovery rate (FDR) was used for a cut-off value at 0.1. The significant markers then

were used to determine candidate genes linked to them within the range of LD and

search in JBrowse in *P. virgatum* v.1.1 in Phytozome (e.g., if the position of the

marker was 77345950 on chromosome 2a and LD was 20 kb,

"Chr02a:77335950..77355950" was used). In case that the candidate gene was not

fully aligned within that LD range, we considered the gene as a candidate if 80% of its

length was within the LD range. Since some of SNPs' positions could not be aligned

to the *P. virgatum* genome assembly v.1.1, they were grouped as U1-U4 chromosome

and were excluded from determining the candidate genes.

Candidate genes based on resistance in rice

Since *B. oryzae* is one of the major pathogens in rice (*Oryza sativa*), major

resistant QTLs have been identified in recombinant inbred lines (RILs) and double

haploids (DH) by using restriction fragment length polymorphism (RFLP), simple-

sequence repeat (SSR) markers, and sequence- tagged site (STS) (Sato et al., 2008,

2015; Katara et al., 2010). Three major QTLs in Chromosomes 1, 4 and 11 were then

studied further in near-isogenic lines (NILs) and 13 SNP markers linked with the re-

sistance (Sato et al., 2015). The NILs are BC3F5 between indica 'Tadukan' (re-

sistance) and temperate japonica 'Koshihikari' (susceptible). According to GWAS in

rice, linkage disequilibrium (LD) are ~ 100 kb in indica and ~200 kb temperate in ja-

ponica (Zhao et al., 2011). Thus, in this study, the candidate resistance genes were

screened from 200 kb around the 13 SNPs from *Oryza sativa* genome assembly v.7.0

(*Oryza sativa* v.7.0, DOE-JGI, http://phytozome.jgi.doe.gov/). The candidate genes

from rice are listed in the table S1. The sequences of these candidate genes were used

to identify potential homologs in switchgrass by using the BLASTP tool on Phyto-

zome (Goodstein et al., 2012).

269   **RESULTS**

Phenotyping and correlation between traits

The results from leaf detachment showed that 'ECS.6' was the most resistant

population (DTVI 24% and DTIA 8%) and 'SW115' was the most susceptible popula-

tion (DTVI 81% and DTIA 60%) (Table S2). However, when comparing each geno-

type, 'SW788.05' was the most resistant genotype (DTVI 6% and DTIA 2%) whereas

'SW63.05' was the most susceptible genotype (Table S3). The result from the leaf

disk assay showed that 'SW803' was the most resistant population (DSVI 20% and

DSIA 48%) and 'SW38' was the most susceptible population (DSVI 98% and DSIA

99%). In the genotype-based comparison, 'High Tide.02' showed the most resistance

(DSVI 10% and DSIA 11%), but the most susceptible were 81 genotypes with DSVI

100% and DSIA 100%. In the field evaluation based on the mean of the severity score

(0 to 5) from two locations, 'Shelter' appeared to be the most resistant population

(0.44) and 'SW787' was the most susceptible. However, when considering each loca-

tion, different populations performed differently due to high GxE effect. In NY,

SW803 showed the most resistant population (0.22) and SW787 appeared to be the

most susceptible one (3.0). Whereas, in PA, Shelter appeared to be the most resistant

one (0.33) and Pathfinder appeared to be the most susceptible population (2.9). As

Silva et al. (2017) phenotyped potato tuber necrotic ringspot disease by using the

highest score for the disease evaluation across replicates to be able to review

significant QTLs, the highest score for BLS was considered to be used in this study

also. At the two locations, 'SW123', 'SW33', 'SW793', 'SW781', High Tide, and

Timber had the lowest MTL at 3 (resistant), and 41 populations had the highest MTL

at 5 (susceptible). In NY, SW123, SW128 and 'ECS.6' had the lowest MNY at 2, and

32 populations had the highest MNY at 5. In PA, 'SW115', 'SW802', 'SW31' and

'SW43' had the lowest MPA at 2, and 19 populations had the highest MPA at 5. The

most resistant and susceptible genotypes in each phenotyping approach are listed in

the Table S3. Based on artificial inoculations, both the leaf detachment and the leaf

disk assay suggested that the lowland ecotypes (DTVI 43%, DTIA 21%, DSVI 43%

and DSIA 70%) appeared to be more resistant than upland ecotypes (DTVI 61%,

DTIA 39%, DSVI 67% and DSIA 89%); however, the field evaluation of means

between two locations did not show the same trend (Figure S1).

Although each trait from different phenotyping approaches had significantly

moderate to high broad-sense heritability (H) (Table 1), H of severity from two loca-

tions was zero. This was suggested by the different trends of the mean of severity from

two locations, NY and PA (Figure S2). Due to zero heritability, the BTL was zero and

cannot be used for conducting GWAS. The different trends were not only present in

the field evaluation, the other phenotyping approaches yielded different trend of re-

sistance as suggested by the trends of DTIA, DSIA and mean from two locations and

correlation among the traits (Figure S2 and Figure 3). For example, from 479 geno-

types, 'KY1625_07' ranked 27 from DTIA but ranked 260 in DSIA, and 100 in mean

from two locations. Such differences led to low correlation among approaches such

that the r^2 between DTIA and DSIA was only 0.1 and between DTIA and mean of two

locations was 0.03 (Figure 3). In contrast to these low correlations, there were high

correlations in DTVI-DTIA ($r^2 = 0.84$) and DSVI-DSIA ($r^2 = 0.85$). There was no

BLS resistance trait correlating to other agronomic or biomass quality traits (Figure

S3).

Linkage disequilibrium and principal component analysis

To determine the gene interval that potentially linked to the significant markers from

GWAS, linkage disequilibrium (LD) decay was estimated by plotting the allele frequency cor-

relations (R^2) against the physical distance in base pairs (Figure 4). In the full set of 479 geno-

types, the LD decayed sharply within 20 kb and reached a plateau background within 50 kb.

The other groups of 4X, 8X, lowland and upland genotypes showed a similar pattern of the

LD despite being slightly different in an 8X group. Therefore, in this study, we focused on

genes within the 20 kb from the significant SNPs.

Based on the principal component analysis, there was a strong population structure in

this association panel (Figure 5). From PC1 and PC2, upland and lowland ecotypes were apart

from each other. Also, within the lowland ecotype, the latitude of lowland can be distin-

guished to lowland north and south groups. The separation of upland ecotype can also be

noticed in PC1 and PC3 in that the upland north was grouped apart from the other upland. Be-

sides ecotypes, ploidy level can be differentiated into groups. Both lowland north and south,

and upland north genotypes were tetraploid (4X) while the upland east and upland west geno-

types were octaploid (8X). In total, PC1, PC2, and PC3 explained the variance due to the pop-

ulation of 15.03%.

Genome-wide association mapping

As we conducted GWAS with five subsets of genotypes including 479 upland, low-

land, 4X and 8X to multiple combinations of assessing resistance to BLS via many phenotyp-

ing approaches, almost all the combinations for GWAS did not provide reliable results based

on flat or inflated QQ-plots. In the full set of 479 genotypes, single traits from artificial inocu-

lation via either leaf detachment or leaf disk assay with vision or image analysis (DTVI,

DTIA, DSVI, and DSIA) provided reliable GWAS results based on inflated QQ-plot (Figure

S4). Although DSIA showed the best QQ-plot (Figure S4d), the convex head and inflated tail

of -log(P-value) suggested confounding of the association analysis. The single-trait GWAS of

field evaluations (MTL, MNY, MPA, BNY, BPA, mean of two locations, mean of NY, and

mean of PA) did not yield any reliable association analysis (Figure S5). The combination of

two traits from artificial inoculations yielded better QQ-plots, but they still showed unreliable

QQ-plots with a convex head and early inflation in the middle of the plot. Among the two-

trait GWAS, DTIA-DSIA analysis provided the better GWAS with the QQ-plot that delayed

the inflated tail from -log(P-value) of 2 to about 2.5. To improve the association analysis, the

GWASs were then conducted with the combination of two artificial inoculations (DTIA-

DSIA) with field evaluations. The only three-trait combination that provided reliable GWAS

was DTIA-DSIA-MNY due to the QQ-plot with the expected distribution (Figure S6d).

In a tetraploid group, the same three-trait combination of DTIA-DSIA-MNY did not

provide the better GWAS than did MNY alone (Figure S7c, d). Among all the trait combina-

tions in the tetraploids, MNY provided the best association analysis with the expected distri-

bution. However, none of the GWAS octaploid group provided significant GWAS with relia-

ble QQ-plots. In the lowland group, the most reliable GWAS was from the combination of

DTIA-DSIA-MNY-MPA (Figure S8d). Interestingly, in the upland group, none of the image

analysis traits of DTIA and DSIA provided a reliable GWAS. The only nearly reliable single-

trait GWAS in the upland group was DTVI with the convex line in the middle of the QQ-plot

(Figure S9a). The two-trait combination of DTVI-MNY provided the most reliable GWAS

with the expected QQ-plot (Figure S9c). In contrast to the full set of 479 genotypes, the inclu-

sion of the trait from the leaf disk assay did not improve the analysis (Figure S9d).

In this study, therefore, we focused on four reliable GWAS analyses including 479-

DTIA-DSIA-MNY, 4X-MNY, low-DTIA-DSIA-MNY-MPA, and up-DTVI-MNY (Figure 6).

However, there was no overlapping region associated with the resistance among the four

GWAS analyses (Table 2). Accumulatively, when considering phenotypic variance explained

(PVE) by all 18 significant SNP markers for each trait, PVE of DSIA was the highest at

26.72% and PVE of DTIA was the lowest at 6.32% (Table 3). In 479-DTIA-DSIA-MNY, the

significant markers were on chromosomes 2b, 6a, 7b, 9a, and 9b. The most significant marker

on chromosome 2a was snp680175 followed by snp151084. Only snp359167 was the signifi-

cant marker on chromosome 6a. The most significant marker on chromosome 7b was

snp938364 followed by snp933838. The most significant marker on chromosome 9a was

snp469523 followed by snp465249. The snp1003871 was the only significant marker on chro-

mosome 9b. These eight SNP markers explained phenotypic variances for DTIA, DSIA and

MNY of 2.1, 24.28, and 5.27%, respectively (Table S4). In 4X-MNY, there were only two

significant markers snp316732 on chromosome 5a and snp444058 on chromosome 8a, accu-

mulatively explaining 1.58% in MNY. In low-DTIA-DSIA-MNY-MPA, the significant mark-

ers were on chromosomes 1b, 3a, 3b and 5b. Only snp594527 was the significant marker on

chromosome 1b. Also, snp197006 was the only significant marker on chromosome 3a, and

snp732235 was the only one on chromosome 3b. The most significant marker on chromosome

5b was snp791318 followed by snp791375. These five SNP markers explained phenotypic

variances for DTIA, DSIA, MNY, and MPA of 2.41, 8.31, 6.86, and 32.48%, respectively. In

up-DTVI-MNY, the significant markers were on only chromosome 7a, in which snp411995

was the most significant marker, followed by snp411929 and snp412960. Among these mark-

ers, only snp938364, on chromosome 7b from 479-DTIA-DSIA-MNY, was linked with two

genes: Pavir.Gb01488.1 and Pavir.Gb01488.2. These two were the only genes that overlapped

with LOC_Os04g39430.1, one of the potential candidate genes for resistance to BLS in rice

on chromosome 4 from AE11005627 marker (Table S1). Also, these three SNP markers ex-

plained phenotypic variances for DTVI and MNY of 7.19 and 0.27%, respectively. The PVEs

of each SNP in each subgroup and each trait are reported in Table S5.

From the QTLs study of Sato et al. (2015), the candidate resistance genes in rice were

reported to be on three chromosomes with 13 SNPs markers within the LD of 200 kb (Table

S1). There were 278 genes linked to these markers. Among them, 172 genes were annotated

with 134 biological functions. There were 106 non-annotated genes. The most frequent

biological functions were Leucine-rich repeat family protein (6), peroxidase precursor (6),

transposon protein (6), nucleotide binding sites – leucine rice repeats (NBS-LRR) type disease

resistance protein (5), MYB family transcription factor (3), dirigent (2), fasciclin domain-

containing protein (2), flavin monooxygenase (2), and heavy metal associated domain (2).

403   **DISCUSSION**

The most resistant population for improvement of resistance to BLS

Since the various phenotyping approaches yielded various resistant populations, it was

challenging to determine the most resistant candidates for further breeding. Based on field

evaluation, BLUPs from two locations were zero due to zero broad-sense heritability. This in-

dicated huge GxE effects between two locations. If the mean of field scores were used, the

upland Shelter should be considered the most resistant population across two locations with a

mean of 0.44. The high resistance of Shelter from field evaluation might explain the reason in

Songsomboon (2018) that it cannot be improved for resistance to BLS in recurrent phenotypic

selection in seedlings. However, the most susceptible SW787 had the mean score only 2.33.

This suggested the low natural inoculation in 2017 which was not conducive to severe disease

development. When comparing each location, the most resistant population in NY was

SW803 at 0.22, but it was almost the most susceptible population in PA at 2.11. Therefore,

this variation needs to be confirmed with more replicates, years, and locations.

To lessen the GxE effect on the BLS, artificial inoculation was conducted in the labor-

atory. In the leaf detachment assay, due to the limitation of sampling, a whole set of 479 gen-

otypes cannot be tested at the same time. The experiment was separated into seven overlap-

ping sets. However, there was significant variation among sets. Such a variation can occur

from the different age of each genotype in the same sampling time. Although the same stage

of leaves was sampled at 30 centimeters from the top, variation existed among leaves from the

same plant. To handle the variation among sets, BLUPs were used by fitting the effect from

the set as a fixed effect. Another artificial inoculation was leaf disk assay, without limitation

of the size, the whole set of 479 genotypes was sampled at the same time with resistant and

susceptible checks in each plate. Although a small effect of plates existed, BLUPs was fitted

by considering plates as a fixed effect. Both artificial inoculations were evaluated via vision

and image analysis. Although vision and image analysis results were highly correlated, they

yielded a different result from association analysis. Image analysis, however, provided a more

significant analysis. It could be explained by the more refined data from image analysis with

decimals than vision analysis (which grouped into ten increments). Moreover, the image anal-

ysis is better regarding repeatability and reanalysis (Stewart and McDonald, 2014). For the

sake of comparison for the most resistant population, mean of percent lesion from image anal-

ysis was considered. In the leaf detachment approach, the most resistant population from

DTIA was ECS.6 at 8%, but the population appeared very susceptible from DSIA at 94%. In

leaf disk assay, the most resistant population from DSIA was SW803 and SW793. These con-

firmed the low correlation among all three disease evaluation approaches. Regardless of low

correlations, SW805 was the most resistant population across the mean of two locations,

DTIA, and DSIA, 10% and 56% resistance, respectively, with mean of 0.78.

The improvement of multi-trait GWAS from single trait GWAS

Although all single traits from a different phenotype approach had moderate to high

heritability, none of them provided reliable GWAS based on QQ-plots that were not

distributed as expected from theory. This unreliability can be explained by the high density of

markers with high collinearity between markers, and too small population size giving the low

power detection (Muir, 2007). The combination of the traits for resistance to BLS from differ-

ent approaches helped improve the association analysis power (Guo et al., 2017). However,

the addition of traits did not always improve the analysis. In the tetraploid groups, for exam-

ple, only MNY yielded the most reliable GWAS. These suggested significant markers linked

with the specific location. The spatial markers of simple sequence repeats (SSRs) in QTLs

were also reported in resistance to BLS in rice (Katara et al., 2010). None of the reliable

GWAS provides the same significant regions linked with the resistance to BLS. As those reli-

able GWASs were from a different subset of the association panel, it suggested different re-

sistance allele distribution in the subgroups. It also suggests the multifaceted defense mecha-

nism to BLS.

456   Despite the improvement of multi-trait GWAS over single-trait GWAS, 18 SNP mark-

457   ers can explain the phenotypic variance ranging from only 6.32 to 26.72% among DTVI,

458   DTIA, DSIA, MNY and MPA. The low PVE from GWAS is a common phenomenon and

459   generally caused by missing heritability and can be improved by more genomic imputation

460   and implementation of a large-scale next-generation sequence data (Fan and Song, 2016).

461   Moreover, each genotype group can show reliable GWAS with specific traits. For ex-

ample, the upland group only provided the reliable GWAS with DTVI-MNY, whereas the full

set of 479 genotypes provided the best fit GWAS with DTIA-DSIA-MNY. To further study

resistance to BLS in switchgrass populations, single-trait GWAS is not suggested due to the

low power of determining significant markers. Multi-trait GWAS from both artificial and nat-

ural inoculation were suggested to improve the power. However, such an improvement cannot

be applied to the octaploid population due to the uncertainty in differentiating heterozygote

genotypes. The model for octaploid should be applied to the population instead of the diploid

model to detect associations.

Dissecting resistance to BLS

As mentioned previously, none of the significant markers overlapped across reliable

GWASs from different groups. Although this suggested the different distribution of re-

sistance genes in different subgroups from the association panel, we considered the biological

functions for the resistance over the fixation of the alleles in each population. In the full set,

on chromosome 2b, Pavir.Ba03861.1 was predicted to function as a copper transport protein

and as an antioxidant protein 1 (ATOX1). There was no study to confirm the function of this

gene related to any plant disease resistance, but its antioxidant capacity may help handle ROS

from the interaction between fungal effectors and plant response. Pavir.Bb02195.1 and

Pavir.Bb02196.1 encoded RING-type E3 ubiquitin-protein ligase and RBR family ring finger,

respectively, which collaboratively regulates cell death and immunity in rice (Smit and

Sixma, 2014; You et al., 2016). On chromosome 6b, Pavir.Fa00433.1 encoded for the EMSY

N-terminal (ENT) domain, which is important to mediate the race-specific NLR gene re-

sistance to *Peronospora parasitica* (*RPP7*) gene in *Arabidopsis thaliana* (Tsuchiya and

Eulgem, 2011). Moreover, Pavir.Fa00434.1 encoded the Myb-like DNA-binding domain.

This gene was confirmed to control many biological functions like biotic and abiotic stress,

development, defense, etc. in many organisms (Ambawat et al., 2013). In rice, the *MYB* gene

controls Jasmonic acid (JA) against rice blast fungus: *Pyricularia greisea* (Lee et al., 2001).

In Arabidopsis interaction with necrotrophic *Alternaria brassicicola*, the response to JA was

the upregulated plant defensin protein (*Pdf*) gene. This gene was negatively controlled by the

heat stress transcription factor B-1 encoded from Pavir.Fa00437.1 (Kumar et al., 2009). The

most interesting region linked with significant markers was on chromosome 7b since

Pavir.Gb01488.1 and Pavir.Gb01488.2 were the only genes that overlapped with the BLAST

result of the resistance to BLS in rice. These two genes encoded for cytochromes P450, which

involved diverse oxidation reactions and triterpene synthesis (Geisler et al.,). The most well-

known triterpene against fungal disease was Avenacins in oat (*Avena* spp.). Also,

Pavir.Gb01298.1 encoded for the polyadenylate-binding protein. Polyadenylation is an im-

portant process to mediate transposon slicing of the RPP7 gene in Arabidopsis (Tsuchiya and

Eulgem, 2013). Pavir.Gb01299.1 encoded for Isocitrate lyase, which was naturally a compo-

nent of the glyoxylate cycle and was required for the pathogenicity of ascomycete *Lepto-*

sphaeria maculans, causing blackleg disease of canola (*Brassica napus*) (Idnurm and

Howlett, 2002). On chromosome 9a, the most interesting gene was Pavir.Ia00351.1 encoding

mitogen-activated kinase (MAP). The mediation of the MAP has confirmed the activation of

expression of ethylene (ET), JA, and salicylic acid (SA) in the response to both abiotic and

biotic stress in Arabidopsis (Brader et al., 2007). Moreover, Pavir.Ia00174.1 encoded tri-

chome Birefringence-like protein responsible for xylan acetylation in rice, which conferred

the resistance to bacterial leaf blight (Gao et al., 2017). On chromosome 9b, Pavir.Ib01315.1,

Pavir.Ib01315.2, Pavir.Ib01317.1 encoded glutathione S-transferase (GST) which was an im-

portant enzyme in redox reactions in cells. Glutathione was the key compound in the signaling

process to trigger plant resistance to pathogens (Dubreuil-Maurizi and Poinssot, 2012).

In 4X-MNY, on chromosome 5a, Pavir.Ea02827.1, encoding Actinidain, cysteine pro-

tease and Cathepsin propeptide inhibitor, was directly involved in programmed cell death in

the plant (Solomon et al., 1999). Additionally, Pavir.Ea02828.1 encoded DEAD/DEAH box

helicase, which showed the up-regulation of defense-related signals like SA and JA and

conferred the resistance again rice blast fungus *Manaporthe grisea* (Li et al., 2008). On chro-

mosome 8a, Pavir.Ha00272.1 encoded for auxilin/cyclin G-associated kinase (GAK). Besides

relating to the kinase signal group conferring innate immunity, auxilin-like protein itself was

confirmed to induce necrotic lesion and confer resistance to *Xanthomonas oryzae* pv. Oryzae

(Xoo) (Park et al., 2017).

In low-DTIA-DSIA-MNY-MPA, on chromosome 1b, Pavir.Ab01193.1,

Pavir.Ab01193.2, and Pavir.Ab01193.3 encoded ginkbilobin-2. This protein was purified

from *Ginkgo biloba* seeds. It is cysteine-rich receptor-like kinase that confirmed the functions

of salt stress and antifungal activity (Sawano et al., 2007; Miyakawa et al., 2009). On chromo-

some 3b, Pavir.Ca02197.1 encodedf ATP-binding cassette (ABC) transporter. There are many

subgroups of ABC. In Arabidopsis, the mutant lacking the ABC transporter of PENETRA-

TION3/ PLEIOTROPIC DRUG RESISTANCE8 (PEN3/PDR8) conferred nonhost suscepti-

bility to *Blumeria graminis* f. sp hordei., suggesting that the ABC transporter was involved in

exporting secreted toxins from the fungus and fungal suppressor from the host (Stein et al.,

2006). Also, in wheat, the ABC suggested the exportation of mycotoxin conferring resistance

to Fusarium head blight (Walter et al., 2015). On chromosome 3b, Pavir.Cb01412.1 encoded

copines, which were diverse protein classes related to calcium-, phospholipid-binding func-

tion. The study in Arabidopsis suggested that copines acted as suppressors of defense re-

sponses in hypersensitive cell death defense against *Pseudomonas syringae* pv *tomato*

(Jambunathan and McNellis, 2003). On chromosome 5b, Pavir.Eb00172.1 encoded ser-

ine/threonine protein kinase (STPK) which were known to be candidates conferring disease

resistance in wheat to tan spot (Juliana et al., 2018) and Arabidopsis to flagellin bacteria

(Gómez-Gómez and Boller, 2000). Pavir.Eb00173.1 encoded BED zing finger which was

classed as NBS-LRR, which played an important role in effector-triggered immunity (ETI).

The zing finger domains suggested the broad-spectrum nature of rice blast resistance gene

Pi54 (Gupta et al., 2012). Pavir.Eb00177.1 encoded chloroplastic membrane protein, which

also can involve mediating ROS from avirulent pathogen invasion (Yaeno et al., 2004).

In up-DTVI-MNY, although there was only one significant peak of markers on chro-

mosome 7, it was unlikely to conclude that there was an R gene in this chromosome due to

multiple significant SNPs with diverse roles. Pavir.Ga00855.1 encoded STPK similarly to

Pavir.Eb00172.1 on chromosome 5. Pavir.Ga00857.1 and Pavir.Ga00858.1encoded for trans-

porter proteins like Pavir.Ca02197.1 on chromosome 3b. Lastly, Pavir.Ga00912.1 encoded

WD40 repeat family protein. Although there was no study directly with disease resistance,

WD40 is involved in many biosynthesis pathways such as Transparent Testa2, which is a

Myb transcription factor related to the resistance as explained earlier, and flavonoid biosyn-

thesis for antioxidation handling ROS.

Although these candidate genes from the switchgrass association panel shared some of

the functions from candidate genes from rice resistance to BLS, some of the potential func-

tions conferring resistance were still missing such as dirigent protein, peroxidase, and Mlo-

like protein. Dirigent protein conferred resistance by lignin and lignan synthesis (Ralph et al.,

2006). The lignin synthesis was expected to relate with resistance to BLS by strengthening

cell walls (Dallagnol et al., 2013). However, since none of the resistance traits for BLS corre-

lated with lignin, the missing dirigent protein from detection can be expected. Peroxidase also

played an important role in Pathogen-associated molecular pattern-triggered immunity (PTI)

(Mammarella et al., 2015). This was unexpected due to the high involvement of ROS in the

infection of BO (Antonious and Jawhar, 2013). Mlo was identified to provide broad resistance

to powdery mildew in barley (Jorgensen, 1992). More powerful model and finer phenotype

combinations were needed to clarify these missing candidates.

CONCLUSIONS AND POTENTIAL APPLICATIONS

In conclusion, the resistance to BLS yielded differently based on different phenotyp-

ing approaches, with low correlations suggesting that resistance was sensitive to phenotyping

methods in different conditions. Although it is difficult to standardize the phenotyping ap-

proach that can take account all conditions, finer and more powerful phenotyping of the re-

sistance is still needed. As such, to the best of our phenotyping combination, SW805 should

be considered the most resistant population. Based on obtained phenotypes, our multi-trait

GWAS with four subsets from 479 genotypes have a handful of markers that can be

considered as potential candidate genes related to resistance to BLS with reasonable biologi-

cal functions. Nevertheless, one needs to be prudent regarding the utilization of these markers.

Due to low correlations between resistance to BLS from different phenotyping approaches,

the combinations of the trait in a different subgroup of genotypes provided no significant

overlapping regions. The application of these markers cannot be directly transferred for ge-

nomics-assisted selection without validation to the choices of phenotyping approaches in new

populations.

When considering biological functions from the GWAS regardless of subgroup and

trait combinations, the defense mechanisms against BLS are complicated and multifaceted,

probably controlled by multiple small-effect alleles. Most of the responses involved ETI via

signaling kinases and NBS-LRR. Moreover, basal defense mechanism and PTI mostly via ox-

idative stress also played an important role in response to BLS. Despite missing expected can-

didate functions and only one overlapping gene for resistance to BLS in rice, most of the

genes from these GWASs in switchgrass encode proteins overlapping with genes in rice. To

my knowledge, this was first research to dissect the resistance to BLS in switchgrass. With the

current power of detection within the population, multi-trait GWAS can provide functional

insight into the resistance within multiple subsets of genotypes.

ACKNOWLEDGEMENTS

The project is supported by PhD funding from the Development and Promotion of Science

and Technology Talents Project (DPST), the Royal Thai Government. The northern

switchgrass association panel and its HAPMAP data were provided by Dr. Michael Casler

and Dr. Paul Grabowski from US Dairy Farage Research Center, USDA-ARS. The genome-

wide association analysis with multiple traits was consulted by Dr. Deniz Akdemir. Visual

and digital disease evaluations were guided by Dr. Shawn Kenaley and Dr. Ethan Stewart.

598   **LITERATURE CITED**

Ambawat, S., P. Sharma, N.R. Yadav, and R.C. Yadav. 2013. MYB transcription factor genes

as regulators for plant responses: an overview. Physiol. Mol. Biol. Plants 19(3): 307–21.

Antonious, A. D., & Jawhar, M. 2013. Proteome analysis of the susceptible barley-Cochliobo-

lus sativus interaction. Research in Biotechnology. 4(2).

Bates, D., Mächler, M., Bolker, B., & Walker, S. 2014. Fitting linear mixed-effects models

using lme4. arXiv preprint arXiv:1406.5823.

Bockhaven, J. V., Spíchal, L., Novák, O., Strnad, M., Asano, T., Kikuchi, S., & De Vleess-

chauwer, D. 2015. Silicon induces resistance to the brown spot fungus *Cochliobolus*

miyabeanus by preventing the pathogen from hijacking the rice ethylene pathway. New

Phytologist. 206(2): 761-773.

Bradbury, P.J., Z. Zhang, D.E. Kroon, T.M. Casstevens, Y. Ramdoss, and E.S. Buckler. 2007.

TASSEL: software for association mapping of complex traits in diverse samples. 23(19):

2633–2635.

Brader, G., A. Djamei, M. Teige, E.T. Palva, and H. Hirt. 2007. The MAP kinase kinase

MKK2 affects disease resistance in arabidopsis. 20(510): 589–596.

Condon, B. J., Leng, Y., Wu, D., Bushley, K. E., Ohm, R. A., Otillar, R., & Xue, C. 2013.

Comparative genome structure, secondary metabolite, and effector coding capacity

across Cochliobolus pathogens. PLoS Genetics, 9(1), e1003233.

Dallagnol, L., F. Rodrigues, and M. Mielli. 2013. Silicon improves the emergence and sanity

of rice seedlings obtained from seeds infected with Bipolaris oryzae. Trop. Plant Pathol.

38(6): 478–484.

Dubreuil-Maurizi, C., and B. Poinssot. 2012. Role of glutathione in plant signaling under bio-

tic stress. Plant Signal. Behav. 7(2): 210–2.

Evans, J., E. Crisovan, K. Barry, C. Daum, J. Jenkins, G. Kunde-Ramamoorthy, A. Nandety,

C.Y. Ngan, B. Vaillancourt, C.-L. Wei, J. Schmutz, S.M. Kaeppler, M.D. Casler, and

C.R. Buell. 2015. Diversity and population structure of northern switchgrass as revealed

through exome capture sequencing. Plant J. 84(4): 800–815.

Evans, J., J. Kim, K.L. Childs, B. Vaillancourt, E. Crisovan, A. Nandety, D.J. Gerhardt, T.A.

Richmond, J.A. Jeddeloh, S.M. Kaeppler, M.D. Casler, and C.R. Buell. 2014. Nucleotide

polymorphism and copy number variant detection using exome capture and next-genera-

tion sequencing in the polyploid grass Panicum virgatum. Plant J. 79(6): 993–1008.

Fajolu, O.L. 2012. Characterization of Bipolaris species, their effects on switchgrass biomass

yield and chemical components.

Fan, Y., & Song, Y. Q. 2016). Finding the missing heritability of genome-wide association

study using genotype imputation. Matters, 2(5), e201604000013.

Friesen, T.L., J.D. Faris, P.S. Solomon, and R.P. Oliver. 2008. Host-specific toxins: effectors

of necrotrophic pathogenicity. Cell. Microbiol. 10(7): 1421–1428

Friesen, T.L., S.W. Meinhardt, and J.D. Faris. 2007. The Stagonospora nodorum-wheat pa-

thosystem involves multiple proteinaceous host-selective toxins and corresponding host

sensitivity genes that interact in an inverse gene-for-gene manner. Plant J. 51(4): 681–

692.

Gao, Y., C. He, D. Zhang, X. Liu, Z. Xu, Y. Tian, X.-H. Liu, S. Zang, M. Pauly, Y. Zhou, and

B. Zhang. 2017. Two Trichome Birefringence-Like Proteins Mediate Xylan Acetylation,

Which Is Essential for Leaf Blight Resistance in Rice. Plant Physiol. 173(1): 470–481.

Geisler, K., R.K. Hughes, F. Sainsbury, G.P. Lomonossoff, M. Rejzek, S. Fairhurst, C.-E. Ol-

sen, M.S. Motawia, R.E. Melton, A.M. Hemmings, S. Bak, A. Osbourn, and J.R. Ecker.

Biochemical analysis of a multifunctional cytochrome P450 (CYP51) enzyme required

for synthesis of antimicrobial triterpenes in plants. Proceedings of the National Academy

of Sciences. 110(35): E3360-E3367.

Gómez-Gómez, L., and T. Boller. 2000. FLS2: an LRR receptor-like kinase involved in the

perception of the bacterial elicitor flagellin in Arabidopsis. Mol. Cell 5(6): 1003–11.

Goodstein, D.M., S. Shu, R. Howson, R. Neupane, R.D. Hayes, J. Fazo, T. Mitros, W. Dirks,

U. Hellsten, N. Putnam, and D.S. Rokhsar. 2012. Phytozome: a comparative platform for

green plant genomics. Nucleic Acids Res. 40(1): 1178–1186.

Grabowski, P.P., J. Evans, C. Daum, S. Deshpande, K.W. Barry, M. Kennedy, G. Ramstein,

S.M. Kaeppler, C.R. Buell, Y. Jiang, and M.D. Casler. 2017. Genome-wide associations

with flowering time in switchgrass using exome-capture sequencing data. New Phytol.

213(1): 154–169.

Guo, Y., Y. Huang, L. Hou, J. Ma, C. Chen, H. Ai, L. Huang, and J. Ren. 2017. Genome-wide

detection of genetic markers associated with growth and fatness in four pig populations

using four approaches. Genet. Sel. Evol. 49(1): 21.

Gupta, S.K., A.K. Rai, S.S. Kanwar, and T.R. Sharma. 2012. Comparative analysis of zinc

finger proteins involved in plant disease resistance. PLoS One 7(8): e42578.

Idnurm, A., and B.J. Howlett. 2002. Isocitrate lyase is essential for pathogenicity of the fun-

gus *Leptosphaeria maculans* to canola (*Brassica napus*). Eukaryot. Cell 1(5): 719–24.

Jambunathan, N., and T.W. McNellis. 2003. Regulation of Arabidopsis COPINE 1 gene ex-

pression in response to pathogens and abiotic stimuli. Plant Physiol. 132(3): 1370–81.

Jorgensen, I.H. 1992. Discovery, characterization and exploitation of Mlo powdery mildew

resistance in barley. Euphytica 63(1–2): 141–152.

Juliana, P., R.P. Singh, P.K. Singh, J.A. Poland, G.C. Bergstrom, J. Huerta-Espino, S. Bha-

vani, J. Crossa, and M.E. Sorrells. 2018. Genome-wide association mapping for re-

sistance to leaf rust, stripe rust and tan spot in wheat reveals potential candidate genes.

Theor. Appl. Genet.: 1–18.

Katara, J. L., Sonah, H., Deshmukh, R. K., Chaurasia, R., & Kotasthane, A. S. 2010. Molecu-

lar analysis of QTLs associated with resistance to brown spot in rice (Oryza sativa L.).

Indian J. Genet. 70(1): 17-21.

Kim, J.Y., J. Wu, S.J. Kwon, H. Oh, S.E. Lee, S.G. Kim, Y. Wang, G.K. Agrawal, R. Rakwal,

K.Y. Kang, I.P. Ahn, B.G. Kim, and S.T. Kim. 2014. Proteomics of rice and Cochliobo-

lus miyabeanus fungal interaction: Insight into proteins at intracellular and extracellular

spaces. Proteomics 14(20): 2307–2318.

Kumar, M., W. Busch, H. Birke, B. Kemmerling, T. Nürnberger, and F. Schöffl. 2009. Heat

shock factors HsfB1 and HsfB2b are involved in the regulation of Pdf1.2 expression and

pathogen resistance in Arabidopsis. Mol. Plant 2(1): 152–65.

Lee, M.-W., M. Qi, and Y. Yang. 2001. A Novel Jasmonic Acid-Inducible Rice myb Gene as-

sociates with fungal infection and host cell death. Mol. Plant-Microbe Interact. 14(4):

527–535.

Li, D., H. Liu, H. Zhang, X. Wang, and F. Song. 2008. OsBIRH1, a DEAD-box RNA hel-

icase with functions in modulating defense responses against pathogen infection and oxi-

dative stress. J. Exp. Bot. 59(8): 2133–46.

Lipka, A. E., Gore, M. A., Magallanes-Lundback, M., Mesberg, A., Lin, H., Tiede, T., & Del-

laPenna, D. 2013. Genome-wide association study and pathway level analysis of to-

cochromanol levels in maize grain. G3: Genes, Genomes, Genetics. g3-113.

Lipka, A.E., F. Lu, J.H. Cherney, E.S. Buckler, M.D. Casler, and D.E. Costich. 2014. Accel-

erating the switchgrass (Panicum virgatum L.) breeding cycle using genomic selection

approaches. PLoS One 9(11): 1–7.

Lu, F., A.E. Lipka, J. Glaubitz, R. Elshire, J.H. Cherney, M.D. Casler, E.S. Buckler, and D.E.

Costich. 2013. Switchgrass Genomic Diversity, Ploidy, and Evolution: Novel Insights

from a Network-Based SNP Discovery Protocol. PLoS Genet. 9(1).

Mammarella, N.D., Z. Cheng, Z.Q. Fu, A. Daudi, G.P. Bolwell, X. Dong, and F.M. Ausubel.

2015. Apoplastic peroxidases are required for salicylic acid-mediated defense against

Pseudomonas syringae. Phytochemistry 112: 110–121.

Martin, E.R., D.D. Kinnamon, M.A. Schmidt, E.H. Powell, S. Zuchner, and R.W. Morris.

2010. SeqEM: an adaptive genotype-calling approach for next-generation sequencing

studies. Bioinformatics 26(22): 2803–2810.

Miyakawa, T., K. Miyazono, Y. Sawano, K. Hatano, and M. Tanokura. 2009. Crystal struc-

ture of ginkbilobin-2 with homology to the extracellular domain of plant cysteine-rich

receptor-like kinases. Proteins Struct. Funct. Bioinforma. 77(1): 247–251.

Muir, W.M. 2007. Comparison of genomic and traditional BLUP-estimated breeding value

accuracy and selection response under alternative trait and genomic parameters. J. Anim.

Breed. Genet. 124(6): 342–355.

Okada, M., C. Lanzatella, M.C. Saha, J. Bouton, R. Wu, and C.M. Tobias. 2010. Complete

switchgrass genetic maps reveal subgenome collinearity, preferential pairing and multi-

locus interactions. Genetics 185(3): 745–760.

Park, C.-J., T. Wei, R. Sharma, and P.C. Ronald. 2017. Overexpression of Rice Auxilin-Like

protein, XB21, induces necrotic lesions, up-regulates endocytosis-related genes, and con-

fers enhanced resistance to *Xanthomonas oryzae* pv. oryzae. Rice. 10(1): 27.

Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M.A.R. Ferreira, D. Bender, J. Maller, P.

Sklar, P.I.W. de Bakker, M.J. Daly, and P.C. Sham. 2007. PLINK: A tool set for whole-

genome association and population-based linkage analyses. Am. J. Hum. Genet. 81(3):

559–575.

Ralph, S., J.-Y. Park, J. Bohlmann, and S.D. Mansfield. 2006. Dirigent proteins in conifer de-

fense: gene discovery, phylogeny, and differential wound- and insect-induced expression

of a family of DIR and DIR-like Genes in spruce (*Picea* spp.). Plant Mol. Biol. 60(1):

21–40.

Sato, H., I. Ando, H. Hirabayashi, Y. Takeuchi, S. Arase, J. Kihara, H. Kato, T. Imbe, and H.

Nemoto. 2008. QTL analysis of brown spot resistance in rice (*Oryza sativa* L.). Breed.

Sci. 58(1): 93–96.

Sato, H., K. Matsumoto, C. Ota, T. Yamakawa, J. Kihara, and R. Mizobuchi. 2015. Confirm-

ing a major QTL and finding additional loci responsible for field resistance to brown

spot (Bipolaris oryzae) in rice. Breed. Sci. 65(2): 170–175.

Sawano, Y., T. Miyakawa, H. Yamazaki, M. Tanokura, and K. Hatano. 2007. Purification,

characterization, and molecular gene cloning of an antifungal protein from Ginkgo bi-

loba seeds. Biol. Chem. 388(3): 273–80.

Silva, W., J. Ingram, C.A. Hackett, J.J. Coombs, D. Douches, G. Bryan, W. De Jong, and S.

Gray. 2017. Mapping loci that control tuber and foliar symptoms caused by PVY in Au-

totetraploid Potato (*Solanum tuberosum* L.). Genes Genomes Genetics: g3-300264.

Smit, J.J., and T.K. Sixma. 2014. RBR E3-ligases at work. EMBO Rep. 15(2): 142–54.

Solomon, M., B. Belenghi, M. Delledonne, E. Menachem, and A. Levine. 1999. The involve-

ment of cysteine proteases and protease inhibitor genes in the regulation of programmed

cell death in plants. Plant Cell 11(3): 431–44.

Songsomboon, K., R. Crawford, J. Crawford, J. Hansen, J. Cummings, N. Mattson, G. Berg-

strom, D. Viands. 2018. Recurrent phenotypic selection for resistance to diseases caused

by Bipolaris oryzae in switchgrass (*Panicum virgatum*). Manuscript submitted for publi-

cation.

Stein, M., J. Dittgen, C. Sánchez-Rodríguez, B.-H. Hou, A. Molina, P. Schulze-Lefert, V.

Lipka, and S. Somerville. 2006. Arabidopsis PEN3/PDR8, an ATP binding cassette

transporter, contributes to nonhost resistance to inappropriate pathogens that enter by di-

rect penetration. Plant Cell 18(3): 731–46.

Stewart, E.L., and B.A. McDonald. 2014. Measuring quantitative virulence in the wheat path-

ogen *Zymoseptoria tritici* using high-throughput automated image analysis. Phytopathol-

ogy 104(9): 985–992.

Stroup, J., M. Sanderson, J.P. Muir, M.J. McFarland, and R.L. Reed. 2003. Comparison of

growth and performance in upland and lowland switchgrass types to water and nitrogen

stress. Bioresour. Technol. 86: 65–72.

Tsuchiya, T., and T. Eulgem. 2011. EMSY-Like Genes Are Required for Full RPP7-Mediated

Race-Specific Immunity and Basal Defense in Arabidopsis. 24(1210): 1573–1581.

Tsuchiya, T., and T. Eulgem. 2013. An alternative polyadenylation mechanism coopted to the

Arabidopsis RPP7 gene through intronic retrotransposon domestication. Proc. Natl. Acad. Sci. U. S. A. 110(37): E3535-43.

Walter, S., A. Kahla, C. Arunachalam, A. Perochon, M.R. Khan, S.R. Scofield, and F.M. Doohan. 2015. A wheat ABC transporter contributes to both grain formation and myco-toxin tolerance. J. Exp. Bot. 66(9): 2583–2593.

Xue, D., Q. Wang, Z. Chen, L. Cai, L. Bao, Q. Qi, L. Liu, X. Wang, H. Jin, J. Wang, H. Wu, H. Liu, and Q. Chen. 2015. 3-Anhydro-6-hydroxy-ophiobolin A, a fungal sesterterpene from Bipolaris oryzae induced autophagy and promoted the degradation of α-synuclein in PC12 cells. Bioorg. Med. Chem. Lett. 25(7): 1464–1470.

Yaeno, T., O. Matsuda, and K. Iba. 2004. Role of chloroplast trienoic fatty acids in plant dis-ease defense responses. Plant J. 40(6): 931–941.

You, Q., K. Zhai, D. Yang, W. Yang, J. Wu, J. Liu, W. Pan, J. Wang, X. Zhu, Y. Jian, J. Liu, Y. Zhang, Y. Deng, Q. Li, Y. Lou, Q. Xie, and Z. He. 2016. An E3 Ubiquitin Ligase-BAG protein module controls plant innate immunity and broad-spectrum disease re-sistance. Cell Host Microbe 20(6): 758–769.

Zhao, K., C. W. Tung, G.C. Eizenga, M.H. Wright, M.L. Ali, A.H. Price, G.J. Norton, M.R. Islam, A. Reynolds, J. Mezey, A.M. McClung, C.D. Bustamante, and S.R. McCouch. 2011. Genome-wide association mapping reveals a rich genetic architecture of complex traits in Oryza sativa. Nat. Commun. 2: 467.

Zhou, X., and M. Stephens. 2012. Genome-wide efficient mixed-model analysis for associa-tion studies. Nat. Genet. 44(7): 821–4.

 TABLES

778   Table 1. Broad-sense heritability (s.e.) of each trait.

Traits	H (s.e.)
Severity from the leaf detachment via vision (DTVI)	0.76 ± 0.02
Severity from the leaf detachment via image analysis (DTIA)	0.74 ± 0.06
Severity from the leaf disk assay via vision (DSVI)	0.56 ± 0.05
Severity from the leaf disk assay via image analysis (DSIA)	0.54 ± 0.04
BLUPs two locations	0.003 ± 0.06
BLUPs NY	0.32 ± 0.05
BLUPs PA	0.61 ± 0.04

 Table 2. Markers significantly associated with resistance in each case.

Cases	SNP	Chromosome	Distance from gene (bp)	p-value	*P. virgatum* gene	Orthologous genes	Identity	Predicted function
479-DTIA-DSIA-MNY	snp151084	Chr02a	419	1.32E-06	Pavir.Ba03861.1	Pahal.B02018.1[Pha]	99.4	copper transport protein ATOX1-related
	snp680175	Chr02b	975	3.17E-06	Pavir.Bb02195.1	Pahal.B02834.1[Pha]	94.8	E3 ubiquitin-protein ligase RNF144 [EC:6.3.2.19]
			3533		Pavir.Bb02196.1	Pahal.J00890.1[Pha]	93.8	RBR family ring finger and IBR domain-containing
	snp359167	Chr06a	2850	2.96E-06	Pavir.Fa00433.1 Pavir.Fa00433.2	Pahal.F00718.1[Pha]	88.9	ENT domain
			716		Pavir.Fa00434.1	Pahal.F00719.1[Pha]	70.9	Myb-like DNA- binding domain, Bromodomain
			2690		Pavir.Fa00435.1	Pahal.F00721.1[Pha]	54.6	uncharacterized C-terminal membrane protein
			8477		Pavir.Fa00437.1	LOC_Os04g48030.1[Osa]	8.5	heat stress transcription factor B-1
	snp933838	Chr07b	1453	2.12E-06	Pavir.Gb01298.1 Pavir.Gb01298.2	Sobic.006G139100.1[Sbi]	96.7	polyadenylate-binding protein
			609		Pavir.Gb01299.1	Pahal.H00668.1[Pha]	93.5	2,3-dimethylmalate lyase, (2R,3S)-2,3-dimethylmalate pyruvate-lyase, isocitrate lyase
	snp938364	Chr07b	597	6.81E-06	Pavir.Gb01488.1* Pavir.Gb01488.2*	LOC_Os04g39430.1[Osa]	85.6	cytochrome P450
			2987		Pavir.Gb01489.1 Pavir.Gb01489.2	Pahal.G00545.1[Pha]	97.4	protection of telomeres 1

Cases	SNP	Chromosome	Distance from gene (bp)	p-value	*P. virgatum* gene	Orthologous genes	Identity	Predicted function
					Pavir.Gb01489.3 Pavir.Gb01489.4			
	snp465249	Chr09a	3464	1.58E-07	Pavir.Ia00174.1	Pahal.I01213.1[Pha]	97.3	protein trichome Birefringence-like 32
			1915		Pavir.Ia00175.1 Pavir.Ia00175.2	Sobic.001G038100.1[Sbi]	92.2	protein FLC expressor
	snp469523	Chr09a	5412	4.77E-06 4.77E-06	Pavir.Ia00350.2	Pahal.I01603.1[Pha]	89.4	protein K03B8.4-related
			2256		Pavir.Ia00351.1	Sobic.001G079200.1[Sbi]	97.3	mitogen-activated kinase kinase kinase (MAP)
	snp1003871	Chr09b	5314	9.85E-06	Pavir.Ib01315.1 Pavir.Ib01315.2	Sevir.9G450400.1[Svi]	95.0	glutathione s-transferase (GST)
			788		Pavir.Ib01317.1	Pahal.I04479.1[Pha]	94.1	glutathione s-transferase (GST)
			3573		Pavir.Ib01318.1	Zm00008a00114T01[Zma]	83.0	ent-kaurenoic acid hydroxylase (KAO) subfamily P450
4X-MNYNY	snp316732	Chr05a	8347	6.08E-07	Pavir.Ea02827.1	Sevir.3G000300.1[Svi]	63.6	Actinidain/Actinidin [3.4.22.14], cysteine protease, Cathepsin pro-peptide inhibitor domain
			2009		Pavir.Ea02828.1 Pavir.Ea02828.2 Pavir.Ea02828.3	Pahal.E01498.1[Pha]	96.3	[3.64.13] RNA helicase, ATP-dependent RNA helicase, DEAD/DEAH box helicase
	snp444058	Chr08a	1739	1.21E-06	Pavir.Ha00272.1	Pahal.H00237.1[Pha]	90.6	auxilin/cyclin g-associated kinase-related
low-	snp594527	Chr01b	7186	2.37E-06	Pavir.Ab01193.1 Pavir.Ab01193.2 Pavir.Ab01193.3	LOC_Os07g35340.1[Osa]	74.7	protein kinase domain (Pkinase), salt stress response,antifungal (stress-antifung)

Cases	SNP	Chromosome	Distance from gene (bp)	p-value	*P. virgatum* gene	Orthologous genes	Identity	Predicted function
			199		Pavir.Ab01194.1	Pahal.A00997.1[Pha]	56.0	component of root nitrate treatment specific coexpression sub-network
	snp197006	Chr03a	1106	1.58E-06	Pavir.Ca02197.1	Sobic.008G093000.1[Sbi]	92.6	ATP-binding cassette (ABC) transporter
	snp732235	Chr03b	1613	1.75E-06	Pavir.Cb01411.1 Pavir.Cb01411.2	Zm00008a03100T01[Zma]	99.8	elongation factor 1-alpha (EEF1A)
			4102		Pavir.Cb01412.1	Sobic.009G118300.1[Sbi]	95.1	copine
	snp791318	Chr05b	7974	2.56E-09	Pavir.Eb00172.1	Sevir.5G014600.1[Svi]	82.6	protein kinase domain (Pkinase), S-locus glycoprotein domain, D-mannose binding lectin (B_lectin), PAN-like domain (PAN_2), serine/threonine protein kinase
			5076		Pavir.Eb00173.1 Pavir.Eb00173.2	Pahal.E04515.1[Pha]	97.7	HAT family dimerization domain-containing protein, BED zinc finger
			1362		Pavir.Eb00174.1	Pahal.E04517.1[Pha]	98.3	Diacylglycerol o-acyltransferase 1 (DGAT1)
			2273		Pavir.Eb00175.1	Sobic.003G013800.1[Sbi]	88.0	uncharacterized
	snp791375	Chr05b	3373	4.16E-06	Pavir.Eb00177.1	Sobic.003G014100.1[Sbi]	64.8	GDT1-like protein 1, chloroplastic, membrane protein
			4141		Pavir.Eb00178.1	Sobic.003G014300.2[Sbi]	99.5	Snoal-like domain, uncharacterized conserved protein
up-	snp411929	Chr07a	2184	3.44E-07	Pavir.Ga00855.1 Pavir.Ga00855.2 Pavir.Ga00855.3	Pahal.G01363.1[Pha]	96.8	RNA polymerase II, C-terminal domain phosphatase-like 1/2

Cases	SNP	Chromosome	Distance from gene (bp)	p-value	*P. virgatum* gene	Orthologous genes	Identity	Predicted function
								[EC:3.1.3.16] (CPL1_2), protein-serine/threonine phosphatase
			6254		Pavir.Ga00856.1	Zm00008a03828T01[Zma]	74.2	lactosylceramide 4-alpha-galacto-syltransferase
			2093		Pavir.Ga00857.1	Sobic.006G156600.2[Sbi]	97.2	transporter family protein
			507		Pavir.Ga00858.1	Sobic.006G156600.2[Sbi]	54.0	sugar (and other) transporter
	snp411995	Chr07a	49	8.66E-07	Pavir.Ga00859.1 Pavir.Ga00859.2 Pavir.Ga00859.3 Pavir.Ga00859.4 Pavir.Ga00859.5 Pavir.Ga00859.6	Zm00008a00013T01[Zma]	76.8	general transcription factor 3c polypeptide 6 (GTF3C6)
			6080		Pavir.Ga00860.1 Pavir.Ga00860.2	Zm00008a00671T01[Zma]	92.2	expansin precursor
	snp412960	Chr07a	673	3.08E-06	Pavir.Ga00912.1 Pavir.Ga00912.2	Sobic.006G156300.1[Sbi]	92.7	WD-40 repeat family protein

* *P. virgatum* genes that overlapping with resistance genes in *Oryza sativa* search

[Osa] gene from *Oryza sativa*

[Pha] gene from Panicum hallii

[Sbi] gene from Sorghum bicolor

[Svi] gene from Setaria viridis

[Zma] gene from Zea may

Table 3 Phenotypic variance explained (PVE) (adjusted PVE) by accumulative 18 significant

SNP markers across all analyses for each trait.

Traits	PVE (adjusted PVE)
DTVI	7.68 (4.07)
DTIA	6.32 (2.65)
DSIA	26.72 (23.85)
MNY	9.93 (6.4)
MPA	11.38 (7.91)

790   Abbreviation: BLUPs of severity from the leaf detachment via vision (DTVI), BLUPs of se-

791   verity from the leaf detachment via image analysis (DTIA), BLUPs of severity from the leaf

792   disk assay via image analysis (DSIA), highest score of BLS in NY (MNY), highest score of

793   BLS in PA (MPA).

FIGURES

Figure 1. Leaf detachment assay; a) sealed plates of detached leaves under 24-h light 25°C; b) control inoculated (left) and inoculated leaves (right) at 7 dpi were taped on white paper with QR code label for scanning for image analysis.

Figure 2. A plate example of leaf disk assay at 7 dpi. Each plate consisted of 28 leaf disks. SW43_09 was used as a resistant check and SW122_02 as a susceptible check. Two control leaf disks (yellow circles) and two inoculated leaf disks (red circles) were included in all plates. The rest of the leaf disks (24) consisted of eight genotypes with three replicates placed in a consecutive row from left to right.

797
798

Figure 3. Correlation plot among BLUPs of severity from the leaf detachment via vison (DTVI), via image analysis (DTIA), from leaf disk assay via vision (DSVI), via image analysis (DSIA), the highest score between two locations (MTL), mean from two location, minerals total ash, *in vitro* dry matter digestibility, and cell wall concentration.

799
800
801

Figure 4. Scatter plot showing the linkage disequilibrium (LD) decay by plotting physical distance in base pairs against the LD estimate as correlation (R^2) in a) 479 genotypes, b) 4X group, c) 8X group, d) lowland ecotype, and e) upland ecotype. From full set of 479 genotypes, the LD (blue curve) decays rapidly within 20 kb (red line) and reaches background levels around 50 kb (dashed red line).

Figure 5. The 479 switchgrass genotypes from the Northern Association Panel showing the

different distributions based on ecotype and region from principal component analysis (PC).

The lowland ecotype can be differentiated into two regions from PC1 and PC2, and the up-

land ecotype can be grouped into three regions from PC1 and PC3.

Figure 6. (Left) Manhattan plot showing genetic associated of a) DTIA-DSIA-MNY in 479 genotypes, b) MNY in 4X genotypes, c) DTIA-DSIA-MNY-MPA in lowland ecotype and d) DTVI-MNY in upland genotypes. The dashed line represents the FDR threshold (0.1) and the line represents Bonferroni correction threshold. On x-axis, the physical positions of the SNPs were aligned in 18 chromosomes of *P. virgatum*; however, there were 16,676 genes that cannot be assigned to the genome and noted as chromosome U1-U4. (right) Quantile-quantile (QQ) plots between the distributions of observed to expected P-values for GWAS of each trait combination in each genotype group.

Abbreviation: BLUPs of severity from the leaf detachment via vision (DTVI), BLUPs of severity from the leaf detachment via image analysis (DTIA), BLUPs of severity from the leaf disk assay via image analysis (DSIA), highest score of BLS in NY (MNY), highest score of BLS in PA (MPA).