A transposable element insertion is the switch between alternative life history strategies

Authors: Alyssa Woronik*[1,2], Kalle Tunström[1], Michael W. Perry[2,3], Ramprasad Neethiraj[1],

Constanti Stefanescu[4,5], Maria de la Paz Celorio-Mancera[1], Oskar Brattström[6], Jason Hill[1,7],

Philipp Lehmann[1], Reijo Käkelä[8], Christopher W. Wheat*[1]

Author affiliations:

[1] Department of Zoology, Stockholm University, S106 91 Stockholm, Sweden

[2] Department of Biology, New York University, New York, New York 10003, USA

[3] Division of Biological Sciences, University of California San Diego, La Jolla, California 92093,

USA

[4] Museum of Natural Sciences of Granollers, Granollers, Catalonia 08402, Spain

[5] CREAF, Cerdanyola del Valles, Catalonia 08193, Spain

[6] Department of Zoology, University of Cambridge, Cambridge CB23EJ, United Kingdom

[7] Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden

[8] Helsinki University Lipidomics Unit, Helsinki Institute for Life Science (HiLIFE) and Molecular

and Integrative Biosciences Research Programme, University of Helsinki, FI00014 Helsinki,

Finland

Correspondence to: AW alyssa.woronik@zoologi.su.se and CWW chris.wheat@zoologi.su.se

Tradeoffs affect resource allocation during development and result in fitness consequences that

drive the evolution of life history strategies. Yet despite their importance, we know little about

the mechanisms underlying life history tradeoffs in wild populations. Many species of *Colias*

butterflies exhibit an alternative life history strategy (ALHS) where females divert resources from

wing pigment synthesis to reproductive and somatic development. Due to this reallocation, a

wing color polymorphism is associated with the ALHS: individuals have either yellow/orange or

white wings. Here we map the genetic basis of the ALHS switch in *Colias crocea* to a

transposable element insertion downstream of the *Colias* homolog of *BarH-1,* a homeobox

transcription factor. Using CRISPR/Cas9 gene editing, antibody staining, and electron

microscopy we find morph-specific specific expression of *BarH-1* suppresses the formation of

pigment granules in wing scales. Lipid and transcriptome analyses reveal physiological

differences associated with the ALHS. These findings characterize a novel mechanism for a

female-limited ALHS and show that the switch arises via recruitment of a transcription factor

previously known for its function in cell fate determination in pigment cells of the retina.

A life-history strategy is a complex pattern of co-evolved life history traits (e.g. number of

offspring, size of offspring, and lifespan[1]), that is fundamentally shaped by tradeoffs that arise

because all fitness components cannot simultaneously be maximized. Therefore, finite

resources are competitively allocated to one life history trait versus another within a single

individual, and selection acts on these allocation patterns to optimize fitness[2]. Evolutionary

theory predicts that positive selection will remove variation from natural populations, as

genotypes with the highest fitness go to fixation[3]. However, across diverse taxa alternative life

history strategies (ALHSs) are maintained within populations at intermediate frequencies due to

balancing selection[4]. Life history theory was developed using methods such as quantitative

genetics, artificial selection, demography, and modeling to gain significant insights into the

causes and consequences of genetic and environmental variation on life history traits. Yet

despite these advances, a key challenge that remains is to identify the proximate mechanisms

underlying tradeoffs, especially for ecologically relevant tradeoffs that occur in natural

populations[5]. Here, we identify the mechanism underlying one such ALHS in the butterfly *Colias*

crocea (Pieridae, Lepidoptera) (Geoffroy, 1785).

Colias butterflies (the "clouded sulphurs") are common throughout the Holarctic and can be

found on every continent except Australia and Antarctica[6]. In approximately a third of the nearly

90 species within the genus, females exhibit two alternative wing-color morphs: yellow or

orange (depending on the species) and white[6,7] (Fig. 1A). The wing color polymorphism arises

because during pupation the white morph, also known as Alba, reallocates larval derived

resources from the synthesis of energetically expensive colored pigments to reproductive and

somatic development[8]. This tradeoff has been well characterized in *Colias crocea*, the Old

World species that we focus upon in this work, via radio-labelled metabolite tracking in pupae*[9]*

as well as in the New World species *Colias eurytheme*[8] (Pieridae, Lepidoptera) (Boisduval,

1852) using ultraviolet spectrophotometry. As a result of the resource reallocation, Alba females

have faster pupal development, a larger fat body, and significantly more mature eggs at

eclosion compared to orange females[10]. However, despite these developmental advantages

and the dominance of the Alba allele, the polymorphism is maintained by several abiotic and

biotic factors [10-14]. For example, males preferentially mate with orange females, as wing color is

an important cue for mate recognition[10,12,13]. This mating bias likely has significant fitness costs

for Alba females because males transfer essential nutrients during mating, and multiply mated

females have more offspring over their lifetime[15,16]. The mating bias against Alba females is

strongest in populations that frequently co-occur with other white Pierid butterfly species due to

interference competition[13]. Also, Alba's development rate advantage is temperature dependent,

with Alba females having faster development in cold temperatures[10]. Field studies confirm Alba

frequency and fitness increases in species that inhabit cold and nutrient poor habitats, where

the occurrence of other white Pierid butterflies is low. While in warm environments with nutrient

rich host plants and a high co-occurrence of other white species, orange females exhibit

increased fitness and frequency[12-14]. Previous work has also suggested Alba females have a

higher sensitivity to viral infections[9]. In all *Colias* species where it has been investigated (n=6),

the switch between the Alba or the orange strategy is controlled by a single, autosomal locus[6].

This fact, along with ancestral state reconstruction[7], has led to the assumption that the Alba

locus is conserved within the genus *Colias*, and potentially across the subfamily Coliadinae. Yet,

despite over a century of research on various aspects of Alba biology the mechanism underlying

this polymorphism remained unknown.

Using a *de novo* reference genome for *C. crocea* that we generated via Illumina and PacBio

sequencing, and three rounds of bulk segregant analyses (BSA) using whole genome

sequencing from a female and two male informative crosses for Alba, we mapped the Alba

locus to a ~3.7 Mbp region (Supplementary Fig.1, & Supplementary Information). Then, with

whole genome re-sequencing data from 15 Alba and 15 orange females from diverse population

backgrounds, a SNP association study fine mapped the Alba locus to a ~430 kb contig that fell

within the ~3.7 Mbp locus identified using the BSA crosses (Fig. 1B and Supplementary

Information). The majority of SNPs significantly associated with Alba (n=70 of 72) were within or

flanking a *Jockey-like* transposable element (TE) (Fig. 1C). We determined that the TE insertion

was unique to the Alba morph in *C. crocea* by assembling orange and Alba haplotypes for this

region, then quantifying differences in read depth between morphs within and flanking the

insertion (Supplementary Information and Supplementary Figs. 2, 3, & 4). We then used PCR to

validate the presence or absence, respectively, of the insertion in 25 Alba and 57 orange wild-

caught females (Supplementary Fig. 7). We also found no evidence of a TE insertion in the

homologous region of other butterfly genomes (*Danaus plexippus* & *Heliconius melpomene*)

(Supplementary Fig. 2).

The Alba-specific insertion was located ~30 kb upstream of a gene encoding a DEAD-box

helicase, and ~6kb downstream of the *Colias* homolog of *BarH-1*, a homeobox transcription

factor (Fig. 1C). *BarH-1* was an intriguing find as it affects color via pigment granule

development within eyes of *Drosophila melanogaster*[17]. To investigate BarH-1 expression in

developing *C. crocea* wings, we used *in situ* hybridization of BarH-1 on wings from two day old

pupae of orange and Alba females. We found the BarH-1 protein is expressed in scale building

cells within the white wing regions in Alba females (Fig. 2B). We did not observe BarH-1 in scale

building cells from orange areas of the wing in orange females (Fig, 2C). Interestingly however,

we found BarH-1 is expressed in scale building cells within black regions for both morphs (Fig,

2A&D). To validate the functional role of *BarH-1* in the Alba phenotype, we generated

CRISPR/Cas9-mediated deletions within exons 1 and 2 using a mosaic knockout (KO)

approach (Supplementary Information). *BarH-1* KO gave rise to a white/orange color mosaic on

the dorsal side of the wings in females with an Alba genotype (i.e. TE insertion +) (Fig. 1D),

while KO males and orange females displayed no white/orange mosaic on the wing. These

results indicate *BarH-1* expression suppresses orange coloration in the wings. We also

observed black and green mosaic coloring of eyes in KO males and females of both morphs,

where green eyes are the wild type color (Fig. 1E). These results indicate *BarH-1* also plays a

role in *Colias* eye development.

We next investigated how the Alba color change manifests within wings. Butterfly wing color can

arise either due to the absorption of light by pigments deposited within the scales, or by the

scattering of light via regularly arranged nanostructures in the scales[18]. *Colias* butterflies have

pteridine pigments. These pigments are synthesized within the wings and previous work using

ultraviolet spectrophotometry in *C. eurytheme* found Alba females exhibit dramatic reductions in

colored pteridine pigments compared to orange[8,9]. In insects, pteridines are synthsized in

pigment granules and pigment granules are concentrated within wing scales of Pierid

butterflies[19,20]. However, whether morphs differed in wing scale morphology was unknown. To

investigate wing morphology, we used scanning electron microscopy and found white scales

from Alba individuals exhibited a dramatic and significant reduction in pigment granules,

compared to orange scales ($t_{5.97}$ = 2.93, p = 0.03) (Fig 3 A&B). These results indicate the color

change to white is caused by reduced pigment granule formation. Congruent with this

interpretation, CRISPR KO Alba individuals exhibited significantly less pigment granules in

scales from the white wild-type region compared to scales in orange *BarH-1* KO regions ($t_{5.45}$ =

10.78, p < 0.001) (Fig. 3C). To further test whether reduction in pigment granule amount alone

was sufficient for the orange to white color change, we chemically removed the pigment

granules from the wing of an orange *C. crocea* female. This resulted in formerly orange regions

turning white (Fig. 3D). Wings likely appear white after granule removal due to the scattering of

light from the remaining non-lamellar nanosctructures[21]. These results demonstrate that *BarH-1*

suppresses pigment granule formation in wing scales, resulting in the white color of Alba

females in *C. crocea*. Thus, we propose the resource tradeoff between color and development

arises due to a classic Y reallocation model, wherein limited resources are competatively

allocated and increased investement in one trait results in a decreased investment to another[22].

Within the energetically closed system of a developing pupa, reduced pigment granule

formation would likely result in reduced pigment synthesis, which would in turn leave more

resources free to be used for other developmental processes. Finally, we also observed scale

building cells in black regions of both morphs express BarH-1 and also lack pigment granules

(Fig 2 A&D and Fig 3 A&B), but these scales appear black due to melanin deposition within the

scale[18]. These results suggest BarH-1 may also repress pigment granule formation within black

scales.

The Alba mechanism is assumed to be conserved across *Colias.* Therefore, we wished to test

whether Alba females from the New World species *Colias eurytheme* also exhibited significantly

less pigment granules than orange females. Indeed, we found orange *C. eurytheme* scales

exhibited abundant pigment granules while Alba scales almost entirely lacked granules (Fig. 3

E&F). These results demonstrate white wing color arises via the same morphological

mechanism within *Colias* and corroborate previous assumptions that Alba is conserved across

the genus. To further validate that other aspects of the Alba/orange alternative life history

strategy are conserved across the genus we tested whether one of the physiological tradeoffs of

Alba reported for New World species was also seen in *C. crocea*. In *C. eurytheme,* Alba females

have larger fat bodies than orange females and the strength of the Alba advantage increased in

cold temperatures[10]. To compare abdominal lipid stores between morphs in *C. crocea*, we

conducted high performance thin layer chromatography on two day old adult females reared

under two temperature treatments (Hot: 27^{o}C vs. Cold: 15^{o}C during pupal development). Adults

were not allowed to feed before samples were taken, therefore these measurements reflect

larval stores, where the putative energetic tradeoff should be more clearly visible. We found

Alba females had larger abdominal lipid stores than orange in both temperature treatments,

though the difference was only significant in the cold treatment (cold: n=32, $t_{29.12}$ = 3.42, P =

0.002, hot: n=25, $t_{22.71}$ = 0.67, P = 0.51) (Fig. 4A). These results are consistent with previous

reports from New World *Colias* species and indicate that the morph-specific tradeoff associated

with the color change is also conserved across the genus.

We then investigated the transcriptome of pupal abdomen and wing tissue at the time of

pteridine synthesis to identify genes that exhibited differential expression between morphs and

therefore may play a role in the morph-specific differences in physiology that arise due to the

resource tradeoff (Fig. 4 B&C, Supplementary Information, Supplementary Tables 5,6,&7). In *C.*

eurytheme Alba females emerge from the pupa with significantly more mature eggs than orange

females[10] and we find evidence that suggests similar dynamics are occurring in *C. crocea.* A

gene set enrichment analysis (GSEA) revealed that 'embryo development ending in birth or egg

hatching' (GO:0009792, p = 0.00072), 'proteasome-mediated ubiquitin-dependent protein

catabolic process' (GO:0043161, p = 0.00073), and 'proteolysis' (GO:0006508, p = 0.00101)

were within the top 5 terms enriched and upregulated within Alba abdomens (Supplementary

Table 6). Additionally our differential expression analysis identified a that gene which encodes a

triacylglycerol lipase was significantly upregulated within Alba abdomen tissue (log fold change

[log FC] of 4.8) (Fig. 4B and Supplementary Table 5). Triacylglycerol composes more than 90%

of the lipids stored in the fat body and during times of energy demand triacylglycerol lipases

mobilize these stores[23]. For example, during embryogenesis there is a massive shift in lipid

distribution from the fat body to ovaries as lipids comprise 30-40% of the dry weight of insect

embryos[23]. Taken together these results suggest that, similar to *C. eurytheme*, Alba females of

C. crocea may be benefitting from increased embryogenesis compared to orange females. We

also observe an enrichment of 'defense response to Gram-positive bacterium' (GO:0050830,

0.00027) for genes upregulated within Alba abdomens. Interestingly, previous work has

suggested that Alba females may have enhanced sensitivity to viral infection[9]. Further

investigation of potential morph-specific tradeoffs between wing color and immunity is of

interest.

For genes downregulated in Alba abdomens the GSEA revealed significant enrichment of

'regulation of nucleoside metabolic process' (GO:0009118, p value < 0.0001) and 'regulation of

purine nucleotide catabolic process' (GO:0033121, p value < 0.0001) (Supplementary Table 6).

Colias wings use purine precursors to synthesize pteridines[24]. Downregulation of these GO

terms in Alba abdomens suggests that the decreased pteridine synthesis observed in Alba

females[8], which likely arises due to the decrease in pigment granules in the wings, leads to a

decrease in purine precursors being shunted from the abdomen to the wings. Additionally,

consistent with previous reports of GTP reallocation from wings to other areas of development
in Alba females[8] we also observed significant enrichment of 'positive regulation of GTPase
activity' (GO:0043547, p value < 0.0001). Additionally, *RIM*, a Rab GTPase effector[25], was one
of the most highly differentially expressed (DE) genes in both tissues (logFC increase in Alba of
3.4 in the abdomen and 5.1 in the wings) (Fig. 4 B&C and Supplementary Table 5). RIM acts as
a molecular switch by converting guanosine diphosphate to guanosine triphosphate (GTP),
thereby activating its associated Rab GTPase, which is in turn involved in synaptic vesicle
exocytosis and secretory pathways[26].
211
Within wings, *BarH-1* was not differentially expressed between morphs at this stage, indicating
that morph specific expression differences are temporal. However, we did observe genes
downregulated in Alba wings were significantly enriched for 'xanthine dehydrogenase activity' (p
= 0.02, GO:0004854) (Supplementary Table 7). Xanthine dehydrogenase is the enzyme that
catalyzes the xanthopterin to leucopterin conversion during pteridine synthesis in *Colias*
butterflies[8]. These results are consistent with previous studies in *C. eurytheme* that reported the
level of xanthopterin in Alba wings was 7-8 fold less than in orange[8]. Additionally we observed
enrichment of 'MAP kinase activity' (GO:0004709, p = 0.00109) in genes downregulated within
Alba wings. In *Drosophila*, BarH-1 represses Decapentaplegic a morphogen that is homolog to
TGFβ[27]. TGFβ can activate signalling cascades, including MAP kinase pathways[28]. Previous
work in *Drosophila* has also suggested an interaction between Bar homeobox genes and
Ras/MAP kinase signalling during eye development[29]. Future functional studies of candidate
genes are needed to better understand their mechanistic roles in wing development and the
tradeoffs associated with the ALHS.
226
Here we identified the proximate mechanism underlying a female-limited ALHS in a natural
population. Historically, the field of life history research has treated mechanistic details as a
black box[5], though recently several genetic mechanisms underlying ecologically relevant ALHSs
have been identified, e.g. in the wall lizard[30], ruff[31,32], white throated sparrow[33], and fire ant[34].
The majority of these studies found that supergenes, large loci that maintain many genes in tight
linkage due to structural variation, gave rise to the alternative morphs[31-35]. Such findings
established that structural variation facilitates the evolution of complex traits. However these
genomic architectures make determining the specific contributions of individual genes to ALHSs
difficult, though there have been significant advances made in the white throated sparrow[33].
Interestingly, our results, and recent work in the wall lizard[30], found that ALHSs arose due to

changes in the regulatory region of either a single or two genes, respectively. This raises the

question of how these regions give rise to the other fitness-related traits associated with the

ALHS. We propose the Alba-associated physiological and developmental traits arise due to a

classic Y reallocation model, where reduced pigment granule formation results in reduced

pigment synthesis, which in turn leaves more resources free to be used for other developmental

processes. Previous literature studying the Alba phenotype had been unable to determine

whether allocation of resources from the fatbody, or pigment biosynthesis within the wing

scales, was the basis of Alba. Here, the mosaicism of the BarH-1 KO documents the cell level

autonomy of the Alba polymorphism, as abdomen level provisioning would affect all wing scales

equally. Nevertheless, there may be other pleiotropic effects of the reallocation, such as the

sensitivity of the Alba to viral disease, or of the TE insertion itself as it may affect BarH-1

expression in tissues other than the wing.

Previous work has shown that *BarH-1* plays a role in the morphogenesis of neurons, leg

segments, and eyes in *Drosophila*[36]. Specifically, *BarH-1* expression is required for the

formation of pigment granules and the deposition of red pteridine pigments in the *Drosophila*

eye[17]. We find that *BarH-1* also plays a role in eye and wing color in *Colias* butterflies. However,

as *BarH-1* expression represses the formation of pigment granules within *Colias* wings, we find

it has a reversed function in *Drosophila* and *Colias.* This may be one of several examples where

either whole or a part of a gene regulatory network that regulates eye development has been

co-opted to give rise to a novel trait in the insect wing[37]. If so, future work could investigate what

aspects of the network have been co-opted and how this lead to BarH-1's contrasting roles in

morphogenesis.

Additionally, recent work in the field of butterfly wing evolutionary-development has found that

several genes are repeatedly involved in wing color variation across distantly related species.

Such genes form a patterning "toolkit" (e.g. *optix*[38], *WntA*[39], and *cortex*[40]). *BarH-1* might serve

as another toolkit gene for patterning wing color in butterflies beyond *Colias* as we found BarH-1

expression in scale building and socket cells of developing wings in *Vanessa cardui*

(Nymphalidae, Lepidoptera) (Linnaeus, 1758) pupae (Supplementary Fig. 10). However, the

functional role of BarH-1 in *V. cardui* wings remains to be determined. BarH-1 may have a novel

function within *V. cardui* wings. Alternatively, the function of BarH-1 as a repressor of pigment

granule formation could be conserved, as *V. cardui* scales do not have pigment pigment

granules[41]

Under the latter assumption, we would expect that BarH-1 is not expressed in closely related

Pierinae species that, despite appearing white, exhibit abundant pigment granules that are

primarily filled with the UV-absorbing pteridine called leucopterin[42]. Future work investigating the

evolutionary history of BarH-1's co-option to the wing and function in other species could shed

light on how complex traits such as ALHSs evolve.

Author Contributions

AW conducted butterfly rearings and lab work, analysed the data, and wrote the manuscript with

CWW and input from the coauthors. AW, MWP, KT, and CWW conducted the CRISPR/Cas9

knockout experiment. AW and KT conducted the electron microscopy. MWP conducted

antibody staining. RN and JH assisted with bioinformatics. PL and RK conducted HPTLC and

AW and PL analyzed the data. AW, CS, CWW and OB conducted fieldwork. MC conducted lab

work. CWW supervised the work at all stages.

Acknowledgements

We would like to thank Lovisa Wennerström, Elishia Harji, Jofre Carnicer, and Christina Hansen

Wheat for help with fieldwork. We thank Marianne Ahlbom for assistance with the SEM. Finally

we would like to thank Karin Kiontke, Christen Bossu, Naomi Keehnen, and Peter Pruisscher for

helpful comments on the manuscript. We thank the Department of Zoology at Stockholm

University, the Swedish Research Council 2012–3715, the Academy of Finland 131155, the

Knut and Alice Wallenberg Foundation 2012.0058 and the Erik Philip-Sörensens foundation for

funding.

References

1   Stearns, S. C. *The Evolution of Life Histories*. (Oxford University Press, 1992).
2   Stearns, S. C. Trade-Offs in Life-History Evolution. *Funct Ecol* **3**, 259-268, doi:Doi
10.2307/2389364 (1989).
3   Fisher, R. A. *The Genetical Theory of Natural Selection*. (Oxford University Press,
1930).
4   Gross, M. R. Alternative reproductive strategies and tactics: Diversity within sexes (vol
11, pg 92, 1996). *Trends Ecol Evol* **11**, 263-263 (1996).
5   Flatt, T. & Heyland, A. *Mechanisms of Life History Evolution: The Genetics and
Physiology of Life History Traits and Trade-Offs*. (Oxford University Press, 2011).
6   Remington, C. L. The genetics of Colias (Lepidoptera). *Adv Genet* **6**, 403-450 (1954).
7   Limeri, L. B. & Morehouse, N. I. The evolutionary history of the "alba" polymorphism in
the butterfly subfamily Coliadinae (Lepidoptera: Pieridae). *Biol J Linn Soc* **117**, 716-724,
doi:10.1111/bij.12697 (2016).

8   Watt, W. B. Adaptive Significance of Pigment Polymorphisms in Colias Butterflies .3.
Progress in Study of Alba Variant. *Evolution* **27**, 537-548, doi:Doi 10.2307/2407188
(1973).
9   Descimon, H. & Pennetier, J. L. Nitrogen-Metabolism in Colias-Croceus (Linne) and Its
Alba Mutant (Lepidoptera, Pieridae). *J Insect Physiol* **35**, 881-885, doi:Doi
10.1016/0022-1910(89)90104-2 (1989).
10   Graham, S. M., Watt, W. B. & Gall, L. F. Metabolic Resource-Allocation Vs Mating
Attractiveness - Adaptive Pressures on the Alba Polymorphism of Colias Butterflies. *P*
Natl Acad Sci-Biol **77**, 3615-3619, doi:DOI 10.1073/pnas.77.6.3615 (1980).
11   Woronik, A., Stefanescu, C., Kakela, R., Wheat, C. W. & Lehmann, P. Physiological
differences between female limited, alternative life history strategies: The Alba
phenotype in the butterfly Colias croceus. *J Insect Physiol* **107**, 257-264,
doi:10.1016/j.jinsphys.2018.03.008 (2018).
12   Nielsen, M. G. & Watt, W. B. Behavioural fitness component effects of the alba
polymorphism of Colias (Lepidoptera, Pieridae): resource and time budget analysis.
Funct Ecol **12**, 149-158, doi:DOI 10.1046/j.1365-2435.1998.00167.x (1998).
13   Nielsen, M. G. & Watt, W. B. Interference competition and sexual selection promote
polymorphism in Colias (Lepidoptera, Pieridae). *Funct Ecol* **14**, 718-730, doi:DOI
10.1046/j.1365-2435.2000.00472.x (2000).
14   Hovanitz, W. The biology of Colias butterflies. II. Parallel geographic variation of
dimorphic color phases in North American species. . *Wasmann Journal of Biology* **8**,
197-219 (1950).
15   Boggs, C. L. & Watt, W. B. Population structure of pierid butterflies IV. Genetic and
physiological investment in offspring by male Colias. *Oecologia* **50**, 320-324,
doi:10.1007/BF00344970 (1981).
16   Wiklund, C., Karlsson, B. & Leimar, O. Sexual conflict and cooperation in butterfly
reproduction: a comparative study of polyandry and female fitness. *Proc Biol Sci* **268**,
1661-1667, doi:10.1098/rspb.2001.1719 (2001).
17   Higashijima, S. *et al.* Dual Bar homeo box genes of Drosophila required in two
photoreceptor cells, R1 and R6, and primary pigment cells for normal eye development.
Genes Dev **6**, 50-60 (1992).
18   Nijhout, H. F. *The Development and Evolution of Butterfly Wing Patterns*. (Smithsonian
Institution Press, 1991).
19   Mackenzie, S. M., Howells, A. J., Cox, G. B. & Ewart, G. D. Sub-cellular localisation of
the white/scarlet ABC transporter to pigment granule membranes within the compound
eye of Drosophila melanogaster. *Genetica* **108**, 239-252 (2000).
20   Morehouse, N. I., Vukusic, P. & Rutowski, R. Pterin pigment granules are responsible for
both broadband light scattering and wavelength selective absorption in the wing scales
of pierid butterflies. *P R Soc B* **274**, 359-366, doi:10.1098/rspb.2006.3730 (2007).
21   Rutowski, R. L., Macedonia, J. M., Morehouse, N. & Taylor-Taft, L. Pterin pigments
amplify iridescent ultraviolet signal in males of the orange sulphur butterfly, Colias
eurytheme. *P R Soc B* **272**, 2329-2335, doi:10.1098/rspb.2005.3216 (2005).
22   van Noordwijk, A. J. & de Jong, G. Acquisition and Allocation of Resources: Their
Influence on Variation in Life History Tactics. *The American Naturalist* **128**, 137-142
(1986).
23   Arrese, E. L. & Soulages, J. L. Insect Fat Body: Energy, Metabolism, and Regulation.
Annu Rev Entomol **55**, 207-225, doi:10.1146/annurev-ento-112408-085356 (2010).
24   Watt, W. B. Pteridine biosynthesis in the butterfly Colias eurytheme. *J Biol Chem* **242**,
565-572 (1967).
25   Pavlos, N. J. & Jahn, R. Distinct yet overlapping roles of Rab GTPases on synaptic
vesicles. *Small GTPases* **2**, 77-81, doi:10.4161/sgtp.2.2.15201 (2011).

| 360 | 26 | Stenmark, H. Rab GTPases as coordinators of vesicle traffic. *Nat Rev Mol Cell Biol* **10**, 513-525, doi:10.1038/nrm2728 (2009). |

26   Stenmark, H. Rab GTPases as coordinators of vesicle traffic. *Nat Rev Mol Cell Biol* **10**,
513-525, doi:10.1038/nrm2728 (2009).
27   Kang, J., Yeom, E., Lim, J. & Choi, K. W. Bar represses dPax2 and decapentaplegic to
regulate cell fate and morphogenetic cell death in Drosophila eye. *PLoS One* **9**, e88171,
doi:10.1371/journal.pone.0088171 (2014).
28   Derynck, R. & Zhang, Y. E. Smad-dependent and Smad-independent pathways in TGF-
beta family signalling. *Nature* **425**, 577-584, doi:10.1038/nature02006 (2003).
29   Hayashi, T., Kojima, T. & Saigo, K. Specification of primary pigment cell and outer
photoreceptor fates by BarH1 homeobox gene in the developing Drosophila eye. *Dev*
Biol **200**, 131-145, doi:10.1006/dbio.1998.8959 (1998).
30   Andrade, P. *et al.* Regulatory changes in pterin and carotenoid genes underlie balanced
color polymorphisms in the wall lizard. *Proc Natl Acad Sci U S A* **116**, 5633-5642,
doi:10.1073/pnas.1820320116 (2019).
31   Lamichhaney, S. *et al.* Structural genomic changes underlie alternative reproductive
strategies in the ruff (Philomachus pugnax). *Nat Genet* **48**, 84-88, doi:10.1038/ng.3430
(2016).
32   Kupper, C. *et al.* A supergene determines highly divergent male reproductive morphs in
the ruff. *Nat Genet* **48**, 79-83, doi:10.1038/ng.3443 (2016).
33   Horton, B. M. *et al.* Estrogen receptor alpha polymorphism in a species with alternative
behavioral phenotypes. *Proc Natl Acad Sci U S A* **111**, 1443-1448,
doi:10.1073/pnas.1317165111 (2014).
34   Wang, J. *et al.* A Y-like social chromosome causes alternative colony organization in fire
ants. *Nature* **493**, 664-668, doi:10.1038/nature11832 (2013).
35   Schwander, T., Libbrecht, R. & Keller, L. Supergenes and complex phenotypes. *Curr*
Biol **24**, R288-294, doi:10.1016/j.cub.2014.01.056 (2014).
36   Reig, G., Cabrejos, M. E. & Concha, M. L. Functions of BarH transcription factors during
embryonic development. *Dev Biol* **302**, 367-375, doi:10.1016/j.ydbio.2006.10.008
(2007).
37   Monteiro, A. Gene regulatory networks reused to build novel traits: co-option of an eye-
related gene regulatory network in eye-like organs and red wing patches on insect wings
is suggested by optix expression. *Bioessays* **34**, 181-186, doi:10.1002/bies.201100160
(2012).
38   Zhang, L., Mazo-Vargas, A. & Reed, R. D. Single master regulatory gene coordinates
the evolution and development of butterfly color and iridescence. *Proc Natl Acad Sci U S*
A **114**, 10707-10712, doi:10.1073/pnas.1709058114 (2017).
39   Mazo-Vargas, A. *et al.* Macroevolutionary shifts of WntA function potentiate butterfly
wing-pattern diversity. *P Natl Acad Sci USA* **114**, 10701-10706,
doi:10.1073/pnas.1708149114 (2017).
40   Nadeau, N. J. *et al.* The gene cortex controls mimicry and crypsis in butterflies and
moths. *Nature* **534**, 106-+, doi:10.1038/nature17961 (2016).
41   Dinwiddie, A. *et al.* Dynamics of F-actin prefigure the structure of butterfly wing scales.
Developmental Biology **392**, 404-418, doi:10.1016/j.ydbio.2014.06.005 (2014).
42   Makino, K., Satoh, K., Koike, M. & Ueno, N. Sex in Pieris rapae L. and the pteridin
content of their wings. *Nature* **170**, 933-934 (1952).
43   Gnerre, S. *et al.* High-quality draft assemblies of mammalian genomes from massively
parallel sequence data. *Proc Natl Acad Sci U S A* **108**, 1513-1518,
doi:10.1073/pnas.1017351108 (2011).
44   Chin, C. S. *et al.* Phased diploid genome assembly with single-molecule real-time
sequencing. *Nat Methods* **13**, 1050-1054, doi:10.1038/nmeth.4035 (2016).
45   Wences, A. H. & Schatz, M. C. Metassembler: merging and optimizing de novo genome
assemblies. *Genome Biol* **16**, 207, doi:10.1186/s13059-015-0764-4 (2015).

46   Woronik, A. & Wheat, C. W. Advances in finding Alba: the locus affecting life history and
color polymorphism in a Colias butterfly. *J Evol Biol* **30**, 26-39, doi:10.1111/jeb.12967
(2017).
47   Sedlazeck, F. J., Rescheneder, P. & von Haeseler, A. NextGenMap: fast and accurate
read mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790-2791,
doi:10.1093/bioinformatics/btt468 (2013).
48   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078-2079, doi:10.1093/bioinformatics/btp352 (2009).
49   Kofler, R., Pandey, R. V. & Schlotterer, C. PoPoolation2: identifying differentiation
between populations using sequencing of pooled DNA samples (Pool-Seq).
Bioinformatics **27**, 3435-3436, doi:10.1093/bioinformatics/btr589 (2011).
50   R: A language and environment for statistical computing. (R Foundation for Statistical
Computing, Vienna, Austria, 2019).
51   Kofler, R. *et al.* PoPoolation: a toolbox for population genetic analysis of next generation
sequencing data from pooled individuals. *PLoS One* **6**, e15925,
doi:10.1371/journal.pone.0015925 (2011).
52   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-2158,
doi:10.1093/bioinformatics/btr330 (2011).
53   Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based
linkage analyses. *Am J Hum Genet* **81**, 559-575, doi:10.1086/519795 (2007).
54   Perry, M. *et al.* Molecular logic behind the three-way stochastic choices that expand
butterfly colour vision. *Nature* **535**, 280-284, doi:10.1038/nature18616 (2016).
55   Zaharia, M. *et al.* Faster and More Accurate Sequence Alignment with SNAP. *arxiv*,
doi:arXiv:1111.5572 (2011).
56   Folch, J., Lees, M. & Sloane Stanley, G. H. A simple method for the isolation and
purification of total lipides from animal tissues. *J Biol Chem* **226**, 497-509 (1957).
57   Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat Biotechnol* **29**, 644-652, doi:10.1038/nbt.1883 (2011).
58   Gilbert, D. in *7th annual arthropod genomics symposium* (Notre Dame, 2013).
59   csvkit (2016).
60   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-
140, doi:10.1093/bioinformatics/btp616 (2010).
61   Huerta-Cepas, J. *et al.* Fast Genome-Wide Functional Annotation through Orthology
Assignment by eggNOG-Mapper. *Mol Biol Evol* **34**, 2115-2122,
doi:10.1093/molbev/msx148 (2017).
62   topGO: Enrichment Analysis for Gene Ontology v. R package version 2.34.0 (2018).
448

Methods

For detailed methods, including all bioinformatic commands, please see the supplementary information.

Data availability: SRA reference numbers for the genome and sequencing data will be included upon acceptance.

Genome assembly: An orange female and male carrying Alba (offspring from wild caught butterflies, Catalonia, Spain) were mated in the lab. DNA from an Alba female offspring of this cross was extracted. Quality and quantity were assessed using a Nanodrop 8000 spectrophotometer (Thermo Scientific) and a Qubit 2.0 fluorometer (dsDNA BR, Invitrogen). A

180 insert size paired end library (101bp reads) was prepared (TruSeq PCR free) and
sequenced on an Illumina Hiseq 4000 at the Beijing Genomics Institute (Shenzhen, China). A
Nextera mate-pair library with a 3 kb insert size was prepared and sequenced on an Illumina
HiSeq 2500 (125bp reads) at the Science for Life Laboratory (Stockholm, Sweden). Raw data
was cleaned and high quality reads were used as input for the AllPaths-LG (v. 50960)[43]
assembly pipeline. High molecular weight DNA was extracted from two more Alba females from
the above mentioned cross (i.e full siblings). Equal amounts of DNA from each individual were
pooled sent to the Science for Life Laboratory (Stockholm, Sweden) for PacBio sequencing on
24 SMRT cells (~17GB of data was produced). A Falcon (v. 0.4.2)[44] assembly was generated by
the Science for Life Laboratory. We then used Metassembler (v. 1.5)[45] to merge our AllPathsLG
and Falcon assemblies, using the AllPathsLG assembly as the primary assembly.

Bulk segregant analyses (BSA): The female informative cross data and mapping protocol
described in Woronik and Wheat, 2017[46] was applied to the high quality reference genome to
identify the contigs that made up the Alba chromosome. Male Informative Cross (MIC) I: DNA
was extracted from a wild caught orange mother (Catalonia Spain) and 26 of her Alba and 24 of
her orange female offspring. DNA quality and quantity of each individual was assessed via a
Nanodrop 8000 spectrophotometer (Thermo Scientific, MA, USA) and a Qubit 2.0 Fluorometer
(dsDNA BR; Invitrogen, Carlsbad, CA, USA) before pooling equal amounts of high-quality DNA
from Alba and orange offspring into two pools, respectively. The library preparation (TruSeq
PCR-free) and Illumina sequencing (101 bp PE HiSeq2500), was performed at the Beijing
Genomics Institute (Shenzhen, China). Raw reads were cleaned and then mapped to the
reference genome using NextGenMap v0.4.10 (-i 0.09)[47]. SAMTOOLS v1.2[48] was used to filter
(view -f 3 -q 20), sort and index the bam files and generate mpileup files for the two pools and
the orange mother. Popoolation2[49] were used to calculate the allele frequency difference
between Alba and orange pools. SNP sites were filtered in R[50], for a read depth ≥ 30 and ≤ 300,
a bi-allelic state, and a minimum minor allele frequency of 3. The orange mother mpileup was
similarly analyzed using Popoolation[51] (read depth ≥ 5 and ≤ 30); but the major and minor allele
frequencies were calculated in R[50] by dividing the major and minor allele count by the read
depth at each site respectively. A SNP site was considered a MIC I Alba SNP when it met the
following expectations: 1) homozygous in the orange mother, 2) homozygous in the orange
pool, 3) the allele frequency difference in the Alba pool compared to the orange was 0.45-0.55.
MIC II: A male carrying Alba mated an orange female in the lab at Stockholm University. DNA
was prepared as described above for 26 Alba and 28 orange female offspring resulting in two
DNA pools. Library preparation (TruSeq PCR-free) and Illumina sequencing (150 bp paired-end
reads with 350bp insert, HiSeqX), was performed at Science for Life Laboratory (Stockholm,
Sweden). The same mapping and SNP calling pipeline used on the MIC I was applied. A site
was considered an Alba SNP if 1) it was homozygous in the orange pool and 2) the allele
frequency difference in the Alba pool compared to the orange was 0.45-0.55. A contig was
considered Alba associated if it had ≥ 3 Alba SNPs in all crosses. Nineteen Alba associated
contig were identified. They total ~3.7Mbp and are considered the Alba BSA locus.

Genome wide association study: DNA for genome re-sequencing was extracted from 15 Alba
and 15 orange females from diverse population backgrounds (Catalonia, Spain and Capri, Italy).
High quality DNA was prepared using Illumina TruSeq and sequenced at the Science for Life
Laboratory (Stockholm, Sweden) (150 bp paired-end reads HiSeqX). Cleaned reads were
mapped to the annotated reference genome using NextGenMap v0.4.10 (-i 0.6 -X 2000)[47]. Bam
files were filtered and sorted using SAMTOOLS v1.2 (view -f 3 -q 20) [48]. A VCF file was
generated using SAMTOOLS v1.2 (-t DP -t SP -Q 15)[48] and bcftools v.1.2 (-Ov -m) [48]. VCFtools
(v0.1.13)[52] was used to call SNP sites with no more than 50% missing data, an average read
depth between 15-50 across individuals, and a minimum SNP quality of 30. An association

analysis was performed with PLINK (v1.07)[53] and a Benjamini & Hochberg step-up FDR control
was applied. SNPs with FDR <0.05 were considered Alba SNPs. We conducted this analysis
both genome wide and only within the BSA locus. Both analyses fine mapped the Alba locus to
the same genomic region.

Antibody Generation and Staining: A Rabbit-anti-Bar antibody was generated against the full
length sequence of the *Vanessa cardui* Bar homolog. Protein was generated by GenScript
(Piscataway, NJ) and purified to >80% purity. DNA sequences to produce this protein were
codon-optimized for bacterial expression and made via gene synthesis. GenScript injected
resultant protein into host animals, collected serum for testing, and affinity purified the product
using additional target protein bound to a column. Antibody staining was performed as
described previously for Drosophila and butterfly tissues[54]. Staged pupal wings and retinas were
dissected and fixed 48 hours post-pupation. The Rabbit-anti-Bar antibody was used at 1:100,
followed by secondary antibody staining with AlexFluor-555-anti-Rabbit secondaries at 1:500
and counterstaining with DAPI. Images were captured using standard confocal microscopy on a
Leica SP5.

CRISPR/Cas9 knockouts: The guide-RNA (gRNA) sequences were generated using the
protocol described in Perry et al. 2016. Viable Cas9 target-sites were located by manually
looking for PAM-sites (NGG) in the exon region of *BarH-1*. Uniqueness of the target regions was
confirmed using a NCBI nucleotide blast (ver. 2.5.0+ using blastn-short flag and filtering for an
e-value of 0.01) against the *C. crocea* reference genome. gRNA constructs were ordered from
Integrative DNA Technologies (Coralville, Iowa, USA) as DNA (gBlocks). Full gRNA constructs
had the following configuration: an M13F region, a spacer sequence, a T7-promotor sequence,
the Target specific sequence, a Cas9 binding sequence, and finally a P505 sequence. Upon
delivery, gBlocks were amplified using PCR to generate single-stranded guide RNA (sgRNA).
For each gBlock, four 50ul reactions were conducted using the M13f and P505 primers and
Platinum Taq (Invitrogen cat. 10966-034). The four reactions were then combined and purified
in a Qiagen Minelute spin column (cat. 28004, Venlo, Netherlands). The resulting template was
transcribed using the Lucigen AmpliScribe T7-flash Transcription Kit from Epicentre/Illumina
(cat. ASF3507, Madison, WI, USA) followed by purification via ammonium acetate precipitation.
Products were resuspended with Qiagen buffer EB, concentrations were quantified by Qubit and
further diluted to 1000 ng/µl. They were then mixed with Cas9-NLS protein (PNA Bio, Newbury
Park, CA, USA) and diluted to a final concentration of 125-250 ng/µl. *C. crocea* females (n > 40)
from Aiguamolls de l'Empordà, Spain were captured and kept in morph-specific flight cages in
the lab at Stockholm University where they oviposited on alfalfa (*Medicago sativa*). Eggs were
collected between 1-7h post-laying and sterilized in 7% benzalkonium chloride for ~5 minutes
before injection. Injections were either at a concentration of 125 or 250 ng/ul and conducted
using a M-152 Narishige micromanipulator (Narishige International Limited, London, UK) with a
50 ml glass needle syringe, with injection pressure applied by hand via a syringe fitting.

CRISPR/Cas9 validation: To validate the mutation, Cas9 cut sites were PCR-amplified and a
~370bp region, centered on the intended cut site were sequenced using Illumina MiSeq 300bp
paired-end sequencing. Primers were designed using Primer3
(http://biotools.umassmed.edu/bioapps/primer3_www.cgi). DNA was isolated from KO-
individuals using KingFisher Cell and Tissue DNA Kit from ThermoFisher Scientific (N11997)
and the robotic Kingfisher Duo Prime purification system. DNA quality and quantity were
assessed via a Nanodrop 8000 spectrophotometer (Thermo Scientific, MA, USA) and a Qubit
2.0 Fluorometer (dsDNA BR; Invitrogen, Carlsbad, CA, USA). Aliquots were then taken and
diluted to 1ng/ul before amplifying the region over the cleavage-site. Sequences were amplified
and ligated with Illumina adapter and indexes in a two-step process following the protocol

provided by Science for Life Laboratories (Stockholm, Sweden) and Illumina. First, we amplified
the ~370bp long sequence around the cut sites and attach the first Illumina adapter, onto which
we later attach Illumina handles and Index using a second round of PCR (Accustart II PCR
Supermix from Quanta Bio [Beverly, MA, USA], settings 94C x 2 min followed by 40 cycles of 94
C x 30 sec + 60 C x 15 sec + 68 C x 1 min followed by 68 C x 5 min). PCR products were
purified using Qiagen Qiaquick (Cat. 28104). Concentration and quality of the product were
assessed via Nanodrop and gel electrophoresis. DNA was diluted to ~0.5ng/ul and then the
unique double indices were attached by the second round of PCR (same protocol as above).
The final PCR products were purified again using Qiaquick spin columns and concentration and
size was assessed using Qubit fluorometer and gel electrophoresis. All samples were then
mixed at equal molarity and sent for sequencing at Science for Life Laboratories (Stockholm,
Sweden). Sequences were aligned to their respective fragments (area surrounding cut site)
using SNAP (ver. 1.0beta18)[55], identical reads were clustered using the collapser utility in Fastx-
Toolkit (http://hannonlab.cshl.edu/fastx_toolkit/). Sequences containing deletions were extracted
and the most abundant sequences containing deletions were selected for confirmation of
deletion in the expected region.

Electron Microscopy: To quantify pigment granule differences between Alba and orange
individuals pieces of the forewing were mounted on aluminum pin stubs (6mm length) with the
dorsal side upwards. Samples were coated in gold for 80 seconds using an Agar sputter coater
and imaged under 5 kV acceleration voltage, high vacuum, and ETD detection using a scanning
electron microscope (Quanta Feg 650, FEI, Hillsboro, Oregon, USA). To quantify pigment
granules within the photos we selected images from the same magnification and randomly
placed three 4 μm^2 squares on the images. We counted the number of pigment granules within
each square and took the average, then conducted a two sample t-test in R. To quantify
pigment granule differences between KO and wild type regions in our CRISPR KO mosaic
individuals, a biopsy hole punch a 2mm in diameter circle was used to cut out one piece mostly
containing white scales and one piece with mostly orange scales. These pieces were first
photographed using a Leica EZ4HD stereo microscope in order to allow us to confirm the color
of each scale once they were covered with gold sputter. Five white and five orange scales were
then selected and the granules from a 4μm^2 square were counted from each of those scales
and a two sample t-test was then conducted in R.

Lipid Analysis: Wild caught *C. crocea* Alba females (Catalonia, Spain) oviposited in the lab.
Eggs were moved into individual rearing cups and split between two temperature treatments
(hot: 27^{o}C and 16 hour day length during larval and pupal development, cold: reared at 22 ^{o}C
with a 16 hour day length during larval development and 15^{O}C with a 16 hour day length during
pupal development). Once pupated, individuals were checked a minimum of every 12 hours.
Upon eclosion adults were stored at 4 ^{o}C until the next day to provide time for meconium
excretion. Butterflies were not allowed to feed before dissection. Body weight was taken using a
Sauter RE1614 scale before dissection. Total lipids were extracted using the Folch method[56]
according to the procedures outlined in Woronik et. al. 2018[11]. HPTLC was conducted as
described in Woronik et. al. 2018[11]. In brief, 5 μl of the sample lipid extract was applied on a
silica plate with a Camag Automatic TLC Sampler 4 (Camag, Muttenz, Switzerland). After the
silica plate developed it was scanned with a Camag TLC plate scanner 3 at 254 nm using a
deuterium lamp with a slit dimension of 6 × 0.45 mm and analyzed with the Win-CATS 1.1.3.0
software. Peaks representing the four major neutral lipid classes (diacylglycerols,
triacylglycerols, cholesterol and cholesterol esters) were identified by comparing their retention
times against known standards. Then the peak areas were integrated and the amount of lipid
within each class was calculated using the formula: $pmol_{sample}$ = ($Area_{sample}$ / $Area_{standard}$) x
$pmol_{standard}$. The total lipid content (nmol per abdomen) was calculated as a sum of pmol

contents of all neutral lipid classes. For the statistical analyses this value was regressed against
abdomen weight and standardized residuals (i.e. mass-corrected storage lipid amount) and
were subsequently used as dependent variable.

Transcriptome assembly and differential expression analysis: Offspring from a wild caught
Alba female from Catalonia, Spain were reared at Stockholm University. When larvae reached
the fifth instar they were checked at least every six hours and the pupation time of each
individual was recorded. Tissue was collected between 82% and 92% of pupal development.
Pupae were dissected in PBS solution, and the abdomen and wings were flash frozen in liquid
nitrogen and stored at -80 °C. RNA was extracted from the abdomen and wing tissues using
Trizol. RNA quality and quantity was assessed using a Nanodrop 8000 spectrophotometer
(Thermo Scientific) and an Experion electrophoresis machine using the manufacturer protocol
(Bio-Rad, Hercules, CA). Library preparation (Strand-specific TruSeq RNA libraries using poly-A
selection) and sequencing (101 bp PE HiSeq2500 - high output mode) was performed at the
Science for Life Laboratories (Stockholm, Sweden). In total 16 libraries were sequenced (4
orange and 4 Alba individuals - wings and abdomen from each individual). Raw data was
cleaned and reads from all libraries were used in a de novo transcriptome assembly (Trinity
version trinityrnaseq_r2013_08_14 with default parameters)[57]. To reduce the redundancy
among contigs and produce a biologically valid transcript set, the tr2aacds pipeline from the
EvidentialGene software package[58] was run on the raw Trinity assembly. The sixteen RNA-Seq
libraries were mapped to the resulting transcriptome using NextGenMap v0.4.10 (-i 0.09)[47].
SAMTOOLS v1.2[48] was then used to filter (view -f 3 -q 20), sort and index the sixteen bam files.
SAMTOOLS v1.2[48] idxstats was then used to calculate the read counts per gene for each of the
sorted bam files. These counts were then joined in a CSV file using an in-house pipeline and
csvjoin[59]. A differential expression analysis was conducted in EdgeR[60]. A Benjamini Hochberg
correction was applied to the raw p values to correct for false discovery rate and differentially
expressed genes were called (adjusted p value <0.05). eggNOG-mapper (v.1)[61] was used with
default settings to functionally annotate the transcriptome. The R package topGo[62] was used to
conduct a gene set enrichment analysis on genes that exhibited > 1 or < -1 log fold change in
the differential expression analysis.

Fig. 1. Color variation in *Colias crocea* and the genetic mechanism of Alba. (A) *Colias crocea* male, orange female, and Alba female (left to right). (B) SNPs significantly associated with the Alba phenotype (red) within the ~3.7 Mbp Alba locus identified via 3 rounds of bulk segregant analysis. Contigs in this region shown as alternating dark and light blue. (C) The location of Alba associated SNPs (red) on the ~430 kb outlier contig identified in the GWAS. Gene models for the DEAD-box helicase*,* the Jockey-like transposable element, and *BarH-1* shown at the top of the panel. (D) Wings of a female with an Alba genotype following CRISPR/Cas9 mosaic knockout of *BarH-1*, wild type regions are white, knockout regions are orange. Orange color is seen on the dorsal forewing (top) and hindwing (bottom). (D) *BarH-1* mosaic knockout also leads to black regions in the eyes, wild type regions are green.

Fig. 2. BarH-1 is expressed in white but not orange regions of the wing in *C. crocea* females. DAPI (nuclei, left, blue) and BarH-1 antibody (right, red) staining of pupal wings. Large nuclei are in scale building cells, small nuclei are in epithelial cells. The right part of the panel shows the approximate location of the stained area and the scales in this region in an adult wing. Scale bars are 2µm. (A) Staining of the forewing of an Alba female in an area at the black margin (top) and a white area (bottom). BarH-1 is expressed in melanic as well as white Alba scale building cells. (B) Antibody staining of the forewing of an orange female. BarH-1 is not expressed in these scale building cells. (C) Antibody staining of the hindwing of an orange female. BarH-1 is heterogeneously expressed in the scale building cells within this region. This staining pattern presumably corresponds to the variation in scale color, with melanic scale building cells expressing BarH-1 but orange lacking expression.

Fig. 3. *Colias* forewings and scanning electron microscope (SEM) images of their wing scale nanostructures. (A) *C. crocea* Alba female wing and wing scale structure. The top panel shows the SEM image of a black scale; pigment granules are absent. The bottom panel shows a white scale, exhibiting near absence of pigment granules. (B) Wing and wing scale structures of a wild type orange *C. crocea* female. The top panel shows a black scale, pigment granules are absent. The bottom panel shows an orange scale with abundant pigment granules. (C) Wing and wing scales of a genetically Alba female exhibiting CRISPR/Cas9 mosaic knockout of *BarH-1*. The top panel shows the wild-type white scale, where pigment granules are mostly absent. The bottom panel shows a scale in an orange *BarH-1* KO region. It exhibits significantly more pigment granules than the white scales. (D) Wing and wing scales of an orange *C. crocea* female where pigment granules have been chemically removed from the distal half of the wing. The SEM image shows a scale from the white region with pigment granules completely missing. The white color of this wing section presumably results from light reflection off the remaining scale nanostructures. (E) Wing and wing scale structure of a *C. eurytheme* Alba female. Wing scales lack pigment granules, similar to the phenotype observed in *C. crocea*. (F) Wing and wing scale structures of a *C. eurytheme* orange female. Orange scales show abundant pigment granules, again consistent with the orange phenotype observed in *C. crocea*.

Fig. 4. Physiological differences between female morphs of *C. crocea*. A) The mass corrected total neutral lipid content for female morphs in two temperature treatments. Alba females, on average, have larger neutral lipid stores than orange females. However there is an interaction between morph and temperature as the difference is only significant in the cold treatment. Error bars are the standard error (cold: n=32, $t_{29.12}$ = 3.42, P = 0.002, hot: n=25, $t_{22.71}$ = 0.67, P = 0.51). B) Volcano plot to visualize gene expression differences between female morphs in pupal abdominal tissue. Each point is a gene. Genes not significantly differentially expressed between morphs are grey, while differentially expressed genes are blue. The black square is the triacylglycerol lipase and the black triangle is *RIM*. The X-axis is the log of the fold change (FC), positive log(FC) indicates the gene is upregulated in Alba individuals. C) Volcano plots to visualize gene expression differences between female morphs in pupal wing tissue. Color coding, shapes, and axes are the same as above.

Figure 1

Bar-H1 expression in scale building cells of developing pupal wings

Corresponding region of adult wing

A black DAPI BarH-1

B white DAPI BarH-1

Alba forewing

C orange DAPI BarH-1

D orange and black DAPI BarH-1

orange forewing

orange hindwing

Figure 2

Alba genotype

orange genotype

C. crocea

C. eurytheme

A

B

C

D

E

F

Figure 3

Figure 4