A homeobox gene, *BarH-1*, underlies a female alternative life-history strategy

Alyssa Woronik*[1], Kalle Tunström[1], Michael W. Perry[2], Ramprasad Neethiraj[1], Constanti Stefanescu[3,4], Maria de la Paz Celorio-Mancera[1], Oskar Brattström[5], Jason Hill[1], Philipp Lehmann[1], Reijo Käkelä[6], Christopher W. Wheat*[1]

Author affiliations:

Department of Zoology, Stockholm University, S106 91 Stockholm, Sweden

Department of Biology, New York University, New York, New York 10003, USA

Museum of Natural Sciences of Granollers, Granollers, Catalonia 08402, Spain

CREAF, Cerdanyola del Valles, Catalonia 08193, Spain

Department of Zoology, University of Cambridge, Cambridge CB23EJ, United Kingdom

Helsinki University Lipidomics Unit, Helsinki Institute for Life Science (HiLIFE) and Molecular and Integrative Biosciences Research Programme, University of Helsinki, FI00014 Helsinki, Finland

Corresponding authors:

AW alyssa.woronik@zoologi.su.se and CWW chris.wheat@zoologi.su.se

Colias butterflies (the "clouded sulphurs") often occur in mixed populations where females exhibit two color morphs, yellow/orange or white. White females, known as the Alba morph, reallocate resources from the synthesis of costly colored pigments to reproductive and somatic development [1]. Due to this tradeoff Alba females develop faster and have higher fecundity than orange females [2]. However orange females, that have instead invested in pigments, are preferred by males who in turn provide a nutrient rich spermatophore during mating [2,3,4]. Thus the wing color morphs represent alternative life history strategies (ALHS) that are female-limited, wherein tradeoffs, due to divergent resource investment, result in distinct phenotypes with associated fitness consequences. Here we map the genetic basis of Alba in *Colias crocea* to a transposable element insertion downstream of the *Colias* homolog of *BarH-1.* To investigate the phenotypic effects of this insertion we use CRISPR/Cas9 to validate *BarH-1*'s functional role in the wing color switch and antibody staining to confirm expression differences in the scale building cells of pupal wings. We then use scanning electron microscopy to determine that *BarH-1* expression in the wings causes a reduction in pigment granules within wing scales,

and thereby gives rise to the white color. Finally, lipid and transcriptome analyses reveal
additional physiological differences that arise due to Alba, suggesting pleiotropic effects beyond
wing color. Together these findings provide the first well documented mechanism for a female
ALHS and support an alternative view of color polymorphism as indicative of pleiotropic effects
with life history consequences.

Evolutionary theory predicts that positive selection will remove variation from natural
populations, as genotypes with the highest fitness go to fixation [5]. However, across diverse taxa
ALHS are maintained within populations at intermediate frequencies due to balancing selection
[6]. Modelling and mechanistic insights have advanced our understanding of ALHS evolution and
maintenance (e.g. negative - frequency dependent selection)[7]. However, the majority of
studies, and consequently our insights, are biased toward male strategies that are
morphologically dramatic (e.g. ruff [8,9] and side-blotched lizards [10]). Whether this bias reflects
true differences in the frequency of alternative strategies between the sexes or is simply an
artifact is unknown [11]. As trade-offs and selection regimes are often sex specific, the lack of
female insights severely limits our understanding of the mechanisms, maintenance, evolution,
and co-evolution of alternative strategies in general [11]. Yet despite calls for further investigation
[11], a well documented mechanism for a female limited ALHS has yet to be identified. Here we
identify one such mechanism in the butterfly *Colias crocea* (Pieridae).

Approximately a third of the nearly 90 species within the butterfly genus *Colias* exhibit a female-
limited ALHS known as Alba[12]. The switch between strategies is controlled by a single,
autosomal locus that causes Alba females to reallocate guanosine triphosphate (GTP),
amounting to several percent of their nitrogen budget, from the synthesis of pteridine pigments
to other areas of development [1]. Consequently, Alba females have white wings, while non-Alba
females are orange/yellow (Fig. 1A). As a result of this trade-off, Alba females gain fitness
advantages over orange females due to faster pupal development, a larger fat body, and
significantly more mature eggs at eclosion [2]. Despite these developmental advantages and the
dominance of the Alba allele, females remains polymorphic due to tradeoffs in abiotic and biotic
factors [2,13-15]. For example, Alba's development rate advantage is higher only in cold
temperatures, also as a result of density-dependent, interference competition with other white
Pierid species and sexual selection, males preferentially mate with orange females [2,3,13,14]. The
mating bias likely has significant fitness costs for Alba because males transfer essential
nutrients during mating and multiply mated females have more offspring over their lifetime [4,16].

Field studies confirm Alba frequency and fitness increases in species that inhabit cold and
nutrient poor habitats, where the occurrence of other white Pierid butterflies is low, while in
warm environments with nutrient rich host plants and a high co-occurrence with other white
species, orange females exhibit increased fitness and frequency [3,14].

Using a *de novo* reference genome for *C. crocea* that we generated via Illumina and PacBio
sequencing (Extended Data Table 1), and three rounds of bulk segregant analyses (BSA) using
whole genome sequencing of a female and two male informative crosses for Alba, we mapped
the Alba locus to a ~3.7 Mbp region (Extended Data Fig.1, & Supplementary Information). Then,
with whole genome re-sequencing data from 15 Alba and 15 orange females from diverse
population backgrounds, a SNP association study was able to fine map the Alba locus to a
single ~430 kb contig that fell within the ~3.7 Mbp BSA locus (Fig. 1B) (Supplementary
Information). The majority of SNPs significantly associated with Alba (n=70 of 72) were within or
flanking a *Jockey-like* transposable element (TE) (Fig. 1B & 1C). We determined the TE
insertion is unique to the Alba morph in *C. crocea* by assembling the orange and Alba
haplotypes for this region, then quantifying differences in read depth between morphs within and
flanking the insertion, and comparing the region to other butterfly genomes (*Danaus plexippus* &
Heliconius melpomene) (Extended Data Figs.2 & 3). Additionally we validated the presence and
absence of the insertion, respectively, across 82 wild females, 25 Alba and 57 orange
(Extended Data Fig.4).

The Alba specific TE insertion was located ~30 kb upstream of a *DEAD-box helicase*, and ~6kb
downstream of *BarH-1*, a homeobox transcription factor (Fig. 1C). *BarH-1* was an intriguing find
as its knockout in *Drosophila melanogaster* causes a dramatic decrease in pigment granules
within the eye, changing eye color from red (wild type) to white [17]. To validate the functional role
of *BarH-1* in the Alba phenotype we generated CRISPR/Cas9-mediated deletions of exons 1
and 2 in a mosaic knockout approach (Extended Data Fig.5 & Supplementary Information).
BarH-1 deletions gave rise to a mosaic lack of pigmentation in the eyes of males and females of
both morphs, consistent with *BarH-1*'s expected role in insect eye development (Fig. 1D).
Additionally, on the dorsal side of the wings, females with an Alba genotype exhibited a
white/orange color mosaic, while males and orange females displayed no wing KO phenotypes,
despite those individuals exhibiting mosaic phenotypes in the eye. (Fig. 1E).

To further investigate the role of BarH-1 in developing wing scales, we used *in situ* hybridization

of BarH-1 on wings from 2 day old pupae of orange and Alba females of *C. crocea*, as well as

Vanessa cardui. The BarH-1 protein is highly expressed in the scale building cells of both

species (Fig. 2), suggesting a previously undescribed role of BarH-1 in the developing wing

scales of butterflies. Comparison between orange and Alba females of *C. crocea* further

documents Alba as a gain of BarH-1 function, as scale building cells in the developing wing

show a BarH-1 expression pattern that is Alba limited (Fig. 2).

In butterflies both pigments and scale morphology can affect wing color [18] and while Alba

females exhibit large reductions in colored pteridine pigments compared to orange [1], whether

morphs differed in wing scale morphology was unknown. Using scanning electron microscopy

we found Alba scales exhibited significantly less pigment granules, the structures that store

pteridine pigments in Pierid butterflies [19], compared to orange ($t_{5.97}$ = 2.93, p = 0.03), suggesting

reduced granule formation as the basis of the Alba color change (Fig. 3 A & B). As expected,

the number of pigment granules were also significantly reduced in the white regions of the

CRISPR/Cas9 *BarH-1* KO individuals ($t_{5.45}$ = 10.78, p < 0.001) (Fig. 3C & D), demonstrating that

BarH-1 is affecting pigment granule formation to give rise to Alba. To further test whether

reduction in pigment granule number alone was sufficient for the orange/white color change, we

chemically removed the pigment granules from the wing of an orange *C. crocea* female,

resulting in formerly orange regions turning white likely due to the scattering of light from

remaining non-lamellar microsctructures on the wing (Fig. 3 E) [20]. Thus, the white wings of Alba

C. crocea (Fig. 1A & 3A) differ from other white Pierid species, as the latter exhibit abundant

pigment granules in their scales (Fig. 3F and Extended Data Fig.6), documenting that there are

multiple routes to white wing color in Pieridae.

We next tested whether the physiological tradeoffs of Alba reported for New World species [1,2],

which were discussed in the introduction, were also seen in *C. crocea*, an Old World species, as

shared tradeoffs would suggest the Alba mechanism is conserved genus wide. To compare

abdominal lipid stores between morphs, we conducted high performance thin layer

chromatography on 2 day old adult females reared under two temperature treatments (Hot:

27^{o}C vs. Cold: 15^{o}C during pupal development). We found Alba females had larger abdominal

lipid stores than orange in both temperature treatments, though the difference was only

significant in the cold treatment (cold: n=32, $t_{29.12}$ = 3.42, P = 0.002, hot: n=25, $t_{22.71}$ = 0.67, P =

0.51) (Fig. 4A), consistent with the known effects of temperature on Alba fitness in New World
Colias species [2].

We then conducted RNASeq on pupal wing and abdomen tissue, at the time of pteridine
synthesis (i.e. when allocation tradeoffs are realized) to identify genes that exhibited differential
expression between morphs (Fig. 4B,C, Supplementary Information). We found that *vitellogenin*
1, which encodes an egg yolk precursor protein synthesized in the fat bodies of insects [21], was
significantly upregulated within Alba abdomen tissue (log fold change [log FC] of 4.8) (Fig. 4B).
Additionally, consistent with previous reports of GTP reallocation in Alba females [1], *RIM*, a Rab
GTPase effector [22], was one of the most highly differentially expressed (DE) genes in both
tissues (logFC increase in Alba of 3.4 in the abdomen and 5.1 in the wings) (Fig. 4B,C). RIM
acts as a molecular switch by converting guanosine diphosphate to GTP, thereby activating its
associated Rab GTPase, which is in turn involved in synaptic vesicle exocytosis and secretory
pathways [23]. These results are consistent with previous qualitative findings of Alba females in
the North American species *C. eurytheme* (Alba females have a larger fat body, emerge from
the pupa with significantly more mature eggs, and reallocate GTP from pigment synthesis to
somatic development [1,2]). Our findings thus quantitatively demonstrate that the trade-offs
associated with the Alba ALHS are likely consistent across the *Colias* genus, suggesting that
Alba may be due to the same genetic mechanism and corroborating previous work that
proposed Alba is homologous across *Colias* [12].

Gene set enrichment analyses identified 85 functional categories that were significantly
enriched in the abdomen tissue of Alba females (Extended Data Fig.7), notably downregulation
of 'positive regulation of GTPase activity' (adjusted p value < 0.0001), 'regulation of Notch
signalling pathway' (adjusted p value = 0.03), and 'canonical Wnt signalling' (adjusted p value <
0.01). While the Wnt pathway is known to regulate wing patterns in several butterfly species
[24,25], these findings are curious as they are observed in abdomen tissue rather than wing,
suggesting potential unexplored pleiotropic effects of these pathways outside of the wing. In
wing tissue, 35 functional categories were significantly enriched and downregulated in Alba
including 'regulation of transcription' (adjusted p value < 0.0001) and 'positive regulation of
GTPase activity' (adjusted p value < 0.0001), while 'protein catabolic process' (adjusted p value
< 0.0001) was upregulated (Supplementary Information). *BarH-1* was not DE between morphs
in our RNASeq data, suggesting that morph specific expression differences are temporal and

likely occur earlier in development. Further functional studies of candidate genes are needed to
better understand their mechanistic roles in the trade-off.

Here we report that the genetic basis of a female-limited ALHS arises from the co-option of the
homeobox transcription factor *BarH-1*, primarily known for its role in the morphogenesis of the
insect eye, neurons and leg segments [26]. We document that *BarH-1* has a similar function in
eye morphogenesis of butterflies, and also find it is expressed during wing scale development in
butterflies from the families Pieridae and Nymphalidae, which last shared a common ancestor
over 70 million years ago. This novel finding suggests a conserved function of *BarH-1* in scale
morphogenesis that warrants further study and suggests a parsimonious route to *BarH-1*'s gain
of function in the ALHS Alba phenotype, with co-option from a role in wing scale development
rather than its previously described functions. *BarH-1*'s well characterized role in determining
cell fate through gene repression [27] suggests it is involved in the repression of pigment granule
formation, providing an explanation for the Alba allele being dominant and a gain of function that
results in the absence of a phenotype (i.e. orange wing color). To what extent *BarH-1* has an
active pleiotropic role in other tissues or developmental stages remains to be determined, as the
extensive physiological responses we document could easily arise from a simple reallocation
following the absence of pigment granule formation. Given the emergence of "toolkit" genes for
butterfly wing patterning, wherein specific genes have been found to be repeatedly involved in
wing color variation across distant species (e.g. *cortex* [28]), determining to what extent *BarH-1* is
involved in other wing phenotypes and ALHS is of interest, especially given the pleiotropic
effects on life history documented here. Finally, our results and others (e.g. side-blotched lizards
[29] & damselflies [30]) suggest that investigating to what extent ALHS are associated with color
variation in other systems is warranted, especially in cases where such variation is female
limited.

Author Contributions
AW conducted butterfly rearings and lab work, analysed the data, and wrote the manuscript with
CWW and input from the coauthors. MWP, KT, CWW, and AW conducted the CRISPR/Cas9
knockout experiment. AW and KT conducted the electron microscopy. MWP conducted
antibody staining. RN and JH assisted with bioinformatics. PL and RK conducted HPTLC and
PL and AW analyzed the data. AW, CS, CWW and OB conducted fieldwork. MC conducted lab
work. CWW supervised the work at all stages.

Acknowledgements

We would like to thank Lovisa Wennerström, Elishia Harji, Jofre Carnicer, and Christina Hansen Wheat for help with fieldwork. We thank Marianne Ahlbom for assistance with the SEM. Finally we would like to thank Christen Bossu, Naomi Keehnen, and Peter Pruisscher for helpful comments on the manuscript. We thank the Department of Zoology at Stockholm University, the Swedish Research Council 2012–3715, the Academy of Finland 131155, the Knut and Alice Wallenberg Foundation 2012.0058 and the Erik Philip-Sörensens foundation for funding.

References

1. Watt, W.B. Adaptive significance of pigment polymorphisms in *Colias* butterflies. III. Progress in the study of the "alba" variant. Evolution **27**: 537–548 (1973).

2. Graham, S.M.S., Watt, W.B., & Gall, L.F.L. Metabolic resource allocation vs. mating attractiveness: Adaptive pressures on the "alba" polymorphism of *Colias* butterflies. Proc Natl Acad Sci U S A **77**: 3615–3619 (1980).

3. Nielsen, M.G. & Watt, W.B. Interference competition and sexual selection promote polymorphism in *Colias* (Lepidoptera, Pieridae). Functional Ecology **14**: 718–730 (2000).

4. Boggs, C.L. & Watt, W.B. Population structure of pierid butterflies IV. Genetic and physiological investment in offspring by male *Colias*. Oecologia **50**: 320–324 (1981).

5. Fisher, R.A. The Genetical Theory of Natural Selection. Oxford University Press (1930).

6. Gross, M.R. Alternative reproductive strategies and tactics: diversity within sexes. Trends Ecol Evol. **11**: 92-98 (1996).

7. Hofbauer, J. & Sigmund, K. Evolutionary Game Dynamics. Bulletin of the American Mathematical Society **40**: 479–519 (2003).

8. Lamichhaney, S. *et al.* Structural genomic changes underlie alternative reproductive strategies in the ruff (*Philomachus pugnax*). Nat Genet **48**: 84–88 (2015).

9. Lank, D.B., Smith, C.M., Hanotte, O., Burke, T. & Cooke, F. Genetic polymorphism for alternative mating-behavior in lekking male ruff *Philomachus pugnax*. Nature **378**: 59–62 (2015).

10. Sinervo, B., & Lively, C. M. The rock-paper-scissors game and the evolution of alternative male reproductive strategies. Nature **380**: 240–243 (1996).

11. Alonzo SH. Oliveira RF, Taborsky M, & Brockmann HJ. Conflict between the sexes and alternative reproductive tactics within a sex, Alternative reproductive tactics: an integrative approach, 2008 Cambridge (UK) Cambridge University Press (pg. 435-450).

12. Limeri, L.B. & Morehouse, N.I. The evolutionary history of the "alba" polymorphism in the
butterfly subfamily Coliadinae (Lepidoptera: Pieridae). Biological Journal of the Linnean
Society **117**: 716–724 (2015).

13. Woronik, A., Stefanescu, C., Käkelä, R., Wheat, C.W., & Lehmann, P. Physiological
differences between female limited, alternative life history strategies: The Alba
phenotype in the butterfly *Colias croceus*. Journal of Insect Physiology **107**: 257-264
(2018).

14. Nielsen, M.G. & Watt, W.B. Behavioural fitness component effects of the alba
polymorphism of *Colias* (Lepidoptera, Pieridae): resource and time budget analysis.
Functional Ecol **12**: 149–158 (1998).

15. Hovanitz, W. The Biology of *Colias* Butterflies. I. The Distribution of the North American
Species. The Wasmann Journal of Biology **8**: 1–28 (1950).

16. Wiklund, C., Karlsson, B., Leimar, B. Sexual conflict and cooperation in butterfly
reproduction: a comparative study of polyandry and female fitness. Proc. R. Soc. B **268**
1661-1667 (2001).

17. Higashijima, S., Kojima, T., & Michiue, T. Dual *Bar* homeo box genes of *Drosophila*
required in two photoreceptor cells, R1 and R6, and primary pigment cells for normal eye
development. Genes & Development **6**: 50–60 (1992).

18. Nijhout, H.F. The Development and Evolution of Butterfly Wing Patterns. Smithsonian
Institution Press (1991).

19. Morehouse, N.I., Vukusic, P., & Rutowski, R. Pterin pigment granules are responsible for
both broadband light scattering and wavelength selective absorption in the wing scales
of pierid butterflies. Proc. R. Soc. B **274**: 359–366 (2007).

20. Rutowski, R.L., Macedonia, J.M., Morehouse, N., & Taylor-Taft, L. Pterin pigments
amplify iridescent ultraviolet signal in males of the orange sulphur butterfly, *Colias*
eurytheme. Proc. R. Soc. B **272**: 2329–2335 (2005).

21. Hagedorn, H.H. & Kunkel, J.G. Vitellogenin and Vitellin in Insects. Annu. Rev. Entomol.
24: 475–505 (1979).

22. Pavlos, N.J. & Jahn, R. Distinct yet overlapping roles of Rab GTPases on synaptic
vesicles. Small GTPases **2**: 77–81 (2011).

23. Stenmark, H. Rab GTPases as coordinators of vesicle traffic. Nat Rev Mol Cell Biol **10**:
513–525 (2009).

24. Gallant, J.R. *et al*. Ancient homology underlies adaptive mimetic diversity across
butterflies. Nature Communications **5**:4817 (2014).

25. Martin, A. *et al*. Diversification of complex butterfly wing patterns by repeated regulatory
evolution of a Wnt ligand. Proc Natl Acad Sci **109**: 12632-12637 (2012).
26. Reig G., Cabrejos, M.E., & Concha. M.L. Functions of BarH transcription factors during
embryonic development. Dev Biol **302**: 367-375 (2007).
27. Kang, J., Yeom, E., Lim, J., & Choi, K. Bar Represses dPax2 and Decapentaplegic to
Regulate Cell Fate and Morphogenetic Cell Death in *Drosophila* Eye. Plos One **9**:
e88171 (2014).
28. Nadeau, N.J. *et al*. The gene *cortex* controls mimicry and crypsis in butterflies and
moths. Nature **534**: 106-110 (2016).
29. Sinervo, B., Svensson, E., & Comendant, T. Density cycles and an offspring quantity
and quality game driven by natural selection. Nature **406**: 985–988 (2000).
30. Sánchez Guillén, R.A. *et al*. Alternative reproductive strategies and the maintenance of
female color polymorphism in damselflies. Ecol Evol **7**: 5592-5602 (2017).

Methods
For detailed methods, including all bioinformatic commands, please see the supplementary
information.
Data availability: SRA reference numbers for the genome and sequencing data will be included
upon acceptance.
Genome assembly: An orange female and male carrying Alba (offspring from wild caught
butterflies, Catalonia, Spain) were mated in the lab. DNA from an Alba female offspring of this
cross was extracted. Quality and quantity were assessed using a Nanodrop 8000
spectrophotometer (Thermo Scientific) and a Qubit 2.0 fluorometer (dsDNA BR, Invitrogen). A
180 insert size paired end library (101bp reads) was prepared (TruSeq PCR free) and
sequenced on an Illumina Hiseq 4000 at the Beijing Genomics Institute (Shenzhen, China). A
Nextera mate-pair library with a 3 kb insert size was prepared and sequenced on an Illumina
HiSeq 2500 (125bp reads) at the Science for Life Laboratory (Stockholm, Sweden). Raw data
was cleaned and high quality reads were used as input for the AllPaths-LG (v. 50960) [31]
assembly pipeline. High molecular weight DNA was extracted from two more Alba females from
the above mentioned cross (i.e full siblings). Equal amounts of DNA from each individual were
pooled sent to the Science for Life Laboratory (Stockholm, Sweden) for PacBio sequencing on
24 SMRT cells (~17GB of data was produced). A Falcon (v. 0.4.2) [32] assembly was generated
by the Science for Life Laboratory. We then used Metassembler (v. 1.5) [33] to merge our
AllPathsLG and Falcon assemblies, using the AllPathsLG assembly as the primary assembly.
Bulk segregant analyses (BSA): The female informative cross data and mapping protocol
described in Woronik and Wheat, 2017 [34] was applied to the high quality reference genome to
identify the contigs that made up the Alba chromosome. ***Male Informative Cross (MIC) I:*** DNA
was extracted from a wild caught orange mother (Catalonia Spain) and 26 of her Alba and 24 of
her orange female offspring. DNA quality and quantity of each individual was assessed via a
Nanodrop 8000 spectrophotometer (Thermo Scientific, MA, USA) and a Qubit 2.0 Fluorometer
(dsDNA BR; Invitrogen, Carlsbad, CA, USA) before pooling equal amounts of high-quality DNA

from Alba and orange offspring into two pools, respectively. The library preparation (TruSeq
PCR-free) and Illumina sequencing (101 bp PE HiSeq2500), was performed at the Beijing
Genomics Institute (Shenzhen, China). Raw reads were cleaned and then mapped to the
reference genome using NextGenMap v0.4.10 (-i 0.09) [35]. SAMTOOLS v1.2 [36] was used to filter
(view -f 3 -q 20), sort and index the bam files and generate mpileup files for the two pools and
the orange mother. Popoolation2 [37] were used to calculate the allele frequency difference
between Alba and orange pools. SNP sites were filtered in R [38], for a read depth ≥ 30 and ≤
300, a bi-allelic state, and a minimum minor allele frequency of 3. The orange mother mpileup
was similarly analyzed using Popoolation [39] (read depth ≥ 5 and ≤ 30); but the major and minor
allele frequencies were calculated in R [38] by dividing the major and minor allele count by the
read depth at each site respectively. A SNP site was considered a MIC I Alba SNP when it met
the following expectations: 1) homozygous in the orange mother, 2) homozygous in the orange
pool, 3) the allele frequency difference in the Alba pool compared to the orange was 0.45-0.55.
MIC II: A male carrying Alba mated an orange female in the lab at Stockholm University. DNA
was prepared as described above for 26 Alba and 28 orange female offspring resulting in two
DNA pools. Library preparation (TruSeq PCR-free) and Illumina sequencing (150 bp paired-end
reads with 350bp insert, HiSeqX), was performed at Science for Life Laboratory (Stockholm,
Sweden). The same mapping and SNP calling pipeline used on the MIC I was applied. A site
was considered an Alba SNP if 1) it was homozygous in the orange pool and 2) the allele
frequency difference in the Alba pool compared to the orange was 0.45-0.55. A contig was
considered Alba associated if it had ≥ 3 Alba SNPs in all crosses. Nineteen Alba associated
contig were identified. They total ~3.7Mbp and are considered the Alba BSA locus.
Genome wide association study: DNA for genome re-sequencing was extracted from 15 Alba
and 15 orange females from diverse population backgrounds (Catalonia, Spain and Capri, Italy).
High quality DNA was prepared using Illumina TruSeq and sequenced at the Science for Life
Laboratory (Stockholm, Sweden) (150 bp paired-end reads HiSeqX). Cleaned reads were
mapped to the annotated reference genome using NextGenMap v0.4.10 (-i 0.6 -X 2000) [35]. Bam
files were filtered and sorted using SAMTOOLS v1.2 (view -f 3 -q 20) [36]. A VCF file was
generated using SAMTOOLS v1.2 (-t DP -t SP -Q 15) [36] and bcftools v.1.2 (-Ov -m) [36]. VCFtools
(v0.1.13) [40] was used to call SNP sites with no more than 50% missing data, an average read
depth between 15-50 across individuals, and a minimum SNP quality of 30. An association
analysis was performed with PLINK (v1.07) [41] and a Benjamini & Hochberg step-up FDR control
was applied. SNPs with FDR <0.05 were considered Alba SNPs. We conducted this analysis
both genome wide and only within the BSA locus. Both analyses fine mapped the Alba locus to
the same genomic region.
Antibody Generation and Staining: A Rabbit-anti-Bar antibody was generated against the full
length sequence of the *Vanessa cardui* Bar homolog:
MTVQRDERDARAPRTRFMITDILDAAPRDLSAHRDSDSDRSATDSPGVKDDSDDVSSKSCGG
DASGLAKKQRKARTAFTDHQLQTLEKSFERQKYLSVQDRMELAAKLGLTDTQVKTWYQNRRT
KWKRQTAVGLELLAEAGNYAAFQRLYGGYWAGVPAYPAQPAPAAADLYYRQAAATAAAAASA
SANTLQKPLPYRLYPGAPLGGVPPLGLGLPGPSAHLGSLGAPGLGALGYYAQARRTPSPDVDP
GSPAPPPRSPREPSIEQRSDDEDDDETIHV. Protein was generated by GenScript (Piscataway,
NJ) and purified to >80% purity. DNA sequences to produce this protein were codon-optimized

for bacterial expression and made via gene synthesis. GenScript injected resultant protein into host animals, collected serum for testing, and affinity purified the product using additional target protein bound to a column. Antibody staining was performed as described previously for *Drosophila* and butterfly tissues [42]. Staged pupal wings and retinas were dissected and fixed between 24-72 hours post-pupation. The Rabbit-anti-Bar antibody was used at 1:100, followed by secondary antibody staining with AlexFluor-555-anti-Rabbit secondaries at 1:500 and counterstaining with DAPI. Images were captured using standard confocal microscopy on a Leica SP5.

CRISPR/Cas9 knockouts: The guide-RNA (gRNA) sequences were generated using the protocol described in Perry et al. 2016. Viable Cas9 target-sites were located by manually looking for PAM-sites (NGG) in the exon region of *BarH-1*. Uniqueness of the target regions was confirmed using a NCBI nucleotide blast (ver. 2.5.0+ using blastn-short flag and filtering for an e-value of 0.01) against the *C. crocea* reference genome. gRNA constructs were ordered from Integrative DNA Technologies (Coralville, Iowa, USA) as DNA (gBlocks). Full gRNA constructs had the following configuration: an M13F region, a spacer sequence, a T7-promotor sequence, the Target specific sequence, a Cas9 binding sequence, and finally a P505 sequence. Upon delivery, gBlocks were amplified using PCR to generate single-stranded guide RNA (sgRNA). For each gBlock, four 50ul reactions were conducted using the M13f and P505 primers and Platinum Taq (Invitrogen cat. 10966-034). The four reactions were then combined and purified in a Qiagen Minelute spin column (cat. 28004, Venlo, Netherlands). The resulting template was transcribed using the Lucigen AmpliScribe T7-flash Transcription Kit from Epicentre/Illumina (cat. ASF3507, Madison, WI, USA) followed by purification via ammonium acetate precipitation. Products were resuspended with Qiagen buffer EB, concentrations were quantified by Qubit and further diluted to 1000 ng/µl. They were then mixed with Cas9-NLS protein (PNA Bio, Newbury Park, CA, USA) and diluted to a final concentration of 125-250 ng/µl. *C. crocea* females (n > 40) from Aiguamolls de l'Empordà, Spain were captured and kept in morph-specific flight cages in the lab at Stockholm University where they oviposited on alfalfa (*Medicago sativa*). Eggs were collected between 1-7h post-laying and sterilized in 7% benzalkonium chloride for ~5 minutes before injection. Injections were either at a concentration of 125 or 250 ng/ul and conducted using a M-152 Narishige micromanipulator (Narishige International Limited, London, UK) with a 50 ml glass needle syringe, with injection pressure applied by hand via a syringe fitting.

CRISPR/Cas9 validation: To validate the mutation, Cas9 cut sites were PCR-amplified and a ~370bp region, centered on the intended cut site were sequenced using Illumina MiSeq 300bp paired-end sequencing. Primers were designed using Primer3 (http://biotools.umassmed.edu/bioapps/primer3_www.cgi). DNA was isolated from KO-individuals using KingFisher Cell and Tissue DNA Kit from ThermoFisher Scientific (N11997) and the robotic Kingfisher Duo Prime purification system. DNA quality and quantity were assessed via a Nanodrop 8000 spectrophotometer (Thermo Scientific, MA, USA) and a Qubit 2.0 Fluorometer (dsDNA BR; Invitrogen, Carlsbad, CA, USA). Aliquots were then taken and diluted to 1ng/ul before amplifying the region over the cleavage-site. Sequences were amplified and ligated with Illumina adapter and indexes in a two-step process following the protocol provided by Science for Life Laboratories (Stockholm, Sweden) and Illumina. First, we amplified the ~370bp long sequence around the cut sites and attach the first Illumina adapter, onto which we later attach Illumina handles and Index using a second round of PCR (Accustart II PCR

Supermix from Quanta Bio [Beverly, MA, USA], settings 94C x 2 min followed by 40 cycles of 94
C x 30 sec + 60 C x 15 sec + 68 C x 1 min followed by 68 C x 5 min). PCR products were
purified using Qiagen Qiaquick (Cat. 28104). Concentration and quality of the product were
assessed via Nanodrop and gel electrophoresis. DNA was diluted to ~0.5ng/ul and then the
unique double indices were attached by the second round of PCR (same protocol as above).
The final PCR products were purified again using Qiaquick spin columns and concentration and
size was assessed using Qubit fluorometer and gel electrophoresis. All samples were then
mixed at equal molarity and sent for sequencing at Science for Life Laboratories (Stockholm,
Sweden). Sequences were aligned to their respective fragments (area surrounding cut site)
using SNAP (ver. 1.0beta18) [43], identical reads were clustered using the collapser utility in
Fastx-Toolkit. Sequences containing deletions were extracted and the most abundant
sequences containing deletions were selected for confirmation of deletion in the expected
region.
Electron Microscopy: To quantify pigment granule differences between Alba and orange
individuals pieces of the forewing were mounted on aluminum pin stubs (6mm length) with the
dorsal side upwards. Samples were coated in gold for 80 seconds using an Agar sputter coater
and imaged under 5 kV acceleration voltage, high vacuum, and ETD detection using a scanning
electron microscope (Quanta Feg 650, FEI, Hillsboro, Oregon, USA). To quantify pigment
granules within the photos we selected images from the same magnification and drew randomly
placed three 4 μm^2 squares on the images. We counted the number of pigment granules within
each square and took the average, then conducted a two sample t-test in R. To quantify
pigment granule differences between KO and wild type regions in our CRISPR KO mosaic
individuals, a biopsy hole punch a 2mm in diameter circle was used to cut out one piece mostly
containing white scales and one piece with mostly orange scales. These pieces were first
photographed using a Leica EZ4HD stereo microscope in order to allow us to confirm the color
of each scale once they were covered with gold sputter. Five white and five orange scales were
then selected and the granules from a 4μm^2 square were counted from each of those scales
and a two sample t-test was then conducted in R.
Lipid Analysis: Wild caught *C. crocea* Alba females (Catalonia, Spain) oviposited in the lab.
Eggs were moved into individual rearing cups and split between two temperature treatments
(hot: 27 ^{O}C and 16 hour day length during larval and pupal development, cold: reared at 22 ^{O}C
with a 16 hour day length during larval development and 15^{O}C with a 16 hour day length during
pupal development). Once pupated individuals were checked a minimum of every 12 hours.
Upon eclosion adults were stored at 4 ^{O}C until the next day to provide time for meconium
excretion. Butterflies were not allowed to feed before dissection. Body weight was taken using a
Sauter RE1614 scale before dissection. Total lipids were extracted using the Folch method [44]
according to the procedures outlined in Woronik et. al. 2018 [13]. HPTLC was conducted as
described in Woronik et. al. 2018 [13]. In brief, 5 μl of the sample lipid extract was applied on a
silica plate with a Camag Automatic TLC Sampler 4 (Camag, Muttenz, Switzerland). After the
silica plate developed it was scanned with a Camag TLC plate scanner 3 at 254 nm using a
deuterium lamp with a slit dimension of 6 × 0.45 mm and analyzed with the Win-CATS 1.1.3.0
software. Peaks representing the four major neutral lipid classes (diacylglycerols,
triacylglycerols, cholesterol and cholesterol esters) were identified by comparing their retention
times against known standards. Then the peak areas were integrated and the amount of lipid

within each class was calculated using the formula: $pmol_{sample}$ = ($Area_{sample}$ / $Area_{standard}$) x
$pmol_{standard}$. The total lipid content (nmol per abdomen) was calculated as a sum of pmol
contents of all neutral lipid classes. For the statistical analyses this value was regressed against
abdomen weight and standardized residuals (i.e. mass-corrected storage lipid amount) and
were subsequently used as dependent variable.
Transcriptome assembly and differential expression analysis: Offspring from a wild caught
Alba female from Catalonia, Spain were reared at Stockholm University. When larvae reached
the fifth instar they were checked at least every six hours and the pupation time of each
individual was recorded. Tissue was collected between 82% and 92% of pupal development.
Pupae were dissected in PBS solution, and the abdomen and wings were flash frozen in liquid
nitrogen and stored at -80 ºC. RNA was extracted from the abdomen and wing tissues using
Trizol. RNA quality and quantity was assessed using a Nanodrop 8000 spectrophotometer
(Thermo Scientific) and an Experion electrophoresis machine using the manufacturer protocol
(Bio-Rad, Hercules, CA). Library preparation (Strand-specific TruSeq RNA libraries using poly-A
selection) and sequencing (101 bp PE HiSeq2500 - high output mode) was performed at the
Science for Life Laboratories (Stockholm, Sweden). In total 16 libraries were sequenced (4
orange and 4 Alba individuals - wings and abdomen from each individual). Raw data was
cleaned and reads from all libraries were used in a *de novo* transcriptome assembly (Trinity
version trinityrnaseq_r2013_08_14 with default parameters) [45]. To reduce the redundancy
among contigs and produce a biologically valid transcript set, the tr2aacds pipeline from the
EvidentialGene software package [46] was run on the raw Trinity assembly. The sixteen RNA-Seq
libraries were mapped to the resulting transcriptome using NextGenMap v0.4.10 (-i 0.09) [35].
SAMTOOLS v1.2 [35] was then used to filter (view -f 3 -q 20), sort and index the sixteen bam files.
SAMTOOLS v1.2 [35] idxstats was then used to calculate the read counts per gene for each of the
sorted bam files. These counts were then joined in a CSV file using an in-house pipeline and
csvjoin. A differential expression analysis was conducted in EdgeR [47]. A Benjamini Hochberg
correction was applied to the raw p values to correct for false discovery rate and differentially
expressed genes were called (adjusted p value <0.05). Babelomics (version 4.2) [48] was used to
conduct a gene set enrichment analysis (Fatiscan, two tailed Fisher's exact test). Revigo [49] was
used to cluster significantly enriched GO terms by semantic similarity (default settings, C = 0.7).
The GO term clusters were named and assigned p-values based on the most significant GO
term in the cluster.
472
473

474   31. Gnerre, S. *et al*. High-quality draft assemblies of mammalian genomes from
475   massively parallel sequence data. Proc Natl Acad Sci **108**:1513-1518 (2011).
476
477   32. Chin, C. *et al*. Phased diploid genome assembly with single-molecule real-time
478   sequencing. Nature Methods **13**: 1050–1054 (2016).
479
480   33. Wences, A.H. & Schatz M.C. Metassembler: merging and optimizing *de novo*
481   genome assemblies. *Genome Biology* **16**: 207 (2015).
482
34. Woronik, A. & Wheat, C.W. Advances in finding Alba: the locus affecting life history
and color polymorphism in a *Colias* butterfly. *Journal of Evolutionary Biology* **30**: 26-39

(2017).

35. Sedlazeck, F.J., Rescheneder, P. & Haeseler, von, A. NextGenMap: fast and
accurate read mapping in highly polymorphic genomes. *J Gerontol* **29**: 2790–2791
(2013).

36. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**:
2078–2079 (2009)

37. Kofler, R., Pandey, R.V. & Schlotterer, C. PoPoolation2: identifying differentiation
between populations using sequencing of pooled DNA samples (Pool-Seq).
Bioinformatics **27**: 3435–3436 (2011b).

38. R Core Team (2015). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

39. Kofler, R. *et al.* PoPoolation: A Toolbox for Population Genetic Analysis of Next
Generation Sequencing Data from Pooled Individuals. *PLoS ONE* **6**: e15925 (2011a).

40. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**: 2156–
2158 (2011).

41. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-
Based Linkage Analyses. *The American Journal of Human Genetics* **81**: 559–575
(2007).

42. Perry, M.W. *et al.* Molecular logic behind the three-way stochastic choices that
expand butterfly colour vision *Nature:* **535**:280–284 (2016).

43. Zaharia, M. *et al.* Faster and More Accurate Sequence Alignment with SNAP.
ArXiv:1111.5572 [Cs, q-Bio], November. http://arxiv.org/abs/1111.5572 (2011).

44. Folch, J.M., Lees, M., & Sloane-Stanley, G.H. A simple method for the isolation and
purification of total lipides from animal tissue. *J. Biol. Chem*. **226**: 497-509 (1957).

45. Grabherr, M.G. *et al*. Full-length transcriptome assembly from RNA-Seq data without
a reference genome. *Nat Biotechnol* **29**: 644–652 (2011).

46. Gilbert, Donald (2013) Gene-omes built from mRNA seq not genome DNA. 7th
annual
arthropod Genomics symposium. Notre Dame.
http://arthropods.eugenes.org/EvidentialGene/about/EvigeneRNA2013poster.pdf   and
http://globalhealth.nd.edu/7th-annual-arthropod-genomics-symposium/
515

47. Robinson, M.D., McCarthy, D.J. & Smyth, G.K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–
140 (2009).

48. Medina, I. *et al*. Babelomics: an integrative platform for the analysis of
transcriptomics, proteomics and genomic data with advanced functional profiling. *Nucleic*

Acids Research **38**: W210–W213 (2010).

49. Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO Summarizes and Visualizes
Long Lists of Gene Ontology Terms. *PLoS ONE* **6**: e21800 (2011).

Figure 1. Color variation in *Colias crocea* and the genetic mechanism of Alba. (A) *Colias crocea* male, orange female, and Alba female (left to right). (B) SNPs significantly associated with the Alba phenotype (red) within the ~3.7 Mbp Alba locus identified via 3 rounds of bulk segregant analysis. Contigs in this region shown as alternating dark and light blue. (C) The location of Alba associated SNPs (red) on the ~430 kb outlier contig identified in the GWAS. Gene models for the DEAD-box helicase, the Jockey-like transposable element, and *BarH-1* shown at the top of the panel. (D) Loss of green color in the *C. crocea* eye following *BarH-1* mosaic KO. KO regions are black. (E) Orange color is seen on the dorsal forewing (upper) and hindwing (lower) of an Alba female following *BarH-1* mosaic KO.

Figure 2. BarH-1 antibody staining in *Colias crocea* and *Vanessa cardui* pupal wings. (A) Depiction of approximate location of antibody images on the *C. crocea* Alba forewing and the scales within the region. (B) DAPI (nuclei) and BarH-1 staining within the scale building cells within the black margin region of the Alba forewing. BarH-1 is expressed in melanic Alba scale building cells. (C) DAPI (nuclei) and BarH-1 staining of white regions of the Alba forewing. Large nuclei are scale building cells while small nuclei are epithelial cells. Antibody staining shows BarH-1 expression in the white Alba scale building cells. (D) Depiction of approximate location of antibody images on the *C. crocea* orange forewing and the scales within the region. (E) DAPI (nuclei) and BarH-1 staining of orange regions of the orange forewing. Large nuclei are scale building cells while small nuclei are epithelial cells. Antibody staining shows a lack of BarH-1 expression in the orange scale building cells. (F) Depiction of approximate location of antibody images on the *C. crocea* orange hindwing and an illustration of the scale heterogeneity found within the region. (G) DAPI (nuclei) and BarH-1 staining of the orange hindwing. Large nuclei are scale building cells while small nuclei are epithelial cells. Antibody staining shows heterogenity in BarH-1 expression in the scale building cells within this region presumably corresponding to the variation in scale color, where melanic scale building cells express BarH-1, but orange do not. (H) Hind and fore wing of *V. cardui.* (I) DAPI (nuclei) and BarH-1 staining of *V. cardui* . Both the scale building and socket cells can be observed and express BarH-1.

Figure 3. Butterfly forewings and scanning electron microscope (SEM) images of their wing scale microstructures. (A) A WT Alba female wing and SEM of one of its wing scales, showing the near absence of pigment granules. (B) A WT orange female wing and SEM, showing an abundance of pigment granules. (C) An Alba female that is a *BarH-1* mosaic KO, SEM images illustrate significant differences in the number of pigment granules in Alba (C) and orange (D) mosaic regions. These differences are consistent with those observed in WT animals. (E) Dorsal forewing of an orange female where pigment granules were chemically removed from distal ½ of wing. SEM inset showing absence of pigments in the white region. (H) A *Pieris brassicae* female

forewing, with SEM showing abundance of pigment granules despite its white color, indicating there are multiple routes to white wing color within Pieridae.

Figure 4. Physiological differences between female morphs. A) The mass corrected total neutral lipid content for female morphs in two temperature treatments. Alba females, on average, have larger neutral lipid stores, however there is an interaction between morph and temperature as the difference is only significant in the cold treatment. Error bars are the standard error (cold: n=32, $t_{29.12}$ = 3.42, P = 0.002, hot: n=25, $t_{22.71}$ = 0.67, P = 0.51). B) Volcano plot to visualize gene expression differences between female morphs in pupal abdominal tissue. Each point is a gene. Grey circles are genes not significantly DE between morphs, while blue circles are significantly DE. The black square is *vitellogenin1* and the black triangle is *RIM*. The X-axis is the log of the fold change (FC), positive log(FC) indicates the gene is upregulated in Alba individuals. C) Volcano plots visualize gene expression differences between female morphs in pupal wing tissue. Color coding, shapes, and axes are the same as above.

A

B

DEAD-box helicase

Jockey-like
transposable element

BarH-1

C

D

E

Figure 1

BarH-1 expression in a matching region of developing pupal wings

A — Alba forewing

B — DAPI / BarH-1

C — DAPI / BarH-1

D — orange forewing

E — DAPI / BarH-1

F — orange hindwing

G — DAPI / BarH-1

H — *Vanessa cardui*

I — DAPI / BarH-1

Figure 2

Figure 3

Figure 4