Type of article : technical report

Title :

Assessing precocious maturation in salmon using ultrasound imaging

Authors:

Marie Nevoux[1,2], Frédéric Marchand[2,3], Jean-Pierre Destouches[2,3], Guillaume Forget[1,2], Dominique

Huteau[1,2], Julien Tremblay[2,3]

Affiliations:

[1] INRA, Agrocampus-Ouest, Ecology and Ecosystem Health, 65 rue de Saint-Brieuc, Rennes, France

[2] Management of Aquatic Ecosystems in Agricultural Watersheds, INRA, Agence Française pour la

Biodiversité, Rennes, France

[3] INRA, Ecologie et Ecotoxicologie Aquatique, 65 rue de Saint-Brieuc, Rennes, France

Corresponding author:

Marie Nevoux,

ESE, Ecology and Ecosystem Health, Agrocampus Ouest, INRA, 35042 Rennes, France

Email: marie.nevoux@inra.fr

Phone: +33(0)2 23 48 52 35

Abstract

Salmonids are characterized by a large diversity of life histories, but their study is often limited by the imperfect observation of the true state of an individual in the wild. Challenged by the need to reduce uncertainty of empirical data, we have taken advantage of recent development in medical imaging techniques to assess precocious maturation in Atlantic salmon. We compared traditional phenotypic (external) examination and ultrasound (internal) examination and provided recommendations on fish handling and ultrasound image interpretation. By seeing the unseen, ultrasound imaging offers great opportunities for ecological studies in the wild.

Keywords

Atlantic salmon *Salmo salar*, ecology, empirical study, fisheries, individual heterogeneity, life history trait, maturation, ultrasound scan

Introduction

Long term monitoring of animal populations in the wild is of great interest for ecology and management (Clutton-Brock and Sheldon 2010). In a context of global changes, such long term monitoring program contributed to highlight major change in population composition, dynamics and abundance over the last decades (Parmesan 2006). Indeed, global changes trigger the need to predict population structure and abundance in the future. However, predictive models are difficult to parameterize and often return high level of uncertainty, which hinder their use by managers and policy makers. This is partly because field records are imperfect observation of the true state of an individual (Genovart et al. 2012). For instance, maturation status and sex is rarely known with certainty in fish monitoring programs. Thus, field studies are challenged by the need to reduce uncertainty in individual data.

Atlantic salmon (*Salmo salar*) is an anadromous species that reproduce in freshwater and migrate at

sea to grow and undergo sexual maturation. However, in this and other salmonid species, some

young males (parr) would mature precociously in freshwater (Fleming 1996). It is proposed that

frequency-dependent selection, as affected by intensity of mate competition and environmental

variability in growth opportunities, determines a polygenic threshold (Hutchings and Myers 1994).

Mature male parrs attempt to 'sneak' access to mating with large anadromous females (Fleming

1996). They can contribute substantially to the reproduction (Morán et al. 1996) but as maturation

seems to be traded against survival, mature parr have a low probability to smoltify (Buoro et al.

2010). The frequency of this alternative reproduction strategy is highly variable over time, space and

age classes (Bagliniere and Maisse 1985; Myers et al. 1986), but precise quantification of this

phenotype remain rare in empirical studies. Furthermore, precocious maturation is virtually ignored

by stock assessment models and fisheries management in Atlantic salmon (e.g. ICES (2013)). This

means that mature parrs are included neither in the stock nor in the recruitment figures. This might

bias stock-recruitment relationships and underestimate freshwater survival and salmon production.

For instance, Myers (1984) estimated that 60% of the stock of adult male salmon was lost due to

precocious maturation. Caswell et al. (1984) demonstrated that the effect of reproduction by parr on

population growth rate is always greater than that of adult reproduction. We suspect that this lack

of consideration for secretive mature parrs owes to the complexity of salmon life cycle. This could

also be partly due to the difficulty of quantifying the proportion of precocious maturation in wild

salmon populations

In the field, precocious maturity in parr is traditionally assessed by pressing the abdomen of the fish

to extract milt. However, not all mature male may be producing milt at the time of capture.

Moreover, based on the general look of the fish, experienced fieldworkers sometimes suspect

maturity in some individuals that do not excrete milt upon abdominal pressure. However,

researchers are often reluctant to use such expert opinion data in their analysis due to unknown

accuracy and potential operator effect.

Ultrasound imaging, which gave rise to significant advances in human medicine, offer reliable

visualization of internal organs. Ultrasound imaging is routinely used in aquaculture as provides an

accurate, non-invasive, non-lethal method that is more accurate that visual methods, e.g. to

determine gender or gonadal growth in fish (Novelo and Tiersch 2012). In adult salmon, although

gonads of immature males may be difficult to discern, gonads of females are always visible and

hence sexing is possible by deduction (Martin et al. 1983; Mattson 1991). However the ovary and

testes of the juvenile (non mature) salmon seem difficult to identify (Martin et al. 1983).

The development of portable devices offer an opportunity to bring ultrasound machines out into the

wild. In this paper, we describe how ultrasound scanner can enhance the range of biological traits

that can be monitored in long-term salmonids population surveys. In particular, we assessed the

maturation status of young salmon in the field using traditional phenotypic examination and

ultrasound examination. We compare the two approaches by testing whether biological or operator

related factors would affect the uncertainty in external phenotypic assessment of maturity in parr.

Finally, we provided recommendation on fish handling and the interpretation of ultrasound images.

Material and methods

Salmon monitoring

Atlantic salmon monitoring takes place every year in September or October in the River Oir,

Normandy, France (Marchand et al. 2018). Young individuals (parr) are captured using a

standardized electrofishing protocol, they are placed in a light anesthetic, measured, weighed,

scanned and returned to the river within 0.5 hour maximum of their capture. We assessed age

through scale reading. In 2015, 2016, and 2017 we examined 850 salmon (366 parrs of age 0 and 484

parrs of age 1) for maturity using traditional phenotypic (external) approach and ultrasound imaging.

We caught all parrs within a period of 8 days in early October. All the animal experimentation in this

study was performed according to French legislation and under licence APAFIS-201602051204637.

Traditional phenotypic examination

First, the field operator assessed the maturation state of parr by gently pressing the side of the fish.

The excretion of milt (sperm) is indicative of a mature male, while the absence of milt can be

encountered in a maturing male whose gonad does not yet produce milt, an immature male, or a

female (females are all immature at this stage). The field operator also reported that some fish *looks*

like mature parr: they display phenotypic traits similar to those of mature fish (e.g. colour pattern,

body shape) but they do not excrete milt. This phenotypic assessment rely less on a fixed set of

quantifiable parameters than on a general and subjective appreciation of the fish, which is specific to

each operator.

Ultrasound examination

Then, the ultrasound operator screened all salmon parr for precocious maturation using a portable

ultrasound scanner M-Turbo (Sonosite) with a 5-10 MHz linear transducer. We selected the default

setting for muscular examination. As water perfectly transmit ultrasounds, we do not need to use

ultrasound transmission gel. We placed fish on their side in a tank of freshwater, and we operated

the transducer in the water keeping the transducer 1-2cm above the belly. We produced sagittal

images of the abdominal cavity, i.e. the transducer was oriented parallel to the lateral line. We

directly assessed the maturation state of each individual from live images on the ultrasound

monitor. Parrs were classified as mature when the gonad was detected, or immature when the

gonad was not detected, assuming that the gonad would be too small at this stage. Images could be

stored on the ultrasound machine and then exported on a USB memory stick for post-treatment.

Comparison of the two approaches

We compared the number of parrs that were recorded as mature or non-mature by one or the other

approach. Then, we conducted a multivariate analysis to investigate which biological factor drove

the probability that a mature male was producing milt at the time of capture. We tested the effect

of year, age, body length in a generalized linear model (GLM) with a binomial distribution of errors.

We also tested whether the probability to miss a mature but parr that does not produce milt was

affected by age, body length, operator identity and time of the day (testing for operator tiredness).

We did not consider the date of capture because of the low variability in this variable. All models

were run in R (R Development Core Team 2004).

Results and discussion

Ultrasound assessment of precocious maturity

The trained ultrasound operator conducted ultrasound examination for maturation assessment in

ca. 5 seconds. In immature individuals, we can identify the liver, the stomach, caecas and the

intestine (Figure 1a). Note that all internal organs appear more clearly on the live images, during

ultrasound examination, than on frozen images. Nevertheless, the heterogeneous filling and uneven

granularity of the abdominal cavity are diagnostic, as well as the concave shape of the cavity, which

appear depressed on both sides. In mature parr (precocious males), as gonads virtually fill the whole

cavity, ultrasound images depict a homogeneous and finely granulated pattern (Figure 1b, c).

Digestive organs are hardly visible and the shape of the cavity is convex. In mature males that does

not produce milt (Figure 1b), gonads appear less developed and fill the cavity to a lesser extent than

in males producing milt (Figure 1c), which is consistent with a lower degree of maturity in the

former. Interestingly, ultrasound imaging provides an objective description of maturation status in

males that did not produce milt (Figure 1b) and the ones that produced milt (Figure 1c). It can also

discriminate between truly mature parr from well-fed individuals, which the operator can mistakenly

confuse with mature male because of their round belly.

Figure 1. Assessing maturity in salmon parr using ultrasound imaging: a) immature one year old parr,

b) mature young of the year parr that did not produce milt (precocious male), and c) mature and

year old parr that produced milt (precocious male) in River Oir. On all images, the head of the fish is

on the left but not shown, the left and right flanks are situated on the lower and upper part of the

image, respectively. Gonads are identified with a "G" letter.

Comparison of phenotypic and ultrasound methods

On average 33.4% of the mature male were not producing milt. The probability to produce milt

increased with age (P = 0.025) and body length (P = 0.041), although this length effect tend to be

weaker in older parrs (age * length: P = 0.055). We also detected a strong year effect (P < 0.001),

which suggests that environmental factors influence the level of maturation of parr at the time of

capture.

In our study, the maturation state of 106 individuals was uncertain. However, following ultrasound

examination, we found that 89.6 % of those parrs that looked like mature parrs were actually

mature. This result acknowledged the high level of expertise of our trained operators to assess the

maturation state of parr from external phenotypic examination. In total, the field operator did not

detect 5.6 % of the mature males (i.e. the maturity was not confirmed nor suspected). This

probability differed between the two operators (P = 0.008), which point out to the difficulty of

assessing parr maturation state from external examination in the field in a fully reliable and

replicable manner. None of the other variable had a significant effect.

The ultrasound operator missed 2.7% of the males producing milt (the sample size is too small to

conduct multivariate analysis). This result pointed out that ultrasound imaging is nevertheless

subject to a low degree of uncertainty in maturity assessment. The interpretation of ultrasound

images requires a good training to improve the accuracy of the information. We also recommend

that all images would be examined and interpreted by two operators.

Table 1. Comparison of the number of records using external phenotypic observation (rows) and

ultrasound imaging (columns) to assess maturation in parr.

Phenotype \ ultrasound	Gonad detected with ultrasound	Gonad not detected with ultrasound
Parr producing milt	221	6
Parr not producing milt that looks like a mature parr	95	11
Parr not producing milt that looks like a non-mature parr	19	498

Conclusion

By seeing the unseen, ultrasound imaging offer great opportunities for ecological studies in the wild.

As illustrated in this paper, key phenotypic traits become accessible and help better characterize

individual's true biological state. Ultrasound imaging is an accessible, easily transferable and

replicable approach to investigate precocious maturity in parr as it relies on key diagnostic features.

However, the accuracy and efficiency of the ultrasound examination will depend on the skill of the

operator. Indeed, some training is required to operate the transducer in a fluent way and navigate

between different anatomic layers. In this regard, we recommend good knowledge of the internal

structure for the species of interest as a prerequisite. This will ensure a quick examination and thus

minimize the impact on fish wellbeing. This will also improve the quality of the images and minimize

misinterpretation issue.

References

Bagliniere, J.L., and Maisse, G. 1985. Precocious maturation and smoltification in wild Atlantic

salmon in the Armorican massif, France. Aquaculture **45**(1): 249–263. doi:10.1016/0044-

8486(85)90274-1.

Buoro, M., Prévost, E., and Gimenez, O. 2010. Investigating Evolutionary Trade-Offs in Wild

Populations of Atlantic Salmon (*Salmo salar*): Incorporating Detection Probabilities and

Individual Heterogeneity. Evolution **64**(9): 2629–2642. doi:10.1111/j.1558-

5646.2010.01029.x.

Caswell, H., Naiman, R.J., and Morin, R. 1984. Evaluating the consequence of reproduction in

complex salmonid life cycles. Aquaculture **43**: 123–134.

Clutton-Brock, T., and Sheldon, B.C. 2010. Individuals and populations: the role of long-term,

individual-based studies of animals in ecology and evolutionary biology. Trends Ecol. Evol.

25(10): 562–573.

Fleming, I.A. 1996. Reproductive strategies of Atlantic salmon: ecology and evolution. Rev. Fish Biol.

Fish. **6**(4): 379–416. doi:10.1007/BF00164323.

Genovart, M., Pradel, R., and Oro, D. 2012. Exploiting uncertain ecological fieldwork data with multi-

event capture–recapture modelling: an example with bird sex assignment. J. Anim. Ecol.: no-

no. doi:10.1111/j.1365-2656.2012.01991.x.

Hutchings, J.A., and Myers, R.A. 1994. The evolution of alternative mating strategies in variable

environments. Evol. Ecol. **8**(3): 256–268.

ICES. 2013. Report of the Working Group on North Atlantic Salmon (WGNAS). Copenhagen,

Denmark.

Marchand, F., Azam, D., Delanoë, R., Destouches, J.-P., Tremblay, J., Baglinière, J.-L., Nevoux, M.,

Prévost, E., and Azam, D. 2018. Abundances and biological traits of fish sampled by

electrofishing (except specific abundance indices) on the River Oir since 1988 (France).

doi:10.15468/yaikgn.

Martin, R.W., Myers, J., Sower, S.A., Phillips, D.J., and Mcauley, C. 1983. Ultrasonic Imaging, a

Potential Tool for Sex Determination of Live Fish. North Am. J. Fish. Manag. **3**(3): 258–264.

doi:10.1577/1548-8659(1983)3<258:UIAPTF>2.0.CO;2.

Mattson, N.S. 1991. A new method to determine sex and gonad size in live fishes by using

ultrasonography. J. Fish Biol. **39**(5): 673–677. doi:10.1111/j.1095-8649.1991.tb04397.x.

Morán, P., Pendás, A.M., Beall, E., and García-Vázquez, E. 1996. Genetic assessment of the

reproductive success of Atlantic salmon precocious parr by means of VNTR loci. Heredity

77(6): 655–660. doi:10.1038/hdy.1996.193.

Myers, R.A. 1984. Demographic Consequences of Precocious Maturation of Atlantic Salmon (Salmo

salar). Can. J. Fish. Aquat. Sci. **41**(9): 1349–1353. doi:10.1139/f84-165.

Myers, R.A., Hutchings, J.A., and Gibson, R.J. 1986. Variation in Male Parr Maturation Within and

Among Populations of Atlantic Salmon, *Salmo salar*. Can. J. Fish. Aquat. Sci. **43**(6): 1242–

1248. doi:10.1139/f86-154.

Novelo, N.D., and Tiersch, T.R. 2012. A Review of the Use of Ultrasonography in Fish Reproduction.

North Am. J. Aquac. **74**(2): 169–181. doi:10.1080/15222055.2012.672370.

Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. Annu. Rev. Ecol.

Evol. Syst. **37**(1): 637–669.

R Development Core Team. 2004. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. Available from http://www.R-

project.org.