Type of article : technical report

Title :

Field assessment of precocious maturation in salmon parr using ultrasound imaging

Authors:

Marie Nevoux[1,2], Frédéric Marchand[2,3], Jean-Pierre Destouches[2,3], Guillaume Forget[1,2], Dominique

Huteau[1,2], Julien Tremblay[2,3]

Affiliations:

[1] INRA, Agrocampus-Ouest, Ecology and Ecosystem Health, 65 rue de Saint-Brieuc, Rennes, France

[2] Management of Aquatic Ecosystems in Agricultural Watersheds, INRA, Agence Française pour la

Biodiversité, Rennes, France

[3] INRA, Ecologie et Ecotoxicologie Aquatique, 65 rue de Saint-Brieuc, Rennes, France

Corresponding author:

Marie Nevoux,

ESE, Ecology and Ecosystem Health, Agrocampus Ouest, INRA, 35042 Rennes, France

Email: marie.nevoux@inra.fr

Phone: +33(0)2 23 48 52 35

Abstract

Salmonids are characterized by a large diversity of life histories, but their study is often limited by the imperfect observation of the true state of an individual in the wild. Challenged by the need to reduce uncertainty of empirical data, recent development in medical imaging techniques offered new opportunities to assess precocious maturation in Atlantic salmon parr. Traditional phenotypic (external) examination and ultrasound (internal) examination were compared and recommendations on fish handling and ultrasound image interpretation were provided. By allowing to see the unseen, ultrasound imaging offers great opportunities for ecological studies in the wild such as individual sexual maturation.

Keywords

Atlantic salmon *Salmo salar*, ecology, empirical study, fisheries, individual heterogeneity, life history trait, maturation, ultrasound scan

Introduction

Long term monitoring of animal populations in the wild is of great interest for ecology and management (Clutton-Brock and Sheldon 2010). In a context of global change, such long term monitoring programs contribute to highlight major changes in population composition, dynamics and abundance over the last decades (Parmesan 2006). In the North Pacific, analyses confirmed climate-related shifts in the abundance of most salmonid species since the 1940's, associated with reported ecosystem regime shifts (Irvine and Fukuwaka 2011). In the North Atlantic, the widespread decline in salmon abundance went with a decrease in both survival and age at maturity (Chaput 2012). This global change makes it particularly important to be able to predict population structure and abundance in the future under different climatic scenarios. However, predictive population dynamics models are difficult to parameterize and often return high level of uncertainty, which hinder their use by managers and policy makers. Stock assessments form the basis for catch advice

for salmon fisheries, but they mask regional differences and annual river-specific stock assessments

are only available for some 25% of the rivers (Chaput 2012). This is partly because field records are

imperfect observation of the true state of an individual (Genovart et al. 2012). For instance,

maturation status and gender is rarely known with certainty in fish monitoring programs without

invasive or lethal methods. Thus, field studies are challenged by the need to reduce uncertainty in

individual-level data.

Atlantic salmon (*Salmo salar*) is an anadromous species that reproduces in freshwater and migrates

at sea to grow and undergo sexual maturation. However, in this and other salmonid species, some

young males (parrs) can mature precociously in freshwater (Fleming 1996). It is proposed that the

intensity of intra-sexual competition for mates and environmental variability in growth

opportunities, determines a polygenic threshold that triggers precocious maturation (Hutchings and

Myers 1994). Mature male parrs attempt to 'sneak' access to large anadromous females for mating

(Fleming 1996). They can contribute substantially to the reproduction (Moran et al. 1996) but, as

maturation seems to be traded against survival, mature parr have a low probability to migrate at sea

(Buoro et al. 2010). The occurrence of this alternative reproduction strategy is highly variable over

time, space, and age classes (Baglinière and Maisse 1985; Myers et al. 1986) and may be driven by

frequency dependent selection, i.e. the reproductive success of precocious parrs increases with

decreasing frequency of precocious maturation in the population (Berejikian et al. 2010). Yet, precise

quantification of this phenotype remains rare in empirical studies and in salmon population dynamic

models. Furthermore, precocious maturation is virtually ignored by stock assessment models and in

the management of Atlantic salmon fisheries (e.g. ICES 2013). This means that mature parrs are

included neither in the stock nor in the recruitment figures. This might bias stock-recruitment

relationships and underestimate freshwater survival as well as salmon production. For instance,

using more than 280,000 records from Little Codroy River parrs, Myers (1984) estimated that 60% of

the stock of adult male salmon in Newfoundland was lost because of increased mortality due to

precocious maturation. Caswell et al. (1984) demonstrated that the effect of reproduction by parrs

on population growth rate was always greater than that of reproduction by adults. It is suspected

that this lack of consideration for mature parrs is due to the complexity of the life cycle of Atlantic

salmon. But it could also be due to the difficulty of quantifying the occurrence of precocious

maturation in wild salmon populations.

In the field, precocious maturity in parrs is traditionally assessed by pressing the abdomen of the fish

to extract milt (sperm). Hence, not all mature males may be producing milt at the time of capture.

Moreover, based on the general aspect of the fish, experienced fieldworkers sometimes suspect

maturity in some individuals that do not excrete milt upon abdominal pressure. However,

researchers are often reluctant to use such expert opinions in their analysis due to the potential for

incorrect diagnosis and strong operator bias.

Ultrasound imaging, which gave rise to significant advances in human medicine, offers reliable

visualization of internal organs. Ultrasound imaging is routinely used in aquaculture as it provides an

accurate, non-invasive, non-lethal method that is more accurate that visual methods, to determine

gender or gonadal growth in fish for example (Novelo and Tiersch 2012). In adult salmon, although

gonads of immature males may be difficult to discern, gonads of females are always visible and

hence sexing is possible by deduction (Martin et al. 1983; Mattson 1991). However the ovaries and

testes of the juvenile (immature) salmons seem difficult to identify (Martin et al. 1983).

The development of portable devices offers an opportunity to bring ultrasound machines in the

field. This paper illustrates how ultrasound scanners can increase the range of biological traits that

can be monitored in long-term salmonid populations surveys. In particular, the maturation status of

salmon parr (i.e. freshwater resident salmon) was assessed in the field using traditional phenotypic

examination and ultrasound examination. The two approaches are compared and it is tested

whether biological or operator-related factors affect the uncertainty in external phenotypic

assessment of parr maturity. Finally, recommendation on fish handling and on the interpretation of

ultrasound images is provided.

Material and methods

Salmon monitoring

Atlantic salmon monitoring takes place every year in September or October in the river Oir,

Normandy, France (Marchand et al. 2018). Young individuals (parrs) are captured using a standard

electrofishing protocol. They are then placed in a light anesthetic solution, measured, weighed,

scanned, and returned to the river within 30 min of their capture. Age is assessed through scale

reading. In 2015, 2016, and 2017, a total of 850 salmon was examined (366 parrs of age 0 and 484

parrs of age 1) for maturity using a traditional, phenotypic (i.e. external), approach and ultrasound

imaging. All parrs were caught within a period of 8 days in early October. All the animal

experimentation in this study was performed according to French legislation and under licence

APAFIS-201602051204637.

Traditional phenotypic examination

First, the field operator assess the maturation state of parrs by gently pressing the side of the fish.

The excretion of milt is indicative of a mature male, while the absence of milt can be encountered in

a maturing male whose gonads do not yet produce milt, an immature male, or a female (females are

all immature at this stage). The field operator also report that some fish *look like* mature parrs: they

display phenotypic traits similar to those of mature fish (e.g. colour pattern, body shape) but do not

excrete milt. This phenotypic assessment relies less on a fixed set of quantifiable parameters than on

a general, and subjective, appreciation of the fish, which is specific to each operator.

Ultrasound examination

Then, the ultrasound operator screens all salmon parrs for precocious maturation using a portable

ultrasound scanner M-Turbo (Sonosite) with a 5-10 MHz linear transducer. The default setting, for

muscular examination, is selected. As water perfectly transmits ultrasounds, the use of ultrasound

transmission gel is not needed. Fish are placed on their side in a tank of freshwater and the

transducer is operated in the water keeping the transducer 1-2cm above the belly of the fish.

Sagittal images of the abdominal cavity are produced, i.e. the transducer is aligned with the lateral

line. The maturation state of each individual is directly assessed from live images on the ultrasound

monitor. Parrs are classified as mature when gonads are detected, or immature when the gonads

are too small to be detected,. Ultrasound snapshots can be stored on the ultrasound machine and

then exported to a USB memory stick for post-processing.

Comparison of phenotypic and ultrasound methods

The comparison of the proportion of parrs that are recorded each year as mature or immature by

each approach is done with a Chi² test. The number of mature parrs is defined as the sum of

individuals whose gonads are detected with the ultrasound and individuals whose gonads are not

detected with the ultrasound but that produce milt. Because the probability to detect precocious

maturation with the phenotypic method depends on the probability for a mature male to produce

milt at the time of capture, an analysis is conducted to investigate which biological factor drive this

probability. Empirical records show a large interannual variability in the proportion of maturing parrs

(Baglinière and Maisse 1985; Myers et al. 1986). Similarly, the year of capture is expected to affect

milting within mature parrs. Thus, a generalized linear model (GLM) with a binomial distribution of

errors is used to test for a year effect. Then, a mixed model with year as a random intercept and a

binomial distribution of errors is used to test for the effect of age, body length, and their interaction.

A GLM is also run to test whether the probability to miss a mature parr that does not produce milt is

affected by age, body length, operator identity, and time of day (testing for operator tiredness). The

date of capture is not considered because of the low variance in this variable. The significance of the

above-mentioned effects is assessed using the z-value. All models are run in R (R Development Core

Team 2018).

Results and discussion

Ultrasound assessment of precocious maturity

The trained ultrasound operator conduct ultrasound examination for maturation assessment in ca. 5

seconds. In immature individuals, the liver, stomach, caeca, and intestine can be identified (Figure

1a). Note that all internal organs appear more clearly on live images, during ultrasound examination,

than on the snapshots displayed here. Nevertheless, the heterogeneous filling and uneven

granularity of the abdominal cavity are diagnostic of immature individuals, as well as the concave

shape of the cavity, which appears depressed on both sides. In mature parrs (precocious males),

gonads virtually fill the whole abdominal cavity and ultrasound snapshots depict a homogeneous

and finely granulated pattern (Figure 1b, c); digestive organs are hardly visible and the cavity is

convex. In mature males that do not produce milt (Figure 1b), gonads appear less developed and fill

the cavity to a lesser extent than in males producing milt (Figure 1c), which is consistent with a lower

degree of maturity in the former. Interestingly, ultrasound imaging provides the same objective

diagnosis of maturation in males that do not produce milt (Figure 1b) and ones that do (Figure 1c). It

also discriminates truly mature parrs from well-fed individuals, which the operator can mistake for

mature males upon visual inspection because of their round belly.

Figure 1. Assessing maturity in salmon parr using phenotypic view (left panel) and ultrasound

imaging (right panel): a) immature one-year-old parr, b) mature young-of-the-year parr that do not

produce milt (precocious male), and c) mature one-year-old parr that produce milt (precocious male)

in river Oir. On all ultrasound snapshots, the head of the fish is on the left but not shown; therefore

the left flank is towards the bottom of the snapshot and the right towards the top. Gonads are

identified by the letter "G".

Comparison of phenotypic and ultrasound methods

The proportion of mature parrs detected by the phenotypic method is much lower and more

variable (63.8 %, SD = 40.7%) than with the ultrasound method (95.4%, SD = 7.1%, p = 0.199). On

average 33.4% of mature males do not produce milt. A strong year effect is detected (p < 0.001) in

the probability to produce milt at the time of capture in mature males. When accounting for this

year effect as a random factor, a strong increase with age (estimate = 5.582, SE = 1.031, p < 0.001)

and a negative interaction with body length within each age class (estimate = -0.031, SE = 0.008, p <

0.001) are detected. This means that a higher proportion of mature males produce milt in autumn in

old parrs than in young ones, and within a given age class in small parrs than in large ones.

Because the probability to detect precocious maturation with the phenotypic method depends on

the probability for a mature male to produce milt at the time of capture, our result highlights that

both intrinsic and environmental factors can affect the level of detection of the precocious state.

Hence, the ultrasound method unveil that only 16.8% and 13.2% of the mature males are identified

as precocious parrs at the time of capture in 2015 and 2017, and this proportion rise to 87.6% in

2016 (Table 1). Empirical evidence have shown an increase in the proportion of mature parrs as the

parrs are getting older (Baglinière and Maisse 1985; Myers et al. 1986). But this study suggests that

mature parrs develop and produce milt earlier in the season than young ones, which may accentuate

even further the above mentioned age-specific pattern reported with the phenotypic method.

In our study, the maturation state of 106 individuals is uncertain according to the phenotypic

method, i.e. individuals that look like mature parrs. However, following ultrasound examination, it is

found that 89.6 % of those parrs are indeed mature. This result acknowledge the high level of

expertise of our trained operators to assess the maturation state of parrs from external phenotypic

examination. Conversely, no gonad is detected in 10.4% of those parrs (false positive) and the field

operators do not detect milt nor suspect maturity in 5.6 % of the mature males (false negative). This

later proportion differ between the two operators (p = 0.008). None of the other variables had a

significant effect. This highlights how difficult it is to assess parr maturation from external

examination in the field in a fully reliable and replicable manner.

Contrary to the phenotypic method, it is assume that the ultrasound method cannot produce false

positive data, because gonads large enough to be detected by the ultrasound equipment are only

found in mature parrs. The ultrasound operator miss 2.7% of the males producing milt (false

negative). The sample size is too small to assess the potential effect of external factors through

multivariate analysis. Still, this shows that ultrasound imaging is nevertheless subject to a low degree

of uncertainty in maturity assessment. The correct interpretation of ultrasound images depends on

the skill of the operator. Indeed, some training is required to operate the transducer in a fluent way

and navigate between different anatomic layers. In this regard, a good knowledge of the internal

structure for the species of interest is a prerequisite. This also improves the quality of the ultrasound

snapshots and minimizes misinterpretation issues. In hindsight, it is also recommended that all

ultrasounds should be examined and interpreted by multiple operators. This ensures a quick

examination and thus minimizes the impact on fish wellbeing.

Table 1. Comparison of the number of records using external phenotypic observation (rows) and

ultrasound imaging (columns) to assess maturation in parrs each year.

Year	Phenotype method \ ultrasound method	Gonads detected with ultrasound	Gonads not detected with ultrasound
2015	Parr producing milt	17	0
	Parr not producing milt that looks like a mature parr	74	8
	Parr not producing milt that looks like an immature parr	10	73
2016	Parr producing milt	181	2
	Parr not producing milt that looks like a mature parr	20	3
	Parr not producing milt that looks like an immature parr	6	251
2017	Parr producing milt	23	4
	Parr not producing milt that looks like a mature parr	1	0
	Parr not producing milt that looks like an immature parr	3	174

Conclusion

Assessing precocious maturation in parrs from phenotypic observation can reach a high level of

accuracy in trained operators. However, this method relies on some subjective criteria and gut

feeling, which render the level of accuracy in the data difficult to quantify. By allowing to see the

unseen, ultrasound imaging offers great opportunities for ecological studies in the wild. As

illustrated in this paper, key phenotypic traits become accessible and help better characterize the

true biological state of individuals. Ultrasound imaging is an objective, easily transferable and

replicable approach to investigate precocious maturity in parrs, as it relies on key diagnostic

features. A naïve operator can gain a good expertise and assess precocious maturation with the

ultrasound method within a day, whereas more training would be required to achieve a similar level

of expertise with the phenotypic method. In the end, this study calls for the assessment of

precocious maturation in Atlantic salmon parrs using the two methods together: 1) testing whether

parrs produce milt is a quick examination in the field and provides an objective and undeniable

evidence of maturity, and 2) ultrasound imaging should remove any ambiguity about the state of

maturation in parrs that do not produce milt. Therefore, reducing uncertainty in empirical data

should offer new opportunities for further research on this alternative breeding strategy in salmon

and improve our understanding of this key biological process.

References

Baglinière, J.-L., and Maisse, G. 1985. Precocious maturation and smoltification in wild Atlantic

salmon in the Armorican massif, France. Aquaculture **45**(1): 249–263. doi:10.1016/0044-

8486(85)90274-1.

Berejikian, B.A., Van Doornik, D.M., Endicott, R.C., Hoffnagle, T.L., Tezak, E.P., Moore, M.E., and

Atkins, J. 2010. Mating success of alternative male phenotypes and evidence for frequency-

dependent selection in Chinook salmon, *Oncorhynchus tshawytscha*. Can. J. Fish. Aquat. Sci.

67(12): 1933–1941. doi:10.1139/F10-112.

Buoro, M., Prévost, E., and Gimenez, O. 2010. Investigating Evolutionary Trade-Offs in Wild

Populations of Atlantic Salmon (*Salmo salar*): Incorporating Detection Probabilities and

Individual Heterogeneity. Evolution **64**(9): 2629–2642. doi:10.1111/j.1558-

5646.2010.01029.x.

Caswell, H., Naiman, R.J., and Morin, R. 1984. Evaluating the consequence of reproduction in

complex salmonid life cycles. Aquaculture **43**: 123–134.

Chaput, G. 2012. Overview of the status of Atlantic salmon (*Salmo salar*) in the North Atlantic and

trends in marine mortality. ICES J. Mar. Sci. **69**(9): 1538–1548. doi:10.1093/icesjms/fss013.

Clutton-Brock, T., and Sheldon, B.C. 2010. Individuals and populations: the role of long-term,

individual-based studies of animals in ecology and evolutionary biology. Trends Ecol. Evol.

25(10): 562–573.

Fleming, I.A. 1996. Reproductive strategies of Atlantic salmon: ecology and evolution. Rev. Fish Biol.

Fish. **6**(4): 379–416. doi:10.1007/BF00164323.

Genovart, M., Pradel, R., and Oro, D. 2012. Exploiting uncertain ecological fieldwork data with multi-

event capture–recapture modelling: an example with bird sex assignment. J. Anim. Ecol.: no-

no. doi:10.1111/j.1365-2656.2012.01991.x.

Hutchings, J.A., and Myers, R.A. 1994. The evolution of alternative mating strategies in variable

environments. Evol. Ecol. **8**(3): 256–268.

ICES. 2013. Report of the Working Group on North Atlantic Salmon (WGNAS). Copenhagen,

Denmark.

Irvine, J.R., and Fukuwaka, M. 2011. Pacific salmon abundance trends and climate change. ICES J.

Mar. Sci. **68**(6): 1122–1130. doi:10.1093/icesjms/fsq199.

Marchand, F., Azam, D., Delanoë, R., Destouches, J.-P., Tremblay, J., Baglinière, J.-L., Nevoux, M.,

Prévost, E., and Azam, D. 2018. Abundances and biological traits of of fish sampled by

electrofishing (except specific abundance indices) on the River Oir since 1988 (France).

doi:10.15468/yaikgn.

Martin, R.W., Myers, J., Sower, S.A., Phillips, D.J., and Mcauley, C. 1983. Ultrasonic Imaging, a

Potential Tool for Sex Determination of Live Fish. North Am. J. Fish. Manag. **3**(3): 258–264.

doi:10.1577/1548-8659(1983)3<258:UIAPTF>2.0.CO;2.

Mattson, N.S. 1991. A new method to determine sex and gonad size in live fishes by using

ultrasonography. J. Fish Biol. **39**(5): 673–677. doi:10.1111/j.1095-8649.1991.tb04397.x.

Moran, P., Pendás, A.M., Beall, E., and García-Vázquez, E. 1996. Genetic assessment of the

reproductive success of Atlantic salmon precocious parr by means of VNTR loci. Heredity

77(6): 655–660. doi:10.1038/hdy.1996.193.

Myers, R.A. 1984. Demographic Consequences of Precocious Maturation of Atlantic Salmon (*Salmo*

salar). Can. J. Fish. Aquat. Sci. **41**(9): 1349–1353. doi:10.1139/f84-165.

Myers, R.A., Hutchings, J.A., and Gibson, R.J. 1986. Variation in Male Parr Maturation Within and

Among Populations of Atlantic Salmon, *Salmo salar*. Can. J. Fish. Aquat. Sci. **43**(6): 1242–

1248. doi:10.1139/f86-154.

Novelo, N.D., and Tiersch, T.R. 2012. A Review of the Use of Ultrasonography in Fish Reproduction.

North Am. J. Aquac. **74**(2): 169–181. doi:10.1080/15222055.2012.672370.

Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. Annu. Rev. Ecol.

Evol. Syst. **37**(1): 637–669.

R Development Core Team. 2018. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. Available from http://www.R-

project.org.