Engineered inter-species amino acid cross-feeding increases population

evenness in a synthetic bacterial consortium

Marika Ziesack[1], Travis Gibson[3], John K.W. Oliver[1,4], Andrew M. Shumaker[1,5], Bryan B.

Hsu[2], David T. Riglar[1,2], Tobias W. Giessen[2], Nicholas V. DiBenedetto[3], Lynn Bry[3],

Jeffrey C. Way[1,7], Pamela A. Silver[1,2,7*], Georg K. Gerber[3,7*]

[1]Wyss Institute for Biologically Inspired Engineering, Harvard Medical School, Boston,

Massachusetts, USA

[2]Department of Systems Biology, Harvard Medical School, 200 Longwood Ave. Boston,

Massachusetts, USA

[3]Massachusetts Host-Microbiome Center, Brigham and Women's Hospital, Harvard

Medical School, 75 Francis Street, Boston, Massachusetts, USA

[4]Present address: ZBiotics, San Francisco, California USA

[5]Current address: Indigo, Boston, Massachusetts USA

[7]These authors contributed equally.

*correspondence: ggerber@bwh.harvard.edu, Pamela_Silver@hms.harvard.edu

Abstract

In nature, microbes interact antagonistically, neutrally or beneficially. To shed light on

the effects of positive interactions in microbial consortia we introduced metabolic

dependencies and metabolite overproduction into four bacterial species. While

antagonistic interactions govern the wildtype consortium behavior, the genetic

modifications alleviated antagonistic interactions and resulted in beneficial interactions.

Engineered cross-feeding increased population evenness, a component of ecological

diversity, in different environments including in a more complex gnotobiotic mouse gut

environment. Our findings suggest that metabolite cross-feeding could be used as a tool

for intentionally shaping microbial consortia in complex environments.

Importance

Microbial communities are ubiquitous in nature. Bacterial consortia live in and on our

body and in our environment and more recently, biotechnology is applying microbial

consortia for bioproduction. As part of our body, bacterial consortia influence us in

health and disease. Microbial consortia function is determined by its composition, which

in turn is driven by the interactions between species. Further understanding of microbial

interactions will help us deciphering how consortia function in complex environments

and may enable us to modify microbial consortia for health and environmental benefits.

Introduction

In nature, microbes occur as conglomerates of various species with diverse sets of genomes and metabolic capabilities (1). Composition of these consortia arises from networks of ecological interactions and determines function and performance in varying environments. Ecological interactions can be antagonistic (e.g. competition for nutrients), neutral (no interaction) or beneficial (e.g. resource sharing) (2). Studies suggest that both antagonistic and beneficial interactions drive microbial consortia composition (3). According to classic evolutionary theory, competitive interactions favoring survival of one species over another should be prevalent, and a body of experimental research supports that hypothesis (4–6). Recent studies of the microbiota also highlight the importance of beneficial interactions, particularly those that provide "public goods" (7, 8). "Public goods" are metabolites that are released into the environment, and not only benefit the producer but also neighboring cells.

A meta-analysis of 77 studies revealed that cross-feeding of metabolites between naturally occurring bacteria is common (7). Metabolite cross-feeding may be explained by comparative advantages of metabolic capabilities in some species over others, and removal of biochemical conflicts in biosynthetic pathways within cells (9, 10). The evolution of such mutually beneficial interactions may occur as follows. Metabolic leakiness in strain A can serve as the first step towards metabolic dependencies as it results in products leaking into the environment where neighboring strain B can take them up (11, 12). Strain B may lose the ability to make the metabolites itself, resulting in auxotrophy. Indeed, microbial auxotrophies are common in natural communities. In a

study of 949 eubacterial genomes, 76% were found to lack at least one of 25 different

metabolic biosynthesis pathways (13).

There is increasing recognition that maintaining species diversity – a measure of

richness and evenness within a population - in microbiomes is important to ecosystem

health (14). A study that analyzed composition of a syntrophic consortium of

methanotrophs found that amino acid cross-feeding may help with retaining species

diversity which leads to higher robustness with regard to limiting loss or dominance of

species in the consortium (15). Despite the fact that bacteria competed for similar

nutritional niches, the consortium composition remained stable. Thus, metabolite cross-

feeding could serve as a mechanism to preserve species diversity by limiting the ability

of some consortium members to dominate and eliminate other species.

Amino acid cross-feeding is also an attractive means to introduce cooperation into

synthetic microbial consortia. Numerous studies have engineered pairwise amino acid

cross-feeding in *E. coli* with native amino acid production levels and generated

quantitative models to describe their behavior (16–21). Most prior synthetic biology

studies that have engineered cooperativity in bacterial communities have used a single

species (21–24). However, natural microbial ecosystems contain a diversity of

interacting species.

In advancing synthetic biology to real applications in complex environments, it will be

useful to expand engineering capabilities to diverse, multi-species consortia.

Importantly, bacteria from naturally occurring ecosystems are likely to have pre-existing

interactions, which are often competitive. In a given bacterial consortium that is

dominated by antagonistic interactions, synthetically introduced positive interactions

need to overcome these antagonistic interactions in order to cause a net beneficial

effect. Here we engineer metabolite cross-feeding in a synthetic consortium of four

different bacterial species and demonstrate net beneficial behavior. We further observe

increased population evenness–a component of species diversity within a population–

in different environments *in vitro* and *in vivo*.

Results

Engineering of metabolic dependency and overproduction between four species.

For our consortium, we chose four bacterial species that naturally reside in the

mammalian gut, a complex environment of intense interest both from the basic scientific

and biomedical perspectives (Figure 1a). The organisms were chosen in part for their

propensity to colonize in the colon, which we hypothesized could allow for sufficient

cross-feeding opportunities. A second crucial criteria for the selected organisms was the

availability of tools for efficient genetic manipulation; to date, a very small subset of

commensal gut microbes are genetically tenable and thus a wider range of organisms

was not feasible. Nonetheless, these organisms represent gut residents with different

spatial niches and carbohydrate utilization preferences: *E. coli* and *S.* Typhimurium tend

to utilize simpler carbohydrates as found in the gut lumen (25, 26), whereas *Bacteroides*

spp. tend to be mucosally associated and utilize more complex carbohydrates (27).

Escherichia coli NGF-1 was originally isolated from BALB/c mice, has been shown to

stably colonize the mouse gut, and can be engineered with standard genetic tools (28–

30). *Salmonella enterica* serovar Typhimurium LT2 (*S.* Typhimurium), an attenuated gut

pathogen, was rendered further harmless by removing the pathogenicity islands SPI1

and SPI2, and thus causes no disease when administered to mice (31). The two

Bacteroides species, *Bacteroides thetaiotaomicron* and *Bacteroides fragilis,* are human

commensals that can achieve high abundance in the mammalian gut, and are also

genetically tractable.

We engineered each of the constituent species to depend on three of the four

metabolites L-methionine, L-histidine, L-tryptophan and L-arginine (hereafter referred to

as Met, His, Trp and Arg) and to overproduce the remaining amino acid (Figure 1a). As

we noted, amino acid cross-feeding has been described as a prominent natural

mechanism (7) and has previously been used extensively in single species engineering

efforts. Our rationale for choosing these specific amino acids for cross-feeding was in

part based on feasibility of engineering overproduction and auxotrophies in the selected

species based on prior work. Additionally, we considered the amino acid composition of

proteins across a variety of bacterial taxa, which indicated lower use of Met, His and

Trp, and higher for Arg (32). We therefore hypothesized that these particular amino

acids would offer a range of nutrient requirements for testing and evaluating consortium

behavior.

The auxotrophies introduced for each strain were: *E. coli*: His, Trp and Arg;

*S.*Typhimurium: Met, Trp, Arg; *B. thetaiotaomicron*: Met, His, Arg; *B.fragilis*: Met, His,

Trp. *E. coli* and *S.* Typhimurium were engineered by sequential phage transduction from

three single auxotroph strains. *E. coli* was transduced with genome fragments from

BW25113 that contained insertions in argA, trpC, hisA (see Methods) and *S.*

Typhimurium with genome fragments of the same parent strain with insertions in argA,

trpC, metA. *Bacteroides spp.* triple knockout generation utilized the pExchange-tdk

vector to precisely delete *metA*, *hisG* and *argF* in *B. thetaiotaomicron* and *metA*, *hisG*

and *trpC* in *B. fragilis*. We introduced the ability to overproduce one amino acid in each

strain by using anti-metabolite selection methods (*E. coli*: Met; *S.* Typhimurium: His; *B.*

thetaiotaomicron: Trp; *B. fragilis*: Arg) and identified putatively causal mutations by

sequencing and SNP analysis (Table 1, Methods).

To assess the auxotrophic strains' requirements for the metabolite, we measured

growth on varying concentrations of each metabolite in the presence of non-limiting

concentrations of all the other metabolites (Figure S1). We found that each strain has

distinct requirements for the amino acids that it was auxotrophic for. In general, the

Bacteroides spp. have higher amino acid requirements compared to *E. coli* and *S.*

Typhimurium, which could in part be due to differences in their capability to

transport/uptake amino acids. We note also that the observed amino acid requirements

correlate with differing amino acid compositions in proteins of these species (33).

Overproduction of metabolites was measured in comparison to a defined amino acid

standard using LC-MS (Figure 1b, filled bars, Figure S6). In order to compare

overproduction with each species' amino acid requirements, we fit a sigmoidal curve to

the growth response data and calculated the EC50 requirement (Figure S1, and 1b,

shaped bars). In this context, the EC50 requirement describes the concentration of

supplemented metabolite that allows for half-maximum growth of the respective strain.

These data on amino acid requirements and overproduction levels provide information

about the expected relative strengths of the engineered beneficial interactions.

Engineered *B. fragilis* overproduced Arg at 362 uM - a concentration exceeding EC50

requirements of any of the other strains. Corresponding supplementation allows for

more than half-maximum growth of either one of the strains. *B. thetaiotaomicron*

overproduced Trp at 34 uM, which exceeded the EC50 requirements of *E. coli* and *S.*

Typhimurium but not *B. fragilis*. Overproduction from *E. coli* (Met at 5.2 uM) and *S.*

Typhimurium (His at 16 uM) were lower than any of the EC50 requirement values. EC50

requirement values exceeded *E. coli* Met overproduction by 4-, 7-, and 6-fold and *S.*

Typhimurium His overproduction by 12- ,51- and 17-fold. Overall, these findings suggest

relatively strong cross-feeding from *Bacteroides spp*, moderate cross-feeding from *E.*

coli, and the weakest cross-feeding from *S.* Typhimurium to other strains. Note that WT

E. coli and *S.* Typhimurium do not produce detectable levels of the respective amino

acids in supernatant, whereas WT *B. theta* produces 5.7 ± 0.8 µM Trp and WT *B.*

fragilis produces 1.7 ± 1.2 µM Arg (Figure S6).

To provide further insight into the ability of each overproducer strain to support the

auxotrophic strains, we performed assays in which each auxotrophic strain was grown

in the presence of supernatant from the overproducer strain or its WT counterpart

(Figure 1c). Overall, our overproduction and requirement assay described above

predicted growth in the supernatant assay. The majority of WT strain supernatants

supported no to minimal growth of the auxotrophs, whereas the overproducer

supernatants variably supported growth depending on the match between

overproduction and metabolite requirements of the respective pair of strains (Figure 1d).

Exceptions to this observation were WT *B. theta* supernatant, which supported some

growth of *E. coli* and *S.* Typhimurium, and WT *B. fragilis* supernatant, which supported

some growth of *B. theta*. This is consistent with our finding that the WT strains of these

organisms produced a small amount of the respective amino acid (Figure S6). Growth

of *E. coli* was rescued by supernatant from overproducing *B. theta* and *B. fragilis* but not

S. Typhimurium. *S.* Typhimurium growth was rescued by all overproducer supernatants.

B. theta growth was partially rescued by *E. coli* and *B. fragilis* but not *S.* Typhimurium

overproducer supernatant. *B. fragilis* growth was not rescued by any of the strains'

overproducer supernatants but is rescued in co-culture (see below).

Engineered cross-feeding introduces beneficial interactions in the consortium.

To characterize the behavior and interaction structure of the consortium as a whole, we

performed a set of experiments in which we cultured all members together while varying

the starting inoculum of one member at a time. By measuring the growth trajectories of

all strains simultaneously, we can analyze how fluctuations in the growth of one

member of the consortium influences changes in the growth of other members over

time, and thus infer the interactions that are occurring. We performed these experiments

under five conditions for both WT and engineered consortia.

In condition 1, which served as the baseline, we inoculated all consortium members at

equal ratios (Figure 2 a, b), and in conditions 2-5 we reduced the starting inoculum of

each of the four species in turn by a factor of 10 (Figure 2 c,d). We grew the consortia

anaerobically at 37 °C for 27 hours in M9 minimal media with specific modifications as

described in Methods, without supplementation of any of the cross-fed amino acids, and

with 0.5% starch and 0.5% glucose as carbon sources. We assessed growth of each

strain in co-culture over time via strain-specific qPCR.

We were able to record growth of each strain in WT consortia indicating that conditions

were permissive for each strain (Figure 2a). We also recorded growth of each strain in

the engineered consortia, which indicates that cross-feeding capabilities were sufficient

to support growth (Figure 2b). As expected based on our previous experiment the

engineered consortia did not fully rescue all of the auxotrophies and hence grew to

about 10-fold lower overall cfu/mL. In the WT consortium, knockdown of one species

persists throughout culturing, whereas in the engineered consortium *S.* Tyhpimurium

and *B. theta* are able to recover to baseline ratios (Figure 2 c,d). Two further

characteristics of the engineered consortia are notable. In all five conditions, *S.*

Typhimurium appears to dominate the culture, and *B. fragilis* knockdown causes a

decrease in overall consortia growth by 2-fold.

We note that no *B. fragilis* growth in supernatant was observed, whereas we do see

growth in the co-culture setting. This difference may be due to several reasons. One

possibility is that in the co-culture setting *B. fragilis* may have time to grow to sufficient

biomass such that the growth inhibition is blunted, whereas in the supernatant setting

the other organisms can accumulate high concentrations of substances that inhibit initial

B. fragilis growth. Alternately, other consortium members may be able to actively buffer

negative effects. For example, we have observed that *S.* Typhimurium acidifies the

media, which, without appropriate buffering, strongly decreased *Bacteroides spp.*

growth (data not shown).

To visualize how lowering the inoculum of each strain ("knockdown") affected growth of

the other strains, we plotted log-fold ratios (logR) comparing the "knockdown" condition

to the equal inoculum condition (Figure S2, S3). Positive logR values for strain A

indicate more growth of that strain in the absence of the "knockdown" strain B,

suggesting a net antagonistic effect of strain B on strain A; conversely, a negative logR

value suggests a net positive effect. We see either negligible or positive logR values for

the WT consortium (Figure S2), suggesting that it is dominated by antagonistic

interactions. In contrast, for the engineered consortium we see negligible or negative

logR values (Figure S3), suggesting antagonistic interactions have been reduced and

positive interactions were introduced.

To quantitate interactions present in the consortium, taking into account both direct and

cascading effects, we used a statistical method based on well-established techniques

for inferring microbial interaction networks from time-series data (33–36). Briefly, the

method models the rate of change in growth of species A as a function of the

concentrations of itself and all other species present in the ecosystem. (See the

Methods section for further details.) Applying our inference method to the data

presented in Figure 2 a-d, we obtained a network of inferred interactions and

accompanying estimates of confidence in each edge in the network (Figure 2 e, f,

Supplemental Figure S4). Consistent with our qualitative visualization, all inferred

interactions for the WT consortium were antagonistic (Figure 2 e). In contrast, we see

two inferred positive interactions in the engineered consortium network, from *B. fragilis*

to *E. coli* and *S.* Typhimurium with high inference confidence, which is also consistent

with our qualitative visualization (Figure 2 f). Further, although some inferred negative

interactions are still present in the engineered consortium network, the confidence in

these predictions is much lower than in the WT network, indicating that the model finds

greatly reduced evidence for antagonistic interactions in the engineered consortium.

Taken together, our findings indicate that introducing metabolic dependencies and

overproduction capacities alleviated antagonistic interactions and introduced some

beneficial interactions in the consortia.

Engineered consortia show higher population evenness than WT counterpart *in*

vitro

To gain insight into how beneficial interactions affect consortia behavior, we compared

consortia compositions in environments with different nutrient levels (Figure 3). We

grew consortia in batch culture for 24 hr anaerobically in modified M9 media as

described in Methods with and without supplementation of 1 mM of the cross-fed amino

acids and 0.2% starch and 0.5% glucose as carbon sources. As described in the

previous trajectory experiment the engineered consortia were able to grow without

amino acid supplementation albeit to lower capacity compared to its WT counterpart.

Addition of cross-fed amino acids rescued growth of the engineered strains to levels

equivalent of the WT counterparts. This is in accordance with previously described

consortia characterization (Figure 1) which shows that not all overproduction levels

were sufficient to fully rescue growth of all species.

An important ecological measure of consortium behavior is its population evenness in

different environments, which is frequently computed using Pielou evenness index. This

measure indicates the extent to which abundances of different species in an ecosystem

are even or balanced; a value of 1.0 indicates equal abundances of all species,

whereas lower values indicate an ecosystem dominated by a subset of the total species

present, and a value of 0 indicates one species completely dominates the community.

The highest population evenness occurred in the engineered consortia without amino

acid supplementation, a setup in which cross-feeding via engineered overproduction

and auxotrophies was expected to be maximized. The WT consortium also exhibited a

similar trend, showing higher population evenness when amino acid supplementation

was omitted. However, the population evenness attained by the WT consortium in this

condition was still considerably less than that of the engineered consortium in the same

condition. Overall, our results suggest that introduction of cross-feeding can increase

population evenness in conditions favoring these beneficial interactions.

Engineered consortia show higher population evenness than WT counterpart in

the mouse gut

We investigated the behavior of our consortium in the mammalian gut, using gnotobiotic

mice as a controlled yet relatively complex environment for evaluation. To investigate

the role of amino acid cross-feeding *in vivo*, we altered amino acid levels in the gut by

changing the animal's diet (37). Four groups of five germfree mice were fed standard or

low protein (3%) chow, and gavaged with either the WT or engineered consortium

(Figure 4). The consortia were allowed to colonize for 10 days, and then stool samples

were collected and interrogated via qPCR with species-specific primers.

We observed consistently higher abundances of all strains in both engineered and WT

consortia in mice that were fed with the standard diet. For the engineered consortium,

species abundances were higher by a factor of approximately 3 for *E. coli*, 8 for *S.*

Typhimurium, 16 for *B. thetaiotaomicron*, and 11 for *B. fragilis* in mice fed a standard

versus a low protein diet. In the case of the WT consortium, *S. Typhimurium*, *B.*

thetaiotaomicron and *B. fragilis* concentrations were similarly higher on standard chow

(fold changes of approximately: 10, 22, 13 respectively), and WT *E. coli* concentrations

were considerably higher (~50-fold higher).

Although overall strain abundance was higher in mice that were fed with standard chow,

the engineered consortium in mice that were fed with a low protein diet showed

significantly greater population evenness (Mann-Whitney test; *p*-values: 0.02, 0.03,

0.02). These findings are consistent with our *in vitro* results showing the greatest

population evenness in the engineered consortium in conditions with low concentrations

of the cross-fed amino acids. We also observed a trend toward greater population

evenness in the engineered consortium in mice fed standard chow compared to the WT

consortium in mice regardless of diet, although this trend was not statistically significant.

Interestingly, in mice that were fed with a low protein diet, *S.* Typhimurium grew about

8-fold better in the engineered consortium compared to the WT consortium. This finding

is consistent with our *in vitro* results, which indicated that engineered *S.* Typhimurium

benefits most from the consortium cross-feeding effects. We observed the same trend

in mice that were fed with standard diet, albeit to a lesser extent

Discussion

We engineered inter-species amino acid cross-feeding in a bacterial consortium and

demonstrated that this increased population evenness in environments of varying

complexity. The ability of an ecosystem to maintain evenness, an important component

of species diversity, in the face of different environments or perturbations to the same

environment is a critical component to its robustness or limiting dominance of some

ecosystem members at the expense of others. We showed that pre-existing

antagonistic interactions in a bacterial consortium can be overcome by engineered

beneficial interactions (Figure 2). The engineered consortium exhibited increased

population evenness *in vitro*, particularly in nutrient sparse conditions when cross-

feeding was favored (Figure 3).

We found that the engineered consortium also showed increased population evenness

in the gnotobiotic mouse gut (Figure 4). Gnotobiotic mice provide a tractable model for

the mammalian gut environment, maintaining the spatial organization and metabolic

composition of the host. However, a limitation is that gnotobiotic mice lack the myriad of

bacterial species present in a naturally occurring commensal microbiota (38). Despite

this limitation, our results provide evidence that engineered bacterial cross-feeding

behavior can function in a complex environment and maintain increased population

evenness. Notably, while the engineered consortium attained the highest population

evenness *in vitro* in conditions lacking the cross-fed amino acids, this was at the

expense of overall biomass. However, in the gnotobiotic mouse gut, the engineered

consortium attained about the same biomass as its WT counterpart in mice receiving

the same diet. Given that the strains in the consortium all originated from the

mammalian gut, these findings reinforce the fact that *in vitro* conditions may not mimic

important aspects of the host environment such as availability of complex nutrients and

chemical environments in the gut as well as differing spatial distribution of bacterial

species, i.e., in large versus small bowel or luminal versus mucosal surfaces. An

interesting direction for future work would be to better characterize the niches of the

consortium member in the gut, such as through *in situ* localization assays, and

determine whether engineering alters spatial localization and other niche behavior.

Engineering diverse, multi-species consortia presents interesting challenges while

holding promise for substantial improvements for applications in human or animal health

and bioproduction. Bacteria from naturally occurring ecosystems will have pre-existing

interactions, which are often competitive (4–6). In a given bacterial consortium that is

dominated by antagonistic interactions, synthetically introduced positive interactions

need to overcome these antagonistic interactions in order to cause a net beneficial

effect. We found that in our consortium there was considerable bias toward naturally

occurring competitive interactions. Thus, any engineering of these organisms had to

overcome this inherent negative bias. We engineered bacteria to overproduce

metabolites at different concentrations from low (*S.* Typhimurium, His) to high (*B.*

fragilis, Arg) (Figure 1). This enabled us to assess the effects of various levels of

overproduction in a single consortium. *B. fragilis* overproduction achieved the most

obvious beneficial interactions and its knockdown *in vitro* had the most profound effect

on consortium growth (Figure 2). Overproduction by the other strains led to reduction of

pre-existing antagonistic interactions, but not to the extent of producing fully beneficial

interactions.

Many studies on engineered cross-feeding in microbial consortia have focused on

knockouts and complementary metabolism but not overproduction (18, 21). Our findings

suggest that relatively high overproduction rates are required in a consortium where

multiple members cross-feed from a single or small number of overproducers. We did

not observe reduced growth of the overproducing strains, with the exception of *B.*

thetaiotaomicron, which showed a 3-fold reduction in growth (Figure S5). Our findings

have implications for naturally occurring systems, and merit further study to understand

the extent to which robust cross-feeding has evolved to depend on multiple producers

versus high output single producers.

We observe that our engineered consortium exhibits increased evenness at the cost of

reduced fitness; the engineered consortium grows at ~10-fold lower concentration than

the WT counterpart (Figure 2). This is likely in part due to the fact that some of the

bacteria overproduce less than is required by the other members (Figure 1). This

imbalance may also make the consortium susceptible toward cheaters; in fact *E. coli*

and *S.* Typhimurium could be viewed as "moochers" that benefit most from *Bacteroides*

spp. production but do not contribute much themselves. Future work could seek to

optimize the balance of overproduction and consumption rates of the consortium

members to improve overall consortium fitness while preserving population evenness,

using approaches such as rationally engineering overproduction pathways and amino

acid uptake transporters in combination with directed evolution via turbidostat.

Our finding that metabolite cross-feeding can lead to increased population evenness,

which is an important aspect of species diversity, may have implications for

understanding naturally occurring microbial ecosystems as well as engineered

communities for biotechnology applications. Indeed, prior work has shown that in a

methanotrophic bacterial consortium, amino acid cross-feeding helps to maintain

species diversity (15). Species diversity is clearly important in the robustness of

microbial ecosystems; an ecosystem with greater genetic diversity as a whole has the

potential to better survive in changing environments (e.g. levels of cross-feed nutrients

for our consortium). The gut microbiota constitutes a complex host-microbial ecosystem

and maintaining species diversity in this context has important implications for human

health. Our internal microbial ecosystems undergo a variety of alterations due to factors

including normal physiologic changes such as exercise or pregnancy, pathologic states

due to disease, modifications to our diets, and medical interventions. Thus, mechanisms

such as cross-feeding that promote maintenance of species diversity in the human

microbiome in the face of perturbing factors may be critical to maintaining host function.

From the perspective of an ecosystem, there is a long-term evolutionary fitness

advantage to maintaining species diversity, which may be supported by naturally

occurring metabolite cross-feeding mechanisms such as those we introduce artificially

here.

Acknowledgement

We thank Laurie Comstock for providing *B. fragilis* 638R and helpful advice on *Bacteroides* handling. This project was funded by the Defense Advanced Research Program Agency (DARPA BRICS HR0011-15-C-0094), NIH T32 HL007627 (to T.G.), the Harvard Digestive Diseases Center P30 DK034854 and The Wyss Institute for Biologically Inspired Engineering. D.T.R. is supported by a Human Frontier Science Program Long-Term Fellowship and a National Health and Medical Research Council (NHMRC) RG Menzies Early Career Fellowship from the Menzies Foundation Australia.

Methods

Auxotroph engineering

For auxotroph generation in the *E. coli* NGF-1 strain we introduced multiple knockouts using

sequential P1 transduction (39) from the Keio knockout collection (40). Flip-out of kanamycin

cassettes was done using pCP20 (41). In brief, for P1 transduction we prepared phage by

diluting an overnight culture of the donor strain 1:100 LB with 0.2% glucose, 5 mM $CaCl_2$ and 25

mM $MgCl_2$ and incubated for 1-2 hours at 37 °C until slightly turbid. We then added 40 µL P1

lysate and continued growth for 1-3 h at 37 °C while shaking until lysed. Lysate was then filtered

with a 20 µm sterile filter and stored in the fridge. For transduction, we harvested 2 mL overnight

culture of recipient strain and re-suspended in 2 mL LB with 5 mM $CaCl_2$ and 100 mM $MgSO_4$.

We then mixed 100 µL donor lysate with 100 µL recipient, incubated 30 min at 37 °C and added

200 µL sodium citrate (1 M, pH 5.5) and 1 mL LB and incubated for another 1 hr at 37 °C. Cells

were harvested, re-suspended in 100 µL LB with 100 mM sodium citrate and plated on LB Kan

plates (75 µg/mL). The transduced kanamycin cassette was then removed using pCP20

according to protocol. We transformed pCP20 via electroporation and transformants were

selected on LB agar plates supplemented with 100 µg/mL carbenicillin grown at 30 °C. Single

colonies were re-streaked on LB without drugs and incubated for 10 hours at 42 °C. From there,

single colonies were re-streaked on LB plates without drugs and grown overnight at 37 °C.

Colonies were checked for Carbenicillin, and Kanamycin sensitivity and further confirmed via

PCR at respective loci. This procedure was repeated until all knockouts were introduced.

Engineering of *S.* Typhimurium LT2 required generation of single knockout strains that

contained pKD46 integrated into the genome, which allowed for linear DNA integration using

lambda red recombination (41). We then introduced the knockouts into the *S.* Typhimurium

strain through sequential P22 transduction and pCP20 flipout analogous to *E. coli* engineering.

Single knockout strains were generated by PCR amplifying a Kanamycin resistance cassette

from pKD13 generating linear fragments that contained upstream and downstream homology to

the gene of interest and the kanamycin cassette with FRT sequences. Fragments were

introduced via electroporation and selected on LB agar plates supplemented with 50 µg/mL

Kanamycin. Sequential P22 transduction and pCP20 flip-out was essentially performed as

described above for P1 transduction but lysis was done overnight.

For knockout generation of both *B. thetaiotaomicron* and *B. fragilis*, we used pExchange KO

vectors as described (42). Briefly, we introduced 750 bp flanking regions for genes of interest

adjacent to each other into the vector. The vector contains an erythromycin resistance positive

marker and a thymidine kinase as counter selection marker. Cloning was done in pir$^+$ *E. coli*

strains and vectors were transferred to MFDpir for conjugation (43). Conjugation was done

according to protocol with minor changes. In brief, five drops of overnight culture of *E. coli* donor

was inoculated in LB supplemented with 300 µM Diamino pimelic acid (DAP) and five drops of

recipient overnight culture was inoculated in 50 mL basal media. Both cultures were grown for

about 2 hr (*E. coli* aerobically, *Bacteroides spp.* anaerobically) until *E. coli* culture was well

turbid and *Bacteroides* culture just slightly turbid. Subsequently, 9 mL recipient and 3 mL donor

were combined and spun down for 10 min at 4000 rpm together. The pellet was re-suspended

in 100 µL fresh basal media with 300 µM DAP and pipetted on basal media agar plates without

cysteine and supplemented with 300 µM DAP. The cells were incubated at 37 °C aerobically

face up for up to 20 hr, scraped off and re-suspended in 10% glycerol. Dilutions were plated on

basal-agar plates supplemented with 10 µg/mL erythromycin and incubated at 37 °C

anaerobically for 2-3 days. Single colonies were re-streaked in the presence of erythromycin

and grown for another 2 days. 10 single colonies were inoculated in basal media without drug

and grown overnight. 500 µL of each culture was mixed, spun down and re-suspended in 10%

glycerol. We then plated different dilutions on basal media plates supplemented with 5-fluoro-2-

deoxy-uridine (FuDR) (200 µg/mL) and incubated at 37 °C anaerobically for 3 days. Knockouts

were verified via PCR. This procedure was repeated multiple times to obtain the multiple

auxotroph strains.

Overproducer selection

Overproducers were generated by selecting for mutants that could grow on minimal media agar

plates supplemented with anti-metabolites (*E. coli*: 5 mg/mL Norleucine for Met overproduction;

S. Typhimurium: > 0.7 mg/mL beta-(2-thiazolyl)-DL-alanine for His overproduction; *B.*

thetaiotaomicron: 50 µg/mL 4-methyl tryptophan for Trp overproduction; *B. fragilis*: 80 µg/mL

Canavinine for Arg overproduction). Single colonies that showed halos were re-streaked and

overproduction was measured using a bioassay. In brief, for screening of overproducing

mutants the isolated strains were grown overnight at 37 °C shaking aerobically (for *E. coli* and

S. Typhimurium) or anaerobically without agitation (for *Bacteroides spp.*). Supernatant was

harvested, diluted 1:1 with fresh media, and *E. coli* auxotrophs were inoculated and their growth

was recorded after 24 hr. For *E. coli* NGF-1 overproducers, we used a *S.* Typhimurium

auxotroph instead, since its colicin production prevented the *E. coli* biosensor from growing.

Confirmed overproducers were further quantified using LC-MS.

LC-MS for Overproduction Measurements

To quantitate amino acid levels in overproducer supernatants, a standard curve was obained

using freshly prepared amino acid standards dissolved in growth media (1mM, 500 uM, 100 uM,

50 uM, 10 uM of L. Methionine, L/Histidine, L-Tryptophan, L-Arginine each). To prepare for

HPLC-MS analysis, 0.5 mL sample or standard were added to 1.5 mL ice-cold methanol and

incubated on ice for 10 min. The mixture was centrifuged for 5 min at 15,000 rpm and 500 µL

supernatant was vacuum concentrated and re-suspended in 50 µL methanol. Samples were

kept on ice or at 4°C. HPLC-MS analysis of standards and extracts was carried out using an

Agilent 1260 Infinity HPLC system equipped with an Agilent Eclipse Plus C18 (100 × 4.6 mm,

particle size 3.5 mm, flow rate: 0.3 mL/min, solvent A: dd.H_2O/0.1% (v/v) formic acid, solvent B:

acetonitrile, injection volume: 4 mL) connected to an Agilent 6530 Accurate-Mass Q-TOF

instrument. The following gradient was used (time/min, %B): 0, 0; 0.5, 0; 14, 100; 19, 100; 20, 0,

25, 0. The mass spectrometer was operated in positive mode and the autosampler was kept at

4°C. After HPLC-MS analysis, extracted ion current (EIC) peaks were automatically integrated

using the MassHunter Workstation Software (version: B.07.00). A plot of peak area versus

amino acid concentration was used to generate a linear fit.

Sequencing

Bacterial cultures were prepared in rich media (basal for *Bacteroides spp.* and LB for *E. coli* and

S. Typhimurium). Genomic DNA (gDNA) extraction was performed using the Wizard Genomic

DNA Purification Kit (Promega) according to protocol. The extracted gDNA was sheared using

Covaris DNA Shearing, and the library was prepared using Kapa Biosystem DNA Hyper Prep

NGS Library (Dana Faber Core MBCFL Genomics). Sequencing was performed on the Illumina

MiSeq instrument, with the 150 bp paired End (PE150) reagents. Sequences were analyzed for

SNPs using Geneious software and published genome sequences (*E. coli*: CP016007.1; *S.*

Typhimurium: NC_003197; *B. thetaiotaomicron*: AE015928; *B. fragilis*: NC_016776).

Growth and Media Conditions

All basal media (20 g/L proteose peptone, 5 g/L yeast extract, 5 g/L sodium chloride, 1 g/L L-

cysteine, 0.5% glucose, 5 mg/mL potassium phosphate, 5 mg/L hemin) and co-culture media

(M9 salts (0.2 g/L Na_2HPO_4, 90 mg/L KH_2PO_4, 30 mg/L NH_4Cl, 15 mg/L NaCl), 1 mM $MgSO_4$, 10

μg/mL heme, 0.1 mM $CaCl_2$, 1 μg/mL Niacinamide, vitamin B12 and thiamine, 400 μg/mL L-

cysteine, 0.3% bicarbonate buffer, 2.5 ng/mL vitamin K, 2 μg/mL $FeSO_4$*$7H_2O$ and carbon

sources and amino acid supplementation as described in Results) were pre-incubated for at

least 24 hr anaerobically before use. In initial experiments we used 0.5% starch, which we later

optimized to 0.2% (resulting in lower viscosity); the observed growth differences were not

statistically significant between those two conditions (Figure S7). *Bacteroides spp.* were

inoculated from glycerol stock into basal media, grown overnight and 400 μL was inoculated in 5

mL basal and grown 2 hr anaerobically. Cells were spun down, washed twice in PBS and

diluted in growth media as described for each experiment in Results. *E. coli* and *S.* Typhimurium

were inoculated from glycerol stock into LB and grown overnight at 37 °C while shaking. 100 μL

of culture was then inoculated into pre-incubated LB and grown anaerobically for 2 hr, diluted,

washed in PBS and diluted into co-culture media as described.

Multiplex qPCR

We designed strain specific primer/probe-fluorophore pairs according to IDT protocol (Table

S1). We chose strain specific genes by multiple genome alignment between the strain of

interest, the other consortia members and closely related strains using Mauve (44). Multiplex

qPCR was used to quantify each strain in co-culture by using a standard curve obtained by

plating late log phase cultures grown in rich media. In brief, each strain was grown from

overnight culture for ~5 hours until about OD of 1. Cells were then counted by plating. Cultures

were mixed, diluted and frozen at -80 °C for use as standard curve. Samples were diluted 1:10

in ddH_2O and snap-frozen in liquid nitrogen and stored at least overnight at -80 °C. Growth

curve and sample were both thawed together and prepared in a 5 μL Primetime Mastermix

(IDT) with 1 μL Primer/Probe mixture (final concentrations: 100 nM for primers and 50 nM for

probes). The qPCR was run with the following program: 20 min at 98 µC (to boil the cells and

denature gDNA), followed by 40 cycles of 60 °C and 98 °C.

Statistical inference of interactions within consortia

We adapted our previously published statistical method for inferring a dynamical systems model

from microbial time-series data (33, 34). Briefly, our model assumes the rate of change over

time of each microbe in the ecosystem is related to its own abundance as well as abundances

of the other microbes in the ecosystem. Our model is fully Bayesian and infers the posterior

probability distribution over the qualitative network of microbe-microbe interactions (e.g.,

probability of an edge being present in the network) as well as the quantitative strengths of the

interactions present. As we have previously described (33), the qualitative network can be

interpreted in terms of Bayes factors (BF), a standard Bayesian alternative for hypothesis

testing that quantifies the evidence for one model over another. For our method, the BF on

network edges quantifies the evidence for existence of the interaction versus absence of the

interaction. A standard interpretation of BFs is: BF between 3 and 10 = substantial evidence, BF

> 10 = strong evidence.

To be precise, our model is based on discrete time stochastic generalized Lotka Volterra (gLV)

dynamics:

$$x_{k+1,i} = x_{k,i} + x_{k,i}\left(r_i + \sum_{j=1}^{4} a_{i,j} z_{i,j} x_{k,j}\right)\Delta_k + \sqrt{\Delta_k}\left(w_{k+1,i} - w_{k,i}\right)$$

548   where $x_{k,i}$ is the abundance of species i at time point k; r_i is the growth rate of species i; $a_{i,j}$ is

549   the interaction coefficient and $z_{i,j}$ is a binary variable indicating the presence or absence of an

550   edge (interaction) between species j and i; Δ_k is the time difference between timepoints $k + 1$

551   and k; and $w_{k,i}$ is a Brownian motion term. The Brownian motion has variance v_i^w. We place

552   prior probability distributions on the variables defined above:

$$z_{i,j} \sim \text{Bernouli}(0.5)$$

$$r_i \mid v^r \sim \text{Normal}(0, v^r)$$

$$a_{i,j} \mid v^a \sim \text{Normal}(0, v^a).$$

553   Note that our prior probability distribution for the indicator variable, $z_{i,j}$, for presence or absence

554   of interactions, expresses maximum uncertainty, e.g., no *a priori* assumption about the

555   presence or absence of an interaction. We place the following prior probability distributions on

556   the variance parameters in our model:

$$\boldsymbol{v}^r \sim \text{Inv} - \chi^2(\eta_r, \theta_r)$$

$$\boldsymbol{v}^a \sim \text{Inv} - \chi^2(\eta_a, \theta_a)$$

$$\boldsymbol{v}_i^w \sim \text{Inv} - \chi^2(\eta_w, \theta_w).$$

In our model $\eta_r, \eta_a, \eta_w = .1$ with $\theta_r = .01$, $\theta_a = 10^{-14}$, and $\theta_w = 10^4$, specifying relatively diffuse

(uninformative) priors. We perform inference using a simplified version of our previously

described Markov Chain Monte Carlo (MCMC) algorithm that omits clustering (34), with 5,000

iterations (1,000 burn in iterations.)

The gLV model described above allows for only pairwise interactions, so we also tested an

extended model with higher order interactions. Specifically, we included 3[rd] order or 4[th] order

interactions, i.e.,

$$+\boldsymbol{z}_i^{(3)} \boldsymbol{a}_i^{(3)} \prod_{j \neq i} x_{k,j} \quad \text{or} \quad +\boldsymbol{z}_i^{(4)} \boldsymbol{a}_i^{(4)} \prod_{j=1}^{4} x_{k,j}$$

appearing on the right-hand side of the gLV dynamics equation above. The superscripts (3) and

(4) denote higher order indicator variables $\boldsymbol{z}_i^{(\cdot)}$ and interaction coefficients $\boldsymbol{a}_i^{(\cdot)}$ for orders 3 and 4

respectively. However, when we applied the model to our data, pairwise interactions dominated,

and the indicator variables associated with the higher order interactions had Bayes factors near

zero. From these analyses, we can conclude that the presence of higher order interactions was

not supported by our data. However, it is possible that such interactions actually exist in the

ecosystem but that our data is insufficient for rigorous inference using our statistical method.

Calculation of Population Evenness (Pielou's Evenness)

Species Diversity (Pielou's evenness) was calculated according to the given formula:

$$Pielou's\ Evenness\ Index = \frac{-\sum_{i=1}^{4} p_i * \ln(p_i)}{\ln(S)}$$

Where p_i refers to the population ratio of a given strain in the consortium of four strains. S is the

number of species.

In vivo **Experiments**

Adult (6-8 weeks) male Swiss Webster germ free mice bred in house at the Massachusetts

Host-Microbiome Center were used. Animals were fed either on standard chow in the facility for

the entire experiment, or on low-protein diet (3% custom diet, envigo, doubly irradiated)

beginning 10 days prior to the experiment and continuing for its duration. To prepare bacteria for

gavage, we grew each strain to mid-log phase, plated for counting and snap-froze aliquots. For

gavage, aliquots were thawed, spun down, combined to achieve concentrations of

approximately 10^7 per bacteria per gavage, and re-suspended in 200 µL 1x PBS with 0.05% L-

cysteine for gavage. After gavage, mice were transferred to Optimice cages and maintained

gnotobiotic for 10 days. Fecal samples were collected prior gavage and at 10 days, and snap-

frozen for storage at -80 °C.

Molecular Analysis of *in vivo* **Samples**

DNA was extracted from fecal samples using the Zymobiomic 96 DNA Kit with the following

modification: we omitted the silicon-ATM-HRC wash. Cells were lysed in a bead beater at speed

20 for 10 min, plates were turned and lysed for another 10 min at the same speed. We added

an additional 3 min incubation step for Binding Buffer and additional 5 min incubation steps

when transferring to Zymo-Spin-I-96-Z plates. Elution was done in 50 µL ZymoBIOMIC

DNase/RNase Free Water.

Direct multiplex probe-based qPCR was done on extracted DNA samples as described above.

For standard curves, we used plated overnight cultures spiked into germfree fecal samples and

extracted them as described above.

References

1.   Hays SG, Patrick WG, Ziesack M, Oxman N, Silver PA. 2015. Better together:
Engineering and application of microbial symbioses. Curr Opin Biotechnol 36:40–49.

2.   Little AEF, Robinson CJ, Peterson SB, Raffa KF, Handelsman J. 2008. Rules of
Engagement: Interspecies Interactions that Regulate Microbial Communities. Annu Rev
Microbiol 62:375–401.

3.   Freilich S, Zarecki R, Eilam O, Segal ES, Henry CS, Kupiec M, Gophna U, Sharan R,
Ruppin E. 2011. Competitive and cooperative metabolic interactions in bacterial
communities. Nat Commun 2:587–589.

4.   Foster KR, Bell T. 2012. Competition, not cooperation, dominates interactions among
culturable microbial species. Curr Biol 22:1845–1850.

5.   Hibbing ME, Fuqua C, Parsek MR, Peterson SB. 2010. Bacterial competition: Surviving
and thriving in the microbial jungle. Nat Rev Microbiol 8:15–25.

6.   Coyte KZ, Schluter J, Foster KR. 2015. The ecology of the microbiome: Networks,
competition, and stability. Science 350:663–6.

7.   D'Souza G, Shitut S, Preussger D, Yousif G, Waschina S, Kost C. 2018. Ecology and
evolution of metabolic cross-feeding interactions in bacteria. Nat Prod Rep 35:455–488.

8.   Rakoff-Nahoum S, Foster KR, Comstock LE. 2016. The evolution of cooperation within
the gut microbiota. Nature 533:255–259.

9.   Hammerstein P, Noë R. 2016. Biological trade and markets. Philos Trans R Soc B Biol
Sci 371.

10.   Werner GDA, Strassmann JE, Ivens ABF, Engelmoer DJP, Verbruggen E, Queller DC,

632  Noe R, Johnson NC, Hammerstein P, Kiers ET. 2014. Evolution of microbial markets.

633  Proc Natl Acad Sci 111:1237–1244.

634 11. Morris JJ. 2015. Black Queen evolution: The role of leakiness in structuring microbial

635  communities. Trends Genet 31:475–482.

636 12. Kallus Y, Miller JH, Libby E. 2017. Paradoxes in leaky microbial trade. Nat Commun 8.

637 13. D'Souza G, Waschina S, Pande S, Bohl K, Kaleta C, Kost C. 2014. Less Is More:

638  Selective Advantages Can Explain the Prevalent Loss of Biosynthetic Genes in Bacteria.

639  Evolution (N Y) 68:2559–2570.

640 14. Reese T, Dunn R. 2018. Drivers of Microbiome Biodiversity: A Review of General Rules,

641  Feces, and Ignorance. MBio 1–14.

642 15. Embree M, Liu JK, Al-Bassam MM, Zengler K. 2015. Networks of energetic and

643  metabolic interactions define dynamics in microbial communities. Proc Natl Acad Sci

644  112:15450–15455.

645 16. Stolyar S, Van Dien S, Hillesland KL, Pinel N, Lie TJ, Leigh JA, Stahl DA. 2007.

646  Metabolic modeling of a mutualistic microbial community. Mol Syst Biol 3:1–14.

647 17. Estrela S, Gudelj I. 2010. Evolution of Cooperative Cross-Feeding Could Be Less

648  Challenging Than Originally Thought. PLoS One 5.

649 18. Wintermute EH, Silver PA. 2010. Emergent cooperation in microbial metabolism. Mol

650  Syst Biol 6:1–7.

651 19. Kerner A, Park J, Williams A, Lin XN. 2012. A programmable escherichia coli consortium

652  via tunable symbiosis. PLoS One 7:1–10.

653 20. Pande S, Merker H, Bohl K, Reichelt M, Schuster S, De Figueiredo LF, Kaleta C, Kost C.

2014. Fitness and stability of obligate cross-feeding interactions that emerge upon gene

loss in bacteria. ISME J 8:953–962.

21.   Mee MT, Collins JJ, Church GM, Wang HH. 2014. Syntrophic exchange in synthetic

microbial communities. Proc Natl Acad Sci U S A 111:E2149-56.

22.   Wintermute EH, Silver PA. 2010. Dynamics in the mixed microbial concourse. Genes Dev

24:2603–2614.

23.   Shou W, Ram S, Vilar JMG. 2007. Synthetic cooperation in engineered yeast

populations. Proc Natl Acad Sci U S A 104:1877–82.

24.   Kong W, Meldgin DR, Collins JJ, Lu T. 2018. Designing microbial consortia with defined

social interactions. Nat Chem Biol 14:821–829.

25.   Aidelberg G, Towbin BD, Rothschild D, Dekel E, Bren A, Alon U. 2014. Hierarchy of non-

glucose sugars in Escherichia coli 1–12.

26.   Spiga L, Winter MG, Santos RL, Hooper L V, Winter SE, Spiga L, Winter MG, Carvalho

TF De, Zhu W, Hughes ER, Beiting DP, Santos RL, Hooper L V, Winter SE. 2017. An

Oxidative Central Metabolism Enables Article An Oxidative Central Metabolism Enables

Salmonella to Utilize Microbiota-Derived Succinate. Cell Host Microbe 22:291-301.e6.

27.   Donaldson GP, Lee SM, Mazmanian SK. 2017. Gut biogeography of the bacterial

microbiota 14:20–32.

28.   Kotula JW, Kerns SJ, Shaket L a, Siraj L, Collins JJ, Way JC, Silver P a. 2014.

Programmable bacteria detect and record an environmental signal in the mammalian gut.

Proc Natl Acad Sci U S A 111:4838–43.

29.   Riglar DT, Giessen TW, Baym M, Kerns SJ, Niederhuber MJ, Bronson RT, Kotula JW,

Gerber GK, Way JC, Silver PA. 2017. Engineered bacteria can function in the

mammalian gut long-term as live diagnostics of inflammation. Nat Biotechnol 35:653–

658.

30. Ziesack M, Karrenbelt MAP, Bues J, Schaefer E, Silver P. 2018. Escherichia coli NGF-1,

a Genetically Tractable, Efficiently Colonizing Murine Gut Isolate. Microbiol Resour

Announc 1–2.

31. Kim S, Kerns SJ, Ziesack M, Bry L, Gerber GK, Way JC, Silver PA. 2018. Quorum

Sensing Can Be Repurposed to Promote Information Transfer between Bacteria in the

Mammalian Gut. ACS Synth Biol 7:2270–2281.

32. Bogatyreva NS, Finkelstein A V, Galzitskaya O V. 2006. Trend of Amino Acid

Composition of Proteins of Different Taxa. J Bioinform Comput Biol 4:597–608.

33. Bucci V, Tzen B, Li N, Simmons M, Tanoue T, Bogart E, Deng L, Yeliseyev V, Delaney

ML, Liu Q, Olle B, Stein RR, Honda K, Bry L, Gerber GK. 2016. MDSINE: Microbial

Dynamical Systems INference Engine for microbiome time-series analyses. Genome Biol

17:121.

34. Gibson TE, Gerber GK. 2018. Robust and Scalable Models of Microbiome Dynamics.

Proc 35th Int Conf Mach Learn 80:1763–1772.

35. Stein RR, Bucci V, Toussaint NC, Buffie CG, Raetsch G, Pamer EG, Sander C, Xavier

JB. 2013. Ecological Modeling from Time-Series Inference□: Insight into Dynamics and

Stability of Intestinal Microbiota. PLoS Comput Biol 9:31–36.

36. Stein RR, Tanoue T, Szabady RL, Bhattarai SK, Olle B, Norman JM, Suda W, Oshima K,

Hattori M, Gerber GK, Sander C. 2018. Computer-guided design of optimal microbial

consortia for immune system modulation 1–17.

37. Ravindran R, Loebbermann J, Nakaya HI, Khan N, Ma H, Gama L, Machiah DK, Lawson

B, Hakimpour P, Wang YC, Li S, Sharma P, Kaufman RJ, Martinez J, Pulendran B. 2016.

The amino acid sensor GCN2 controls gut inflammation by inhibiting inflammasome

activation. Nature 531:523–527.

38.   Luczynski P, Neufeld KAMV, Oriach CS, Clarke G, Dinan TG, Cryan JF. 2016. Growing

up in a bubble: Using germ-free animals to assess the influence of the gut microbiota on

brain and behavior. Int J Neuropsychopharmacol 19:1–17.

39.   Thomason LC, Costantino N, Court DL. 2007. E. coli genome manipulation by P1

transduction. Curr Protoc Mol Biol Chapter 1:Unit 1.17.

40.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko K a, Tomita M,

Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12 in-frame, single-gene

knockout mutants: the Keio collection. Mol Syst Biol 2:2006.0008.

41.   Cherepanov PP, Wackernagel W. 1995. Gene disruption in Escherichia coli: TcR and

KmR cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance

determinant. Gene 158:9–14.

42.   Mimee M, Tucker AC, Voigt CA, Lu TK. 2015. Programming a Human Commensal

Bacterium, Bacteroides thetaiotaomicron, to Sense and Respond to Stimuli in the Murine

Gut Microbiota. Cell Syst 18:386–392.

43.   Ferrières L, Hémery G, Nham T, Guérout AM, Mazel D, Beloin C, Ghigo JM. 2010. Silent

mischief: Bacteriophage Mu insertions contaminate products of Escherichia coli random

mutagenesis performed using suicidal transposon delivery plasmids mobilized by broad-

host-range RP4 conjugative machinery. J Bacteriol 192:6418–6427.

44.   Darling ACE, Mau B, Blattner FR, Perna NT. 2004. Mauve□: Multiple Alignment of

Conserved Genomic Sequence With Rearrangements 1394–1403.

45.   Veeravalli K, Laird MW, Fedesco M, Zhang Y, Yu XC. 2014. Strain Engineering to

724   Prevent Norleucine Incorporation During Recombinant Protein Production in Escherichia

725   coli.

726   46.   Malykh EA, Butov IA, Ravcheeva AB, Krylov AA, Mashko S V. 2018. Specific features of l

727   ⬚ histidine production by Escherichia coli concerned with feedback control of AICAR

728   formation and inorganic phosphate / metal transport. Microb Cell Fact 1–15.

729   47.   Fang M, Zhang C, Yang S, Cui J, Jiang P, Lou K, Wachi M, Xing X. 2015. High crude

730   violacein production from glucose by Escherichia coli engineered with interactive control

731   of tryptophan pathway and violacein biosynthetic pathway 1–13.

732   48.   Ginesy M, Belotserkovsky J, Enman J, Isaksson L, Rova U. 2015. Metabolic engineering

733   of Escherichia coli for enhanced arginine biosynthesis. Microb Cell Fact 14:1–11.

Table 1 Engineered strains, engineered genotypes and a subset of identified SNPs

previously implicated as causal mutations. [a]prevents feedback inhibition (45), [b]decouples

from histidine feedback inhibition (46); [c]trpE, removes feedback inhibition (47); [d]arginine

repressor, nonfunctional (48)

Figure 1 Engineered Metabolite Overproduction and Growth Requirements. **a.** Consortia

design. Each strain is auxotrophic for three amino acids and overproduces one. **b.**

Quantification of overproduction and growth requirement (EC50). Overproduction in each

engineered strain was measured via LC-MS with appropriate amino acid standard. Metabolite

growth requirement EC50 was measured in media with full supplementation of two amino acids

and varying amounts of one other. **c.** Cross-feeding capabilities of each strain were assessed

by testing for rescue of auxotroph growth in supernatants obtained from overproducers. **d.**

Growth assay of auxotrophic strains. Auxotroph strains were grown in media without and with

addition of all cross-fed amino acids (left side of graphs, -/+) which served as a reference point.

Each strain was also grown in fresh media without supplementation and supernatant of

engineered overproducers or WT counterpart at a 1:1 ratio. Shown are three biological

replicates with median indicated as horizontal lines.

Figure 2 Engineered consortium exhibits added beneficial interactions and reduced

antagonistic interactions. a./b. Growth trajectories in anaerobic batch co-cultures for WT (**a.**)

and engineered (**b.**) consortia. All strains were inoculated at equal ratios and grown

anaerobically at 37 °C for 27 hours in M9 minimal media with specific modifications as

described in Methods, without supplementation of any of the cross-fed amino acids, and with

0.5% starch and 0.5% glucose as carbon sources. The engineered consortium grows to an

order of magnitude lower density than the WT counterpart. **c./d.** Growth trajectories of consortia

with each strain inoculum reduced by 10-fold of WT (**c.**) and engineered (**d.**) consortia. These

data combined were used to infer a network interaction using the MDSINE algorithm. **e./f.**

Inferred interaction networks for WT (**e.**) and engineered (**f.**) consortia. WT consortia show

mainly negative interactions with large Bayes factors. Engineered consortia's negative

interactions have reduced Bayes factors and positive interactions with strong Bayes factors

occurred between *B. fragilis* and *E. coli* and *S.* Typhimurium.

Figure 3 Engineered Consortia exhibit increased population evenness *in vitro.* WT and

engineered consortia were grown anaerobically in batch co-culture for 24 hr anaerobically in

modified M9 media as described in Methods with and without supplementation of 1 mM of the

cross-fed amino acids and 0.2% starch and 0.5% glucose as carbon sources. Upper panel:

average population composition after 24 hr. Lower panel: Quantified Pielou Evenness. The

engineered consortium has higher population evenness with or without amino acid

supplementation compared to WT consortium.

Figure 4. Consortia engineering increases population evenness in the mammalian gut in

a diet dependent manner. Four groups of germfree mice (n=5, except second group from the

left which is n=4) were either fed low protein diet or standard diet, and inoculated with either the

engineered or WT bacterial consortia. Fecal samples 10 days post-inoculation were analyzed

via strain-specific qPCR to assess concentrations of each consortia species. Population

evenness of consortia was calculated. Bar indicates Median. Mann-Whitney test showed

significantly increased population of the engineered consortia in mice that were fed low protein

diet compared to the consortia in the three other groups (*p*-values: 0.024, 0.032, 0.015). yellow:

E. coli; purple: *S.* Typhimurium; green: *B. thetaiotaomicron*; blue: *B. fragilis*.

 Supplemental Material

Figure S1 Growth response. Each auxotroph was grown in media supplemented with varying

concentrations of one amino acid and saturating concentration of the two others. Depicted is the

average of three biological replicates; error bars indicate standard deviation. A sigmoidal curve

was fit using GraphPad Prism 8.

Figure S2 Log fold ratios (LogR) trajectories in WT consortium in all four conditions compared

to condition one in Figure 2. Legend: Ec = *E. coli*, ST = *S.* Typhimurium, BT = *B. theta*, BF = *B.*

fragilis.

Figure S3 Log fold ratios (logR) trajectories in engineered consortium in all four conditions

compared to condition one in Figure 2. Legend: Ec = *E. coli*, ST = *S.* Typhimurium, BT = *B.*

theta, BF = *B. fragilis*.

Figure S4 Growth rates, Bayes Factors (for microbial interactions), and Microbial Interaction

Strengths for unengineered (A) and engineered (B) consortia learned from *in vitro* growth data.

The inferred growth rates for the engineered consortia are smaller by about an average factor of

3 for any species. The Bayes Factors quantify the evidence for an interaction being present

verses being absent (Bayes Factor between 3 and 10 = substantial evidence, Bayes Factor >

10 = strong evidence). The inference is highly confident in 7 of the 12 unenginereerd consortia

interactions. Note that an infinite Bayes Factor means that the edge was included in every

single step when learning the posterior distribution. For the engineered consortia two of the 12

interactions exist with at least substantial evidence with 5 of the interactions having very low

evidence for the edge being present in the model but still greater than evidence for no

interaction (Bayes Factor of 1 is the scenario where evidence for and against an interaction

being present are equal). The interaction strengths for the engineered consortia are of higher

magnitude when compared to the unengineered consortia. This occurs because the carrying

capacity of the unengineeered consortia is about 1.5 orders of magnitude larger than the

engineered consortia and microbe-microbe interactions are inversely proportional to the carrying

capacity in a gLV model.

Figure S5 Growth of WT and engineered strains after 24 hr, at which point supernatant was

collected for cross-feeding experiment (see Figure 1). Overproduction does not affect growth

with the exception of *B. theta* (3-fold reduction). Shown are three biological replicates with

median indicated as horizontal line.

Figure S6 Metabolite overproduction and WT and engineered strains. Amino acid

production was measured using LC-MS. The average of three biological replicates is shown.

Figure S7 Growth after 24 hr in media supplemented with 0.5% or 0.2% starch. Growth did

not differ significantly for any of the strains (Mann-Whitney test, p-values > 0.1).

Table S1 qPCR probes and primers.

a.

equal inocula

E. coli   S. Typhimurium   B. fragilis   B. theta

b.

equal inocula

E. coli   S. Typhimurium   B. fragilis   B. theta

c.

Condition 2: *E. coli* knockdown

Condition 3: *S.* Typhim. knockdown

Condition 4: *B. theta* knockdown

Condition 4: *B. fragilis* knockdown

d.

Condition 2: *E. coli* knockdown

Condition 3: *S.* Typhim. knockdown

Condition 4: *B. theta* knockdown

Condition 5: *B. fragilis* knockdown

e.

Wild Type Inferred Interaction Network

E coli   S typhimurium

B fragilis   B theta

Bayes Factor
15 or greater

f.

Engineered Inferred Interaction Network

E. coli   S. Typhimurium

B. fragilis   B. theta

Species	Strain	Auxotroph Genotype	Other Genotype	Overproduction Mutations
E. coli	NGF	ΔargA, ΔtrpC, ΔhisA	ΔthiE	metA(I296S)[a]
S. Tyhpimurium	LT2	ΔargA, ΔtrpC, ΔmetA	ΔthiE, ΔSPI1, ΔSPI2	hisG(E271K)[b]
B theta	VPI5482	ΔmetA, ΔhisG, ΔargF,	ΔthiSEG, Δtdk	BT_0532 (A306V; N63D)[c]
B. fragilis	368R	ΔmetA, ΔhisG, ΔtrpC	ΔthiSEG, Δtdk	BF638R_0532 (L26R)[d]

Species	Strain	Auxotroph Genotype	Other Genotype	Overproduction Mutations
E. coli	NGF	ΔargA, ΔtrpC, ΔhisA	ΔthiE	metA(I296S)[a]
S. Tyhpimurium	LT2	ΔargA, ΔtrpC, ΔmetA	ΔthiE, ΔSPI1, ΔSPI2	hisG(E271K)[b]
B theta	VPI5482	ΔmetA, ΔhisG, ΔargF,	ΔthiSEG, Δtdk	BT_0532 (A306V; N63D)[c]
B. fragilis	368R	ΔmetA, ΔhisG, ΔtrpC	ΔthiSEG, Δtdk	BF638R_0532 (L26R)[d]