High-speed atomic force microscopy visualizes mobility of photosynthetic proteins in grana thylakoid membranes

Bibiana Onoa[1,†,*], Shingo Fukuda[1,†,‡], Masakazu Iwai[2,3], Carlos Bustamante[1,2,4,5,6,7], Krishna K. Niyogi[1,2,3,*]

[1]Howard Hughes Medical Institute, University of California, Berkeley, CA 94720, USA.

[2]Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA.

[3]Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA.

[4]Jason L. Choy Laboratory of Single Molecule Biophysics, University of California, Berkeley, CA 94720, USA.

[5]Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA.

[6]Department of Physics, University of California, Berkeley, CA 94720 USA.

[7]Kavli Energy NanoSciences Institute at the University of California, Berkeley and the Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA.

* For correspondence: bibianaonoa@berkeley.edu (BO) and niyogi@berkeley.edu (KKN)

† These authors contributed equally to this work.

Present address: ‡ Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kanazawa 920-1192, Japan.

SUMMARY

Thylakoid membranes in chloroplasts contain photosynthetic protein complexes that convert light energy into chemical energy. Under fluctuating light conditions, photosynthetic protein complexes are considered to undergo structural reorganization to maintain the efficiency of photochemistry. Electron and atomic force microscopes have been broadly used to visualize the macromolecular architecture of thylakoid membranes as well as individual photosynthetic protein complexes at high spatial resolution. However, real-time observation of the dynamics driving structural reorganization in the membrane is challenging with these techniques. Here, we used high-speed atomic force microscopy (HS-AFM) to visualize and characterize the mobility of individual protein complexes in grana thylakoid membranes isolated from *Spinacia oleracea*. Our results indicate that grana thylakoid membranes can be grouped into two classes according to protein mobility. A large fraction of grana membranes contained proteins with quasi-static mobility, exhibiting molecular displacements smaller than 10 nm^2. In the remaining fraction, the protein mobility appeared to be variable with molecular displacements of up to 100 nm^2. The calculated average diffusion coefficient was ~1 nm^2 s^{-1}. Both confined and Brownian diffusion models could describe the protein mobility of the second group of membranes. The HS-AFM temporal resolution also allows

detecting rotational protein diffusion in grana membranes. Thus, HS-AFM provides both visualization of individual photosynthetic protein complexes in grana membranes and direct measurement of their mobility and dynamics *in situ*. This dynamic characterization of the diffusion of individual photosynthetic complexes opens the door to in-depth study of the structural responses of thylakoid membranes to environmental fluctuations.

Keywords: high-speed atomic force microscopy, thylakoid membrane, high-spatiotemporal resolution imaging, PSII dynamics, *Spinacia oleracea*

SIGNIFICANCE STATEMENT

High-speed atomic force microscopy enables the characterization of the dynamics of individual photosynthetic proteins in grana thylakoid membranes from *Spinacia oleracea*. The high-spatiotemporal resolution imaging unveils that the mobility of photosynthetic proteins is heterogeneous but governed by the confinement effect imposed by the high protein density in the thylakoid membrane.

INTRODUCTION

Photosynthesis is a fundamental process that sustains virtually all life on earth. The two photosystems (PSI and PSII), cytochrome b_6f complex (Cyt b_6f), and ATP synthase are the major multisubunit membrane protein complexes that catalyze light-driven chemical reactions to generate ATP and NADPH in chloroplasts (Nelson and Yocum, 2006). Light energy is funneled into the reaction centers of each photosystem through light-harvesting complex (LHC) proteins. LHC proteins of PSII (LHCII) are the most abundant membrane proteins in chloroplast thylakoid membranes, and they are also known to play an essential role in photoprotection (Muller, 2001; Wobbe *et al.*, 2016; Ruban, 2016). Given the complexity of the light reactions of photosynthesis and their regulation, investigation of thylakoid membrane structure and function has long been a central topic in the field of photosynthesis research.

Thylakoid membranes in plant chloroplasts are organized into intricate structures comprised of highly stacked and non-stacked membrane regions called grana and stroma lamellae, respectively (Anderson, 1981; Austin and Staehelin, 2011). It is well established that PSII and LHCII proteins are predominantly localized in grana, whereas PSI and ATP synthase are exclusively located in stroma lamellae (Andersson and Anderson, 1980). Photo-regulatory processes such as photosynthesis, photoprotection, protein repair, etc. are thought to trigger architectural reorganization in the thylakoids (Ruban and Johnson, 2015). However, the molecular displacement of the photosynthetic complexes responsible for those structural rearrangements is poorly characterized. To investigate the dynamics of photosynthetic proteins in the membrane at high spatial-temporal resolution, a microscopy technique has to meet at least four conditions: (1) The spatial resolution is such that individual complexes can be resolved. (2) The observation is performed in physiological conditions. (3) The temporal

resolution is high enough to track molecular displacements. (4) The detection method is not invasive to preserve native-like organizations.

Electron microscopy (EM) meets the first and fourth conditions, indeed, EM studies have shown that grana are highly packed with membrane proteins, where PSII and LHCII form a protein supercomplex (Boekema *et al.*, 2000; Yakushevska *et al.*, 2001; Dekker *et al.*, 2002). It has also been suggested that the macroorganization of PSII-LHCII supercomplexes plays a critical role during photoprotection, which implies a high significance of their diffusion rates within the thylakoid membrane (Goral *et al.*, 2010; Johnson *et al.*, 2011; Goral *et al.*, 2012). Optical microscopy based on fluorescence detection, on the other hand, fulfills conditions (2) and (3). By following chlorophyll (Chl) fluorescence emitted from photosynthetic membrane proteins, it is possible to infer dynamics of thylakoid membranes at high temporal resolution. However, this strategy remains blind to photosynthetic proteins themselves, which provide the binding sites for Chl. In spite of the efforts to improve the spatial resolution using super-resolution optical microscopy (Willig *et al.*, 2006; Betzig *et al.*, 2006; Sigal *et al.*, 2018; Iwai *et al.*, 2018), the information obtained is still somewhat indirect due to the high abundance of Chl pigments in the membrane which limits the specific assignment of the actual photosynthetic complex.

Atomic force microscopy (AFM) is a unique technique that enables direct observation of macromolecules at high spatial resolution (XY < 1 nm and Z < 0.1 nm) in aqueous conditions. AFM has been used to characterize the structure and organization of thylakoid membranes (Kirchhoff *et al.*, 2008; Sznee *et al.*, 2011; Johnson *et al.*, 2014; Onoa *et al.*, 2014; Stoichev *et al.*, 2015; Phuthong *et al.*, 2015). These studies support and complement EM reports about how PSII and LHCII organization is affected by illumination (Sznee *et al.*, 2011; Onoa *et al.*, 2014; Stoichev *et al.*, 2015). Despite the scope of AFM in the characterization of the thylakoid membrane, studying its dynamic behavior was elusive because of the AFM's intrinsic low imaging speed. Thanks to the development of high-speed AFM (HS-AFM) (Kodera *et al.*, 2005; Kodera *et al.*, 2006; Fantner *et al.*, 2006; Ando *et al.*, 2008; Fukuda *et al.*, 2013), it is now possible to visualize real-time dynamics of biological macromolecules. Thus, HS-AFM has the potential to meet all four conditions stated above. Indeed, HS-AFM has been applied to study dynamic events of purified molecules (Kodera and Ando, 2018; Uchihashi *et al.*, 2011; Katan *et al.*, 2015; Milhiet *et al.*, 2010; Banerjee *et al.*, 2017), membrane proteins embedded in lipid bilayers (Casuso *et al.*, 2012; Ruan *et al.*, 2017), and even biological tissues (Colom *et al.*, 2013; Yamashita *et al.*, 2012; Sakiyama *et al.*, 2016). Here, we employed HS-AFM to characterize the *in situ* dynamic behavior of the photosynthetic protein complexes in grana thylakoid membranes isolated from *Spinacia oleracea*. Our HS-AFM observations indicated that the fraction of mobile membrane proteins is less than 10% of the total population. The high temporal resolution permitted herein showed that the diffusion of the protein complexes is mainly rotational. Our results also indicated that the diffusion of the photosynthetic membrane proteins is heterogeneous not only between different grana membranes, but also within a single granum. This characterization provides a foundation for studies of thylakoid structural response to different environments.

RESULTS

HS-AFM visualizes dimeric photosynthetic complexes without altering macromolecular organization in grana membranes.

We isolated grana membranes from spinach using digitonin as described previously (Anderson and Boardman, 1966) (Figure S1). This preparation yielded small stacks of grana discs with high content of PSII, LHCII and Cyt b$_6$f complexes as previously shown (Johnson *et al.*, 2014). We optimized the HS-AFM setup for imaging photosynthetic proteins in grana membranes, such that deflection of the AFM cantilever was detected using a near-infrared laser (830 nm wavelength) to minimize Chl excitation. The membranes were scanned at one frame per second in buffer solution (see Methods for details).

HS-AFM observations indicated that macroorganization of grana membranes and associated protein structures were well preserved (Figures 1 and S3). As shown previously (Kirchhoff *et al.*, 2008; Phuthong *et al.*, 2015, Onoa *et al.*, 2014; Kouřil *et al.*, 2013), grana membranes were packed with dimeric complexes with an overall density of ~1456 ± 230 particles/μm^2. Dimeric structures were distributed throughout the membranes, but their structural arrangement appeared to be disordered. We observed a bimodal distribution of height of the dimeric structures, which is consistent with previous results (Figure S2a, b) (Johnson *et al.*, 2014). Immunoblot analysis detected the existence of PSII and Cyt b$_6$f in our grana membranes prepared using digitonin (Johnson *et al.*, 2014) (Figure S1a). Certainly, the unambiguous distinction of PSII and Cyt b$_6$f is highly desirable whenever both of them are present in the membranes. However, imaging by itself is insufficient to discriminate these two dimers. Comparison of available structures of Cyt b$_6$f and PSII indicates that their dimensions are very similar (Kurisu *et al.*, 2003; Umena *et al.*, 2011). Both Cyt b$_6$f and PSII have similar shapes and lateral sizes of ~16 and ~20 nm, respectively. Moreover, in both cases, their membrane lumenal protrusions are separated by ~8 nm and their expected heights differ by only ~1 nm (Johnson *et al.*, 2014). These characteristics challenge an unambiguous discrimination of each individual complex but are sufficient to estimate the presence of distinct populations of complexes. There is a structural report that succeeded in distinguishing both dimers by affinity-mapping AFM (Johnson *et al.*, 2014), which specifically recognizes plastocyanin-Cyt b$_6$f interactions. It is worth noting that we did not observe ATP synthase by HS-AFM in our grana membranes. It is known that solubilization by digitonin cannot remove ATP synthase completely from the grana preparation, as observed in our immunoblot analysis (Figure S1a) and a previous study (Johnson *et al.*, 2014). For the purpose of this study, it was more important to prepare the membrane samples closer to native conditions than to remove ATP synthase completely by using stronger detergents. In fact, the ATP synthase content in our grana membranes was reduced 2-fold from leaves, which was sufficient for our observation. For HS-AFM observation, the samples were 5-to-10-fold further diluted to adjust their density on the surface, such that individual grana discs could be isolated.

Furthermore, we avoided scanning membranes with large protruding structures, and thus we minimized the probability to study membranes containing ATP synthase complexes.

We also calculated the nearest neighbor distance (NND) distribution function (Figure S2c). A main peak centered at ~20 nm flanked by shorter (~16 nm) and longer (~25 nm) distances was well fitted, which is also consistent with previous results (Onoa *et al.*, 2014; Kirchhoff *et al.*, 2004). These results observed by HS-AFM are qualitatively comparable with those observed by conventional AFM performed in air (Figure S1b-d). The protein density and NND distribution observed by HS-AFM are also consistent with those observed by EM (e.g. the samples with no light treatment (Kouřil *et al.*, 2013; Betterle *et al.*, 2009). Taken together, these results demonstrated that HS-AFM is a suitable technique to image the thylakoid protein complexes at high spatial resolution *in situ* without altering their intrinsic macroorganization.

HS-AFM revealed heterogeneous protein diffusion in individual grana membranes.

To analyze the dynamics of photosynthetic complexes in grana membranes, we performed HS-AFM observation for 60 s or more per sample. HS-AFM images of representative grana discs are shown in Figure S3e. HS-AFM was performed on a total of 19 grana discs from two biological replicate preparations. We tracked individual protruding dimeric structures to calculate the mean square displacement (MSD, Eq. 1). Based on the level of lateral displacement, grana membranes that we observed here could be divided into two groups. The first group, which comprises approximately 90% of the total grana membranes observed in this study, was termed QSM membranes because they contained dimeric structures with quasi-static mobility (Figure 2, Movie S1). The distinct dimeric structure of each particle in a QSM membrane was still apparent after averaging 50 frames of the HS-AFM images (Figure 2a), which indicates that the lateral displacement was confined to a few nanometers or within the mechanical drift in HS-AFM apparatus. The MSD values of 53 dimeric structures in this representative QSM membrane show that the molecular displacement was less than 10 nm^2 (Figure 2b). The average MSD trace of all structures (thick line in Figure 2b) was well fitted to a confined diffusion model (Eq. 2). Because the grana membranes used in this study showed preserved macroorganization (Figures 1, S1 and S2), we consider that the quasi-static mobility observed in the QSM group does not indicate an aberrant state of the membranes. In the second group, which comprises about 10% of the total grana membranes observed in this study, most dimeric structures showed quasi-static mobility and sometimes appeared to be clustered (arrowheads in Figure S4). However, there was a subpopulation of dimeric structures that displayed larger displacements. Figure 3a shows representative time-lapse HS-AFM images revealing such dimeric structures (see also Movie S2). The individual MSD traces were variable among the selected dimeric structures (thin gray lines), and some of them showed larger displacements. (Figure 3b). Therefore, the second group of grana membranes was termed VPM for variable protein mobility. The average MSD trace of all observed dimeric structures in the VPM membranes (yellow line in Figure 3b) fitted well to a confined diffusion model (Eq. 2, black dashed line, R^2 = 0.996) as compared to a

Brownian diffusion model (Eq. 3, orange dotted line, $R^2 = 0.981$). Using Equations 2 and 4, we obtained the average diffusion coefficient of ~1 nm^2 s^{-1} that was consistent with previous Monte Carlo simulation reports (see Discussion). This result indicates that, overall, the dimeric structures in the VPM grana membranes still exhibited dynamics that can be explained by a confined diffusion model, similar to those observed in the QSM group.

The average MSD trace of the entire population of particles in the VPM membranes may underestimate a certain subpopulation, which apparently showed an unconfined diffusion. To carefully determine the diffusion of each individual dimeric structure, we first fitted each MSD trace to the confined diffusion model. Next, we extracted each confined domain (L) and correlated it with the diffusion coefficient during the first 4 s ($D_{initial}$). We then applied k-means clustering criteria (Hartigan and Wong, 1979). The results showed that most of the dimeric structures (84%) (Figure 3c, blue circles) were each confined to a small region (average L = 5.8 ± 0.04 nm). In contrast, the remaining population (Figure 3d, magenta squares) displayed larger displacements (average L = 44.3 ± 17.6 nm). We re-calculated the average MSD trace separately for these two populations. The average MSD trace (Figure 3d, blue trace) from the larger population was better fitted to a confined diffusion model (light blue dashed line; Eq. 2, $R^2 = 0.997$) than to a Brownian diffusion model (blue dotted line; Eq. 3 $R^2 = 0.939$). On the other hand, the average MSD trace from the smaller population (Figure 3d, magenta trace) was well fitted to either a confined (light pink dashed line; Eq. 2, $R^2 = 0.983$) or Brownian diffusion model, (magenta dotted line; Eq. 3, $R^2 = 0.983$). This result illustrates that the diffusion characteristics of this particular population are heterogeneous even within the same VPM grana membrane.

In an attempt to explain the nature of this heterogeneous mobility and/or to distinguish PSII from Cyt b_6f by their mobility, we interrogated if there was any correlation of the complexes' mobility and their height (i.e. smaller complexes might diffuse faster than taller/larger ones). We measured the height of dimeric proteins in the VPM membranes and correlated them to their MSD. There was no correlation between complexes' heights and MSD (Figure S5). From this result, we consider that the complexes' mobility was inconclusive to distinguish whether the mobile fraction corresponded specifically to PSII or Cyt b_6f. We also noticed that the protein density (1594 ± 364 particles/μm^2) in the VPM membranes was not statistically different from that in QSMs (P = 0.7, Student's t-test). Interestingly, the median height distribution of the particles in VPM membranes was broader compared to that of QSMs, and it was best fitted with a single Gaussian based on Akaike information criterion (Figure S6a). The NNDs were also modestly shifted towards shorter distances compared to those of QSMs (Figure S6b), which is consistent with the presence of protein clusters. Protein clustering might favor molecular unconfinement by creating unpopulated domains within the membrane. Furthermore, we detected modest fluctuations of the complexes' height throughout the observation. We compared the heights and NNDs of the particles (first and last frame) in the same VPM membrane. There are visible differences between both height and NND distributions (Figures S6c and S6d), suggesting that the global organization of the membrane, as well as the level of protrusion of the complexes, were slightly changed during our observation. It is worth mentioning that we

occasionally observed grana membranes that contained larger populations of mobile dimeric structures (>80%) than immobile ones (Movie S3). In such cases, dimeric structures tended to collide with each other more frequently, which made it difficult to properly identify individual structures and to track their lateral diffusion.

HS-AFM visualized rotational displacement of dimeric structures in grana

High temporal resolution of HS-AFM in this study also enabled us to analyze rotational displacement of the dimeric structures. In the VPM grana membranes, which contained the dimeric structures with heterogeneous diffusion, we measured changes in the angle (θ) of two adjacent structures during a 1-min observation period (magenta and blue arrows in Figure 4a; Movie S4). The results indicated that the angle changed from 20 to 80 degrees within a few seconds (Figure 4b). To further illustrate the rotational diffusion observed here, we calculated a 2D-correlation coefficient of variation (CCV; Eq. 5) of the dimeric structures as shown in Figure 4c. The dimeric structure showing rotational diffusion exhibited constant fluctuations with an average CCV value of 0.65 throughout the observation (Figure 4c, red trace). By averaging the total time-lapse images, the dimeric morphology disappeared because of the rotational displacement (red profile, Figure 4d). In contrast, the other dimeric structure showed an average CCV value of 0.83 (Figure 4c, blue trace), and the dimeric morphology of this structure stayed the same after averaging the time-lapse images (blue profile in Figure 4d). It is known that PSII-LHCII supercomplexes in grana can be organized into a higher order of associations (e.g. megacomplexes or two-dimensional crystalline arrays (Dekker and Boekema, 2005). Intriguingly, it has been suggested that there are two types of PSII megacomplexes, in which 80% of the supercomplexes show parallel associations, whereas the other 20% interact in a non-parallel manner with variable associations between the two supercomplexes (Nosek *et al.*, 2017). The rotational displacement of the dimeric structures detected here is in line with the idea that PSII-LHCII supercomplexes might associate with their neighbors at slightly different orientations.

DISCUSSION

This contribution aims to establish that HS-AFM is a powerful technique to study the dynamic behavior of photosynthetic protein complexes in plants under biologically relevant conditions. Indeed, we were able to characterize the mobility of individual complexes in highly crowded grana membranes. Despite extensive previous investigation, there is currently a lack of consensus about the actual diffusion properties of photosynthetic complexes in thylakoids. This is, in part, because the experimental results are indirectly obtained from the total pool of Chl-binding proteins in the membranes, and because the large size of the supercomplexes limits the scope of molecular simulations (Goral *et al.*, 2010; Ruban and Johnson, 2015; Consoli *et al.*, 2005; Drepper *et al.*, 1993; Kirchhoff, 2014; Kirchhoff *et al.*, 2008; Van Eerden *et al.*, 2017). Our HS-AFM observations provide insight into diffusion of photosynthetic proteins in grana thylakoid membranes, which is

essential information for understanding relevant biological processes such as photosynthesis, photoprotection, photodamage, and repair.

The HS-AFM scanning of grana membranes in solution at a rate of one frame per second was sufficient to resolve the dimeric structure of the photosynthetic protein complexes (PSII or Cyt b_6f). In our preparation, the diameter of grana discs varied from 250 to 600 nm, so in order to observe the entire disc, the scanning area was set at 300-500 nm^2 (~ 150×150 pixels). However, higher temporal resolution can be achieved by scanning smaller areas (Ando *et al.*, 2008).

Our HS-AFM observation revealed the presence of at least two groups of isolated grana membranes (QSM and VPM) according to the diffusion behavior of observed dimeric protein structures. QSM represents the majority of grana membranes observed (~90% of ~19 grana discs) and contains dimeric structures with quasi-static mobility that fits a confined diffusion model (Figure 2). As the remaining 10%, VPM contains dimeric structures showing larger displacements and a higher diffusion rate than the first group. The diffusion model for VPM appears to be fitted with both confined and Brownian models, but there are individual trajectories that fit only a Brownian model because of their high diffusion rates (Figure 3). The architecture of both types of membranes are qualitatively similar (lack of crystalline arrays and similar complex densities), nevertheless it appears that the global organization is slightly different (i.e., more protein clustering). We do not have evidence to suggest the exact reason for this heterogeneity. As we observed a similar density of dimeric structures in both QSM and VPM grana membranes but shorter NNDs, a possible reason for this heterogeneity could be different lipid compositions in the membranes, which might originate from different parts of thylakoid membranes. Also, it has been previously shown that, under fluctuating light, the organization of PSII-LHCII supercomplexes could undergo reversible transitions from crystalline to fluid phases (Onoa *et al.*, 2014; Kouřil *et al.*, 2013). Therefore, we speculate that the heterogeneous protein mobility observed in this study might partially reflect different physiological conditions or transient events related to photoacclimation mechanisms in the leaves from which the grana were isolated. Future experiments using plants acclimated to different light environments will provide a detailed connection between our HS-AFM observation and physiological mechanisms. In this study, we provide a proof of concept that HS-AFM is a suitable technique to study the protein dynamics in thylakoid membranes.

Molecular confinement is well-described in highly crowded membranes such as thylakoid membranes (Kirchhoff, 2014), which would have a significant impact on protein mobility (e.g. diffusion paths and velocities). Fluorescence recovery after photobleaching (FRAP) experiments both in intact or broken chloroplasts and isolated grana membranes have shown that about 80% of chlorophyll-binding proteins are immobile (Goral *et al.*, 2010; Kirchhoff *et al.*, 2008). HS-AFM allows us to visualize the displacement of individual dimeric photosynthetic complexes in thylakoid membranes and to characterize their diffusion. Our HS-AFM observation indicated that 90% of observed grana contain immobile proteins, similar to the results reported by FRAP

experiments. This suggests that the overall protein immobility in chloroplasts observed by FRAP might reflect the confined protein mobility occurring in grana.

Additionally, our HS-AFM observation revealed that protein diffusion can be segregated even within the same grana disc (Figures 3 and 4). It has been shown by using *in vitro* reconstituted lipid membranes with a protein surface density of ~50% that local protein density is correlated with protein mobility, demonstrating that molecular confinement has a significant effect on protein diffusion in membranes (Casuso *et al.*, 2012). The grana thylakoid is however a special type of membrane expected to restrict diffusion of its proteins more than plasma membranes as it is highly packed with a ratio of protein:lipid of 86:14 (w/w) (Kaňa, 2013). AFM scanning can only visualize those domains protruding from the lipid bilayer (e.g. the oxygen evolving complex in PSII). In the thylakoid there are many more proteins embedded in the lipid bilayer that are invisible to AFM (darker areas in the micrographs) that could also affect the mobility of the visible ones. LHCII proteins are most abundant in grana and are suggested to be both immobile and mobile as they can associate with PSII and interact with other LHCII proteins, reorganizing different protein complexes in response to light fluctuations (Ruban and Johnson, 2015). The heterogeneity of protein mobility in a single granum observed here might indicate such different situations of LHCII, some of which are strongly associated with PSII, whereas others diffuse freely between PSII supercomplexes and thereby affect the apparent mobility of dimeric structures. To fully understand diffusion of dimeric proteins (PSII and Cyt b_6f) in thylakoid membranes, it will be necessary to consider the effect of these embedded membrane proteins.

Our HS-AFM observation indicated that the average diffusion coefficient of dimeric structures in grana is approximately 1 $nm^2 s^{-1}$. The results from FRAP measurements estimated a diffusion coefficient of ~100 $nm^2 s^{-1}$ (Kirchhoff *et al.*, 2008), 100-fold higher than our observation, which is most likely due to the fact that our HS-AFM tracks individual dimeric structures in grana, whereas FRAP measures the ensemble of Chl-binding proteins. Coarse-grained simulations of individual PSII complexes calculated a diffusion coefficient of 100,000 $nm^2 s^{-1}$ (Van Eerden *et al.*, 2017). However, this simulation did not account for the molecular crowding effect. Monte Carlo simulations based on FRAP experimental data that include the effect of molecular crowding calculated a diffusion coefficient of 1 $nm^2 s^{-1}$ (Kirchhoff *et al.*, 2004), which agrees well with our direct observation of these complexes. These results emphasize that it is essential for understanding diffusion of membrane proteins to measure not only the mobility of individual molecules but also to consider the effect of molecular crowding (Kirchhoff, 2014; Kirchhoff *et al.*, 2008), which is only achievable experimentally by HS-AFM.

HS-AFM imaging of thylakoid membranes also provided direct evidence of rotational diffusion of photosynthetic complexes (Figure 4). Rotational mobility of membrane proteins have been detected by many different techniques, including HS-AFM (Casuso *et al*., 2012, Jacobson *et al*., 2019). The rotational diffusion is a consequence of combined effects: the protein's structure, its location, its function, protein-protein and

protein-lipid interactions, as well as the local environment (Guigas and Weiss, 2006, Rahman *et al.*, 1992, Wang *et al.*, 2010, Wong and Case, 2008). In a much more simplified, reconstituted membrane (50% of protein density and ~10-fold smaller than PSII reaction core), Casuso *et al.* were also able to detect by HS-AFM the rotational motion of OmpF homotrimers (Casuso *et al.*, 2012). In both cases, photosynthetic and reconstituted membranes, the rotation coefficient of the proteins could not be determined, because of the heterogeneous mobility of the proteins due to the confinement effect. It is conceivable that during the photosynthetic reactions, protein rotation could affect the efficiency of electron transfer process. For example, a rotation of Cyt b_6f with respect to PSII could potentially speed up or slow down the diffusion and binding of the plastoquinone/plastoquinol molecular carriers in the membrane.

In conclusion, we demonstrate that HS-AFM is a powerful technique for characterizing the dynamics of photosynthetic protein complexes in grana thylakoid membranes. Our real-time HS-AFM observation showed heterogeneous mobility of individual proteins. With our current HS-AFM setup, the molecular displacement of PSII and Cyt b_6f was indistinguishable. Evidently, this methodology can be adapted to study the effects of light, pH, ionic strength, inhibitors, mutations, etc. on the mobility of the complexes in thylakoid membranes. Our successful application of HS-AFM to photosynthetic proteins in grana membranes opens a much-needed avenue to address long-standing questions regarding the dynamics of these protein complexes during photoacclimation and photoprotection mechanisms. A follow-up aim is to visualize intact thylakoids containing the complete photosynthetic electron transport chain, which will allow us to characterize molecular rearrangements induced by light fluctuations in real time.

EXPERIMENTAL PROCEDURES

Grana sample preparation

Grana membranes were prepared from spinach (*Spinacia oleracea*) according to the previous method (Anderson and Boardman, 1966), except for the following modification. Spinach leaves were obtained from a local store and kept in the dark overnight at 4 °C. Digitonin (the final concentration at 0.7% w/v; MilliporeSigma, St. Louis, MO) was used to solubilize chloroplasts (0.4 mg Chl/mL) at 4 °C for 30 min in the buffer containing 50 mM phosphate (pH 7.2), 300 mM sucrose, and 10 mM KCl. Crude grana fractions were removed by centrifugation at 1,000 × g for 3 min at 4 °C. The supernatant was centrifuged at 1,000 × g for 5 min at 4 °C to sediment taller grana. The supernatant was further centrifuged at 1,000 × g for 10 min at 4 °C. The pellet containing shorter-height grana was resuspended in the same buffer and immediately used for AFM observation. Different preparations of grana membranes were considered to be biological replicates. Repeated measurements on the same grana preparation were considered to be technical replicates. Prior to initiating HS-AFM experiments, several preparations (~6) of grana membranes were routinely inspected by conventional AFM to assure reproducibility and uniformity in appearance and dispersion of individual grana discs.

Immunoblot analysis

Membrane samples were solubilized with standard Laemmli sample buffer and separated by electrophoresis using sodium dodecyl sulfate polyacrylamide gel prepared using Any kD TGX precast protein gels (Bio-Rad, Hercules, California). Separated proteins in gel were electroblotted onto a polyvinylidene difluoride membrane using Trans-Blot Turbo transfer system according to the manufacturer's instruction (Bio-Rad, Hercules, California). Primary antibodies specific for D1 (PSII), PsaD (PSI), Lhcb2 (LHCII), Cyt *f*, and AtpB were obtained commercially (Agrisera, Vännäs, Sweden) and used according to their recommendations.

Conventional AFM in air

Grana membranes were deposited on freshly cleaved mica in high ionic strength buffer (10 mM Tris-HCl, pH 8.0, 150 mM KCl, and 25 mM $MgCl_2$) (Scheuring *et al.*, 2004) and incubated at room temperature for 1–3 h. Mica was rinsed with water ten times and dried under N_2 gas flow for 2 min. We used a Multimode AFM Nanoscope V (Bruker Co. California) and performed the observation as described previously (Onoa *et al.*, 2014).

HS-AFM

Grana membranes were diluted 5- to 10-fold in high ionic strength adsorption buffer. Two microliters of the diluted sample was deposited on freshly cleaved mica and incubated for 1 h in the dark. Weakly bound membranes were removed by rinsing 10 times with imaging buffer (10 mM Tris-HCl, pH 8.0, 5 mM $MgCl_2$, 150 mM KCl) followed by a gentle, brief (2 s) puff of high purity Argon gas. The sample was immediately immersed in 2 μL of imaging buffer. For scanning, the HS-AFM bath contained the same imaging buffer. We optimized data acquisition and analyses to deconvolute the noise intrinsic to our HS-AFM (e.g. drift, probe, and scan artifacts) to minimize over-interpretation and maximize unbiased observations. We used the Ando-model HS-AFM (Research Institute of Biomolecule Metrology, Tsukuba, Japan) (Ando *et al.*, 2008) equipped with a near-infrared laser (830 nm wavelength) to minimize chlorophyll excitation during observation. All optical components were adjusted to near infrared region except for an objective lens. We used a wide area scanner (maximum range: 6×6 μm^2 in XY-directions and 1 μm in Z-direction). First, we set the scan range between 1×1 μm^2 to 4×4 μm^2 in order to find appropriate membranes. Then, we moved the stage to place the membrane at the cantilever position and observed it with scan range of 150~500 nm^2 at 1 frame s^{-1}. The samples were scanned in liquid using tapping mode at room temperature. The deflection of micron-sized cantilever (AC10DS, Olympus, Japan, spring constant ~0.1 N/m, resonance frequency 400~500 kHz in liquid) was detected using an optical beam detector. The free-oscillation amplitude of the cantilever (A_0) was set to ~2 nm, and the set point of feedback amplitude was set to about $0.9A_0$. The detailed procedure of HS-AFM observation was described elsewhere (Uchihashi *et al.*, 2012).

Data analysis

Individual frames from HS-AFM movies selected from a second biological replicate experiment were processed using customized algorithms written in Igor Pro (Wave Metrics Inc. Oregon). First, noise was reduced by Gaussian filtering followed by a flattening filter to accurately measure heights. Second, entire patches were tracked using a 2D correlation method to correct and minimize lateral drift (Uchihashi *et al.*, 2011). Finally, the corrected images were aligned to the first frame or average image (QSM membranes) to remove residual artifactual displacements. Movies whose center of mass from individual frames was satisfactorily aligned were selected for further analysis. Particle dimensions (median heights, diameters, center of mass, etc.) were obtained from those selected by thresholding segmentation (package features) from the particle and pore analysis module included in SPIP™ (Hørsholm, Denmark). Dimensional fluctuations and spatial displacements were tracked, plotted, and fit using customized scripts written in Wolfram Mathematica® (Illinois) or Igor Pro. The goodness of fit for normal distributions was done using the Akaike information criterion. The contrast of high resolution images was digitally adjusted to facilitate the visual detection of dimeric structures chosen for particle analysis; therefore, small and membrane-embedded proteins appear invisible. Particle MSD was calculated according to:

$$MSD = \frac{1}{T}\sum (x - x_0)^2 + (y - y_0)^2 \qquad (1)$$

where x and y are the particle's center of mass coordinates at different time points; x_0 and y_0 represents the initial *x,y* center of mass coordinate; T = total duration of observation. Each MSD trace was then fit with two diffusion models: confined (Eq. 2) (Ishihama and Funatsu, 2009; Daumas *et al.*, 2003) and Brownian (Eq. 3).

$$MSD = \frac{L^2}{3}\left\{1 - e^{-t/\tau}\right\} + 4D_{macro}t \qquad (2)$$

$$MSD = 4Dt \qquad (3)$$

where D_{macro} = macroscopic diffusion coefficient, L = confined domain, τ = equilibration time, D = the diffusion coefficient of natural diffusion and t = time interval. The microscopic diffusion coefficient (D_{micro}) in the confined diffusion model can be obtained from Eq. 4 (Ishihama and Funatsu, 2009; Daumas *et al.*, 2003).

$$D_{micro} = \frac{L^2}{12\tau} \qquad (4)$$

To establish whether or not the particle's diffusion properties were identical, we determined the relationship between L and the initial diffusion coefficient ($D_{initial}$ at ≈ 4 s) or D_{micro} obtained from Eq. 4. We confirmed an apparent particle's segregation from this correlation by applying the k-means clustering criteria. The diffusion coefficients reported in this study were obtained from the best fit (Brownian or confined) to the average trace resulted from each subgroup. The goodness of the fitting was evaluated by determining the squared correlation coefficient R^2.

To dissect the molecular rotational movement, we calculated the correlation coefficient of 2D image (Uchihashi *et al.*, 2011). After tracking a selected molecule to eliminate lateral diffusion effects, we defined it into a rectangular region of interest (ROI) to calculate the 2D correlation coefficient frame by frame with Eq. 5.

$$CCV = \frac{\sum_m \sum_n (H_{m,n} - \overline{H})(I_{m,n} - \overline{I})}{\sqrt{(\sum_m \sum_n (H_{m,n} - \overline{H})^2)}\sqrt{(\sum_m \sum_n (I_{m,n} - \overline{I})^2)}} \qquad (5)$$

H and I denote the height values at a pixel point (m, n) for the targeted ROI at different time points and the initial one (from the first frame), respectively. $\overline{H}$ and $\overline{I}$ are their respective height mean values of the matrix.

ACKNOWLEDGMENTS

We thank Daniel Westcott and Graham Fleming for critical reading of the manuscript.

COMPETING OF INTEREST

No competing interests declared.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

Figure S1. Grana membranes used for HS-AFM observation.

Figure S2. HS-AFM characterization of the dimeric protein structures in grana.

Figure S3. HS-AFM micrographs of grana membranes.

Figure S4. Time-lapse HS-AFM observation of a representative grana membrane showing large protein displacements.

Figure S5. Correlation between displacement and height of dimeric proteins in the membrane.

Figure S6. HS-AFM characterization of the dimeric protein structures in VPM membranes.

Movie S1. HS-AFM movie of photosynthetic proteins in the first group of grana membranes, showing quasi-static mobility (QSM).

Movie S2. HS-AFM movie of photosynthetic proteins in the second group of grana membranes, showing variable protein mobility (VPM).

Movie S3. HS-AFM movie of photosynthetic proteins in the second group of grana membranes (VPM), showing that a large fraction of proteins is mobile.

Movie S4. HS-AFM movie of photosynthetic proteins in a VPM grana membrane, showing a rotational movement of a dimeric structure with respect to its neighbor.

REFERENCES

Anderson, J.M. (1981) Consequences of spatial separation of photosystem 1 and 2 in thylakoid membranes of higher plant chloroplasts. *FEBS Lett.*, **124**, 1-10.

Anderson, J.M. and Boardman, N.K. (1966) Fractionation of the photochemical systems of photosynthesis I. Chlorophyll contents and photochemical activities of particles isolated from spinach chloroplasts. *BBA - Biophys. Incl. Photosynth.*, **112**, 403–421.

Andersson, B. and Anderson, J.M. (1980) Lateral heterogeneity in the distribution of chlorophyll-protein complexes of the thylakoid membranes of spinach chloroplasts. *BBA - Bioenerg.*, **593**, 427-440.

Ando, T., Uchihashi, T. and Fukuma, T. (2008) High-speed atomic force microscopy for nano-visualization of dynamic biomolecular processes. *Prog. Surf. Sci.*, **83**, 337–437.

Austin, J.R. and Staehelin, L.A. (2011) Three-Dimensional Architecture of Grana and Stroma Thylakoids of Higher Plants as Determined by Electron Tomography. *Plant Physiol.*, **155**, 1601-1611.

Banerjee, S., Sun, Z., Hayden, E.Y., Teplow, D.B. and Lyubchenko, Y.L. (2017) Nanoscale Dynamics of Amyloid β-42 Oligomers As Revealed by High-Speed Atomic Force Microscopy. *ACS Nano*., **11**, 12202-12209.

Betterle, N., Ballottari, M., Zorzan, S., Bianchi, S. de, Cazzaniga, S., Dall'Osto, L., Morosinotto, T. and Bassi, R. (2009) Light-induced dissociation of an antenna hetero-oligomer is needed for non-photochemical quenching induction. *J. Biol. Chem.*, **284**, 15255–15266.

Betzig, E., Patterson, G.H., Sougrat, R., Lindwasser, O.W., Olenych, S., Bonifacino, J.S., Davidson, M.W., Lippincott-schwartz, J. and Hess, H.F. (2006) Imaging Intracellular Fluorescent Proteins at Nanometer Resolution. *Science*, **313**, 1642–1646.

Boekema, E.J., Breemen, J.F.L. Van, Roon, H. Van and Dekker, J.P. (2000) Arrangement of photosystem II supercomplexes in crystalline macrodomains within the thylakoid membrane of green plant chloroplasts. *J. Mol. Biol.,* **301**, 1123-1133.

Casuso, I., Khao, J., Chami, M., Paul-Gilloteaux, P., Husain, M., Duneau, J.P., Stahlberg, H., Sturgis, J.N. and Scheuring, S. (2012) Characterization of the motion of membrane proteins using high-speed atomic force microscopy. *Nat. Nanotechnol.*, **7**, 525–529.

Colom, A., Casuso, I., Rico, F. and Scheuring, S. (2013) A hybrid high-speed atomic force-optical microscope for visualizing single membrane proteins on eukaryotic cells. *Nat. Commun.*, **4**: 2155.

Consoli, E., Croce, R., Dunlap, D.D. and Finzi, L. (2005) Diffusion of light-harvesting complex II in the thylakoid membranes. *EMBO Rep.*, **6**, 782-786.

Daumas, F., Destainville, N., Millot, C., Lopez, A., Dean, D. and Salomé, L. (2003) Confined diffusion without fences of a G-protein-coupled receptor as revealed by single particle tracking. *Biophys. J.*, **84**, 356–366.

Dekker, J.P. and Boekema, E.J. (2005) Supramolecular organization of thylakoid membrane proteins in green plants. *Biochim. Biophys. Acta - Bioenerg.*, **1706**, 12–39.

Dekker, J.P., Germano, M., Roon, H. Van and Boekema, E.J. (2002) Photosystem II solubilizes as a monomer by mild detergent treatment of unstacked thylakoid membranes. *Photosynth. Res.*, **72**, 203-210.

Drepper, F., Haehnel, W., Carlberg, I. and Andersson, B. (1993) Lateral Diffusion of an Integral Membrane Protein: Monte Carlo Analysis of the Migration of Phosphorylated Light-Harvesting Complex II in the Thylakoid Membrane. *Biochemistry.*, **32**, 11915-11922.

Eerden, F.J. Van, Berg, T. Van Den, Frederix, P.W.J.M., Jong, D.H. De, Periole, X. and Marrink, S.J. (2017) Molecular Dynamics of Photosystem II Embedded in the Thylakoid Membrane. *J. Phys. Chem. B*, **121**, 3237–3249.

Fantner, G.E., Schitter, G., Kindt, J.H., *et al*. (2006) Components for high speed atomic force microscopy. *Ultramicroscopy*, **106**, 881–887.

Fukuda, S., Uchihashi, T., Iino, R., Okazaki, Y., Yoshida, M., Igarashi, K. and Ando, T. (2013) High-speed atomic force microscope combined with single-molecule fluorescence microscope. *Rev. Sci. Instrum.*, **84**., 073706.

Goral, T.K., Johnson, M.P., Brain, A.P.R., Kirchhoff, H., Ruban, A. V. and Mullineaux, C.W. (2010) Visualizing the mobility and distribution of chlorophyll proteins in higher plant thylakoid membranes: effects of photoinhibition and protein phosphorylation. *Plant J.*, **62**, 948–959.

Goral, T.K., Johnson, M.P., Duffy, C.D.P., Brain, A.P.R., Ruban, A. V and Mullineaux, C.W. (2012) Light-harvesting Antenna Composition Controls the Macrostructure and Dynamics of Thylakoid Membranes in Arabidopsis. *Plant J.*, **69**, 289–301.

Guigas, G. and Weiss, M. (2006) Size-dependent diffusion of membrane inclusions. *Biophys. J.*, **91**, 2393-2398.

Hartigan, J.A. and Wong, M.A. (1979) A k-means clustering algorithm. *Appl. Stat.*, **28**, 100-108.

Ishihama, Y. and Funatsu, T. (2009) Single molecule tracking of quantum dot-labeled mRNAs in a cell nucleus. *Biochem. Biophys. Res. Commun.*, **381**, 33–38.

Iwai, M., Roth, M.S. and Niyogi, K.K. (2018) Subdiffraction-resolution live-cell imaging for visualizing thylakoid membranes. *Plant J.*, **96**, 233-243.

Jacobson, K., Liu, P. and Lagerholm, B.C. (2019) The Lateral Organization and Mobility of Plasma Membrane Components. *Cell*, **177**, 806-819

Johnson, M.P., Goral, T.K., Duffy, C.D.P., Brain, A.P.R., Mullineaux, C.W. and Ruban, A. V. (2011) Photoprotective Energy Dissipation Involves the Reorganization of Photosystem II Light-Harvesting Complexes in the Grana Membranes of Spinach Chloroplasts. *Plant Cell*, **23**, 1468–1479.

Johnson, M.P., Vasilev, C., Olsen, J.D. and Hunter, C.N. (2014) Nanodomains of Cytochrome $b_6 f$ and Photosystem II Complexes in Spinach Grana Thylakoid Membranes. *Plant Cell*, **26**, 3051–3061.

Kaňa, R. (2013) Mobility of photosynthetic proteins. *Photosynth. Res*, **116**, 465-479

Katan, A.J., Vlijm, R., Lusser, A. and Dekker, C. (2015) Dynamics of nucleosomal structures measured by high-speed atomic force microscopy. *Small*, **11**, 976–984.

Kirchhoff, H. (2014) Diffusion of molecules and macromolecules in thylakoid membranes. *Biochim. Biophys. Acta - Bioenerg.*, **1837**, 495–502.

Kirchhoff, H., Haferkamp, S., Allen, J.F., Epstein, D.B.A. and Mullineaux, C.W. (2008) Protein Diffusion and Macromolecular Crowding in Thylakoid Membranes. *Plant Physiol.*, **146**, 1571–1578.

Kirchhoff, H., Lenhert, S., Bchel, C., Chi, L., Nield, J., Chi, L. and Nield, J. (2008) Probing the Organization of Photosystem II in Photosynthetic Membranes by Atomic Force Microscopy. *Biochemistry*, **47**, 431–440.

Kirchhoff, H., Tremmel, I., Haase, W. and Kubitscheck, U. (2004) Supramolecular photosystem II organization in grana thylakoid membranes: Evidence for a structured arrangementt. *Biochemistry*, **43**, 9204–9213.

Kodera, N. Yamamoto, D. Isikawa, R and Ando, T. (2010) Video imaging of walking myosin V by high-speed atomic force microscopy. *Nature*, **468**, 72–76.

Kodera, N., Sakashita, M. and Ando, T. (2006) Dynamic proportional-integral-differential controller for high-speed atomic force microscopy. *Rev. Sci. Instrum.*, **77**, 083704.

Kodera, N., Yamashita, H. and Ando, T. (2005) Active damping of the scanner for high-speed atomic force microscopy. *Rev. Sci. Instrum.*, **76**, 1–5.

Kouřil, R., Wientjes, E., Bultema, J.B., Croce, R. and Boekema, E.J. (2013) High-light vs. low-light: Effect of light acclimation on photosystem II composition and organization in Arabidopsis thaliana. *Biochim. Biophys. Acta - Bioenerg.*, **1827**, 411–419.

Kurisu, G., Zhang, H., Smith, J.L. and Cramer, W.A. (2003) Structure of the Cytochrome b6f Complex of Oxygenic Photosynthesis: Tuning the Cavity. *Science*, **302**, 1009-1014.

Milhiet, P.E., Yamamoto, D., Berthoumieu, O., Dosset, P., Grimellec, C. le, Verdier, J.M., Marchal, S. and Ando, T. (2010) Deciphering the structure, growth and assembly of amyloid-like fibrils using high-speed atomic force microscopy. *PLoS One*, **5**: e13240.

Miyagi, A., Ando, T. and Lyubchenko, Y.L. (2011) Dynamics of nucleosomes assessed with time-lapse high-speed atomic force microscopy. *Biochemistry*, **50**, 7901–7908.

Muller, P. (2001) Non-Photochemical Quenching. A Response to Excess Light Energy. *Plant Physiol.*, **125**, 1558–1566.

Nelson, N. and Yocum, C.F. (2006) Structure and function of photosystems I and II. *Annu. Rev. Plant Biol.*, **57**, 521-565.

Nosek, L., Semchonok, D., Boekema, E.J., Ilík, P. and Kouřil, R. (2017) Structural variability of plant photosystem II megacomplexes in thylakoid membranes. *Plant J.*, **89**, 104–111.

Onoa, B., Schneider, A.R., Brooks, M.D., Grob, P., Nogales, E., Geissler, P.L., Niyogi, K.K. and Bustamante, C. (2014) Atomic force microscopy of photosystem II and its unit cell clustering quantitatively delineate the mesoscale variability in Arabidopsis Thylakoids. *PLoS One.*, **9**, e101470.

Phuthong, W., Huang, Z., Wittkopp, T.M., Sznee, K., Heinnickel, M.L., Dekker, J.P., Frese, R.N., Prinz, F.B. and Grossman, A.R. (2015) The Use of Contact Mode Atomic Force Microscopy in Aqueous Medium for Structural Analysis of Spinach Photosynthetic Complexes. *Plant Physiology*. **169**, 1318-1332

Rahman, N.A., Pecht, I., Roess, D.A. and Barisas, B.G. (1992) Rotational dynamics of type I Fc epsilon receptors on individually-selected rat mast cells studied by polarized fluorescence depletion. *Biophys. J.*, **61**, 334-346

Ruan, Y., Miyagi, A., Wang, X., Chami, M., Boudker, O. and Scheuring, S. (2017) Direct visualization of glutamate transporter elevator mechanism by high-speed AFM. *Proc. Natl. Acad. Sci.*, **114**, 1584–1588.

Ruban, A. V. (2016) Nonphotochemical Chlorophyll Fluorescence Quenching: Mechanism and Effectiveness in Protecting Plants from Photodamage. *Plant Physiol.*, **170**, 1903–1916.

Ruban, A. V. and Johnson, M.P. (2015) Visualizing the dynamic structure of the plant photosynthetic membrane. *Nat. Plants*, **1**, 1–9.

Sakiyama, Y., Mazur, A., Kapinos, L.E. and Lim, R.Y.H. (2016) Spatiotemporal dynamics of the nuclear pore complex transport barrier resolved by high-speed atomic force microscopy. *Nat. Nanotechnol.*, **11**, 719-724.

Scheuring, S., Bernadac, A., Sturgis, J.N., Prima, V., Rigaud, J.-L. and Levy, D. (2004) Watching the photosynthetic apparatus in native membranes. *Proc. Natl. Acad. Sci.*, **101**, 11293-11297

Sigal, Y.M., Zhou, R. and Zhuang, X. (2018) Visualizing and discovering cellular structures with super-resolution microscopy. *Science*, **361**, 880-887.

Stoichev, S., Krumova, S.B., Andreeva, T., Busto, J. V., Todinova, S., Balashev, K., Busheva, M., Goñi, F.M. and Taneva, S.G. (2015) Low pH modulates the macroorganization and thermal stability of PSII supercomplexes in grana membranes. *Biophys. J.*, **108**, 844–853.

Suzuki, Y., Shin, M., Yoshida, A., Yoshimura, S.H. and Takeyasu, K. (2012) Fast microscopical dissection of action scenes played by Escherichia coli RNA polymerase. *FEBS Lett.*, **586**, 3187–3192.

Sznee, K., Dekker, J.P., Dame, R.T., Roon, H. Van, Wuite, G.J.L. and Frese, R.N. (2011) Jumping mode atomic force microscopy on grana membranes from spinach. *J. Biol. Chem.*, **286**, 39164–39171.

Uchihashi, T., Iino, R., Ando, T. and Noji, H. (2011) High-speed atomic force microscopy reveals rotary catalysis of rotorless F 1-ATPase. *Science*, **333**, 755–758.

Uchihashi, T., Kodera, N. and Ando, T. (2012) Guide to video recording of structure dynamics and dynamic processes of proteins by high-speed atomic force microscopy. *Nat. Protoc.*, **7**, 1193–1206.

Umena, Y., Kawakami, K., Shen, J.R. and Kamiya, N. (2011) Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9Å. *Nature*, **473**, 55-60.

Wang, Y., Li, C. and Pielak, G.J. (2010) Effects of proteins on protein diffusion. *J. Am. Chem. Soc.,* **132**, 9392-9397

Willig, K.I., Rizzoli, S.O., Westphal, V., Jahn, R. and Hell, S.W. (2006) STED microscopy reveals that synaptotagmin remains clustered after synaptic vesicle exocytosis. *Nature*, **440**, 935–939.

Wobbe, L., Bassi, R. and Kruse, O. (2016) Multi-Level Light Capture Control in Plants and Green Algae. *Trends Plant Sci.*, **21**, 55-68.

Wong, V. and Case, D.A. (2008) Evaluating rotational diffusion from protein MD simulations. *J. Phys. Chem. B*, **112**, 6013-6024.

Yakushevska, A.E., Jensen, P.E., Keegstra, W., Roon, H. Van, Scheller, H. V., Boekema, E.J. and Dekker, J.P. (2001) Supermolecular organization of photosystem II and its associated light-harvesting antenna in Arabidopsis thaliana. *Eur. J. Biochem.*, **268**, 6020-6028.

Yamashita, H., Taoka, A., Uchihashi, T., Asano, T., Ando, T. and Fukumori, Y. (2012) Single-molecule imaging on living bacterial cell surface by high-speed AFM. *J. Mol. Biol.*, **422**, 300–309.

Figure 1

Figure 2

Figure 3

Figure 4

Figure Legends

Figure 1. HS-AFM micrographs of the representative grana membrane. (a) HS-AFM image of a grana membrane. Z-scale, 94.4 nm. (b) Magnified averaged HS-AFM image of a grana membrane. Z-scale, 6.0 nm. The image was acquired at 1 frame per second and averaged over 50 frames. The contrast of the membrane was adjusted to enhance the dimeric structures; small, membrane-embedded proteins such as LHCII are invisible.

Figure 2. HS-AFM characterization of the protein mobility observed in the first (QSM) group of membranes. (a) Time-average image of a representative HS-AFM movie. Each image was acquired at 1 frame per second and averaged over 50 frames. (b) Mean square displacement (MSD) trajectories of 53 proteins; gray thin lines are MSD trajectories of each tracked protein in this membrane, and the superimposed black thick line is the average trajectory of all proteins selected in this grana membrane. The confined diffusion model agrees well with the average displacement of proteins in the QSM group of membranes. Inset shows the enlarged MSD trajectories.

Figure 3. HS-AFM characterization of the protein mobility observed in the second (VPM) group of membranes. (a) Time-lapse HS-AFM images showing the heterogeneous protein mobility in the VPM group of membranes. Images were acquired at 1 frame per second. Corresponding illustrations of few proteins enclosed in the red squares are presented to facilitate the visualization of the protein dynamics. The particles in the cartoon were drawn freehand by visually tracing the protein's contour in the HS-AFM images. Z-scale, 6.7 nm. (b) MSD trajectories of the proteins selected in this membrane (N = 50). Gray lines are individual proteins' MSD trajectories; superimposed yellow line is the MSD average of all proteins; dashed black line is a fit to a confined diffusion model (Eq. 2); and orange dotted line is a fit to the Brownian diffusion model (Eq. 3). (c) Relationship between each protein's confined length and its initial diffusion coefficient (4 s). A protein's mobility can be segregated into two distinct groups as shown in blue circles and magenta squares, according to the k-means criteria. (d) Average MSD traces of two distinct groups (blue and magenta in C); dotted and dashed lines are the fits to Brownian and confined diffusion models, respectively.

Figure 4. HS-AFM characterization of the rotational displacement of a dimeric structure. (a) Time-lapse HS-AFM images illustrating the rotation of a dimeric protein (magenta arrow) with respect to its neighbor protein (blue arrow) captured at 1 fps. Corresponding illustrations are shown in the right (the particles were drawn by following the protein's contour as described in Fig. 3). Z-scale, 4.5 nm. (b) Variation of the angle (θ) in panel A over 50 s. (c) Time-course examples of the 2D correlation coefficient of variation (CCV) for two different proteins. Non-rotational symmetry will produce CCV values closer to 1. Rotational displacement is depicted by the red trace. Negligible rotational displacement is illustrated by the blue trace (blue arrow in the bottom right inset). (d) Left, the average HS-AFM image. Right, height profiles indicated in averaged AFM images (left).