A novel rotifer-derived alkaloid paralyzes schistosome larvae and prevents

infection

Jiarong Gao[1,2,9], Ning Yang[3,9], Fred A. Lewis[4^], Peter Yau[5], James J. Collins III[2†], Jonathan V.

Sweedler[3*], Phillip A. Newmark[2,6,7,8*]

[1]Cellular and Molecular Biology Program, University of Wisconsin-Madison, Madison, WI,

USA. [2]Department of Cell and Developmental Biology, University of Illinois at Urbana-

Champaign, Urbana, IL, USA. [3]Department of Chemistry, University of Illinois at Urbana-

Champaign, Urbana, IL, USA. [4]Biomedical Research Institute, Rockville, MD, USA. [5]Roy J.

Carver Biotechnology Center, University of Illinois at Urbana-Champaign, Urbana, IL, USA.

[6]Howard Hughes Medical Institute, USA. [7]Morgridge Institute for Research, Madison, WI, USA.

[8]Department of Integrative Biology, University of Wisconsin-Madison, Madison, WI, USA.

[9]These authors contributed equally: Jiarong Gao, Ning Yang.

[*]e-mail: jsweedle@illinois.edu; pnewmark@morgridge.org

[^]Present address: Bernville, PA, USA

[†]Present Address: Department of Pharmacology, UT Southwestern Medical Center, Dallas, TX,

USA

Schistosomiasis, caused by the blood fluke *Schistosoma*, is a devastating and

widespread disease affecting more than 200 million people and threatening over 700

million people worldwide[1,2]. Treatment relies exclusively on one drug, praziquantel, for

which there are concerns of emerging drug resistance[3,4] and which has shown some

limitations in mass drug administration programs[5]. Efforts to develop an effective

vaccine to combat schistosomiasis have been unsuccessful; thus, there is an urgent

need for alternative drugs and better prevention strategies. Here we report the

discovery and purification of a novel tetracyclic alkaloid secreted by the rotifer *Rotaria*

rotatoria, which lives commensally on snails that serve as intermediate hosts for *S.*

mansoni[6]. This compound effectively paralyzes cercariae, the infective larvae of

schistosomes, at nanomolar concentrations*.* Furthermore, we show that worm burden

and disease pathology are significantly reduced when cercariae are pre-treated with this

Schistosome Paralysis Factor (SPF). Identifying this molecule provides new directions

for understanding motility of schistosome cercariae and new strategies for preventing

schistosomiasis.

Schistosomes have a complex life cycle that alternates between an intermediate host

(snail) and a definitive host (mammal) via two free-living, water-borne forms called

miracidia and cercariae, respectively[7] (Fig. 1a). For decades, inconsistency in cercarial

production by snails and infectivity of mammalian hosts has been observed in most

schistosome laboratories[8]. Intriguingly, Stirewalt and Lewis reported that rotifer

colonization on snail (*Biomphalaria glabrata*) shells significantly reduced cercariae

output, motility, and infectivity[6]. Furthermore, they observed that cercarial motility was

affected not only by the presence of rotifers, but also by rotifer-conditioned water,

indicating that rotifers released water-soluble molecules with paralytic activity. Almost

40 years have passed since this important finding, yet this factor's identity has remained

a mystery.

Encouraged by this anti-cercarial effect and its potential to prevent schistosome

infection, we sought to purify this paralyzing agent. We isolated individual rotifers from

snail shells and found two species, *Rotaria rotatoria* (Fig. 1b) and *Philodina acuticornis*

(Fig. 1c), as previously reported[6]. To identify which rotifer was responsible for the

paralytic effect, we grew clonal isolates of each species, producing rotifer-conditioned

artificial pond water (APW). Adding *Rotaria*-conditioned APW to freshly collected

cercariae resulted in gradual paralysis within five minutes (Fig. 1d). Most cercariae

stopped swimming and sank to the bottom of the dish. Tapping the dish could stimulate

their movement, but their response was limited to writhing on the dish bottom or short-

distance swimming before becoming paralyzed again. In contrast, *P. acuticornis*-

conditioned water had no effect (Fig. 1e).

To purify the paralyzing agent, we performed molecular weight cut-off filtration (MWCO)

of rotifer-conditioned water and found that the activity was present in the <650 Da

fraction. The <650 Da filtrate was fractionated by reversed-phase high-performance

liquid chromatography (RP-HPLC) (Fig 2a) and each fraction was tested on cercariae.

Paralysis was only observed following treatment with a peak eluting at 25-27 min (Fig.

2b). As expected, this peak was detected only in *R. rotatoria*- but not *P. acuticornis*-

conditioned water (Fig. 2b). A second round of HPLC on this peak revealed one peak

(eluting at 24-26 min) with paralytic activity (Fig. 2c). A predominant signal of 273.16 Da

(M+H) in this peak was revealed by matrix-assisted laser desorption/ionization mass

spectrometry (MALDI-MS) (Fig. 2d). Consistent with the paralysis assay, this signal (m/z

273.16) was detected exclusively in the fraction eluting at 24-26 min but not in the

fractions before or after (Fig. 2e). These results suggested that the component with m/z

273.16 was the paralyzing agent, which we named "Schistosome Paralysis Factor"

(SPF). We then determined the monoisotopic mass for protonated SPF using high-

resolution quadrupole time-of-flight (Q-TOF) MS, 273.1595 Da (Fig. 2f), suggesting

$C_{16}H_{20}N_2O_2$ as the best-fitting formula for SPF.

To elucidate its structure, we purified SPF from *R. rotatoria*-conditioned water, obtaining

0.126 mg from 25 L. Nuclear magnetic resonance (NMR) spectroscopy revealed a novel

tetracyclic structure, which conformed to the best-fitting formula and contained one

exchangeable proton, consistent with the results of H/D-exchange MS (Fig. 2g;

Extended Data Fig. 1-6; Extended Data Table 1). Nuclear Overhauser effect

spectroscopy (NOESY) (Extended Data Fig. 7) and 3D minimized steric-energy models

favored the (R, S, S) configuration (steric energy 35.6 kcal/mol) over the (S, R, R)

configuration (110.9 kcal/mol) at the three chiral centers, C12-14. Based on energy-

minimized models, the proton-proton spatial distance in the (R, S, S) configuration

matched the relative intensity of cross-peak signals in NOESY, whereas the distances

in the (S, R, R) configuration did not (Extended Data Table 2; Extended Data Fig. 7).

To test its dose dependency, we examined the paralytic effect of serially diluted SPF on

cercariae by quantifying their movement over time. In the absence of SPF, over 82% of

cercariae were free-swimming over three minutes (Fig. 3a). In 2.5 nM SPF, the

percentage of free-swimming cercariae dropped to 67% at three-minutes post drug

treatment. As the concentration of SPF increased, so did the rate of paralysis, and more

cercariae were paralyzed at the end of treatment. We observed maximum effects in 250

nM and 2.5 μM SPF, with the majority of cercariae paralyzed within 30 s.

Two natural compounds isolated from *Streptomyces sp.*, ht-13-A and ht-13-B[9], are

structurally related to SPF. All three alkaloids share a novel oxepineindole framework

fused with a pyrrolidine ring (Fig. 3a-c). Although synthesis of SPF has not been

achieved, total syntheses of ht-13-A and ht-13-B have been reported[10-12]. To test

whether this shared tetracyclic scaffold is responsible for the paralytic effect, we

analyzed structure-activity relationships by using ht-13-A, ht-13-B, three ht-13-A

derivatives, and one epimer in cercarial paralysis assays. Importantly, ht-13-A, although

not as potent as SPF, also had a paralytic effect on cercariae (Fig. 3b). In contrast, ht-

13-B did not paralyze cercariae, suggesting that the extra methyl group disrupts

interaction with the target (Fig. 3c). Of the three ht-13-A analogs, only ht-13-A-pr

effectively paralyzed cercariae; it was more potent than ht-13-A, indicating that the

nature of the side chain is important for proper target interaction (Fig. 3d-e). In contrast

to ht-13-A, the epimer was unable to paralyze cercariae, supporting the (R, S)

configuration of SPF at C12, 13.

Since motility of the cercarial tail is essential for swimming and provides force for skin

penetration[13-15], we examined whether SPF prevented infection. We treated ~200

cercariae with different concentrations of SPF for 10 mins, and tested their ability to

infect mice after a 30-min exposure to their tails (N=6 for each condition). Six-weeks

post infection, we euthanized the mice, counted schistosomes recovered after hepatic

portal vein perfusion, and examined liver pathology. From controls, we recovered 83

adult worms on average (Fig. 4a), consistent with typical recoveries of ~40%[16]. Livers

from these mice appeared dark and contained extensive granulomas (Fig. 4a). In

contrast, we did not recover any adult worms from mice after treatment with 250 nM or

2.5 μM SPF (Fig. 4b) and no granulomas were observed (Fig. 4a). Histological

examination confirmed that these livers were free of schistosome eggs (Fig. 4e),

suggesting complete inhibition of infection. These data are consistent with the full

paralysis observed after treatment with 250 nM or 2.5 μM SPF (Fig. 3a). Although 25

nM SPF paralyzed most cercariae *in vitro*, the effects on mouse infection were not as

severe (Fig. 4a). Mechanical and/or chemical stimuli from mouse tails may overcome

SPF-induced paralytic effects at low SPF concentrations. Notably, neither Ht-13-A nor

Ht-13-A-pr blocked infection as completely as 250 nM SPF, even at 25 μM (Fig. 4a, c, d,

f, g). Under more realistic infection conditions, in which mouse tails were lifted 1-2 cm

from the bottom of the test tube containing cercariae, so they had to swim actively

towards the tail to infect the mouse, Ht-13-A and Ht-13-A-pr were still not as effective as

SPF, which completely blocked infection (Extended Data Fig. 8).

This work has identified a novel tetracyclic alkaloid, produced by the rotifer *R. rotatoria,*

that paralyzes the infective larvae of schistosomes. Although its mechanism of action

remains unknown, its chemical structure provides important clues. SPF contains a

serotonin backbone, suggesting that SPF might antagonize serotonin signaling, perhaps

via G protein-coupled receptors (GPCRs) or serotonin-gated channels. Consistent with

this idea, the structurally related compounds, ht-13-A and ht-13-B, bind several human

serotonin receptors[9]. In schistosomes, serotonin has been implicated in neuromuscular

functions in multiple life stages[17-20]; knocking down a serotonergic GPCR (Sm5HTR) in

schistosomulae and adult worms led to decreased movement[21]. Interestingly,

praziquantel partially activates human HT2BR, suggesting that it may also target

schistosome serotonergic GPCRs[22].

The chemical ecology underlying *R. rotatoria*'s production of SPF is also unclear.

Whether SPF is used naturally to combat other aquatic creatures (e.g., to prevent other

rotifers from colonizing areas where *R. rotaria* live) and, thus, the effect on schistosome

cercariae is indirect, or whether SPF benefits the rotifer's commensal host will require

further study. In the past few decades, the discovery and development of natural

products have helped combat parasitic diseases[23]. Based on its ability to block infection,

SPF holds great promise as an anti-schistosomal agent. Identifying the biologically

active chemical scaffolds and understanding SPF's mode of action are expected to

provide important clues for preventing schistosomiasis.

Methods

Artificial pond water

Four stock solutions were prepared to make artificial pond water[24]: 1) 0.25 g/L $FeCl_3 \bullet$
$6H_2O$, 2) 12.9 g/L $CaCl_2 \bullet 2H_2O$, 3) 10g/L $MgSO_4 \bullet 7H_2O$, and 4) 34 g/L KH_2PO_4 1.5 g/L
$(NH_4)_2SO_4$, pH 7.2. For 1L artificial pond water, we added 0.5 mL of $FeCl_3$ solution, 2.5
mL $CaCl_2$ solution, 2.5 mL $MgSO_4$ solution and 1.25 mL phosphate buffer.

Obtaining *S. mansoni* cercariae

Infected *B. glabrata* snails provided by Biomedical Research Institute (BRI, Rockville,
MD) were maintained in artificial pond water and fed Layer Crumbles (chicken feed)
(Rural King, Mattoon, IL). To obtain *S. mansoni* cercariae, *B. glabrata* snails were
exposed to light at 26°C for 1-2 hrs. Artificial pond water containing cercariae was
passed through 100 μm cell strainer (Falcon) to remove snail food and feces. Cercariae
were then collected using custom-made 20 μm cell strainers.

Rotifer culture

Since both rotifer species reproduce parthenogenetically, we clonally expanded each
species into one-liter cultures from a single rotifer. Individual rotifers (*R. rotatoria* and *P.
acuticornis*) were initially isolated from the shell of *B. glabrata* and cultured in artificial
pond water in 24-well plates. Each individual colony was expanded into ever-larger
culture volumes and ultimately maintained in two-liter flasks. Both species were fed
Roti-rich liquid invertebrate food (Florida Aqua Farms Inc). Rotifer-conditioned water
was collected every month by filtering out the rotifers using a 20 μm cell strainer.

166   Filtered rotifers were then passaged to fresh artificial pond water to propagate the

167   cultures.

Crude rotifer-conditioned water preparation

One liter rotifer media was lyophilized, reconstituted with 50 mL dH_2O and filtered

through 10,000 and 650 (MWCO) Pall Minimate TFF Capsules with Omega membrane

(Ann Arbor, MI). Filtrate (<650 Da) was freeze dried. For RP-HPLC, 300 mg of the dried

material was dissolved in dH_2O and run on a RP-HPLC – Merck Chromolith semi-prep

RP-18e column (Darmstadt, Germany) at 5 ml/min using a gradient of 100% A to 60% B

in 60 min. 10 mL fractions were collected and assayed for biological activity. Fractions

containing biological activity were saved for further study.

Further purification of rotifer media

The bioactive fractions were pooled, freeze dried with SpeedVac (Savant, MA),

reconstituted with 500 μL dH_2O and injected into a 4.6mm diameter × 25cm Symmetry

column (Waters, MA). Breeze2 analytical LC system (Waters, MA) was employed for

separation at 0.5 ml/min with the following solvents and gradients: Solvent A, 0.1%

formic acid (FA); solvent B, methanol with 0.1% FA; 0—10 min 0—10% B, 10—30 min

10—35% B, 30—33 min 35—80% B, 33—37 min 80—80% B, 37—40 min 80—0% B.

Eluents were collected manually based on peak elution. All fractions were lyophilized,

reconstituted with water and analyzed with MALDI-MS. Fractions containing biological

activity were saved for future use.

Matrix-assisted laser desorption/ionization mass spectrometry (MALDI-MS) analysis

For each collected fraction, 1 μL of sample solution was spotted on ground steel MALDI target and mixed with 1 μL of alpha-cyano-4-hydroxy-cinnamic acid (CHCA, Sigma-Aldrich, MO) solution (10 mg/mL CHCA in 50% acetonitrile solution with 0.005% trifluoroacetic acid). Mass calibration, spectra acquisition and analysis were performed under conditions as previously described[25].

High-resolution quadrupole time-of-flight mass spectrometry (Q-TOF MS) analysis

1 μL of the bioactive fraction was separated on a Magic 0.1 × 150mm column (Michrom, CA) and analyzed with maXis 4G mass spectrometer (Bruker, MA) using previously established methods for metabolite study[26]. The separation was performed at 300 nl/min by use of solvent A (95% water, 5% acetonitrile with 0.1% FA) and solvent B (5% water, 95% acetonitrile with 0.1% FA) with the following gradient conditions: 0—5 min 4% B, 5—50 min 4—50% B, 50—52 min 50—90% B, 52—60 min 90% B, 60—70 min 90—4% B, 70—90 min 4% B.

Hydrogen/deuterium (H/D) exchange analysis

Acidified deuterated methanol (CD_3OD, methanol—d4, Sigma—Aldrich, MO) was made by adding 1 μL of deuterated FA into 1 mL of CD_3OD. 2 μL of the bioactive fractions were added into 18 μL of acidified methanol above. 15 μL of the mixture were analyzed by direct infusion into a modified 11 Tesla FTMS (Thermo Scientific, MA) through NanoMate robot (Advion, NY)[27]. Full spectra were acquired with resolution set at 100k.

Nuclear magnetic resonance (NMR) analysis

Purified bioactive materials were dissolved in 250 µL of CD_3OD and transferred into a 5 mm Shigemi NMR tube with a glass magnetic plug with susceptibility matched to CD_3OD on the bottom. All NMR data were collected at 40°C on an Agilent VNMRS 750 MHz spectrometer equipped with a 5 mm Varian indirect detection probe with z gradient capability. Collected NMR data included 1H spectrum, gradient selected correlation spectroscopy (gCOSY), total correlation spectroscopy (TOSCY), nuclear overhauser enhancement spectroscopy (NOESY) with a mixing time of 500 ms, heteronuclear single quantum coherence spectroscopy (^1H—^{13}C HSQC) and heteronuclear multiple-bond correlation spectroscopy (^1H—^{13}C HMBC). The NMR spectra were analyzed using Mnova NMR software (Mestrelab Research, Spain).

Determination of SPF concentration

The proton quantification experiments were performed at 23^oC on an Agilent 750 MHz VNMRS NMR spectrometer equipped with a 5mm triple-resonance (^1H/^{13}C/^{15}N) indirect-detection probe with XYZ PFG gradient capability. The probe was calibrated using the qEstimate tool in the Agilent VnmrJ4.2 software with a known standard. The proton spectrum of the sample was collected with a 90-degree pulse angle of 8.5 ms, 16 scans and 10.4 s delay between scans. The Agilent VnmrJ4.2 software was used to determine the concentration of the sample based on the integration values of proton peaks. A total of 5 well-resolved proton peaks (7.12ppm (1H), ~6.89ppm (2H), 4.41 (1H), 3.83 (3H), and ~3.58 (2H)) was used, and the concentration of the sample was 1.55 ± 0.07 mM. All concentrations used in the cercarial paralysis assay were calculated based on this value.

Cercarial paralysis assay

To capture the whole field while avoiding excess reflected light in a well, we used the lid

of 96-well plate (Costar). 40 μL of artificial pond water containing ~50 cercariae were

added to each shallow well on the lid. 10 μL of SPF (dissolved in APW) was then added

to reach the final concentration indicated. Using a high-speed camera (Olympus i-

SPEED TR), attached to a stereomicroscope (Leica MZ125), we recorded cercariae

movement at 20-60 fps at 1.25X magnification just prior to addition of test compounds

until 3-4 min after treatment started. Raw movies were converted to .avi files using i-

SPEED Viewer and compressed into JPEG format using ImageJ (addition of compound

is considered time 0). We then counted the numbers of free swimming/paralyzed

cercariae every 5 s for 1 min and every 30 s thereafter for 3 min. The number of dead

cercariae (those that never swim before and after SPF treatment) were subtracted from

data. Experiments were performed in biological triplicate.

Mouse infectivity assay

Mouse infections were performed by exposing mouse tails to *S. mansoni* cercariae

according to standard protocol from BRI[16] with slight modifications. Briefly, we secured

mice in rodent restrainers (Thomas Scientific, Cat #551-BSRR) and put them vertically

on top of a rack with grids. We pipetted 100 μL of each drug at proper concentration into

a skinny glass tube (Fisher Scientific, Cat #14-958A) inside a 12 X 75 mm holding glass

tube (VWR, Cat # 47729-570). 300 μL of APW containing ~200 cercariae were pipetted

into each skinny tube and incubated for 10 mins before we inserted the mouse tail.

Mouse tails were wiped with APW-moistened Kimwipes, inserted into the skinny tube,

and exposed to cercariae for 30 mins. The mouse tail was touching the bottom of the

test tube unless otherwise specified. We euthanized and perfused these mice six week-

post infection according to standard protocols[16]. For each drug, we initially used three

mice for controls (APW only) and three mice for each concentration tested except for 25

nM Ht-13-A and Ht-13-A-pr. We then repeated the experiments again with three mice

for each condition. In addition to that, we included six mice for 25 nM Ht-13-A and Ht-

13-A-pr.

Adult worms were recovered by hepatic portal vein perfusion and briefly incubated in

2.5% Tricaine (Sigma) to separate males and females. We counted total numbers of

adult worms under a stereomicroscope (Leica MZ75). Livers from infected mice were

fixed in 4% formaldehyde in PBS overnight. Largest liver lobes (left lobe) were

submitted to University of Wisconsin-Madison Histology Core Facility for sectioning and

Hematoxylin and Eosin staining. Each left lobe was evenly cut into 4-6 pieces and

paraffin embedded on a large cassette. One slide (4-6 liver sections) for each liver was

used for histological examination, which provided a representative view throughout the

whole liver lobe. We took a tiled image of the whole slide using a Zeiss Axio Zoom

microscope and used ImageJ to determine the area of each section. Total numbers of

eggs in each section were counted and normalized to the area.

In adherence to the Animal Welfare Act and the Public Health Service Policy on

Humane Care and Use of Laboratory Animals, all experiments with and care of mice

were performed in accordance with protocols approved by the Institutional Animal Care

and Use Committee (IACUC) of the University of Wisconsin-Madison (protocol approval

number M005569).

Statistical analysis

GraphPad Prism (Version 7) was used for all statistical analyses. One-way ANOVA test

followed by Dunnett's multiple comparison test was used. Mean ± S.D. is shown in all

figures.

1   Steinmann, P., Keiser, J., Bos, R., Tanner, M. & Utzinger, J. Schistosomiasis and water
resources development: systematic review, meta-analysis, and estimates of people at
risk. *The Lancet. Infectious diseases* **6**, 411-425, doi:10.1016/S1473-3099(06)70521-7
(2006).
2   WHO. Schistosomiasis and soil-transmitted helminthiases: number of people treated in
2016. *Wkly Epidemiol Rec* **92**, 749-760 (2017).
3   Doenhoff, M. J., Cioli, D. & Utzinger, J. Praziquantel: mechanisms of action, resistance
and new derivatives for schistosomiasis. *Current opinion in infectious diseases* **21**, 659-
667, doi:10.1097/QCO.0b013e328318978f (2008).
4   Wang, W., Wang, L. & Liang, Y. S. Susceptibility or resistance of praziquantel in human
schistosomiasis: a review. *Parasitol Res* **111**, 1871-1877, doi:10.1007/s00436-012-3151-z
(2012).
5   Assare, R. K. *et al.* Sustaining Control of Schistosomiasis Mansoni in Western Cote
d'Ivoire: Results from a SCORE Study, One Year after Initial Praziquantel Administration.
PLoS Negl Trop Dis **10**, e0004329, doi:10.1371/journal.pntd.0004329 (2016).
6   Stirewalt, M. & Lewis, F. A. Schistosoma mansoni: effect of rotifers on cercarial output,
motility and infectivity. *Int J Parasitol* **11**, 301-308 (1981).
7   Basch, P. F. Schistosomes : development, reproduction, and host relations. *New York :*
Oxford University Press, 248 p (1991).
8   Lewis, F. A., Stirewalt, M. A., Souza, C. P. & Gazzinelli, G. Large-scale laboratory
maintenance of Schistosoma mansoni, with observations on three schistosome/snail
host combinations. *J Parasitol* **72**, 813-829 (1986).
9   Kamigauchi, T. & Yasui, M. Heterocyclic compounds having affinities for serotonin
receptors. WO/2000/059909 (2000).
10   Zhang, Y. L., McArdle, I. W., Hubbard, J. W., Akhmedov, N. G. & Soderberg, B. C. G. Total
synthesis of the tetracyclic indole alkaloid ht-13-A. *Tetrahedron Lett* **57**, 2865-2867,
doi:10.1016/j.tetlet.2016.05.060 (2016).
11   Tao, P., Chen, Z. & Jia, Y. A concise gram-scale synthesis of ht-13-A via a rhodium-
catalyzed intramolecular C-H activation reaction. *Chem Commun (Camb)* **52**, 11300-
11303, doi:10.1039/c6cc05930a (2016).
12   Zhang, Y., Hubbard, J. W., Akhmedov, N. G., Petersen, J. L. & Soderberg, B. C. Total
synthesis of the tetracyclic indole alkaloid ht-13-B. *J Org Chem* **80**, 4783-4790,
doi:10.1021/acs.joc.5b00433 (2015).
13   Haas, W. Physiological analysis of cercarial behavior. *J Parasitol* **78**, 243-255 (1992).
14   Haas, W. Parasitic worms: strategies of host finding, recognition and invasion. *Zoology*
(Jena) **106**, 349-364, doi:10.1078/0944-2006-00125 (2003).
15   Whitfield, P. J., Bartlett, A., Khammo, N. & Clothier, R. H. Age-dependent survival and
infectivity of Schistosoma mansoni cercariae. *Parasitology* **127**, 29-35 (2003).
16   Lewis, F. Schistosomiasis. *Curr Protoc Immunol* **Chapter 19**, Unit 19 11,
doi:10.1002/0471142735.im1901s28 (2001).
17   Ribeiro, P., Gupta, V. & El-Sakkary, N. Biogenic amines and the control of neuromuscular
signaling in schistosomes. *Invertebrate neuroscience : IN* **12**, 13-28, doi:10.1007/s10158-
012-0132-y (2012).

18   Boyle, J. P., Zaide, J. V. & Yoshino, T. P. Schistosoma mansoni: effects of serotonin and
serotonin receptor antagonists on motility and length of primary sporocysts in vitro. *Exp*
Parasitol **94**, 217-226, doi:10.1006/expr.2000.4500 (2000).
19   Boyle, J. P. & Yoshino, T. P. Serotonin-induced muscular activity in Schistosoma mansoni
larval stages: importance of 5-HT transport and role in daughter sporocyst production. *J*
Parasitol **91**, 542-550, doi:10.1645/GE-432R (2005).
20   Chan, J. D. *et al.* A Miniaturized Screen of a Schistosoma mansoni Serotonergic G
Protein-Coupled Receptor Identifies Novel Classes of Parasite-Selective Inhibitors. *PLoS*
Pathog **12**, e1005651, doi:10.1371/journal.ppat.1005651 (2016).
21   Patocka, N., Sharma, N., Rashid, M. & Ribeiro, P. Serotonin signaling in Schistosoma
mansoni: a serotonin-activated G protein-coupled receptor controls parasite movement.
PLoS Pathog **10**, e1003878, doi:10.1371/journal.ppat.1003878 (2014).
22   Chan, J. D. *et al.* The anthelmintic praziquantel is a human serotoninergic G-protein-
coupled receptor ligand. *Nat Commun* **8**, 1910, doi:10.1038/s41467-017-02084-0 (2017).
23   Shen, B. A New Golden Age of Natural Products Drug Discovery. *Cell* **163**, 1297-1300,
doi:10.1016/j.cell.2015.11.031 (2015).
24   Collins, J. J., 3rd, King, R. S., Cogswell, A., Williams, D. L. & Newmark, P. A. An atlas for
Schistosoma mansoni organs and life-cycle stages using cell type-specific markers and
confocal microscopy. *PLoS Negl Trop Dis* **5**, e1009, doi:10.1371/journal.pntd.0001009
(2011).
25   Tillmaand, E. G. *et al.* Peptidomics and Secretomics of the Mammalian Peripheral
Sensory-Motor System. *J Am Soc Mass Spectrom* **26**, 2051-2061, doi:10.1007/s13361-
015-1256-1 (2015).
26   Aerts, J. T. *et al.* Patch clamp electrophysiology and capillary electrophoresis-mass
spectrometry metabolomics for single cell characterization. *Anal Chem* **86**, 3203-3208,
doi:10.1021/ac500168d (2014).
27   Lee, J. E. *et al.* Endogenous peptide discovery of the rat circadian clock: a focused study
of the suprachiasmatic nucleus by ultrahigh performance tandem mass spectrometry.
Mol Cell Proteomics **9**, 285-297, doi:10.1074/mcp.M900362-MCP200 (2010).
28   Collins, J. J., 3rd & Newmark, P. A. It's no fluke: the planarian as a model for
understanding schistosomes. *PLoS Pathog* **9**, e1003396,
doi:10.1371/journal.ppat.1003396 (2013).
355

Acknowledgements

B. glabrata snails were provided by the NIAID Schistosomiasis Resource Center of the

Biomedical Research Institute (Rockville, MD) through NIH-NIAID Contract

HHSN272201700014I for distribution through BEI Resources. We thank: Melanie

Issigonis, Umair Khan, Jayhun Lee, and Tania Rozario for helpful comments on the

manuscript; Tracy Chong and Jayhun Lee for help maintaining the schistosome life

cycle; Björn Söderberg and Yanxing Jia for providing Ht-13-A, -B, and derivatives;

Lingyang Zhu for expert assistance with NMR; Brian Imai for assistance with SPF

purification; as well as Peg Stirewalt, James Leef, Tom Nerad, and Paul Mazzocchi for

their early efforts to help solve this puzzle. This work was supported by the National

Institutes of Health P30 DA018310 and R01 NS031609 to JVS. JG was supported by an

HHMI International Student Research Fellowship. PAN is an investigator of Howard

Hughes Medical Institute. The content is solely the responsibility of the authors and

does not necessarily represent the official views of the National Institutes of Health.

Figure 1 | *R. rotatoria*-conditioned water paralyzes *S. mansoni* cercariae. a, Life

cycle of *S. mansoni* adapted from [28]. **b,c,** Bright-field microscopy images of *R. rotatoria*

and *P. acuticornis*. **d, e,** Maximum intensity projection (5 s, 150 frames) of cercariae

motility after treatment with *R. rotatoria*- or *P. acuticornis*- conditioned water.

375
Figure 2 | SPF is a novel tetracyclic alkaloid. a, Flowchart for SPF purification. **b,** 1st
HPLC plots of *R. rotatoria*- and *P. acuticornis*-conditioned water. Red arrowhead
indicates the peak containing bioactivity. **c,** 2nd HPLC plot of the bioactive fraction (red
arrowhead in **b**). Blue arrowhead indicates the peak containing bioactivity. **d,** MS
showing the dominant signal of m/z 273.1601 from the peak (blue arrowhead). **e,** MS
plots showing this signal (asterisk, m/z 273.1601) was only detected in the fraction
eluting at 24-26 min. **f,** Tandem MS acquired from high-resolution Q-TOF analysis. **g, h,**
NOESY resolved the relative stereochemistry of three chiral centers and narrowed it
down to two possible configurations. Predicted steric energy for the R,S,S (**g**) and
S,R,R (**h**) stereoisomers is shown.

386
387 **Figure 3 | Structure-activity relationships of SPF and related compounds as**
388 **measured by cercarial motility assays. a-g,** Percentage of cercariae (~50) continuing
389 to swim over three minutes after addition of each compound at specified final
390 concentrations. Triplicates were performed. Data are mean ± S.D.

Figure 4 | Treating cercariae with SPF, Ht-13-A or Ht-13-A-pr blocks schistosome infection and alleviates pathology. a, Representative livers (post perfusion) from mice (N=6) infected with drug-treated cercariae. Livers from mice treated with control and lower drug concentrations were darker in color and contained more granulomas (white spots). With higher drug concentrations, livers had normal morphologies with few or no granulomas. 25 μM SPF treatment was not determined (N.D.) due to limited amounts of purified SPF. **b-d,** Numbers of adult worms recovered from infected mice (two experiments for each drug, 6 mice total for each condition). **e-g,** Numbers of schistosome eggs per area (/mm²) from liver sections (4-6 sections per mouse). Data (**b-g**) are mean ± S.D. Statistics: One-way ANOVA, post Dunnett's test.