Leveraging single cell RNA sequencing experiments

to model intra-tumor heterogeneity

Meghan C. Ferrall-Fairbanks (1), Markus Ball (2,3),

Eric Padron (2), and Philipp M. Altrock* (1)

1: Department of Integrated Mathematical Oncology, Moffitt Cancer Center and Research

Institute

2: Department of Malignant Hematology, Moffitt Cancer Center and Research Institute, Tampa,

FL 33612, USA

3: present address: Institute of Pathology, University Hospital Cologne, 50937 Köln, Germany

*Corresponding author: Philipp M. Altrock, philipp.altrock@moffitt.org

Moffitt Cancer Center, 12902 USF Magnolia Drive, SRB 24007, Tampa, FL 33612, USA

Phone: +1 813 745-5897 Fax: +1 813 745- 8357

<u>Running Title</u>: General intra-tumor heterogeneity can be calculated from scRNA-seq data.

<u>Key Words</u>: Acute Myeloid Leukemia. Tumor Heterogeneity. Mathematical Oncology. Diversity.

Single Cell RNA sequencing.

The authors declare no competing financial interests.

ABSTRACT

PURPOSE: Many cancers can be treated with targeted. Almost inevitably, tumors develop resistance to targeted therapy, either from preexistence or by evolving new genotypes and traits. Intra-tumor heterogeneity serves as a reservoir for resistance, which often occurs due to selection of minor cellular sub-clones. On the level of gene expression, the 'clonal' heterogeneity can only be revealed by high-dimensional single cell methods. We propose to use a general diversity index (GDI) to quantify heterogeneity on multiple scales and relate it to disease evolution.

METHODS: We focused on individual patient samples probed with single cell RNA sequencing to describe heterogeneity. We developed a pipeline to analyze single cell data, via sample normalization, clustering and mathematical interpretation using a generalized diversity measure, and exemplify the utility of this platform using single cell data.

RESULTS: We focused on three sources of RNA sequencing data: two healthy bone marrow (BM) samples, two acute myeloid leukemia (AML) patients, each sampled before and after BM transplant (BMT), four samples of pre-sorted lineages, and six lung carcinoma patients with multi-region sampling. While healthy/normal samples scored low in diversity overall, GDI further quantified in which respect these samples differed. While a widely used Shannon diversity index sometimes reveals less differences, GDI exhibits differences in the number of potential key drivers or clonal richness. Comparing pre and post BMT AML samples did not reveal differences in heterogeneity, although they can be very different biologically.

CONCLUSION: GDI can quantify cellular heterogeneity changes across a wide spectrum, even when standard measures, such as the Shannon index, do not. Our approach offers wide applications to quantify heterogeneity across samples and conditions.

INTRODUCTION

In many cancers, there still exists a critical need to understand the mechanisms of therapy resistance evolution. For example, Acute Myeloid Leukemia (AML) is an aggressive hematologic malignancy that is hallmarked by proliferation of immature myeloid cells in the bone marrow and life-threatening ineffective hematopoiesis[1]. AML is the most common adult leukemia, with an incidence of about 20,000 cases yearly and a 5-year survival of only 26%[2,3]. The diagnosis of AML requires greater than 20% of myeloid immature cells (myeloblasts) in the peripheral blood or bone marrow. The median survival of untreated AML is measured in weeks[4]. Several AML targeted therapies have been recently approved, e.g. midostaurin for FLT3 mutated patients and enasidenib for those with mutations in IDH2[5,6]. These mutations occur at rates of 25% (FLT3) and 5% (IDH2) of all AML patients and their targeted therapies are generally well tolerated relative to chemotherapeutic counterparts[7]. However, midostaurin (and even more potent FLT3 inhibitors in clinical trial[8]) does not fully eradicated the disease, leading to refractory or relapsing AML in most patients[9]. The complete response rate for enasidenib in relapse/refractory IDH2 mutated AML is less than 20%. Further refinements in patient selection are required to realize mutationally-directed therapy[5]. Little is known regarding the emerging resistance mechanism and whether targeted therapies (single or combination) against AML alone can ever be successful.

The conventional dogma postulates that therapeutic resistance occurs via the acquisition of mutations that result in clonal evolution. Emerging data suggests that these mutations are either subclonally present or present at frequencies detectable using digital PCR or ultra-deep sequencing technologies at diagnosis or prior to progression. Very low level somatic mutations are also detected in pre-leukemic states[10-13]. Somatic mutations are often present years before

the diagnosis of therapy related myeloid neoplasms[14,15]. Interestingly, these mutations are

commonly associated with disease progression and transformation[16]. The presence of such low-

frequency genetic markers suggests that high levels of intra-tumor heterogeneity (ITH) persist

over long periods of time, and that pre-existing ITH is a primary driver of future therapy

resistance, while variation in transcription over time shapes the disease phenotype. A clinically

relevant summary metric to describe ITH on the transcriptional level has not been developed.

scRNA-sequencing technologies can present a cost-effective method to identify

transcriptomic heterogeneity and directly measure ITH. Proof of concept studies have been

performed in AML, using DROP-seq that yields potentially cost-effective single cell annotations

of thousands of transcripts per cell. In triple negative breast cancer, intercellular heterogeneity

of gene expression programs within tumors is variable and correlates with genomic clonality[17].

A study in Chronic Myeloid Leukemia (CML) demonstrated that scRNA-seq was capable of

segregating patients with discordant responses to targeted tyrosine kinase inhibitor therapy[18].

These data provide rationale to explore ITH in scRNA-seq data, and to determine whether

defined measures of ITH can be predictive of progression, and eventually leveraged to mitigate

progression and relapse.

Our goal is to quantify ITH in cancer such that it has maximal predictive value, in particular

in hematologic malignancies. To this end we here present a platform that uses a generalized

diversity index that characterizes cell population heterogeneity across a spectrum of scales

(orders of diversity)[19]. These scales range from clonal richness (low order of diversity reveals

the number of distinct subpopulations), to more classical measures such as Shannon or

Simpson indices (intermediate order of diversity), to the number of most abundant cell types that

possibly act as the key drivers of heterogeneity before transformation or perturbation by therapy

(high order of diversity).

MATERIALS AND METHODS

We created a computational and modeling approach to develop a robust statistical picture

of the persistent and emerging variability in scRNA-seq data, chiefly based on drop-seq

technologies; the 10x Genomics platform offered a variety of datasets that were linked to disease

and treatment dynamics[20]. We specifically used the datasets of two healthy/control bone marrow

mononuclear cell samples (BMMCs), two individuals with AML BMMCs sampled pre and post

bone marrow transplant (BMT), to develop and test our ITH pipeline. pipelines.

First, we ran publicly available FASTQ-format files (a typical output from a drop -seq

experiment) through the cellranger count pipeline and then through the cellranger aggr pipeline,

to pool the samples together for comparisons during cluster analysis, interrogated through the

10x Genomics Loupe™ Cell Browser (Fig. S1). To test the robustness and valid our diversity

metrics and the ITH pipeline, we extended our analysis to include additional publicly available

datasets for other hematopoietic cell types (CD34+, CD14+, CD19+, and CD4+)[20], as well as six

patient normal-tumor matched lung cancer samples[21], for which we used the same approaches

and pipelines. To calculate summary metrics (outlined in **Fig. 1**), first the transcript expression

data was clustered into groups of cells with similar transcript expressions (cellranger aggr). Next,

we quantified the distance between each of the clusters to determine if clusters separated based

on healthy or disease status (healthy vs AML). A Euclidean distance was calculated between

the mean expression values for each gene, of each cluster, to establish a distance metric (**Fig.**

2).

A: schematic for quantifying diversity of an individual malignant sample

B: schematic for quantifying and comparing diversity across multiple states or individuals

Figure 1. Schematic of our single cell RNA sequencing-based approach. A: Workflow of calculating a generalized diversity index (GDI) for a single sample. After sequencing and library preparation, normalization to reduce the number of false negatives or false positives is applied, for example using 10x-genomics platform, or. Clustering can then be applied (Loupe™ Cell browser, or other platforms, see Supplement), from which diversity can be calculated. **B**: A similar approach can be used when multiple samples are compared. Data normalization and clustering now have to be implemented considering all samples (see Supplement), from which diversity scoring can inform a ranking of intratumor-heterogeneity across samples. Single dots in the tSNE plots represent single cells, which might either be marked according to their cluster classification or according to their sample of origin.

Second, we sought to characterize across-sample differences by calculating the Kolmogorov-Smirnov (KS) distance[22] of the cell count distributions in each cluster, in order to compare samples or pooled samples of the same condition (e.g. disease v. healthy) in terms of the cellular distribution over the identified clusters (**Fig. 3 A-G**).

Third, we calculated an ecological diversity index[23] using the cellular frequencies over clusters, across a range of order of diversity (**Fig. 3H, I**). To assess the robustness of our diversity metric, we performed down sampling of the original datasets and found the relative change in diversity index, across a range of order of diversity, to determine the sensitivity of our diversity metric (**Fig. 4**).

Last, we applied our ITH pipeline and diversity metric to two additional datasets, (1) a hematopoietic cell type datasets comparing CD34+ cells with CD4+, CD14+, and CD19+ cell populations[20], and (2) a lung cancer dataset with tumor-normal matched tissue sites taken from

six different lung cancer patients[21] (**Fig. 5**, Supplement). Further specific details of our methods,

such as cells per sample, are described in the Supplement and is available online (including all

code used to generate our results): https://github.com/mcfefa/scRNAseq.

RESULTS AND DISCUSSION

We established proof of concept that we can generate clinically relevant summary metrics

of ITH by analyzing publicly available scRNA-seq data[20,21]. Within BMMC samples from

diagnosed AML and healthy control groups, we sought to establish how to summarize both inter-

and intra-sample ITH. First, we clustered transcript expression of two healthy individuals, two

AML patients, each sampled twice—once before and once after allogenic bone marrow

transplant.

As a verification we sought to distinguish between healthy and AML samples based on

the mean expression values across cells, across clusters (**Fig. 2A)**. With the 23 clusters

identified (**Fig. 2B, C)**, a network of clusters emerged, displayed as an undirected graph where

the distance between mean UMI counts determines the thickness and length of the edges (**Fig.**

2D). The size of the node was chosen to indicate of the total number of cells in that cluster. We

colored each node according to the condition (health, pre BMT AML, post BMT AML) that was

in the majority (breakdown of actual proportions per cluster shown in **Fig. 2E**). This showed that

indeed the large clusters with mostly healthy cells are most similar in average gene expression,

while the large clusters with mostly AML cells cluster separately (to the right). The post BMT

cells clustered more closely to the healthy than pre BMT AML samples. This supports the idea

that these patients were potentially still transitioning but closer to a healthy phenotype. However,

some AML-dominant clusters still grouped near the healthy/post BMT super cluster. Based on

this "bulk" measure alone, one may not be able to easily distinguish between healthy and

diseased cells. Therefore, other quantifications and metrics to describe the gene expression

differences may better discriminate clinical and stages.

Figure 2. Mean cellular gene expression across clusters within patients can separate disease conditions to some
degree. Here we built a network based on mean differences in overall expression. **A**: The geometric mean of UMI counts across
samples and genes was calculated for each cluster. Then a Euclidean distance was calculated between clusters. Here we used
publicly available scRNA-seq data[20]: two healthy donor BMMCs, two AML patient BMMCs pre bone marrow transplant (preBMT),
and post bone marrow transplant (postBMT). These six samples were then clustered (using 10x-genomics Loupe™ Cell browser;
for alternative clustering methods see Supplement), for which we show the sample-based (**B**) and cluster-based (**C**) tSNE-plots
out of Loupe™ browser. Each dot represents a single cell, which is colored either according to sample of origin or its assigned
cluster. The cluster-based differences in mean gene expression over unique molecular identifier (UMI) counts the gave rise to a
"clustering of the clusters" (**D**). The nodes in the resulting graph were colored based on the dominant cell type from each condition
present in each cluster; gray for healthy, red for AML pre BMT and purple for AML post BMT, the distance between nodes was
chosen inversely proportional to the difference in mean gene expression level, the individual distributions of cells from a specific
condition in each cluster are shown in **E**.

To determine metrics better at discriminating between healthy and disease AML, we

analyzed and summarized inter- and intra-heterogeneity in two different ways. First, we

considered each samples' grouping of cells into clusters of similar gene expression. To this end,

we used the Kolmogorov-Smirnov (KS) distance, which compares two discrete probability mass

functions (the fraction of cells per cluster, **Fig. 3A**). We identified rather similar distributions within

the same condition, and rather different distributions between conditions, with post BMT being

a notable exception (**Fig. 3B-D**). The KS distance between the two healthy samples was 0.139,

between the two pre BMT AML samples it was 0.174, but between the two post BMT samples it

was 0.551. Since we had clustered all six samples together, we could also compare them pooled

by condition, which revealed that indeed conditions distribute differently across the identified

cellular subpopulations in high-dimensional gene expression space (**Fig. 3E-G**).

Second, we calculated a general diversity index (GDI) for each condition (healthy, pre

BMT AML, post BMT). The mathematical definition of GDI, ^qD, is shown in **Fig. 3H**. We

established segregation of the different clinical conditions according to this ecology-based

diversity index[23]. The pre BMT AML samples had consistently higher diversity index compared

to the healthy sample and this held true across the entire order of diversity range, q (**Fig. 3I**).

Interestingly, on this level the post BMT samples also scored unanimously higher in GDI. This

could indicate that post BMT settings may require a certain amount of time after transplant to

evolve toward a healthy spectrum of intra-leukemic diversity. Also, in a comparison of the

individual samples within each condition (**Fig. 4, A-C**), the post BMT samples were most different

from each other.

To interrogate the robustness of GDI further and to establish confidence in the metric, we

down sampled the dataset and then re-clustered and calculated the ^qD spectrum (**Fig. 4D-I**).

A: definition of the Kolmogorov-Smirnov (KS) distance between two discreet distributions

B: healthy, KS-dist=0.139

C: AML pre BMT, KS-dist=0.174

D: AML post BMT, KS-dist=0.551

E: KS-dist=0.372

F: KS-dist=0.360

G: KS-dist=0.369

H: mathematical definition of cluster diversity

number of clusters in the data set = n
frequency of each cluster = $p_1, p_2, ..., p_n$
order of diversity = q

$$^q D = \left(\sum_{i=1}^{n} p_i^q \right)^{\frac{1}{1-q}}$$

I

Figure 3. Cluster-based diversity scoring reveals strong differences between healthy individuals and cancer patients.
In our analysis, using data from Zheng, et. al.[20], we evaluated our ability to score significant differences in "cluster diversity" across healthy and AML samples pre and post bone marrow transplant (BMT). As a first indicator of between-sample or between-condition differences we used the Kolmogorov-Smirnov (KS) distance for discrete probability mass functions (**A**). Little difference was found in the KS distance within condition-differences (**B**, **C**), except in post BMT samples (**D**). Between condition-differences were larger when comparing pooled samples across conditions (**E**, **F**, **G**). We calculated a general diversity index (GDI) qD (**H**) to quantify diversity across "orders of diversity" q. For all orders of diversity measure, AML patients (pre and post BMT) had a higher diversity index compared to healthy individuals (**I**, 2 samples per condition), suggesting GDI can be used as a metric for stratification.

During down sampling, we analyzed each sample individual by randomly removing 50% of the

cells, then calculating the number of clusters identified for that individual's transcript expression,

and finally calculating the diversity index for specific q values of interest, including $q=10^{-2}$, $q=10^{-1}$,

$q=1$ (which relates to the Shannon index), $q=2$ (which defines the inverse of the Simpson

index), $q=10$, and $q=10^2$. The distributions shown were obtained from 1000 runs of independent

down sampling. Intriguingly, these distributions showed that with removing half of the cells, the

diversity scores did not change more than a unit or two in either direction. When comparing to

the diversity spectrum shown in **Fig. 3I**, this suggests that if healthy diversity spectrum were

shifted up by two units (10% of the maximum) and the AML samples diversity spectrum were

shifted down by two units, there would still be visible separation between the healthy and AML

conditions.

To further validate our metric, we implemented on our approach with two other datasets.

One dataset described different hematopoietic cellular subtypes, CD34+, CD4+, CD14+, and

CD19+[20]. CD34+ is a hematopoietic progenitor cell marker and represent a polyclonal population

that includes many different subtypes (hematopoietic stem cells, multipotent progenitor cells,

common myeloid progenitor cells, common lymphoid progenitor cells, megakaryocytes erythroid

progenitor cells and granulocytes macrophage progenitor cells) all of which express CD34[24-26].

The CD34+ polyclonal population contrasted the CD4+, CD14+, and CD19+ populations, which

represent more homogenous cellular populations (helper T-cells, monocytes, and B-cells,

respectively). This clonality pattern was recovered by GDI (**Fig. 5A**), where the CD34+

population had a considerable higher diversity score across the spectrum. Interestingly, lower

values of q seem to separate differentiated cells or robustly.

Figure 4. AML patients have consistently higher diversity compared to healthy individuals. Individual diversity spectrums were reported for healthy (**A**), AML pre BMT (**B**), and AML post BMT (**C**) samples (each line is one from one sample). Cell-gene matrices were down sampled to 50% of the cells 1000 times, and qD scores were calculated (using the full pipeline, see Supplement) for specific values of q=0.01, 0.1, 1, 2, 10, 100. The distributions of relative qD changes for healthy (**D**, **G**), AML (**E**, **H**), and postBMT AML (**F**, **I**) samples showed that generally, lower q values lead to less change in measured diversity. Across all cases the diversity score did not change by more than two units (relative change is measured by dividing the entire distribution by the distribution mean). For sample sizes, see Supplement. BMT: Bone marrow transplant.

Finally, we quantified ITH using lung cancer scRNA-seq samples[21]. We analyzed six

different patients, each with up to three different tumor sites (core, middle, and edge) and a

patient-matched adjacent normal lung tissue sample. Using our GDI metric, we see that the

diversity spectrum of the normal lung tissue was much lower than any of the tumor site diversity

scores (pooled conditions, **Fig. 5B**). Interestingly here, more clear separation of conditions was

achieved at high orders of diversity q, indicating differences in the number of driver clones at

different sites within the tumors. These additional datasets further support the ability of quantified

diversity metric to discriminated between healthy and diseases states, which can be applied in

a clinical setting.

Figure 5. Higher diversity indicates higher clonality in in normal tissues and solid tumors. (**A**) Additional available data
from Zheng, et. al.[20] for CD34+ cells, CD4+ helper T-cells, CD14+ monocytes and CD19+ B-cells were run through pipeline S
(see Supplement), and the continuum of diversity was calculated for each population. The naturally polyclonal population
(CD34+) shows the highest diversity score. Each of the other differentiated immune cell compartments are more homogeneous
across orders of diversity. (**B**) In solid tumors, location matters: normal-tumor matched lung carcinoma samples were obtained
from publicly available data for six lung cancer patients[21] (individual patients: see Supplement). The diversity metric across q
shows an increase in diversity within tumors across different tumor locations.

SUMMARY AND CONCLUSIONS

Single-cell RNA sequencing efforts have greatly helped to uncover population structures

and mapping to specific cellular population patterns[27]. Although these methods can also

elucidate tumorigenesis[28], immune-profiles[29], and detect and track genomic profiles of

clones[30,31], the overall utility of scRNA-seq for cancer progression survival metrics has been elusive[32]. Here, we demonstrated the potential utility of two scRNA-seq based scores of cellular heterogeneities, using a generalized diversity index that may be elevated in disease. Remarkably, using previously published data without further processing, this quantification of intra-tumor heterogeneity was able to accurately distinguish AML from healthy individuals, as well as from post-transplant conditions. These data suggest that ITH can be estimated by diversity-based summary statistics, and that these summary statistics can be leveraged to predict clinical outcomes.

Our work here aimed at optimizing and identifying a clinically relevant summary index for ITH in the context of AML, for which targeted single-cell genome sequencing was also able to sensitively uncovered complex clonal evolution[33]. We anticipate that our ILH metric will be prognostic for leukemia-free survival (LFS) and potentially overall survival (OS), even after correction for known clinical prognostic variables. We have also shown how this metric can also be used to effectively describe heterogeneity in other malignancies, including solid tumors such as lung carcinomas.

From a clinical perspective in terms of tumor heterogeneity and emergence of resistance clones during targeted therapy[34,35], we expect our metric can discriminate patients clinically. We hypothesize that these heterogeneity metrics would be elevated *independently* (at least *a priori*) in potentially highly resistant patients. The advantage of a more general metric used here is that it allows us to look across many orders of diversity, and potentially pick a desired range of heterogeneity quantification. For example, one might be interested especially in lower values of q, where a higher diversity score may indicate an individual (sample) more at risk of resistance evolution, as it shows high standing variation. On the other hand, differences at high values of q

point to key differences in the number of important driver clones, which might uncover distinct

vulnerabilities that can be targeted in combination or adaptively.

Diversity measures have long received attention in ecology and evolution[19,36]. We here

approached to measure diversity, and thus tumor heterogeneity, using a general definition of

non-spatial diversity[37], in form of the quantity ^qD (**Fig. 3H-I**). This approach considers all possible

orders of diversity q, but also allows to compare disease stages according to a specific diversity

index (fixed choice of q), which emerge as special cases of ^qD. The species (clonal) richness of

a sample is given by q=0. The Shannon index (log scale) can be found when q approaches 1.

The Simpson index, which approximated the probability that any two cells are identical emerges

from the case q=2. Both Shannon and Simpson index have been used in mathematical and

statistical models of cancer evolutionary dynamics to quantify tumor heterogeneity as it

potentially changes during tumor growth, with disease progression, or during treatment[38-40].

Shannon entropy-based statistics have also been used to quantify single cell heterogeneity, to

deliver insights into emerging or disappearing clones during transitions between clinical

conditions[41].

Single cell RNA sequencing experiments give a snapshot of the cell population state on

the level of gene expression, and it can characterize how individual cell's transcriptomes

compare to the bulk. In contrast to mass cytometry, drop-seq is fast and extremely high-

throughput. Other single cell technologies, like flow cytometry can be used to generate single

cell data for a relatively small subset of potential markers that distinguish between normal and

disease, and it requires that the researcher/clinician knows what these markers are in advance.

Among a variety of outcomes that may be distinguished by our metric, one can be the study for

segregating samples based on disease severity, for which additional follow-up knowledge will

be needed.

Further extending our results to the potential impact in a clinical setting in leukemias,

intraleukemic heterogeneity (ILH) is a known reservoir for tumor resistance and clinical

refractoriness to targeted therapies[9]. For current targeted therapies to treat AML, clinical

responses have been modest. Specific means to change ILH can be of particular appeal in such

cases, as they might help render tumors less aggressive and hinder their ability to rapidly evolve

resistance. In particular, hypomethylating agents (HMA, cytosine analogs that irreversibly bind

to DNA methyltransferase, an enzyme required for methylation of CpG-rich DNA) have the

potential to diminish ILH[42]. Transcriptome changes upon treatment with HMA therapy have not

been analyzed at single cell resolution. Our analyses here provide a quantitative basis to

understand and reliably track these changes.

Clear separation of diversity metrics by condition, as we show it, might not be expected

in general. A weakness of our approach is that it does not consider any meaning of the

associated phenotypes or genotypes. Therefore, as it stands, our method cannot be transferred

to improve the predictive power of existing bulk signatures. Hence, existing survival data is

unlikely to be useful to prove that GDI is predictive of survival, and novel databased that uniquely

connect high throughput single cell experiments with clinical outcomes are needed.

Once the appropriate cohorts are established, however, changes in an individual's

diversity score could indicate unique features of disease progression. In the context of adaptive

therapy[43,44], which aims at tumor burden control rather than difficult tumor eradication, it might

be critical to identify the appropriate scale of diversity that best predicts outcomes. One could

speculate that there is an optimal window of diversity that should be maintained—very low

diversity could indicate fast disease progression and very high diversity could mean that the

tumor could adapt to the treatment schedule too quickly. The concept we introduced here is

sufficiently flexible in its ability to quantify optimally predictive windows of diversity that should

be maintained during adaptive therapy.

AUTHORSHIP CONTRIBUTIONS

P.M.A., E.P. conceived the study. M.C.F., M.B. and P.M.A. performed computational

experiments and statistical analyses. M.C.F., and P.M.A. performed mathematical modeling.

M.C.F., M.B., E.P. and P.M.A. analyzed the data. M.C.F., M.B., E.P. and P.M.A. wrote the

manuscript. P.M.A. and E.P. supervised the project.

339
340

REFERENCES

1. Perl AE: The role of targeted therapy in the management of patients with AML. Hematology Am Soc Hematol Educ Program 2017:54-65, 2017

2. Percival ME, Tao L, Medeiros BC, et al: Improvements in the early death rate among 9380 patients with acute myeloid leukemia after initial therapy: A SEER database analysis. Cancer 121:2004-12, 2015

3. Medeiros BC, Satram-Hoang S, Hurst D, et al: Big data analysis of treatment patterns and outcomes among elderly acute myeloid leukemia patients in the United States. Annals of Hematology 94:1127-1138, 2015

4. Arber DA, Orazi A, Hasserjian R, et al: The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. Blood 127:2391-2405, 2016

5. Stein EM, DiNardo CD, Pollyea DA, et al: Enasidenib in mutant IDH2 relapsed or refractory acute myeloid leukemia. Blood 130:722-731, 2017

6. Stone RM, Mandrekar SJ, Sanford BL, et al: Midostaurin plus Chemotherapy for Acute Myeloid Leukemia with a FLT3 Mutation. N Engl J Med 377:454-464, 2017

7. Papaemmanuil E, Gerstung M, Bullinger L, et al: Genomic Classification and Prognosis in Acute Myeloid Leukemia. N Engl J Med 374:2209-2221, 2016

8. Cooper TM, Cassar J, Eckroth E, et al: A Phase I Study of Quizartinib Combined with Chemotherapy in Relapsed Childhood Leukemia: A Therapeutic Advances in Childhood Leukemia & Lymphoma (TACL) Study. Clinical Cancer Research 22:4014-4022, 2016

9.   Smith CC, Paguirigan A, Jeschke GR, et al: Heterogeneous resistance to
quizartinib in acute myeloid leukemia revealed by single-cell analysis. Blood 130:48-58, 2017

10.   Paguirigan AL, Smith J, Meshinchi S, et al: Single-cell genotyping demonstrates
complex clonal diversity in acute myeloid leukemia. Sci Transl Med 7:281re2, 2015

11.   Corces-Zimmerman MR, Hong WJ, Weissman IL, et al: Preleukemic mutations in
human acute myeloid leukemia affect epigenetic regulators and persist in remission. Proc Natl
Acad Sci U S A 111:2548-53, 2014

12.   Ding L, Ley TJ, Larson DE, et al: Clonal evolution in relapsed acute myeloid
leukaemia revealed by whole-genome sequencing. Nature 481:506-10, 2012

13.   Padron E, Yoder S, Kunigal S, et al: ETV6 and signaling gene mutations are
associated with secondary transformation of myelodysplastic syndromes to chronic
myelomonocytic leukemia. Blood 123:3675-7, 2014

14.   Wong TN, Ramsingh G, Young AL, et al: Role of TP53 mutations in the origin and
evolution of therapy-related acute myeloid leukaemia. Nature 518:552-555, 2015

15.   Gillis NK, Ball M, Zhang Q, et al: Clonal haemopoiesis and therapy-related myeloid
malignancies in elderly patients: a proof-of-concept, case-control study. Lancet Oncol 18:112-
121, 2017

16.   Zuffa E, Franchini E, Papayannidis C, et al: Revealing very small FLT3 ITD
mutated clones by ultra-deep sequencing analysis has important clinical implications in AML
patients. Oncotarget 6:31284-94, 2015

17.   Karaayvaz M, Cristea S, Gillespie SM, et al: Unravelling subclonal heterogeneity
and aggressive disease states in TNBC through single-cell RNA-seq. Nature Communications
9:3588, 2018

18.   Giustacchini A, Thongjuea S, Barkas N, et al: Single-cell transcriptomics uncovers
distinct molecular signatures of stem cells in chronic myeloid leukemia. Nat Med 23:692-702,
2017

19.   Hill MO: Diversity and evenness: a unifying notation and its consequences.
Ecology 54:427-432, 1973

20.   Zheng GX, Terry JM, Belgrader P, et al: Massively parallel digital transcriptional
profiling of single cells. Nat Commun 8:14049, 2017

21.   Lambrechts D, Wauters E, Boeckx B, et al: Phenotype molding of stromal cells in
the lung tumor microenvironment. Nat Med 24:1277-1289, 2018

22.   Carruth J, Tygert M, Ward R: A comparison of the discrete Kolmogorov-Smirnov
statistic and the Euclidean distance. arxiv.org:arXiv:1206.6367, 2012

23.   Lou J: Entropy and diversity. Oikos 113:363-375, 2006

24.   Greaves MF, Brown J, Molgaard HV, et al: Molecular features of CD34: a
hemopoietic progenitor cell-associated molecule. Leukemia 6 Suppl 1:31-6, 1992

25.   Krause DS, Fackler MJ, Civin CI, et al: CD34: structure, biology, and clinical utility.
Blood 87:1-13, 1996

26.   Zhang Y, Gao S, Xia J, et al: Hematopoietic Hierarchy - An Updated Roadmap.
Trends Cell Biol, 2018

27.   Trapnell C, Cacchiarelli D, Grimsby J, et al: The dynamics and regulators of cell
fate decisions are revealed by pseudotemporal ordering of single cells. Nat Biotechnol 32:381-
386, 2014

28.   Ellsworth DL, Blackburn HL, Shriver CD, et al: Single-cell sequencing and
tumorigenesis: improved understanding of tumor evolution and metastasis. Clin Transl Med
6:15, 2017

29.   Schelker M, Feau S, Du J, et al: Estimation of immune cell content in tumour tissue
using single-cell RNA-seq data. Nat Commun 8:2032, 2017

30.   Pellegrino M, Sciambi A, Treusch S, et al: High-throughput single-cell DNA
sequencing of acute myeloid leukemia tumors with droplet microfluidics. Genome Res 28:1345-
1352, 2018

31.   Navin NE: The first five years of single-cell cancer genomics and beyond. Genome
Res 25:1499-507, 2015

32.   Wang L, Livak KJ, Wu CJ: High-dimension single-cell analysis applied to cancer.
Mol Aspects Med 59:70-84, 2018

33.   Pellegrino M, Sciambi A, Treusch S, et al: High-throughput single-cell DNA
sequencing of acute myeloid leukemia tumors with droplet microfluidics. Genome research 28:1-
8, 2018

34.   Gatenby RA, Silva AS, Gillies RJ, et al: Adaptive therapy. Cancer Res 69:4894-
903, 2009

35.   Staňková K, Brown JS, Dalton WS, et al: Optimizing cancer treatment using game
theory: A review. JAMA Oncology, 2018

36.   MacArthur RH: Patterns of species diversity. . Biological Reviews 40:510-533,
1965

37.   Tuomisto H: A consistent terminology for quantifying species diversity? Yes, it
does exist. Oecologia 164:853-60, 2010

38.   Almendro V, Cheng YK, Randles A, et al: Inference of tumor evolution during
chemotherapy by computational modeling and in situ analysis of genetic and phenotypic cellular
diversity. Cell Rep 6:514-27, 2014

39.   Marusyk A, Tabassum DP, Altrock PM, et al: Non-cell-autonomous driving of
tumour growth supports sub-clonal heterogeneity. Nature 514:54-8, 2014

40.   Altrock PM, Liu LL, Michor F: The mathematics of cancer: integrating quantitative
models. Nat Rev Cancer 15:730-45, 2015

41.   Li J, Smalley I, Schell MJ, et al: SinCHet: a MATLAB toolbox for single cell
heterogeneity analysis in cancer. Bioinformatics 33:2951-2953, 2017

42.   Kaminskas E, Farrell AT, Wang YC, et al: FDA drug approval summary: azacitidine
(5-azacytidine, Vidaza) for injectable suspension. Oncologist 10:176-82, 2005

43.   Zhang J, Cunningham JJ, Brown JS, et al: Integrating evolutionary dynamics into
treatment of metastatic castrate-resistant prostate cancer. Nat Commun 8:1816, 2017

44.   Gallaher JA, Enriquez-Navas PM, Luddy KA, et al: Spatial Heterogeneity and
Evolutionary Dynamics Modulate Time to Recurrence in Continuous and Adaptive Cancer
Therapies. Cancer Res 78:2127-2139, 2018
446
447

SUPPLEMENTAL MATERIAL

Leveraging single cell RNA sequencing experiments

to model intra-tumor heterogeneity

Meghan C. Ferrall-Fairbanks (1), Markus Ball (2,3),

Eric Padron (2), and Philipp M. Altrock* (1)

1: Department of Integrated Mathematical Oncology, Moffitt Cancer Center and Research Institute

2: Department of Malignant Hematology, Moffitt Cancer Center and Research Institute, Tampa, FL 33612, USA

3: present address: Institute of Pathology, University Hospital Cologne, 50937 Köln, Germany

*Corresponding author: Philipp M. Altrock, philipp.altrock@moffitt.org

Moffitt Cancer Center, 12902 USF Magnolia Drive, SRB 24007, Tampa, FL 33612, USA

Phone: +1 813 745-5897 Fax: +1 813 745- 8357

SUPPLEMENTAL MATERIALS AND METHODS

DROPLET-BASED scRNA$_{SEQ}$ SAMPLES

The 10X Genomics platform offered a variety of datasets[1] that were used to create our

pipeline for quantifying intraleukemic heterogeneity. The major focus of our analysis and pipeline

development was of Healthy/Control and AML patient bone marrow mononulear cells (BMMCs).

The Healthy Controls 1 and 2 BMMCs had been sequenced on Illumina Hiseq 2500 Rapid Run

V2 with 90-135 thousand reads per cell and 2000-2400 cells detected. The AML027 and

AML035 pre-transplant BMMCs had been sequenced on Illumnia NextSeq 500 High Output with

16.6-58 thousand reads per cell and 3500-3900 cells detected. The AML027 and AML035 post-

transplant BMMCs had also been sequenced on Illumina NextSeq 500 High Output with 41-51

thousand reads per cell and 900-3900 cells detected. Additional datasets[1] for CD34+, CD14+,

CD19+, and CD4+ (hematopoietic cell type set) cells as well as patient normal-tumor matched

lung cancer samples[2] available in ArrayExpress under accessions E-MTAB-6653 and E-MTAB-

6149 were used to test the robustness of the pipeline. The CD34+ dataset was CD34+ cells

enriched from peripheral blood mononuclear cells (PBMCs), sequenced on an Illumnia NextSeq

500 High Output with 24.7 thousand reads per cell with 9000 cells detected. The CD14+ dataset

was enriched from PBMCs, sequenced on Illumnia NextSeq 500 Output with 100 thousand

reads per cell with 2600 cells detected. The CD19+ dataset was enriched from PBMCs,

sequenced on Illumnia NextSeq 500 Output with 25 thousand reads per cell with 10000 cells

detected. The CD4+ dataset was enriched from PBMCs, sequenced on Illumnia NextSeq 500

High Output with 21 thousand reads per cell with 11000 cells detected. All lung tissue samples

were prepared as single-cell suspensions, sequenced on Illumina HiSeq4000, and aimed for an

estimated 4000 cells per library[2]. There were six patients analyzed, each patient had 4 distinct

tissues locations sampled: an adjuacent normal lung sample (normal), a sample from the tumor

core (core), a sample from the tumor margin (edge), and a sample between the tumor core and

margin/edge (middle).

INITIAL DATA PROCESSING

First, the all samples were run through the 10x Genomics cellranger count pipeline for

transcriptome alignment, calling individual cell barcodes, estimating multiplet rates, and finally

clustering of normalization corrected FPKM (fragments per kilobase million; expression level)

values (**Fig. S1**, pipeline X). In our analysis of the Healthy versus AML high-throughput scRNA-

seq, we used cellranger aggr to pool together the individual dataset results, so we could compare

intra-tumor heterogeneity (ITH) between samples with the same clustering. Using a graph-based

clustering algorithm, 23 different clusters were identified based on clustering cells by expression

similarity (shown in **Fig. 2B-C**). The clustering was performed by forming a graph with cells as

vertices and edges indicating pairs of cells that are sufficiently similar (implemented by

CellRanger). The CellRanger graph-based clustering with Louvain Modularity Optimization[3] was

used to partition this graph into clusters of similar cells. For analysis with the hematopoietic cell

set and the lung cancer samples the cell/gene matrices produced by cellranger count pipeline

were then loaded into R using the Seurat package[4] (designed for QC, analysis, and exploration

of single cell RNA-seq data; **Fig. S1**, pipelines M and S). With Seurat, cells were removed that

had either fewer than 200 UMIs and greater than 6000 UMIs and more than 10% mitochondrial

DNA[2], then scaled and normalized before clustering using graph-based clustering.

QUANTIFYING SUMMARY DIVERSITY METRICS

After the data was clustered, we then sought to quantify whether the clustering could

disentangle healthy and AML samples by average gene expression. To approach within-sample

differences in overall gene expression, we computed Euclidean distance matrices of the mean

expression values in each cluster, to establish a distance metric of cluster differences and a

subsequent diversity of distances across samples (standard deviation, ANOVA) (**Fig. 2A**). For

each cluster, the geometric mean of each unique molecular identifier (UMI) across all genes

were computed, then the Euclidean distance was computed between clusters. This was plotted

as a graph where each node represents each clusters identified in the leukemic dataset (**Fig.**

2D). The size of the node indicates the total number of cells in that cluster and the color identifies

the major species (AML, Healthy, or postBMT) present in that cluster. The distribution of cells

per cluster (**Fig. 2E**) was also included to show which other conditions were also present in each

cluster and how the number of cells compared to the major species used to dictate the color of

the nodes in graph describing the connectivity of the clusters in **Fig. 2D**.

Next, we sought to characterize across-sample differences by calculating the

Kolmogorov-Smirnov (KS) distance[5] of the cell count distributions in each cluster (**Fig. 3A**). The

KS distance is a non-parametric measure that is calculated as the supremum of paired

differences of two empirical probability mass functions[5]. Last, we calculated a continuum of the

ecological diversity index[6] based on the individual cell frequencies in each of the clusters

identified by the graph-based clustering (**Fig. 3H**), across all clusters, which can be written as:

$$^{q}D = \left(\sum_{l=1}^{n} p_i^q \right)^{\frac{1}{1-q}}$$

where n is the number of clusters in the data set, p_n is the frequency of each cluster, and q is

the order of diversity. q is a hyperparameter that would be optimized in a clinical setting. The two

distributions for AML were joined together and the two distributions of the healthy indiviuals were

joined together and plotted across orders of diversity (**Fig. 4A-C**). The same technique was used

to group the lung cancer samples based on location as reported in (**Fig. 5B**).

DOWNSAMPLING

We sought to test the robustness of our metrics by downsampling the number of cells and

re-calculating the number of clusters identified as well as changes in key diversity scores. This

downsampling was preformed removing increments of 10% of the data at a time and the results

were reported as the summary statistics of 1000 runs at each downsampling percentile (**Fig.**

S2). The downsampling results were also used to quantify confidence in the diversity index

spectrum and was calculated for the following orders of diversity: 10^{-2}, 10^{-1}, 10^0, 2, 10^1, 10^2. The

relative change in diversity score was reported for the downsampling results with 50% the initial

amount of cells in **Fig. 4D-I**. Relative change in the diversity score was calculated by subtracting

the mean of the diversity score from all 1000 and reported as the distributions of scores around

that mean score.

CODE AVAILABILITY

The code used in the pipelines described in **Fig. S1** were uploaded to a GitHub repository

(https://github.com/mcfefa/scRNAseq).   These   pipelines   were   implemented   using   data   run

through at least cellranger counts, and then post-processed with CellRanger Loupe Browser,

Mathematica, and R. Code available includes:

- Mathematica code and R scripts and data described to implement all pipelines (described

  in **Fig. S1**)

- Mathematica code for calculating the geometric mean of UMIs and Euclidean distances

  between clusters (for **Fig. 2**)

- R code to draw the graph of the clusters (**Fig. 2D**)

•   Mathematica code for calculating the KS distance and diversity spectrum across all

clusters for the leukemic dataset (**Fig. 3**)

•   R code used to downsample datasets, then cluster and calculate diversity indices (**Fig. 4**

and **S2**)

•   R code for calculating the diversity spectrum for the hematopioteic subtype and matched

lung carcinoma datasets (**Fig. 5**)

Supplemental Figure 1. Workflow diagram for the scRNAseq quantifying intraleukemia

heterogeneity for an example comparing two samples. Cells were processed through the

10X Genomics Chromium and ultimately sequenced, producing a raw base call (BCL) file, which

was demultiplexed for each flowcell directory and converted to a FASTQ file using cellranger

mkfastq. Then cellranger count was run separately for each library to align, filter, and count

barcodes and UMIs. These instances were then aggregated into a single instance using

cellranger aggr to normalize runs to the same sequencing depth and recompute analysis on

combined dataset. Following along pipeline X, from the aggregated analysis, the Loupe Browser

contains the clustered data, which were exported and run through a Mathematic script to

calculate the diversity metrics. This pipeline was used to generate Figs. 3 and 4. This pipeline

was expanded to give the user increased flexibility by using the gene-barcode matrices output

by cellranger and then processing the data in R using the Seurat package, which allows users

to set the filtering criteria (M) and allows data to be combined without running through the

cellranger aggr pipeline (S) as well as both new pipelines allow users to implement additional

clustering algorithms. Pipeline S was used to generate Fig. 5 from gene-barcode matrices

available from additional data from Zheng, et. al.[18].

Supplemental Figure 2. AML populations converge to consistent number of clusters that

describe the population sooner than healthy populations. graph-based clustering. Each

original population's gene-barcode matrix obtain from publicly available data from Zheng, et.

al.[18] was then downsampled to contain 10% to 90% of the cells originally in the matrix. This data

was loaded into R using the Seurat package, downsampled, and then clustered to determine

how sensitive the cluster metrics was to the starting number of cells. Downsampling was

performed 1000 times per cutoff and the violin plots show the distribution of clusters identified.

The red-dashed lined indicates the number of clusters identified with no cells removed from the

dataset. Gray numbers indicate the number of cells present in each downsampling condition.

AML populations (**A**, **B**) showed quickest convergence to number of clusters after about 1500

cells were present in clustering. Healthy populations (**C**, **D**) showed that as more cells were

added, an additional cluster could be found. AML post-bone marrow transplant populations (**E**,

F) behaved more similar to healthy than AML populations.

Supplemental Figure 3: Individual lung cancer patient diversity scores show same trends as

aggregated clustering results with tumor samples have greater diversity scores than their

matched normal counterpart. Individual patients were run through pipeline S to aggregate

normal, core, middle, and edge tissue samples.

REFERENCES

1. Zheng GX, Terry JM, Belgrader P, et al: Massively parallel digital transcriptional profiling of single cells. Nat Commun 8:14049, 2017

2. Lambrechts D, Wauters E, Boeckx B, et al: Phenotype molding of stromal cells in the lung tumor microenvironment. Nat Med 24:1277-1289, 2018

3. 10xGenomics: https://support.10xgenomics.com/single-cell-gene-expression/software/pipelines/latest/algorithms/overview, 2018

4. Butler A, Hoffman P, Smibert P, et al: Integrating single-cell transcriptomic data across different conditions, technologies, and species. Nat Biotechnol 36:411-420, 2018

5. Carruth J, Tygert M, Ward R: A comparison of the discrete Kolmogorov-Smirnov statistic and the Euclidean distance. arxiv.org:arXiv:1206.6367, 2012

6. Lou J: Entropy and diversity. Oikos 113:363-375, 2006