Title: Dual mechanism of TRKB activation by anandamide through CB1 and TRPV1 receptors

Authors: Diniz CRAF[1], Biojone C[2,3], Joca SRL[2,5], Rantamäki T[4], Castrén E[3], Guimarães FS[1], Casarotto PC*[1,3]

Affiliations:

1: Department of Pharmacology, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil
2: Department of Physics and Chemistry, Ribeirão Preto School of Pharmaceutical Sciences, University of São Paulo, Ribeirão Preto, São Paulo, Brazil
3: Neuroscience Center - HILIFE, University of Helsinki, Helsinki, Finland
4: Division of Pharmacology and Pharmacotherapeutics, Faculty of Pharmacy, University of Helsinki, Helsinki, Finland
5: Department of Clinical Medicine, Translational Neuropsychiatric Unit, University of Aarhus, Denmark

*correspondence: plinio.casarotto@helsinki.fi
Plinio Casarotto, PhD
Neuroscience Center - HILIFE
Haartmaninkatu 8, B233b, PO Box 63
University of Helsinki, Finland

Abstract

Administration of anandamide (AEA) or 2-arachidonoylglycerol (2AG) induces CB1 coupling and activation of TRKB receptors, regulating the neuronal migration and maturation in the developing cortex. However, at higher concentrations AEA also engages vanilloid receptor - TRPV1, usually with opposed consequences on behavior. Using primary cell cultures from the cortex of rat embryos (E18) we determined the effects of AEA on phosphorylated TRKB (pTRK). We observed that AEA (at 100 and 200nM) induced a significant increase in pTRK levels. Such effect at AEA 100nM was blocked by pretreatment with the CB1 antagonist AM251 (200nM) and, at the higher concentration of 200nM, by the TRPV1 antagonist capsazepine (20nM). CB1 and TRKB co-immunoprecipitated with TRPV1 in cortical cultures. Increase in pTRK levels, after AEA 200nM treatment, was positively correlated with the levels of BDNF (brain-derived neurotrophic factor) in the culture medium. Interestingly, the effect of AEA at 200nM on pTRK in cortical cells was blocked by TRKB.Fc (a soluble form of TRKB able to bind BDNF), while the effects of BDNF were not prevented by AM251. The increase in pTRK levels by the TRPV1 agonist capsaicin (200nM) was also inhibited by pretreatment with TRKB.Fc, suggesting a mechanism dependent on BDNF release. Using the marble-burying test (MBT) in mice, we observed that the local administration of ACEA (CB1 agonist) into the PFC was sufficient to reduce the burying behavior, while CPS (capsaicin, a TRPV1 agonist) or BDNF exerted the opposite effect by increasing the number of buried marbles. In addition, both ACEA and CPS effects were blocked by previous administration of k252a into PFC. The effect of systemically injected CB1 agonist WIN55,212-2 was blocked by previous administration of k252a, an antagonist of TRK receptors. We also observed a partial colocalization of CB1/TRPV1/TRKB in the prefrontal cortex, and the co-localization of TRPV1 in CaMK2+ cells. Taken together, our data indicate that cannabinoids engage a coordinated activation of TRKB, via CB1 and TRPV1. Endocannabinoids, acting upon CB1 and TRPV1, could regulate the TRKB-dependent plasticity in both pre- and postsynaptic compartments in a coordinated manner and based on BDNF concentration.

Keywords: BDNF; CB1; TRPV1; marble-burying

Introduction

Repetitive behaviors are part of the normal background of animal's learning [1,2]. These stereotypic motor procedures are regulated by the cortical-striatal-thalamo-cortical (CSTC) circuitry, a series of multiple reverberatory loops, with the prefrontal cortex, striatum and thalamic nuclei playing a core role as integrative hubs [3]. Changes in the CSTC can trigger abnormal repetitive behaviors. These aberrant behaviors can be related to several human neuropsychiatric conditions such as Huntington's disease and obsessive-compulsive disorder (OCD). In a simplified version, the CSTC is divided into two major pathways: direct and indirect [4]. The direct pathway comprises the cortical projections to *globus pallidus* (*pars internalis*) and *substantia nigra pars reticulata,* while the indirect pathway engages a more complex set of relay structures, involving the *globus pallidus pars externalis* and subthalamic nucleus – (for review see Canales et al. 2000 [5]). The resultant activity between these two pathways regulates the output of the basal ganglia in favor of one of the two possible effects: increase in the repetitive movements (favored by the direct pathway activity) or inhibition of such programs, a consequence of activation of the indirect pathway. Although highly simplified, this model of the CSTC circuit provides a useful framework for understanding circuit physiology and putative dysfunctions [4].

Multiple neurotransmitters/neuromodulators systems act in coordination to regulate the balance in the CSTC circuitry. Among them, endocannabinoids play a central role regulating not only glutamatergic, but also GABAergic, serotonergic, and dopaminergic transmission (for review see López-Moreno et al. 2008 [6]). Briefly, the activation of CB1 receptors by AEA (N-arachidonoylethanolamine) or 2-AG (2-arachidonoylglycerol), produced as 'on-demand retrograde messengers', decreases the activity of presynaptic neurons via Gi/0 and modulation of calcium and potassium channels [for review see Chevaleyre et al. 2006 [7]]. Due to these effects, endocannabinoids are putatively able to regulate excessive neurotransmission in the CSTC system. Besides, the synthesis and release of endocannabinoids usually is triggered by depolarization-induced calcium influx, as well as by activated phospholipase-C-beta following activation of Gq-protein coupled receptors [8,9].

Endocannabinoids can also act on TRPV1 receptors. These receptors increase calcium influx, facilitating a rapid depolarization of the neuronal cells [10,11]. In preclinical anxiety models, high doses of AEA are usually ineffective or cause anxiogenic rather than anxiolytic effects. This bell-shaped dose-effect curve has been associated with TRPV1 activation and is reversed by pretreatment with antagonists of these receptors [10] (for review see Moreira and Wotjak, 2010; Aguiar et al. 2014 [12,13]). Accordingly, the anxiolytic effect of capsaicin, an agonist of TRPV1 receptors, observed after intracerebral administration was attributed to a desensitization of the channels [14].

CB1 receptors are highly expressed in the anterior cingulate cortex, striatum and *substantia nigra* [15,16], major hubs of CSTC circuitry. TRPV1 has a less broad expression when compared to CB1 [17,18]. However, these two receptors are colocalized in the periaqueductal grey matter (PAG) and prefrontal cortex [10,19], playing opposite functional roles.

CB1 receptors can couple to and transactivate tyrosine kinase receptors [20], including the tropomyosin-related kinase B - TRKB [21]. TRKB is the receptor of brain-derived neurotrophic factor (BDNF), a member of the neurotrophic family that includes NGF, NT-3 and NT4/5 [22]. The activation of TRKB is a crucial factor in the regulation of brain plasticity [23]. Animals lacking CB1 receptors (CB1.KO) display behavioral despair and a reduction in BDNF levels in hippocampus [24], suggesting an interplay between these two systems during development and in adulthood. CB1.KO animals also exhibit impaired extinction of aversive memories [25], a feature usually facilitated by the activation of TRKB following antidepressant treatment, and prevented by reduced BDNF expression [26].

Thus, the initial aim of the present study was to investigate if CB1 and TRKB systems interact in a mouse model of anxiety/repetitive behavior, trying to replicate the *in vitro* mechanism proposed by Berghuis and colleagues [21]. Following the results obtained in the present study, we then speculate on the participation of TRPV1 in the AEA effects on TRKB.

Material and methods

Animals. Male and female C57BL6/j were used (weight 25-30g, 12-18 weeks old at the start of the experiments), with free access to food and water, except during experimental procedures. All procedures were approved by the University of São Paulo (protocol: 146/2009) and for the experiments conducted at the University of Helsinki (protocol: ESAVI/10300/04.10.07/2016), in accordance with international guidelines for animal experimentation.

Drugs. Anandamide (AEA, #1017, Tocris, USA), arachidonyl-2'-chloroethylamide (ACEA, #1319, Tocris, USA), WIN 55,212-2 mesylate (WIN, #1038, Tocris, USA), capsaicin (CPS, #0462, Tocris, USA), AM251 (#1117, Tocris, USA), TRKB.Fc (#688-TK-100, R&D systems, USA), capsazepine (CZP, #0464, Tocris, USA), k252a (#05288, Sigma-Aldrich, USA) and recombinant-human brain-derived neurotrophic factor (rhBDNF, #450-02, Peprotech, USA) were used. AEA and ACEA were dissolved in Tocrisolve (#1684, Tocris, USA). WIN and AM251 were suspended in 2%Tween 80, and k252a was diluted in DMSO 0.2%, and dissolved in sterile saline for ip injections. CSP, CZP or AM251 were diluted in DMSO (at 1000x the final concentration used in the assays) for *in vitro* experiments. BDNF and TRKB.Fc were dissolved in phosphate buffered saline (PBS).

Primary cultures. Cortex of E18 rat embryos were dissected and the cells plated in poly-L-lysine coated wells at a 0.5×10^6 cells/ml density in Neurobasal medium [27]. The cells were left undisturbed, except for medium change, for 8 days before drug treatment and sample collection.

Apparatus and procedure

Marble burying test (MBT). Test was performed using a square box (30x20x12cm) with 5cm sawdust layer covered floor. Twelve green clear glass marbles (1.5cm in diameter) were evenly spaced over the floor. One hour before test the animals were left undisturbed in the experimental room. Mice were initially placed on the box´s center containing marbles. Twenty-five min later they were taken from the box, and the number of buried marbles was counted. Criteria for buried marbles included only those with at least two-thirds under sawdust [28].

Locomotor activity. Male mice were submitted to a 25min session in a circular arena (40cm in diameter with a 50cm high plexiglass wall). Each session was videotaped, and the total activity (distance travelled by the mice in meters) was analyzed with the help of the Anymaze software (version 4.5, Stoelting, Ireland). The apparatus was previously cleaned with 70% alcohol solution before each animal assessment.

Stereotaxic surgery. Briefly, male mice were anesthetized with tribromoethanol and fixed in a stereotaxic frame. Next, stainless steel guide cannulas (0.7mm OD, 9mm length) were implanted into the prelimbic ventromedial prefrontal cortex (PL; coordinates: AP= -2.8mm from bregma, L= 1.5mm, DV= 2mm; lateral inclination of 20°) and then attached to skull bone with steel screws and acrylic cement. Obstruction was prevented with a stylet inside guide cannula. About five days after surgical procedure, bilateral intracerebral infusions in a volume of 200nl were performed with a dental needle (0.3mm OD) during 1min. Hamilton micro-syringe (Sigma-Aldrich) and an infusion pump (KD Scientific) were used. Following the behavioral tests, mice were anesthetized with chloral hydrate and 200nL of methylene blue was infused 1.5mm deeper than the guide cannulas end. Brains were withdrawn and the injection sites verified. Data obtained from injections outside the aimed area were discarded. Coordinates for stereotaxic surgery and the histological location of the injections represented in figure 5 are based on the Paxinos' mouse brain atlas [29].

Sample collection, immunoprecipitation, and western blotting. Cell cultures at DIV8 were washed with ice-cold PBS and lysed using the following buffer [20mM Tris-HCl; 137mM NaCl; 10% glycerol;

0.05M NaF; 1% NP-40; 0.05mM Na3VO4] containing a cocktail of protease and phosphatase inhibitors (Sigma-Aldrich, #P2714 and #P0044, respectively). The samples were centrifuged (16000xg) at 4°C for 15min and the supernatant was collected and stored -80°C until use.

For the immunoprecipitation procedure, samples were incubated overnight at 4°C with anti-TRKB (#AF1494, R&D systems, United Kingdom), anti-CB1 (#CB1-Rb-Af380, Frontier Institute, Japan) or anti-TRPV1 (#sc12498, Santa Cruz, USA) at 1ug of antibody / 500ug of total protein ratio or no antibody (negative control). Following incubation with 30ul of Protein G-sepharose (#101242, Life Technologies, USA) the samples were centrifuged (1000xg/3min) and the supernatant discarded. Precipitated was washed 3 times with lysis buffer and stored at -80°C until use. Samples were then added to the sample buffer, heated to 80°C for 15 min, centrifuged at 1000 xg for 3min and the supernatant resolved by SDS-PAGE, and finally transferred to PVDF membranes. The membranes were blocked with 3%BSA in TBST buffer (20mM Tris-HCl; 150mM NaCl; 0.1% Tween-20; pH 7.6) and incubated with anti-TRPV1 for 48h at 4°C. The detection was performed using HRP-conjugated rabbit anti-goat IgG (1:5000, #2768, Santa Cruz, USA), followed by incubation with ECL and exposure to a CCD camera.

Determination of pTRK levels. Levels of phosphorylated TRK were determined by ELISA [30]. Briefly, 120ug of total protein content from each sample was incubated in white 96-well plates previously coated with anti-TRKB (rabbit, 1:500; Santa Cruz, #sc-8316; in carbonate buffer; pH 9.6) and blocked with 3% BSA in PBST buffer supplemented with Na3VO4 [0.1% Tween-80 in PBS; pH 7.6]. Following incubation with biotin-conjugated anti-phosphotyrosine (mouse, 1:2000; AbD Serotec, #MCA2472B) and streptavidin-HRP conjugate (1:10000; Thermo Scientific, #21126) the amount of luminescence was determined by incubation with ECL using a plate reader (Varioskan Flash, Thermo-Fisher). The signal obtained from each sample was discounted of blank and normalized by the average of the control groups, thus expressed as percentage from control.

Determination of BDNF levels. levels of BDNF in culture medium was determined by ELISA [31]. Briefly, samples from cell culture medium were incubated in precoated 96-well plates with anti-BDNF (provided by Roche, 0.8ng/ml) for 48h. Following overnight incubation with the secondary antibody against BDNF (supplied by Roche, 2.2ng/ml) the peroxidase activity was detected with 3,3'-5,5'-tetramethylbenzidine (TMB). The reaction was stopped with 1M H2SO4 and the color intensity read at 450nm (Varioskan Flash, Thermo-Fisher). The BDNF concentration in each sample was determined based on a concentration curve built with the rhBDNF (6.25-500pg/ml) in Neurobasal

medium. The absorbance obtained from each sample was discounted of blank and normalized by the average of the control groups, thus expressed as percentage from control.

Co-localization of CB1, TRPV1 and TRKB in the prefrontal cortex. Three female C57BL6/j mice were deeply anesthetized and perfused with saline followed by 4% PFA in PBS. The brains were removed from skull and post-fixated in 4% PFA at 4ºC for 48h, and then cryoprotected in 30% sucrose. Coronal sections (40um thick) were obtained in a vibratome (Leica VT1000-S). For immunostaining, the slices were washed in PBS, blocked for 30min at room temperature (3% BSA, 10% donkey normal serum in PBST containing 0.04% sodium azide), followed by incubation for 72h at 4ºC with the primary antibodies, as follows: anti-CB1 (1:1000), anti-TRPV1 (1:100, #sc12498 or #sc398417) and anti-TRKB (1:1000). Another set of samples were labelled for CB1 or TRPV1, and CaMK2 (1:100, Abcam, UK, #AB22609). Additionally, a negative control staining without the primary antibody was included. After washing, sections were incubated 1h at room temperature with 1:200 fluorophore-conjugated secondary antibodies (AlexaFluor donkey anti-goat - 647nm #A21447; anti-rabbit - 568nm #A11077; anti-mouse - 488nm #A21202; Life Technologies, USA). Prelimbic prefrontal cortex was identified based on mouse brain atlas of Paxinos and Franklin [32]. Images were acquired at Biomedicum Imaging Unit at the University of Helsinki using a laser scanning confocal microscope (Zeiss LSM 780) [33] equipped with 63x objective lens (Plan-Apochromat 63/1.40 oil DIC M27) and imaging Software ZEN (Zeiss). A z-stack consisting of at least 8 consecutive images was obtained from each brain section, and images were later analyzed in ImageJ Software (NIH).

Experimental design

Dose-response effects of AEA, CPS, and BDNF on TRKB phosphorylation and co-immunoprecipitation. Cortical cultures (DIV8) were treated with AEA (CB1/TRPV1 agonist; 0, 2.5, 10, 50, 100 or 200nM), CPS (TRPV1 agonist; 0, 10, 100 or 200nM) or BDNF (0, 5, 20 or 50ng/ml, as positive control). Thirty min after drug administration the level of pTRK was determined by ELISA as described (n= 4-6/group). Next, lysates from cortical cells were pre-incubated with anti-CB1, anti-TRPV1 or anti-TRKB, and the levels of TRPV1 determined in the precipitated by western-blotting.

CB1/TRPV1-dependent mechanism of AEA-activated TRKB. Cortical cells received AM251 (CB1 antagonist; 0 or 200nM) followed 10min later by AEA (0 or 100nM). Another batch of cells received AM251 (0 or 200nM), but now followed 10min later by BDNF (0 or 20ng/ml). Next, cells were incubated with CZP (TRPV1 antagonist; 0 or 20nM) followed 10min later by AEA (0, 100 or 200nM). Concerning

each experimental set above described, cells were lysed 30min after the last drug administration for pTRK ELISA assay (n=5-6/group).

TRPV1-dependent mechanism of BDNF release by AEA. In another experimental set, cortical cells received TRKB.Fc (200ng/ml) followed, 10min later, by CPS (0 or 200nM), AEA (0 or 200nM) or BDNF (0 or 20ng/ml, used only as positive control but not part of the statistical analysis). Cells were lysed 30min after the last drug administration for pTRK ELISA assay (n=5-6/group). Finally, for determination of BDNF release, the medium of cortical cultures was reduced from 500ul to 200ul 2h before the drug administration. This procedure facilitates the detection of released BDNF. Following the incubation with AEA (0 or 200nM/30min), the medium was collected for BDNF ELISA and the cells lysed for pTRK (n=5/group).

Effect of systemic and PFC local drug administration. Male mice received an ip injection of K252a (TRK antagonist; 0 or 80ug/kg) followed, 10min later, by another ip injection of WIN (CB1 agonist; 0 or 1mg/kg). MBT test was performed 30min after the last injection (n=8/group). Next, mice with implanted cannulae aimed at the prefrontal cortex received a bilateral injection of ACEA (CB1 agonist; 0.05pmol), ACEA+k252a (10pmol), CPS (5nmol), CPS+k252a, BDNF (200pg) or vehicle (DMSO) and were submitted to the MBT 20min later (n=5-15/group). An independent group of animals received the same pharmacological treatments and was subjected to the locomotor activity test (n=5-9/group).

Statistical analysis. Data was analyzed by one- or two-way ANOVA with drug treatments as factors followed by Fisher's LSD *post-hoc* test when appropriate. Correlation analysis was performed using Spearman's test. Values of $p<0.05$ were considered significant. All the data used in the statistical analysis, as well as background controls for immunostaining, in the present study is publicly available at figshare under CC-BY license [34].

Results

In vitro **dose-response effects of AEA, CPS, and BDNF on TRKB phosphorylation**

As depicted in figure 1a, one-way ANOVA indicates a significant effect of AEA treatment in pTRK levels [$F_{(5,18)}= 16.53$, $p<0.0001$]. *Post-hoc* analysis indicates a difference between the vehicle- and AEA 100nM- and 200nM-treated groups (Fisher, $p=0.0003$ and <0.0001, respectively). Administration of CPS also increased the levels of pTRK [$F_{(3,19)}= 9.28$, $p=0.0005$]. *Post-hoc* analysis indicates a difference between vehicle- and CPS 200nM-treated groups (Fisher, $p<0.0001$; figure 1b). Treatment with BDNF,

used as a positive control, was also able to increase the levels of pTRK [$F_{(3,16)}$= 134.40, $p < 0.0001$], with difference between vehicle- and 20ng/ml- or 50ng/ml-treated groups (Fisher, $p < 0.0001$ for both; figure 1c). As seen in figure 1d, TRPV1 (VR1: band at 100kDa) was detected in samples from primary cultures after precipitation of CB1 or TRKB, the ratio of CB1- and TRKB-coupled TRPV1 compared to total TRPV1 were: 27.0% and 13.2%, respectively.

Figure 1. *In vitro* **dose response effects of anandamide (AEA), capsaicin (CPS) and BDNF on TRKB phosphorylation.** Cortical cultures were treated with (a) AEA 0, 2.5, 10, 50, 100 or 200nM, (b) CPS 0, 10, 100 or 200nM and (c) BDNF 0, 5, 20 or 50ng/ml and 30min after the pTRK levels of cell homogenate was measured (n=4-6/group). Data is expressed as mean/SEM. Dashed lines represent the non-linear regression curve for the dose-response effect. *$p < 0.05$ compared to veh (0) group. (d) coIP of CB1 or TRKB and TRPV1 (VR1) in primary cultures of cortical cells.

CB1/TRPV1-dependent mechanism of AEA-activated TRKB

As observed in figure 2a-b, two-way ANOVA indicates a significant interaction between AM251 and AEA effects [$F_{(1,20)}$= 16.01, $p = 0.0007$], but not between AM251 and BDNF [$F_{(1,16)}$= 0.85, $p = 0.37$] on pTRK levels. *Post-hoc* analysis indicates a difference between veh/AEA100nM- and AM251/AEA100nM-treated groups (Fisher, $p < 0.0001$). In figure 2c, two-way ANOVA indicates a significant interaction between AEA and CZP on pTRK levels [$F_{(2,30)}$= 17.04, $p < 0.0001$]. *Post-hoc* test indicates a significant difference between the veh/AEA200nM- and CZP/AEA200nM-treated groups, but not between veh/AEA100nM- and CZP/AEA100nM-treated groups (Fisher, $p < 0.0001$).

TRPV1-dependent mechanism of AEA-released BDNF

As seen in figure 2d, two-way ANOVA indicates a significant interaction between TRKB.Fc and AEA treatment [$F_{(2,18)}$= 15.01, $p < 0.0001$]. *Post-hoc* analysis indicates difference between veh/veh- and veh/AEA-treated groups (Fisher, $p < 0.0001$), but only a marginal difference between TRKB.Fc/veh- and

TRKB.Fc/AEA-treated groups (Fisher, p=0.065). Figure 2e, two-way ANOVA depicts an interaction between CPS and TRKB.Fc treatments [$F_{(1,20)}$= 9.28, p=0.0064], and *post-hoc* test indicates a difference between veh/veh- and veh/CPS-treated groups (Fisher, p=0.0001). Treatment with AEA at 200nM concentration was able to increase pTRK cell levels [t(8)= 3.75, p=0.0065; data not shown] inasmuch as increased BDNF levels in the medium [t(8)= 4.76, p=0.0014; data not shown]. Both parameters were positively correlated [Spearman's r= 0.77, p=0.011; figure 2f].

Figure 2. CB1/TRPV1-dependent mechanism of AEA-induced activation of TRKB. Cortical cultures received AM251 (0 or 200nM) followed 10min later by (a) AEA (0 or 100nM) or (b) BDNF (0 or 20ng/ml). Cortical cells were incubated with (c) CZP (0 or 20nM) followed 10min later by AEA (0, 100 or 200nM). Cortical cultures received (d) TRKB.Fc (0 or 200ng/ml) followed 10min later by (d) AEA (200nM), BDNF (20ng/ml); or (e) by CPS (200nM). (f) cortical cultures were incubated with AEA (0 or 200nM) and 30min later the medium was collected for BDNF measure and cells were lysed for pTRK measure (n=5/group). Cells were lysed and homogenized 30min after the last drug administration for pTRK measure (n=5-6/group). Data is expressed as mean/SEM. *p<0.05 compared to veh/veh group. #p<0.05 compared to the respective veh/drug group.

Effect of systemic and PFC local drug administration

As seen in figure 3a, two-way ANOVA indicates a significant interaction between drug treatments [$F_{(1,28)}$= 4.52; p=0.0425]. *Post-hoc* analysis depicted the pretreatment with k252a preventing the WIN-induced decrease in the number of buried marbles (veh/WIN vs k252a/WIN, Fisher, p=0.0125). In figure 3b, one-way ANOVA indicates a significant effect of drugs treatment [$F_{(5,53)}$= 10.32; p<0.0001]. *Post-hoc* test indicates that ACEA were able to decrease (Fisher, p=0.0095), while CPS (5nmol) and BDNF (200pg) increased (Fisher, p=0.0253 and =0.0003, respectively) the number of buried marbles, such effect of ACEA and CPS were counteracted by previous administration of k252a (Fisher, p=0.63 and =0.37, respectively). No effect of drugs was observed on locomotor activity, as sees in figure 3c [$F_{(4,23)}$= 0.11; p=0.97]. Figure 3d depicts the injection sites in the PL-PFC of mice according to Paxinos' atlas [29].

Figure 3. Effect of systemic and PFC local drug administration. (a) Mice received ip injection of K252a (80ug/kg) followed, 10min later, by ip injection of WIN (1mg/kg) and were submitted to the MBT 30min after the last drug injection (n=8/group). (b) Mice implanted with a guide cannula aiming the prefrontal cortex received bilateral infusion of ACEA 0.05pmol, ACEA+k252a, CPS 5nmol, CPS+k252a or BDNF 200pg, and were submitted to the MBT 20min after drug administration (n=5-15/group). Data is expressed as mean/SEM. (c) An independent cohort of animals received the same pharmacological treatment and was submitted to the locomotor activity test (n=5-9/group). Data are expressed as mean ± SEM of distance traveled in meters. (d) Infusions are represented by points in the Paxinos´s diagrams (2012). The number of points are fewer than the number of animals used due to overlap. *p<0.05 compared to veh-treated (ctrl) group.

Immunolabeling of CB1, TRPV1, and CaMK2 or TRKB

As observed in figure 4a-c, the majority of TRPV1+ cells are also CaMK2+, while CB1 positive labels surround CaMK2+ cells, figure 4d-f. Figure 5, depicts the triple labeling of TRPV, CB1 and TRKB. Again, the majority of TRPV1+ cells were also positive for TRKB.

Figure 4. Co-staining of (a-c) CaMK2 (from mouse - ms) and TRPV1 (from goat - gt), and (d-f) CaMK2 and CB1 (from rabbit - rb) in prelimbic area of prefrontal cortex in mice. a) cells positive to CaMK2 (green); b) cells positive to TRPV1 (blue); c) cells merged to both CaMK2 and TRPV1. TRPV1+/CaMK2- cells are also observed, as indicated by white arrows. d) cells positive to CaMK2 (green); e) cells positive to CB1 (red); cells merged to both CaMK2 and CB1. scale bar: 20um.

Figure 5. Co-staining of (a) TRKB (from goat - gt), (b) CB1 and (c) TRPV1 (from mouse - ms) in prelimbic area of prefrontal cortex in mice. a) cells positive to TRKB (blue); b) cells positive to CB1 (red); c) cells positive to TRPV1 (green); d) cells merged to TRKB, CB1 and TRPV1. TRKB-/TRPV1+ are also observed, as indicated by white arrows.

Discussion

In the present study we described a coordinated activation of TRKB (pTRK) by AEA in a CB1- and TRPV1-dependent manner. Lower doses of AEA administered to cultured cortical cells were able to increase pTRK levels through CB1 activation. On the other hand, our pharmacological approach has led us to suggest that BDNF effect on pTRK levels does not engage endocannabinoids production, at least in the timeframe used in the present study. Higher doses of AEA increased the pTRK levels in a TRPV1-dependent manner, an effect mimicked by capsaicin (CPS). Both AEA (at 200nM) and CPS required BDNF release to increase pTRK levels. However, CB1 and TRPV1 agonists, infused into PL-PFC produced opposite effects on burying behavior. CB1 agonist ACEA decreased, and TRPV1 agonist CPS increased the number of buried marbles. Further, BDNF infused into PL-PFC mimicked the behavioral effect of TRPV1 agonist. Data from total distance traveled indicates that the effects observed in the burying behavior could not be confounded by changes in locomotion.

Endocannabinoids (eCBs) exert multiple effects both at cellular and behavioral levels [35]. These molecules usually act as 'on-demand' retrograde messengers, following the activation of glutamatergic or gabaergic receptors in the postsynaptic terminal. Activation of CB1 receptors by eCBs increases the presynaptic terminal permeability to potassium while reducing calcium influx. As a result, the presynaptic terminal is hyperpolarized and the neurotransmission blunted [36]. On the other hand, higher concentrations of eCB also act on vanilloid receptor TRPV1 [37]. Given that TRPV1 is a ligand-gated calcium channel, its activation by eCB increases the calcium permeability in the presynaptic terminal, therefore counteracting the activation of CB1 [37]. Indeed, the behavioral data available indicates that in most cases eCB exhibit a bell-shaped curve, with lower concentrations acting mainly through CB1 and higher quantities exerting its effects through TRPV1 [19].

Previous studies indicate that, upon activation by AEA or 2AG, CB1 can couple to TRKB, and triggers its activity, underlying the maturation of cortical neurons [21]. The observed CB1:TRKB coupling did not involve the increase of BDNF production, both at translation or transcription level. In line with these findings, we observed that AEA administration in cultured cortical cells of rat embryo was indeed able to increase the levels of pTRK in a CB1-dependent manner. To verify other possible aspects of CB1:TRKB interaction, we tested if activation of TRKB by BDNF requires previous activation of CB1 or it is, at least partially, dependent on eCB production. Thereby, we found that TRKB activation by BDNF in our experimental conditions (cultivated embryonic cortical cells) did not demand any level of CB1 activation. On the other hand, in preparations of cerebellar granule neurons, the activation of TRKB by BDNF increased the neuronal sensitivity to eCB by increasing CB1 and decreasing the expression of the degrading enzyme monoacylglycerol lipase [38]. Other studies, using preparations of somatosensory cortex,

suggest that GABAergic synapses are inhibited by a TRKB/PLCgamma1-dependent production of eCB from the postsynaptic pyramidal cell [39–41]. Finally, the probability of glutamate release in somatosensory cortex is decreased by BDNF/TRKB-induced eCB release [42]; and eCBs are suggested to be produced and released to act also on midbrain dopamine neurons via previous activation of TRKB/PLC-gamma1 signaling [43]. Moreover, in organotypic hippocampal slice cultures, BDNF is a key downstream intermediary in CB1 protection against kainic acid-dependent excitotoxicity [44]. Thus, as previously described, TRKB activity modulates eCBs levels and the sensitivity of CB1 in response to eCB binding. Then, both eCBs and BDNF can modulate each other but, concerning the direct activation of TRKB, a 'hierarchical' relationship between these two receptors is established.

The higher dose of AEA was more effective to increase the levels of pTRK through a TRPV1-dependent mechanism. Confirming this observation, CPS administration also increased pTRK levels in the same system. Our data suggest that AEA acts through TRPV1 receptors to release BDNF and increase pTRK levels. Interestingly, the levels of pTRK were decreased from 325% to 200% above the control levels (same level reached with AEA 100nM) in cells treated with AEA 200nM following the TRPV1 antagonist capsazepine or TRKB.Fc. Therefore, even if AEA has a higher affinity for CB1 than for TRPV1, the related efficacy appears to be inverted in favor of TRPV1. Thus, it is plausible that lower doses of eCB could induce modest activation of TRKB via CB1 coupling whereas higher doses, acting in both receptors CB1 and TRPV1, would boost BDNF release and pTRK levels. Accordingly, after treatment with AEA 200nM a positive correlation was observed between levels of BDNF released in culture medium and cellular levels of pTRK, thus corroborating the proposal that pTRK levels is increased by BDNF released, possibly in a TRPV1-dependent manner.

Next, we used the marble burying test (MBT) to further investigate *in vivo* the relevancy of BDNF/TRK modulation by CB1 and TRPV1. This test was initially proposed as a model of neophobic anxiety [28], being sensitive to a variety of antidepressants, anxiolytics and antipsychotics [45,46]. However, this proposal has been questioned by recent studies – for more detailed information see Thomas et al. 2009 [47]. The burying behavior seems to reflect natural motor programs presented in the CSTC circuitry, rather than a novelty-induced anxious state. In accordance with prior data described by our group [48], mice systemically treated with the CB1 agonist WIN display a reduced burying behavior. Such effect of WIN was counteracted by the previous administration of AM251 (a CB1 antagonist); and, in the present study, by the TRK blocker k252a. Therefore, as suggested by *in vitro* results, *in vivo* data supports BDNF/TRKB signaling as a downstream trigger of CB1-induced behavioral changes.

In this scenario, given that the PFC is one of the major hubs of CSTC regulation of the burying behavior, we tested the effects of CB1, TRKB and TRPV1 activation in this structure. Our data indicates

that ACEA administration into PFC decreases the number of buried marbles. Surprisingly, the infusion of BDNF into PL-PFC increased the number of marbles buried. A similar facilitatory effect was also observed with CPS, and both ACEA and CPS effects were blocked by previous administration of k252a. Opposed behavioral effects are usually observed when, after increasing concentrations of AEA, there is a target displacement from preferential CB1 to TRPV1 signaling. For example, whereas CB1 is usually responsible for the anxiolytic-like effect of low AEA doses, the activation of TRPV1 signaling by higher doses is associated with null or even anxiogenic-like effect [10,19]. Thus, our behavioral results suggest that in the PL-PFC BDNF and TRPV1 play a similar role.

Taken together, our results indicate that in the PL-PFC, upon activation CB1 couples to TRKB in a BDNF-independent manner (as suggested by Berghuis and colleagues) to induce a modest increase of pTRK levels and an anticompulsive-like effect. TRPV1 activation by high AEA concentrations further boosts BDNF release counteracting the CB1-mediated effect. Exogenous BDNF administration, by producing high local concentrations of this growth factor, would mimic the effects observed after TRPV1 activation. One possible explanation for these results would be the existence of two distinct populations of neurons in the PL-PFC that colocalize TRKB:CB1 and TRKB:TRPV1, modulating the direct and indirect pathways of the CSTC circuitry, respectively. The second possibility is that the CB1- and TRPV1-dependent activation of TRKB occurs in pre and postsynaptic neurons, respectively. In this scenario, CB1 and TRPV1 would play opposed effects in the same pathway of the CSTC. These two scenarios, however, are not mutually exclusive. In agreement with this proposal, we observed a low level, although non-negligible, of co-immunoprecipitation between TRPV1 and TRKB or CB1. TRPV1, CB1, and TRKB immunolabeling in the PL-PFC slices indicate a partial overlap of these three receptors. The majority of the cells in the PL-PFC slices were double-positive to TRPV1/CaMK2, while CB1 surrounds CaMK2+ cells.

In addition, both CB1 and TRKB are mutually expressed in cortical layers 2/3 and 5 [42]. CB1 receptors are also found in different neuronal populations. For instance, Marsicano and Lutz [49] observed that nearly all CB1 positive cells in multiple cortical layers are also GAD65- or CCK-positive, suggesting a GABAergic phenotype. On the other hand, TRPV1 is present in hippocampal pyramidal neurons [50]. However, some studies point also to the occurrence of co-localization between both CB1 and TRPV1 receptors, thus indicating a more complex interaction [10,19].

In conclusion, our *in vitro* and *in vivo* data suggests the presence of a complex interaction between CB1, TRPV1, and TRKB culminating in a final balance between BDNF-dependent and independent signaling to induce or inhibit burying behavior, respectively. Regulation of the amount of BDNF released in the synaptic cleft would be an additional mechanism, beyond inherent receptor

properties, by which CB1 and TRPV1 activation usually produce opposite effects. Therefore, endocannabinoids acting upon CB1 and TRPV1, could regulate the TRKB-dependent plasticity in both pre- and postsynaptic compartments of different neuronal populations in a coordinated manner. More studies, however, are required to investigate this possibility.

Figure 6. Graphical abstract for the interaction between CB1, TRPV1 and TRKB regulating burying behavior. The activation of CB1 by endocannabinoids (eCB) leads to TRKB coupling, activation and decrease in burying, putatively in GABAergic cells (dashed blue). On the other hand, the activation of TRPV1 receptors by eCB triggers the release of BDNF and activation of TRKB, possibly in pyramidal (dashed red) and/or neighbor cells (dashed green) leading to increase in burying.

Acknowledgments

We thank Flávia Salata (USP), Senem Merve Fred (UH), and Sulo Kolehmainen (UH) for their technical support.

Funding and disclosure

This work was supported by CNPq (#471382/2011-6) and FAPESP (#2011/02746-4, #2013/01029-2, #2013/02549-0 and #2018/04250-5) for experiments conducted in Brazil, and by ERC (#322742) for experiments conducted in Finland. The authors declare no conflict of interest.

References

1. Langen, M., Durston, S., Kas, M. J. H., van Engeland, H. & Staal, W. G. The neurobiology of repetitive behavior: …and men. *Neurosci. Biobehav. Rev.* **35,** 356–365 (2011).
2. Langen, M., Kas, M. J. H., Staal, W. G., van Engeland, H. & Durston, S. The neurobiology of repetitive behavior: of mice…. *Neurosci. Biobehav. Rev.* **35,** 345–355 (2011).
3. Alexander, G. E., DeLong, M. R. & Strick, P. L. Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu. Rev. Neurosci.* **9,** 357–381 (1986).
4. Casarotto, P. C., Gomes, F. V. & Guimarães, F. S. Cannabinoids and obsessive-compulsive disorder. in *Cannabinoids in Neurologic and Mental Disease* (ed. Fattore, L.) 365–387 (Academic Press, 2015).
5. Canales, J. J. & Graybiel, A. M. A measure of striatal function predicts motor stereotypy. *Nat. Neurosci.* **3,** 377–383 (2000).
6. López-Moreno, J. A., González-Cuevas, G., Moreno, G. & Navarro, M. The pharmacology of the endocannabinoid system: functional and structural interactions with other neurotransmitter systems and their repercussions in behavioral addiction. *Addict. Biol.* **13,** 160–187 (2008).
7. Chevaleyre, V., Takahashi, K. A. & Castillo, P. E. Endocannabinoid-mediated synaptic plasticity in the CNS. *Annu. Rev. Neurosci.* **29,** 37–76 (2006).
8. Hashimotodani, Y., Ohno-Shosaku, T., Watanabe, M. & Kano, M. Roles of phospholipase Cβ and NMDA receptor in activity-dependent endocannabinoid release. *J. Physiol.* **584,** 373–380 (2007).
9. Castillo, P. E., Younts, T. J., Chávez, A. E. & Hashimotodani, Y. Endocannabinoid signaling and synaptic function. *Neuron* **76,** 70–81 (2012).
10. Casarotto, P. C. *et al.* Opposing roles for cannabinoid receptor type-1 (CB_1) and transient receptor potential vanilloid type-1 channel (TRPV1) on the modulation of panic-like responses in rats. *Neuropsychopharmacology* **37,** 478–486 (2012).
11. Moreira, F. A., Aguiar, D. C., Terzian, A. L. B., Guimarães, F. S. & Wotjak, C. T. Cannabinoid type 1 receptors and transient receptor potential vanilloid type 1 channels in fear and anxiety—two sides of one coin? *Neuroscience* **204,** 186–192 (2012).
12. Moreira, F. A. & Wotjak, C. T. Cannabinoids and anxiety. *Curr. Top. Behav. Neurosci.* **2,** 429–450 (2010).
13. Aguiar, D. C. *et al.* Modulation of defensive behavior by Transient Receptor Potential Vanilloid Type-1 (TRPV1) channels. *Neurosci. Biobehav. Rev.* **46 Pt 3,** 418–428 (2014).
14. Terzian, A. L. B., Aguiar, D. C., Guimarães, F. S. & Moreira, F. A. Modulation of anxiety-like behaviour by Transient Receptor Potential Vanilloid Type 1 (TRPV1) channels located in the dorsolateral periaqueductal gray. *Eur. Neuropsychopharmacol.* **19,** 188–195 (2009).
15. Harkany, T. *et al.* The emerging functions of endocannabinoid signaling during CNS development. *Trends Pharmacol. Sci.* **28,** 83–92 (2007).
16. Díaz-Alonso, J., Guzmán, M. & Galve-Roperh, I. Endocannabinoids via CB1 receptors act as neurogenic niche cues during cortical development. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **367,** 3229–3241 (2012).
17. Tóth, A. *et al.* Expression and distribution of vanilloid receptor 1 (TRPV1) in the adult rat brain. *Brain Res. Mol. Brain Res.* **135,** 162–168 (2005).
18. Menigoz, A. & Boudes, M. The expression pattern of TRPV1 in brain. *J. Neurosci.* **31,** 13025–13027 (2011).
19. Fogaça, M. V., Aguiar, D. C., Moreira, F. A. & Guimarães, F. S. The endocannabinoid and endovanilloid systems interact in the rat prelimbic medial prefrontal cortex to control anxiety-like behavior. *Neuropharmacology* **63,** 202–210 (2012).
20. Dalton, G. D. & Howlett, A. C. Cannabinoid CB1 receptors transactivate multiple receptor tyrosine kinases and regulate serine/threonine kinases to activate ERK in neuronal cells. *Br. J. Pharmacol.* **165,** 2497–2511 (2012).
21. Berghuis, P. *et al.* Endocannabinoids regulate interneuron migration and morphogenesis by transactivating the TrkB receptor. *Proc. Natl. Acad. Sci. U. S. A.* **102,** 19115–19120 (2005).
22. Bothwell, M. NGF, BDNF, NT3, and NT4. *Handb. Exp. Pharmacol.* **220,** 3–15 (2014).
23. Castrén, E. & Antila, H. Neuronal plasticity and neurotrophic factors in drug responses. *Mol. Psychiatry* (2017). doi:10.1038/mp.2017.61
24. Aso, E., Ozaita, A., Serra, M.-À. & Maldonado, R. Genes differentially expressed in CB1 knockout mice: involvement in the depressive-like phenotype. *Eur. Neuropsychopharmacol.* **21,** 11–22 (2011).
25. Marsicano, G. *et al.* The endogenous cannabinoid system controls extinction of aversive memories. *Nature* **418,** 530–534 (2002).
26. Karpova, N. N. *et al.* Fear erasure in mouse requires synergy between antidepressant drug treatment and exposure therapy. *Science* **334,** 1731–1734 (2011).
27. Rantamäki, T. *et al.* Antidepressant drugs transactivate TrkB neurotrophin receptors in the adult rodent brain independently of BDNF and monoamine transporter blockade. *PLoS One* **6,** e20567 (2011).
28. Njung'e, K. & Handley, S. L. Evaluation of marble-burying behavior as a model of anxiety. *Pharmacol. Biochem. Behav.* **38,** 63–67 (1991).
29. Paxinos, G. Paxinos and Franklin's the mouse brain in stereotaxic coordinates. (2013).

30. Antila, H. *et al.* Utilization of in situ ELISA method for examining Trk receptor phosphorylation in cultured cells. *J. Neurosci. Methods* **222,** 142–146 (2014).

31. Karpova, N. N. *et al.* Darkness reduces BDNF expression in the visual cortex and induces repressive chromatin remodeling at the BDNF gene in both hippocampus and visual cortex. *Cell. Mol. Neurobiol.* **30,** 1117–1123 (2010).

32. Paxinos, G. & Franklin, K. The mouse brain atlas in stereotaxic coordinates. *San Diego, CA: Academic* (2001).

33. Umemori, J., Winkel, F., Castrén, E. & Karpova, N. N. Distinct effects of perinatal exposure to fluoxetine or methylmercury on parvalbumin and perineuronal nets, the markers of critical periods in brain development. *Int. J. Dev. Neurosci.* **44,** 55–64 (2015).

34. Casarotto, P. Complete experimental data. (2018). doi:10.6084/m9.figshare.6292901.

35. Benarroch, E. E. Synaptic effects of cannabinoids: complexity, behavioral effects, and potential clinical implications. *Neurology* **83,** 1958–1967 (2014).

36. Piomelli, D. The molecular logic of endocannabinoid signalling. *Nat. Rev. Neurosci.* **4,** 873–884 (2003).

37. Starowicz, K., Nigam, S. & Di Marzo, V. Biochemistry and pharmacology of endovanilloids. *Pharmacol. Ther.* **114,** 13–33 (2007).

38. Maison, P., Walker, D. J., Walsh, F. S., Williams, G. & Doherty, P. BDNF regulates neuronal sensitivity to endocannabinoids. *Neurosci. Lett.* **467,** 90–94 (2009).

39. Lemtiri-Chlieh, F. & Levine, E. S. BDNF evokes release of endogenous cannabinoids at layer 2/3 inhibitory synapses in the neocortex. *J. Neurophysiol.* **104,** 1923–1932 (2010).

40. Zhao, L. & Levine, E. S. BDNF-endocannabinoid interactions at neocortical inhibitory synapses require phospholipase C signaling. *J. Neurophysiol.* **111,** 1008–1015 (2014).

41. Zhao, L., Yeh, M. L.-W. & Levine, E. S. Role for Endogenous BDNF in Endocannabinoid-Mediated Long-Term Depression at Neocortical Inhibitory Synapses. *eNeuro* **2,** (2015).

42. Yeh, M. L., Selvam, R. & Levine, E. S. BDNF-induced endocannabinoid release modulates neocortical glutamatergic neurotransmission. *Synapse* **71,** (2017).

43. Zhong, P. *et al.* BDNF interacts with endocannabinoids to regulate cocaine-induced synaptic plasticity in mouse midbrain dopamine neurons. *J. Neurosci.* **35,** 4469–4481 (2015).

44. Khaspekov, L. G. *et al.* Involvement of brain-derived neurotrophic factor in cannabinoid receptor-dependent protection against excitotoxicity. *Eur. J. Neurosci.* **19,** 1691–1698 (2004).

45. Broekkamp, C. L., Rijk, H. W., Joly-Gelouin, D. & Lloyd, K. L. Major tranquillizers can be distinguished from minor tranquillizers on the basis of effects on marble burying and swim-induced grooming in mice. *Eur. J. Pharmacol.* **126,** 223–229 (1986).

46. Bruins Slot, L. A., Bardin, L., Auclair, A. L., Depoortere, R. & Newman-Tancredi, A. Effects of antipsychotics and reference monoaminergic ligands on marble burying behavior in mice. *Behav. Pharmacol.* **19,** 145–152 (2008).

47. Thomas, A. *et al.* Marble burying reflects a repetitive and perseverative behavior more than novelty-induced anxiety. *Psychopharmacology* **204,** 361–373 (2009).

48. Gomes, F. V., Casarotto, P. C., Resstel, L. B. M. & Guimarães, F. S. Facilitation of CB1 receptor-mediated neurotransmission decreases marble burying behavior in mice. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **35,** 434–438 (2011).

49. Marsicano, G. & Lutz, B. Expression of the cannabinoid receptor CB1 in distinct neuronal subpopulations in the adult mouse forebrain: CB1 expression in murine forebrain. *Eur. J. Neurosci.* **11,** 4213–4225 (1999).

50. Cristino, L. *et al.* Immunohistochemical localization of cannabinoid type 1 and vanilloid transient receptor potential vanilloid type 1 receptors in the mouse brain. *Neuroscience* **139,** 1405–1415 (2006).