Trait-based approach to bacterial growth efficiency

Mario E. Muscarella[1,2], Xia Meng Howey[1], Jay T. Lennon[1,*]

[1]Department of Biology, Indiana University, Bloomington, Indiana, USA

[2]Current Address: Département des Sciences Biologiques, Université du Quebéc à Montréal, Montréal, Quebec, Canada

* Corresponding author, Tel 812-856-0962, Fax 812-855-6082, Email lennonj@indiana.edu

Running Title: Growth Efficiency Traits

Key Words: Bacterial Production, Bacterial Respiration, Physiological Trade-offs

16 **ABSTRACT**

Bacterial growth efficiency (BGE) is the proportion of assimilated carbon that is converted

18   into biomass and reflects the balance between growth and energetic demands. Often measured as

an aggregate property of the community, BGE is highly variable within and across ecosystems.

20   To understand this variation, we focused on how species and resource identity affect BGE using

a trait-based approach with 20 bacterial isolates enriched from north temperate lakes. We used

22   phenotypic and genomic approaches to characterize the metabolism of each isolate and test for

predicted trade-offs between growth rate and efficiency. Regardless of resource identity, 20 % of

24   the variation in BGE could be attributed to the taxonomic class of the bacterial species, and 58 %

of the variation could be explained by species identity. Meanwhile, resource identity explained a

26   relatively small amount of variation (7 %) in BGE across species but accounted for > 60 % of the

variation within a species. Metabolic trade-offs and genomic features suggest that BGE is a

28   predictable trait that is independent of resource type. As such, genomic and phylogenetic

information from microbiomes may help predict aggregate community functions like BGE and

30   the fate carbon in ecosystems.

 INTRODUCTION

In most ecosystems, heterotrophic bacteria play a pivotal role in determining whether

36 organic carbon is respired as CO_2 or converted into biomass and retained in food webs (Pomeroy

et al. 1998; Bardgett et al. 2008; Ducklow 2008). Many factors control how bacteria process

38 carbon, but perhaps the most important is reflected by bacterial growth efficiency (BGE). BGE is

the proportion of assimilated organic carbon that is converted into bacterial biomass (del Giorgio

40 and Cole 1998). When BGE is high, more carbon is turned into bacterial biomass where it can be

retained for longer periods of time while also serving as a resource for other members of the food

42 web. In contrast, when BGE is low, microbially assimilated carbon has a shorter residence time

and is released into the environment as CO_2. When measured at the community scale, BGE is

44 notoriously variable among habitats and has proven difficult to predict (del Giorgio and Cole

1998).

46 While BGE can be influenced by a range of chemical and physical properties (Apple and

del Giorgio 2007; Hall and Cotner 2007; del Giorgio and Newell 2012; Sinsabaugh et al. 2013;

48 Geyer et al. 2016), it may also reflect taxon-specific differences in microbial metabolism. For

example, BGE results from the physiological balance between cellular growth and energetic

50 demands. As such, bacterial growth strategy is predicted to constrain BGE via physiological

trade-offs (Litchman et al. 2015), and it has been hypothesized that oligotrophs have higher

52 maximum growth efficiency than copiotrophs (Roller and Schmidt 2015). In addition, species

that specialize on only a few resources are predicted to be more efficient at using those resources

54 than generalist species (Dykhuizen and Davies 1980; Glasser 1984). As such, traits such as

maximum growth rate and the number of resources used (i.e., niche breadth) could underlie

56 species-specific differences in BGE.

Variation in BGE may also be influenced by the properties of resources that are used to

58   meet energetic and growth demands. Different resources can affect cellular ATP yield depending

on the metabolic pathways used (Fuhrer et al. 2005; Flamholz et al. 2013). For example, glucose

60   is metabolized via glycolysis, but growth on aromatic compounds, such as protocatechuate,

requires the β-ketoadipate pathway which yields less ATP (Gottschalk 1986). Furthermore,

62   energy-producing catabolic processes and biomass-producing anabolic processes are not

independent (Russell and Cook 1995; Kempes et al. 2012). For example, cells have the potential

64   to produce > 30 ATP from a single glucose molecule if it is completely oxidized, but there would

be no remaining carbon to yield new biomass. Instead, cells must use the intermediate products

66   of glycolysis to form proteins and other cellular material, which diminishes the maximum ATP

yield (Gottschalk 1986; Flamholz et al. 2013). In addition, biomass production requires materials

68   and energy. For example, the synthesis of proteins, which constitute ~70 % of cellular dry mass,

requires amino acid building blocks and 4 ATP per peptide bond (Tempest and Neijssel 1984;

70   Gottschalk 1986; Lynch and Marinov 2015). Therefore, because resources differ in their

potential energy yield and bacteria differ in their ability to extract energy and form biomass from

72   a given resource, BGE should vary based on the resources available to bacteria.

In this study we measured BGE in a set of bacterial isolates growing on three different

74   carbon resources that varied in chemical structure and metabolic pathway (Fig. 1). In addition to

partitioning variation in BGE based on species and resource identity, we tested for hypothesized

76   trade-offs with growth rate and niche breadth. Furthermore, using the genomes of each isolate,

we evaluated whether metabolic pathways could explain differences in BGE among diverse

78   representatives of aquatic bacteria from north temperate lakes. Last, to test if resources have

differential effects on the metabolic traits that underlie BGE (i.e., production and respiration), we

80   tested for resource-specific relationships between respiration and production rates for each

resource. Together, our trait-based approach provides a framework for understanding linkages

82   between community structure and function due to the physiological constraints on BGE and

suggest that large changes in community composition or available resources may alter BGE and

84   therefore carbon cycling in predictive ways.

RESULTS

88   **Bacterial Growth Efficiency** — Using measures of bacterial productivity (BP) and respiration

(BR), we calculated bacterial growth efficiency (BGE) for 20 isolates growing on three different

90   resources: glucose, succinate, and protocatechuate (Fig. 1). All isolates belonged to the

Proteobacteria phylum with representatives from the Alpha-, Beta-, and Gammaproteobacteria

92   subphyla (Fig. 2, Fig. S1). Across isolates, we found that BGE ranged from <0.01 to 0.32 (Fig.

2). Based on linear mixed-effects models we found that isolate and resource identity explained

94   substantial amount of variation in BGE. Across resources, isolate identity explained 58 % of the

variation in BGE, and 67 % of the variation within resource, and the taxonomic order of each

96   isolate explained 20 % of the variation in BGE across resources, and 28 % of the variation within

resource. Across isolates, resource identity only explained 7 % of the variation in BGE across all

98   isolates, but 63 % of variation within isolate.

        Next, we determined if there was any evidence of phylogenetic signal for BGE among

100   our isolates. Based on Blomberg's K (a measure of phylogenetic signal strength), there was weak

phylogenetic signal for BGE. We did not detect phylogenetic signal for BGE when isolates used

102   succinate ($K = 0.002$, $p = 0.24$) or protocatechuate ($K = 0.001$, $p = 0.146$), but there was a

significant phylogenetic signal when isolates used glucose ($K = 0.002$, $p = 0.04$). However, the

104   low K value suggests that BGE is over-dispersed (i.e., less phylogenetic signal than expected

under Brownian motion). Furthermore, when we use Pagel's λ (a measure of phylogenetic signal

106   compared to Brownian motion), we found no evidence of a signal (Glucose: $\lambda = 0.10$, $p = 0.76$;

Succinate: $\lambda = 0.13$, $p = 0.66$; Protocatechuate: $\lambda = < 0.01$, $p = 0.99$).

108   Last, we determined if the values of BGE observed across isolates and resources were

unimodal. Based on Hartigan's dip test, we found that there was a bimodal distribution of BGE

110   among our isolates ($D = 0.07$, $p = 0.58$, Fig. S2). Using this distribution, we split isolates into

two groups, which we define as the "high BGE", and "low BGE" groups.

Phenotypic Comparisons — Using linear models, we found phenotypic differences between

114   isolates that were related to BGE (Fig. 3). While there was no relationship between BGE and

maximum growth rate (μ_{max}; $F_{1,7} = 0.035$, $r^2 < 0.01$, $p = 0.86$) in the low-BGE group of bacteria,

116   we identified a significant negative relationship between BGE and μ_{max} for the high-BGE group

($F_{1,7} = 7.79$, $r^2 = 0.53$, $p = 0.027$). Based on this model we would predict a 2.6 % decrease in

118   BGE for each per minute increase in μ_{max} in the high-BGE group. In contrast to our predictions,

there was no relationship between niche breadth (Levin's Index) and BGE for the low-BGE

120   group ($F_{1,7} = 1.42$, $r^2 = 0.17$, $p = 0.27$) or the high-BGE group ($F_{1,7} = 0.92$, $r^2 = 0.11$, $p = 0.37$).

122   **Genomic Comparisons** — We found genomic differences between isolates that were related to

BGE. First, isolates in the high-BGE group had 26 % more annotated metabolic pathways (based

124   on an 80 % module completion ratio cutoff) than isolates in the low BGE-group ($m_{high} = 66$,

$m_{low} = 652$, t-test: $t_{18} = -2.64$, $p = 0.02$). Second, we found that the number of metabolic

126   pathways was related to the observed BGE when grown on glucose, but the relationship

depended on the group. For the high-BGE group there was a negative relationship between BGE

128   and the number of pathways ($\beta = -0.006$, r2 $= 0.48$, $p = 0.04$), but for the low-BGE group there

was a positive relationship ($\beta = 0.0003$, r2 $= 0.37$, $p = 0.05$). Next, we found that differences in

130   the metabolic pathway composition of a genome could help explain which group an isolate

belongs, and pathway composition was related to the observed value of BGE. Specifically, we

132   found three pathways that were indicators of an isolate being in the high-BGE group (Table 1).

Likewise, for the high-BGE group we found that we could explain 24 % of the variation in BGE

134   based on the composition of metabolic pathways (dbRDA: $F_{1,7} = 2.17$, $R^2 = 0.24$, $p = 0.05$). We

found eight pathways with significant positive or negative correlations ($|\rho| > 0.7$) with BGE

136   (Table 2). However, we not find a relationship between pathway composition and BGE for the

low-BGE group ($p = 0.45$).

Resource Effects — Indicator variable linear regression revealed a positive relationship between

140   respiration and production rates (Fig. 4, $F_{9,42} = 8.07$, $R^2 = 0.63$, $p < 0.001$) with there being a

higher y-intercept for the high-BGE group of isolates ($\beta_{Group} = 2.7$, $p < 0.001$). Resource identity,

142   however, had no effect on the BR-BP relationship or the effect of group (all $p > 0.25$). Last, we

did not find evidence that the slope of the BR-BP relationship was different from one ($t_{42} = 0.76$,

144    $p = 0.45$); therefore, the two measures of bacterial metabolism appear to scale proportionately

with one another.

DISCUSSION

148   We measured bacterial growth efficiency (BGE) in 20 environmental bacterial isolates grown on

three resources that varied in their bioavailability, structure, and pathways required for

150   metabolism (Fig. 1). While BGE varied across isolates, there was no evidence for a strong

phylogenetic signal in efficiency. However, a substantial amount (20 %) of the variation in BGE

152   could be explained by an isolate's taxonomic order while a much smaller amount of the variation

(7 %) could be attributed to resource identity (Fig. 3). We found evidence for the predicted trade-

154   off between maximum growth rate and efficiency, but only on the most labile resource (glucose)

(Fig. 3). Even though we found that resource identity explained 63 % of the variation in BGE

156   within an isolate, resource identity did not alter the relationship between respiration and

production rate observed across isolates (Fig. 4) suggesting that resource identity has a stronger

158   effect on BGE within a species than across species. Together, we propose that growth efficiency

is a physiological trait independent of resource identity, but resource characteristics may modify

160   species-specific physiological performances. We propose that taxonomic groups of bacteria may

have fundamentally different growth efficiencies such that changes in community composition

162   may alter the fate of carbon resources (i.e., biomass versus CO_2) within the ecosystem.

164   **Bacterial Growth Efficiency as a Trait**

Our results indicate that there are species-specific properties regulating BGE, which may be

166   conserved at higher taxonomic levels. This conclusion is consistent with the view that BGE

represents a complex bacterial trait (i.e., aggregate property of numerous cellular functions) with

168   ecological significance, and that different groups of bacteria have fundamentally different

strategies for carbon allocation. Our phylogenetic analyses suggest that BGE may be an over-

170   dispersed trait (at least with glucose) such that the efficiency of closely related bacteria may be

less similar than expected. One potential explanation for this pattern of over-dispersion is that

172   our culture collection lacked phylogenetic resolution within some of our taxonomic groups (e.g.,

Betaproteobacteria) or that the variation in BGE within a taxonomic group (e.g., order) may not

174   be the same across taxonomic groups. Alternatively, BGE may not be a phylogenetically

conserved trait. Though some traits such as phosphorus acquisition, photosynthesis, and

176   methanogenesis are phylogenetically conserved deep in the microbial tree of life (Martiny et al.

2006, 2013), others such as complex carbon metabolism are not (Zimmerman et al. 2013).

178   Therefore, it is possible that BGE may be similar to traits such as complex carbon metabolism

that are not deeply conserved, which appears to be common among complex traits (Martiny et al.

180   2015). Regardless, our data reveal that BGE is a complex bacterial trait that is influenced by

taxonomic affiliation. As such, it may be possible to make predictions about BGE and other

182   ecosystem functions given information about composition of resident microbiomes (Goberna and

Verdú 2016).

Bacterial Growth Efficiency on Different Resources

186   Differences in resource complexity and the metabolic pathways required for degradation may

explain species-specific differences in BGE due to resource identity. Within an isolate, resource

188   identity accounted for 63 % of the variation in BGE. Given that different resources are processed

via different metabolic pathways, resource-based variation in BGE within a species is expected.

190   For example, BGE was higher when isolates were supplied with glucose compared to when they

were supplied with protocatechuate. Glucose is a simple sugar that is able to be metabolized by

192   numerous pathways and converted to acetyl-CoA (Neidhardt 2007). Protocatechuate, on the

other hand, is a complex aromatic compound that requires a specific metabolic pathway to be

194   converted to acetyl-CoA. Furthermore, because protocatechuate is chemically more complex, it

requires more energy (i.e., ATP) to be degraded than more labile resources such as glucose

196   (Harwood and Parales 1996). Therefore, resource complexity and the metabolic pathways

required may explain the within-isolate variation in BGE. However, across isolates, we did not

198   find resource-specific differences in the relationship between respiration and production rate.

Such findings suggest that energetic demands required to use different resources may also be

200   species-specific trait. That is, the energetic demands for individual species may be highly

constrained and therefore not change much when growing on different resources. Together, these

202   findings suggest that the effect of resources on the efficiency of entire microbiomes may depend

on the composition of bacteria consuming those resources.

Bacterial Growth Efficiency Groups

206   Across all isolates, we found a bimodal distribution of BGE suggesting that there were

two distinct groups with contrasting efficiencies. One group had low BGE (< 5 %) across all

208   treatments, and the other group ranged in BGE from 7-30 % (Fig. 2 & 3). Although, the range of

BGE measured across isolates is similar to the range observed in many ecosystems (del Giorgio

210   and Cole 1998), our results suggest that some species of bacteria grow relatively inefficiently,

irrespective of resource quality. One explanation is that the minimum cellular energetic demand

212   (i.e., cellular maintenance costs) is higher in some bacteria than others (Russell and Cook 1995).

Furthermore, energetic demand may be higher when bacteria are grown in minimal media where

214   they must produce all cellular components from a single carbon resource (Tao et al. 1999).

Alternatively, nutrient concentrations (e.g., phosphorus) and other physical properties (e.g.,

216   temperature) may regulate efficiency (Smith and Prairie 2004; Frey et al. 2013) and the effects of

these properties may be species-specific. As such, it is possible that maintenance costs, resource

218 imbalances, and the physical growth conditions affected BGE of our isolates. Differences in low-

BGE and high-BGE isolates was also reflected in genomic content, including the number and

220 presence-absence of metabolic pathways. Together, these findings suggest that there are

fundamental differences between bacterial species that determine BGE. However, these

222 genomic features seem to best explain large-scale rather than fine-scale differences in BGE.

Physiological Trade-Offs

We found evidence to support a trade-off between maximum growth rate and BGE (Fig. 4),

226 which is predicted in microbial and non-microbial systems (Glasser 1984; Roller and Schmidt

2015). For example, theoretical models of microbial communities predict a rate-efficiency trade-

228 off (Allison 2014), and this trade-off has been observed across microbial taxa (Lipson 2015).

Physiologically, the trade-off is based on allocation constraints imposed by the balance between

230 energy requirements and biomass yield: organisms with higher maximum growth rates may have

more energetic requirements and thus lower BGE (Russell and Baldwin 1979; Russell and Cook

232 1995). Furthermore, processes that limit respiration, such as oxygen availability, have been

shown to suppress bacterial growth rate (Meyenburg and Andersen 1980). Therefore, respiration

234 rate is likely a major control on biomass production and BGE. Consistent with this, we observed

a scaling relationship between respiration and production rates (Fig. 4). Furthermore, this

236 relationship between respiration and production and the non-zero intercept suggest that there is a

minimum respiration rate required before any biomass can be produced, which is commonly

238 interpreted as the cellular maintenance requirement. Therefore, it is possible that the maintenance

energy demand of bacterial species explains the physiological trade-off between maximum

240   growth rate and growth efficiency.

Theory also predicts a trade-off between resource niche-breadth and growth efficiency

242   (Glasser 1984). This trade-off is based on the assumption that there is an energetic cost to

maintaining numerous metabolic pathways (Johnson et al. 2012). As such, species with more

244   metabolic pathways should have more energetic requirements and thus lower BGE; although, the

effects of genome reduction has been debated (Giovannoni et al. 2005; Livermore et al. 2014). In

246   this study, we did not find evidence of a trade-off between resource niche breadth and BGE (Fig.

4). One possible explanation is that the resources used in our phenotypic assay (i.e. Ecolog

248   plates) did not reflect the full metabolic potential of our isolates. Alternatively, there may not be

a strong trade-off between niche breadth and efficiency, but further experiments with additional

250   isolates and resources would be required to test this prediction more rigorously.

252   **Genomic Signatures**

In addition to the physiological differences documented among our isolates, we found genomic

254   evidence of metabolic pathways that are associated with BGE. Specifically, we found genomic

differences between isolates that belong to low-BGE and high-BGE groups. We discovered that

256   isolates in the high-BGE group had 26 % more metabolic pathways than the low-BGE group.

Furthermore, we identified three pathways that were unique to the high-BGE group (Table 1)

258   and a number of pathways that were correlated with the observed BGE (Table 2). Together, our

findings suggest that there are genomic features that may contribute to or regulate BGE.

260   In general, the genomic composition of BGE groups appear to reflect differences in cellular

biosynthesis. It is possible that species with particular biosynthesis pathways may generate

262   essential cellular components with less energetic demand. For example, the low-BGE isolates

lacked some metabolic pathways, including pyridoxal biosynthesis and histidine degradation,

264   which were present in the high-BGE group. The pyridoxal biosynthesis pathway produces

vitamin B_6 from erythrose-4-phosphate (Mukherjee et al. 2011). Because vitamin B_6 is essential

266   for growth, the isolates lacking the pyridoxal pathway must use alternatives such as uptake from

the environment if they are auxotrophic (i.e., unable to synthesize) or other synthesis pathways

268   such as the Deoxyxylulose-5-phosphate synthase (DXS) pathway (Mukherjee et al. 2011).

However, the DXS pathway requires pyruvate (a precursor for Krebs cycle) and thus may limit

270   central metabolism and possibly lead to lower BGE. Likewise, the histidine degradation pathway

is used to breakdown histidine into ammonium and glutamate (Bender 2012). Alternatively,

272   glutamate can by synthesized from α-ketoglutarate; however, because α-ketoglutarate is an

intermediate component of Krebs cycle this may limit central metabolism and possibly lead to

274   reduced BGE.

Conclusion

276   At the cellular level, BGE reflects the balancing energetic and cellular growth demands.

We find evidence of this based on physiological trade-offs (i.e., maximum growth rate) as well

278   as metabolic pathways. As such, changes in community composition and resource availability

have to potential to alter food web and ecosystem function due to changes in BGE. For example,

280   communities dominated by species with low BGE should yield a net release of CO_2 from the

ecosystem. Alternatively, communities comprised of individuals with high BGE should yield a

282   net increase in ecosystem productivity. However, variation in BGE can arise within a species due

to the ways in which it processes different resources. Therefore, changes in the resource supply

284   will alter the performance of individual taxa, but we predict that these changes will not be as

strong as changes in BGE that arise owing to differences in community composition. A trait-

286 based approach can be used to provide a mechanistic link between the structure and function of

bacterial communities.

METHODS

290 **Bacterial Isolates** — Using a novel cultivation approach, we isolated 20 bacterial strains from

lakes in the Huron Mountain Research Preserve (Powell, MI, USA) by incubating inert carbon

292 beads (Bio-Sep Beads) in the water column for one week. Prior to the incubations, the beads

were saturated with a sterile complex carbon substrate, i.e., Super Hume (CropMaster, United

294 Agricultural Services of America, Lake Panasoffkee, Florida, USA). Super Hume is a lignin-rich

resource comprising 17 % humic and 13 % fulvic acids, and has been shown to be an analog of

296 terrestrial DOC in aquatic ecosystems that can be used by diverse bacteria (Lennon et al. 2013).

We used this enrichment technique to select for bacteria with a range of metabolic potentials

298 (Ghosh et al. 2009). After the incubation, beads were rolled on R2 agar plates (BD Difco, Sparks

Maryland, USA) and incubated at 25 °C. We picked random colonies from plates and serially

300 transferred until axenic. All isolates were preserved in 25 % glycerol at -80 ºC.

We identified each bacterial strain by direct sequencing the 16S rRNA gene. We obtained

302 genomic DNA from log phase cultures using the FastPrep DNA extraction kit according to the

manufacturer's specifications (MP Biomedical). We used 10 ng of genomic DNA to amplify the

304 16S rRNA gene using the 27F and 1492R bacterial primers (See Supplemental for primer

sequences and PCR conditions). We sequenced the PCR products at the Indiana Molecular

306 Biology Institute (IMBI) at Indiana University (Bloomington, Indiana, USA). Raw sequence

reads were quality-trimmed based on a Phred quality score of 25. Forward and reverse reads

308   were manually merged after aligning sequences to the Silva 16S SSU rRNA reference alignment

(release 132) using SINA v. 1.2.11 and the Bacteria variability profile (Pruesse 2011). After

310   merging into full length 16S rRNA sequences, we used mothur (Schloss et al. 2009) to check the

quality of sequences and alignments were checked using the ARB software package (Ludwig et

312   al. 2004). Finally, sequences were compared to the Silva All-Species Living Tree Project

database (Yilmaz et al. 2014) for taxonomic identification (Fig. S1).

Bacterial Growth Efficiency — To test for differences in bacterial growth efficiency (BGE), we

316   measured BGE for each isolate when grown on each of three different carbon substrates:

glucose, succinate, or protocatechuate (Fig. 2). These carbon sources (i.e., resources) were

318   chosen based on differences in their bioavailability and structure but also the required pathways

for metabolism (see Fig. 2).

320   To determine BGE we measured bacterial respiration and production rates and then

calculated BGE as BP/(BP + BR), where BP is bacterial productivity and BR is bacterial

322   respiration (del Giorgio and Cole 1998). BP and BR were measured using triplicate cultures of

each isolate. Cultures of each isolate were grown in R2 broth (BD Difco, Sparks Maryland,

324   USA) until mid-log phase. We then transferred 100 µL of culture into 10 mL of M9 broth (Green

and Sambrook 2012) with the appropriate carbon source (25 mM C) and allowed 24 h for the

326   cultures to acclimate. We then transferred 100 µL of culture into 10 mL of fresh carbon-amended

M9 broth and incubated 1-3 h to replenish nutrients. Using these transfers, we were able to

328   establish populations of each isolate at target cell densities between 10^4 and 10^5 cells mL^{-1}. We

used the populations to measure BP and BR, which were normalized to cell density using plate

330   counts of colony forming units. We measured BP using the ^{3}H-Leucine assay (Smith and Azam

1992) with 1.5 mL of culture. We added ^{3}H-Leucine to a final concentration of 50 mM and

332   incubated for 1 h. Following incubation, we terminated production with trichloroacetic acid

(final concentration 3 mM) and measured leucine incorporation using a liquid scintillation

334   counter. We measured BR using an automated O_2 measurement system (PreSens Sensor Dish

System, PreSens, Regensburg, Germany) on 5 mL of culture based on the rate O_2 consumption

336   during three-hour incubations. We estimated BR as the slope of O_2 concentration during the

incubation using linear regression. We used theoretical respiratory quotients for each resource to

338   convert O_2 depletion into C respiration assuming aerobic growth.

340   **Taxonomic and Phylogenetic Relationships** — We compared differences in BGE across

isolates and resources using linear models. First, we used a taxonomic framework to compare

342   BGE between isolates (Lennon et al. 2012). Isolates were classified into taxonomic groups based

on the species tree constructed in ARB. We then used mixed linear models to compare BGE

344   across taxonomic groups and resources. To test the hypothesis that taxonomy (i.e., at the class

level) affects BGE, we nested resource identity within isolate in the linear model. To test the

346   hypothesis that resource identity affects BGE, we nested isolate identity within resource. We

identified the best statistical models based on the variation explained (R^2) and AIC values.

348   Second, we tested if phylogenetic relationships between isolates explained differences in BGE

across isolates. We created a phylogenetic tree based on the full-length 16S rRNA gene

350   sequences. We aligned sequences using the SINA aligner (Pruesse et al. 2012) and checked

alignments using ARB. We generated a phylogenetic tree using the CIPRES science gateway

352   (Miller et al. 2010). The phylogenetic tree was created using RAxML v.8.2.12 (Stamatakis

2006). We used the GTRGAMMA DNA substitution model and the rapid hill-climbing

354   algorithm to build our maximum likelihood trees, and we used the extended majority rule to find

the consensus tree. We used Blomberg's K and Pagel's Lambda to compare trait variation across

356   the tree and test if phylogenetic relationships between isolates could explain differences in traits

(Pagel 1999; Blomberg et al. 2003). Blomberg's K is a test for phylogenetic signal that

358   determines if trait variation is better explained by phylogenetic relationships or Brownian

motion. Pagel's Lambda is a test of phylogenetic signal that determines if trait variation differs

360   from Brownian motion. Last, to determine if the distribution of BGE across isolates was

unimodal, we used Hartigan's dip test for unimodality (Hartigan and Hartigan 1985). Hartigan's

362   dip test is used to determine if a distribution is unimodal by testing the null hypothesis that there

is a dip in the distribution. A significant Hartigan's dip test would suggest that the distribution is

364   unimodal. Alternatively, the distribution has an internal "dip" (reported as D). All statistical tests

were conducted in the R statistical environment (R Core Development Team 2013). We used the

366   nlme package (Pinheiro and Bates 2011) for the mixed linear models, the picante package

(Kembel et al. 2015) for the phylogenetic methods, and the diptest package (Maechler 2015) for

368   Hartigan's dip test.

370   **Phenotypic Comparisons and Trade-offs** — To test the hypothesis that phenotypic differences

and physiological trade-offs underlie BGE variation, we compared the maximum growth rate

372   (μmax) and niche breadth of each isolate. First, to test whether BGE was affected by growth

strategy (i.e., copiotrophs vs. oligotrophs), we measured the maximum growth rate of each

374   isolate. Bacterial growth rates were measured based on changes in optical density during 18-h

incubations. Bacterial isolates were grown in R2 broth in 48-well plates. We incubated plates

376   with continuous shaking and measured optical density every 15 min using a plate reader (BioTek

MX). Growth curves were analyzed by fitting a modified Gompertz growth model (Zwietering et

378   al. 1990; Lennon 2007) to the observed growth curves using maximum likelihood fitting. We

used the model fit as our estimate of μmax.

380   Second, to test whether BGE was affected by niche breadth, we generated carbon usage

profiles using BioLog EcoPlates[TM] (Garland and Mills 1991). The EcoPlate is a phenotypic

382   profiling tool consisting of 31 unique carbon sources. In addition to the carbon source, each well

contains a tetrazolium dye, which in the presence of NADH will be reduced and change color.

384   We used this colorimetric assay to generate carbon usage profiles for each strain. We

standardized profiles for each strain by subtracting water blanks (average water blank + 1 SD),

386   and relativizing across substrates. Using these data, we calculated resource niche breadth using

Levin's Index (Colwell and Futuyma 1971).

Genomic Comparisons — To test the hypothesis that variation in metabolic pathways could

390   explain differences in BGE, we compared the genomes of each isolate. First, we determined the

metabolic pathways found in the genome of each isolate. We characterized each isolate using

392   whole genome sequencing. Genomic DNA libraries for each isolate were prepared using the

Illumina TruSeq DNA sample prep kit using an insert size of 250 base pairs (bp). Libraries were

394   sequenced on an Illimina HiSeq 2500 (Illumina, San Diego, GA) using 100-bp paired-end reads

at the Michigan State University Research Technology Support Facility. We processed raw

396   sequence reads (FASTQ) by removing the Illumina TruSeq adaptors using Cutadapt (Martin

2011), interleaving reads using Khmer (McDonald and Brown 2013), and quality-filtering based

398   on an average Phred score of 30 using the FASTX-toolkit (Hannon Lab 2010). Finally, we

normalized coverage to 25 based on a k-mer size of 25 using Khmer. We assembled the genomes

400   using Velvet (Zerbino and Birney 2008) after optimizing assembly parameters for each isolate

with Velvet Optimizer (Gladman and Seemann 2012). We annotated contigs larger than 200 bp

402   using Prokka (Seemann 2014), and predicted metabolic and physiological functions using

MAPLE with bidirectional best-hit matches (Takami et al. 2012). We identified functional

404   pathway based on the presence of intermediate genes within a pathway. We scored pathways as

functional if more than 80 % of the intermediate genes were recovered in the genomes based on

406   module completion ratios.

To test the hypothesis that metabolic pathways affect BGE, we used multivariate methods

408   to compare the pathways of each isolate. First, we used PERMANOVA to determine if there

were differences in pathways associated with different levels of BGE. When significant

410   differences were found, we used indicator species analysis (Dufrene and Legendre 1997) to

determine which metabolic pathways contributed to group differences in BGE. Next, to

412   determine if metabolic pathways could explain differences in BGE within a group, we used

distance-based redundancy analysis (dbRDA) which is a multivariate technique that tests if a

414   quantitative predictor can explain differences in multivariate datasets (Legendre and Legendre

2012). Because we scored pathways as present or absent, metabolic distances between isolates

416   were calculated using the Jaccard Index. We tested for significance using a permutation test. If

the dbRDA model was significant, we used Spearman's rank-order correlation to test for

418   correlations between BGE and individual metabolic pathways. We used the vegan R package

(Oksanen et al. 2012) for multivariate analyses.

Resource Effects — To test the hypothesis that resources have different effects on components

422   of metabolism that affect BGE, we used a linear model to test for a relationship between BR and

BP. Because BP required energy through respiration, we used production rate as the dependent

424   variable and respiration rate as the independent variable. We used an indicator variable linear

regression to test for changes in BP rate due to BR. We included resource identity and group

426   (high- versus low-BGE) as the categorical predictors and BR as the continuous predictor

(Lennon and Pfaff 2005). In addition, we included all interactions terms. Respiration and

428   production rates were $\log_{10}$-transformed to meet model assumptions. Last, to determine if the

relationship between BR and BP rates was isometric (proportional scaling, slope = one) or

430   allometric (disproportional scaling, slope ≠ one), we used a one-sample t-test to determine if the

slope was different from one. All statistical tests were conducted in the R statistical environment.

ACKNOWLEDGEMENTS

434   We thank BK Lehmkuhl and MA Carrison for technical assistance and JB McKinlay and

members of the Lennon Lab for critical feedback on an earlier version of this manuscript. This

436   work was supported by the Huron Mountain Wildlife Foundation (MEM & JTL), the National

Science Foundation (DEB-0842441, DEB-1442246, and DEB-1501164), and US Army Research

438   Office (W911NF-14-1-0411). All code and data used in this study can be found in a public

GitHub repository (https://www.github.com/LennonLab/MicrobialCarbonTraits). Isolate

440   genomes are available on NCBI (BioProject PRJNA420393).

REFERENCES

444   Allison, S. D. 2014. Modeling adaptation of carbon use efficiency in microbial communities. Frontiers in Microbiology 5:1–9.

446   Apple, J. K., and P. A. del Giorgio. 2007. Organic substrate quality as the link between bacterioplankton carbon demand and growth efficiency in a temperate salt-marsh estuary. The

448   ISME Journal 1:729–742.

Bardgett, R. D., C. Freeman, and N. J. Ostle. 2008. Microbial contributions to climate change

450   through carbon cycle feedbacks. The ISME Journal 2:805–14.

Bender, R. A. 2012. Regulation of the histidine utilization (Hut) system in bacteria.

452   Microbiology and Molecular Biology Reviews 76:565–584.

Blomberg, S. P., T. Garland, and A. R. Ives. 2003. Testing for phylogenetic signal in

454   comparative data: behavioral traits are more labile. Evolution 57:717–745.

Buchan, A., L. S. Collier, and E. L. Neidle. 2000. Key aromatic-ring-cleaving enzyme,

456   protocatechuate 3,4-dioxygenase, in the ecologically important marine *Roseobacter* lineage. Applied and Environmental Microbiology 66:4662–4672.

458   Colwell, R. K., and D. J. Futuyma. 1971. On the Measurement of Niche Breadth and Overlap. Ecology 52:567–576.

460   del Giorgio, P. A., and J. J. Cole. 1998. Bacterial growth efficiency in natural aquatic systems. Annual Review of Ecology and Systematics 29:503–541.

462   del Giorgio, P. A., and R. E. I. Newell. 2012. Phosphorus and DOC availability influence the partitioning between bacterioplankton production and respiration in tidal marsh ecosystems.

464   Environmental Microbiology 14:1296–307.

Ducklow, H. W. 2008. Bacterial Production and Biomass in the Oceans. Pages 1–47 *in*Microbial

466   Ecology of the Oceans.

Dufrene, M., and P. Legendre. 1997. Species assemblages and indicator species: the need for a

468   flexible asymmetrical approach. Ecological Monographs 67:345–366.

Dykhuizen, D., and M. Davies. 1980. An Experimental Model: Bacterial Specialists and

470   Generalists Competing in Chemostats. Ecology 61:1213–1227.

Flamholz, A., E. Noor, A. Bar-Even, W. Liebermeister, and R. Milo. 2013. Glycolytic strategy as

472   a tradeoff between energy yield and protein cost. Proceedings of the National Academy of

Sciences 110:10039–44.

474   Frey, S. D., J. Lee, J. M. Melillo, and J. Six. 2013. The temperature response of soil microbial

efficiency and its feedback to climate. Nature Climate Change 3:395–398.

476   Fuhrer, T., E. Fischer, and U. Sauer. 2005. Experimental identification and quantification of

glucose metabolism in seven bacterial species. Journal of Bacteriology 187:1581–1590.

478   Garland, J. L., and A. L. Mills. 1991. Classification and characterization of heterotrophic

microbial communities on the basis of patterns of community-level sole-carbon-source

480   utilization. Applied and Environmental Microbiology 57:2351–9.

Geyer, K. M., E. Kyker-Snowman, A. S. Grandy, and S. D. Frey. 2016. Microbial carbon use

482   efficiency: accounting for population, community, and ecosystem-scale controls over the fate of

metabolized organic matter. Biogeochemistry 127:173–188.

484   Ghosh, D., K. Roy, V. Srinivasan, T. Mueller, O. H. Tuovinen, K. Sublette, A. Peacock, et al.

2009. In-situ enrichment and analysis of atrazine-degrading microbial communities using

486   atrazine-containing porous beads. Soil Biology and Biochemistry 41:1331–1334.

Giovannoni, S. J., H. J. Tripp, S. Givan, M. Podar, K. L. Vergin, D. Baptista, L. Bibbs, et al.

488   2005. Genome streamlining in a cosmopolitan oceanic bacterium. Science 309:1242–5.

Gladman, S., and T. Seemann. 2012. Velvet Optimizer.

490   Glasser, J. W. J. 1984. Evolution of efficiencies and strategies of resource exploitation. Ecology 65:1570–1578.

492   Goberna, M., and M. Verdú. 2016. Predicting microbial traits with phylogenies. The ISME Journal 10:959–967.

494   Gottschalk, G. 1986. Bacterial Metabolism (2nd ed.). Springer-Verlag, New York, NY.

Green, M. R., and J. Sambrook. 2012. Molecular Cloning: A Laboratory Manual (4th ed.). Cold

496   Spring Harbor Laboratory Press, Cold Spring Harbor, New York.

Hall, E. K., and J. B. Cotner. 2007. Interactive effect of temperature and resources on carbon

498   cycling by freshwater bacterioplankton communities. Aquatic Microbial Ecology 49:35–45.

Hannon Lab. 2010. FASTX Toolkit.

Hartigan, J. A., and P. M. Hartigan. 1985. The dip test of unimodality. The Annals of Statistics 13:70–84.

Harwood, C. S., and R. E. Parales. 1996. The beta-ketoadipate pathway and the biology of self-identity. Annual Review of Microbiology 50:553–90.

Johnson, D. R., F. Goldschmidt, E. E. Lilja, and M. Ackermann. 2012. Metabolic specialization and the assembly of microbial communities. The ISME Journal 6:1985–1991.

Kembel, S. W., D. D. Ackerly, S. P. Blomberg, W. K. Cornwell, M. R. Helmus, M. Helene, and C. O. Webb. 2015. Picante: R tools for integrating phylogenies and ecology.

Kempes, C. P., S. Dutkiewicz, and M. J. Follows. 2012. Growth, metabolic partitioning, and the size of microorganisms. Proceedings of the National Academy of Sciences 109:495–500.

Legendre, P., and L. F. J. Legendre. 2012. Numerical Ecology (Vol. 24). Elsevier.

Lennon, J. T. 2007. Diversity and metabolism of marine bacteria cultivated on dissolved DNA.

512   Applied and Environmental Microbiology 73:2799–2805.

Lennon, J. T., Z. T. Aanderud, B. K. Lehmkuhl, and D. R. J. Schoolmaster. 2012. Mapping the

514   niche space of soil microorganisms using taxonomy and traits. Ecology 93:1867–1879.

Lennon, J. T., S. K. Hamilton, M. E. Muscarella, A. S. Grandy, K. Wickings, and S. E. Jones.

516   2013. A source of terrestrial organic carbon to investigate the browning of aquatic ecosystems.

PloS One 8:e75771.

518   Lennon, J. T., and L. E. Pfaff. 2005. Source and supply of terrestrial organic matter affects

aquatic microbial metabolism. Aquatic Microbial Ecology 39:107–119.

520   Lipson, D. A. 2015. The complex relationship between microbial growth rate and yield and its

implications for ecosystem processes. Frontiers in Microbiology 6:1–5.

522   Litchman, E., K. F. Edwards, and C. A. Klausmeier. 2015. Microbial resource utilization traits

and trade-offs: implications for community structure, functioning, and biogeochemical impacts at

524   present and in the future. Frontiers in Microbiology 06:1–10.

Livermore, J. A., S. J. Emrich, J. Tan, and S. E. Jones. 2014. Freshwater bacterial lifestyles

526   inferred from comparative genomics. Environmental Microbiology 16:746–758.

Ludwig, W., O. Strunk, R. Westram, L. Richter, H. Meier, Yadhukumar, A. Buchner, et al. 2004.

528   ARB: a software environment for sequence data. Nucleic Acids Research 32:1363–71.

Lynch, M., and G. K. Marinov. 2015. The bioenergetic costs of a gene. Proceedings of the

530   National Academy of Sciences 112:201514974.

Maechler, M. 2015. diptest: Hartigan's dip test statistic for unimodality.

532   Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal 17:10.

534   Martiny, A. C., M. L. Coleman, and S. W. Chisholm. 2006. Phosphate acquisition genes in

Prochlorococcus ecotypes: evidence for genome-wide adaptation. Proceedings of the National Academy of Sciences 103:12552–7.

Martiny, A. C., K. Treseder, and G. Pusch. 2013. Phylogenetic conservatism of functional traits in microorganisms. The ISME Journal 7:830–838.

Martiny, J. B. H., S. E. Jones, J. T. Lennon, and A. C. Martiny. 2015. Microbiomes in light of traits: A phylogenetic perspective. Science 350:aac9323–aac9323.

McDonald, E., and C. T. Brown. 2013. Khmer: working with big data in bioinformatics. arXiv 1303.2223:1–18.

Meyenburg, K. V. O. N., and K. B. Andersen. 1980. Are growth rates of *Escherichia coli* in batch cultures limited by respiration? Journal of Bacteriology 144:114–123.

Miller, M. A., W. Pfeiffer, and T. Schwartz. 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. 2010 Gateway Computing Environments Workshop.

Mukherjee, T., J. Hanes, I. Tews, S. E. Ealick, and T. P. Begley. 2011. Pyridoxal phosphate: biosynthesis and catabolism. Biochimica et Biophysica Acta - Proteins and Proteomics 1814:1585–1596.

Neidhardt, F. C. 2007. The Physiology and Biochemistry of Prokaryotes, 3rd ed. Microbe Magazine (4th ed., Vol. 2). Oxford University Press, New York, NY.

Oksanen, A. J., F. G. Blanchet, R. Kindt, P. Legen-, P. R. Minchin, R. B. O. Hara, G. L. Simpson, et al. 2012. Community Ecology Package. … Ecology Package ….

Pagel, M. 1999. Inferring the historical patterns of biological evolution. Nature 401:877–884.

Pinheiro, J., and D. Bates. 2011. Linear and nonlinear mixed effects models (nlme). Package of the R software for statistical computing on the CRAN repository.

Pomeroy, L. R., P. J. Williams, F. Azam, and J. E. Hobbie. 1998. The microbial loop.

558   Oceanography 20:28–33.

Pruesse, E. 2011. SINA: high throughput multiple sequence alignment.

560   Pruesse, E., J. Peplies, and F. O. Glöckner. 2012. SINA: accurate high-throughput multiple

sequence alignment of ribosomal RNA genes. Bioinformatics 28:1823–1829.

562   R Core Development Team. 2013. A language and environment for statistical computing.

Roller, B. R., and T. M. Schmidt. 2015. The physiology and ecological implications of efficient

564   growth. The ISME Journal 9:1481–1487.

Russell, J. B., and R. L. Baldwin. 1979. Comparison of maintenance energy expenditures and

566   growth yields among several rumen bacteria grown on continuous culture. Applied and

Environmental Microbiology 37:537–543.

568   Russell, J. B., and G. M. Cook. 1995. Energetics of bacterial growth: balance of anabolic and

catabolic reactions. Microbiological Reviews 59:48–62.

570   Schloss, P. D., S. L. Westcott, T. Ryabin, J. R. Hall, M. Hartmann, E. B. Hollister, R. A.

Lesniewski, et al. 2009. Introducing mothur: open-source, platform-independent, community-

572   supported software for describing and comparing microbial communities. Applied and

Environmental Microbiology 75:7537–7541.

574   Seemann, T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–9.

Sinsabaugh, R. L., S. Manzoni, D. L. Moorhead, and A. Richter. 2013. Carbon use efficiency of

576   microbial communities: stoichiometry, methodology and modelling. (J. Elser, ed.)Ecology

Letters 16:930–939.

578   Smith, D. C., and F. Azam. 1992. A simple, economical method for measuring bacterial protein

synthesis rates in seawater using 3H-leucine. Marine Microbial Food Webs 6:107–114.

580   Smith, E. M., and Y. T. Prairie. 2004. Bacterial metabolism and growth efficiency in lakes: the

importance of phosphorus availability. Limnology and Oceanography 49:137–147.

582   Stamatakis, A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with

thousands of taxa and mixed models. Bioinformatics 22:2688–2690.

584   Takami, H., T. Taniguchi, Y. Moriya, T. Kuwahara, M. Kanehisa, and S. Goto. 2012. Evaluation

method for the potential functionome harbored in the genome and metagenome. BMC Genomics

586   13:699.

Tao, H., C. Bausch, C. Richmond, F. R. Blattner, and T. Conway. 1999. Functional genomics:

588   expression analysis of *Escherichia coli* growing on minimal and rich media. Journal of

Bacteriology 181:6425–6440.

590   Tempest, D. W., and O. M. Neijssel. 1984. The status of YATP and maintenance energy as

biologically interpretable phenomena. Annual Reviews in Microbiology 459–486.

592   Yilmaz, P., L. W. Parfrey, P. Yarza, J. Gerken, E. Pruesse, C. Quast, T. Schweer, et al. 2014. The

SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks. Nucleic Acids

594   Research 42:643–648.

Zerbino, D. R., and E. Birney. 2008. Velvet: algorithms for de novo short read assembly using de

596   Bruijn graphs. Genome Research 18:821–9.

Zimmerman, A. E., A. C. Martiny, and S. D. Allison. 2013. Microdiversity of extracellular

598   enzyme genes among sequenced prokaryotic genomes. The ISME Journal 7:1187–1199.

Zwietering, M. H., I. Jongerburger, F. M. Rombouts, and K. van't Riet. 1990. Modeling of the

600   bacterial growth curve. Applied and Environmental Microbiology 56:1875–1881.

604 **Table 1**: Genetic pathways unique to the high BGE isolates. Pathways are functional metabolic

pathways identified from genome sequencing and predicted using Maple. Prob. = probability

606 statistic from indicator species analysis: the probability that the "species" (i.e., pathway) is not

unique to the group.

Group	Prob.	Pathway	Reference Function
	0.02	M00045	Histidine degradation (histidine → N-formiminoglutamate → glutamate)
High BGE	0.02	M00060	Lipopolysaccharide biosynthesis (Kdo2-lipid A biosynthesis)
	0.03	M00565	Trehalose biosynthesis (D-glucose-1P → trehalose)

612 **Table 2**: Genetic pathways correlated with BGE in the high-BGE group. Correlations are

spearman's rank correlations between BGE and the pathway presence. Pathways with

614 correlations greater than 0.70 or less than -0.70 were considered significant.

Relationship	ρ	Pathway	Reference Function
Positive	0.82	M00025	Tyrosine biosynthesis (chorismate → tyrosine)
	0.72	M00034	Methionine salvage pathway
Negative	-0.73	M00117	Ubiquinone biosynthesis (chorismate → ubiquinone)
	-0.82	M00044	Tyrosine degradation (tyrosine → homogentisate)
	-0.82	M00053	Pyrimidine deoxyribonuleotide biosynthesis (CDP/CTP → dCDP/dCTP,dTDP/dTTP)
	-0.82	M00549	Nucleotide sugar biosynthesis (glucose → UDP-glucose)
	-0.82	M00568	Catechol ortho-cleavage (catechol → 3-oxoadipate)
	-0.82	M00637	Anthranilate degradation (anthranilate → catechol)

626   **Fig. 1:** Carbon resources used to study BGE variation in environmental isolates. **A**: Glucose –

the baseline resource used to compare BGE across isolates. Glucose can be degraded by the

628   Embden-Meyerhof-Parnas, pentose phosphate, or Entner-Doudoroff pathway. Ultimately, these

pathways produce pyruvate (and then acetyl-CoA), which enters Krebs cycle and is used to

630   produce energy and intermediate for biomass synthesis, when cells are grown aerobically.

Alternatively, glucose can be fermented into organic acids (e.g., lactate), but these reactions yield

632   less energy (Gottschalk 1986). **B**: Succinate – is a simple organic acid. Succinate is an

intermediate of Krebs cycle and thus it does not require previous degradation. Additionally,

634   succinate can be used directly to produce energy via succinate dehydrogenase (Neidhardt 2007).

C: Protocatecuate – is a complex resource with an aromatic core. Typically, it is degraded to

636   acetyl-CoA and Succinyl-CoA via the β-ketoadipate pathway (Harwood and Parales 1996).

Protocatechuate is commonly used to study aromatic resource degradation in ecosystems, and the

638   β-ketoadipate pathway is commonly found in bacteria across the phylum Proteobacteria (Buchan

et al. 2000).

642 **Fig. 2:** Bacterial growth efficiency (BGE) of each isolate for each resource. BGE was calculated

based on measured production (BP) and respiration (BR) rates using the following equation:

644 BGE = BP / (BP + BR). Cladogram is based on the consensus phylogeny. Taxonomic class and

order are included based on the ribosomal database taxonomy: α = Alphaproteobacteria. β =

646 Betaproteobacteria, γ = Gammaproteobacteria, *Xan.* = Xanthomonadales, *Aero.* = Aeromondales,

Pseu. = Pseudomondales.

Fig. 3: Phenotypic traits associated with BGE. A: Maximum growth rate, a measure of growth

650 strategy, demonstrates a trade-off (negative relationship) with BGE in the high BGE group ($F_{1,7}$

= 9.52, r^2 = 0.54, p = 0.015), but not the low BGE group ($F_{1,7}$ = 0.51, r^2 = 0.06, p = 0.50). B:

652 Levin's Index, a measure of niche breadth, does not demonstrate a trade-off with BGE in either

the high or low BGE groups (high: $F_{1,7}$ = 0.92, r^2 = 0.11, p = 0.37; low: $F_{1,7}$ = 1.47, r^2 = 0.17, p =

654 0.27). High and low BGE groups were determined based on bimodal distribution of BGE.

656 **Fig. 4**: Relationship between respiration and production rates. Respiration and production rates

were compared using an indicator variable linear regression ($F_{9,42}$ = 4.92, R^2 = 0.51, $p < 0.001$).

658 According to the regression model, production rate increases with respiration rate proportionally

(i.e., slope not significantly different from one, t_{42} = -0.26, p = 0.79). In addition, group (high vs.

660 low BGE) was a significant factor and isolates in the high BGE group had a greater y-intercept

(p = 0.02). Symbols indicates isolate group (high and low BGE), and color indicates the resource

662 being used. Symbol size is scaled by growth efficiency.

FIGURES

Fig. 1:

A

Glucose

B

Succinate

C

Protocatecuate

Fig. 2:

672 **Fig. 3:**

Fig. 4:

676