Shared and Distinct Genetic Risk Factors

for Childhood Onset and Adult Onset Asthma

Milton Pividori, PhD[1,^], Nathan Schoettler, MD, PhD[1,2,^], Dan L. Nicolae, PhD[1,2,3], Carole

Ober, PhD[2,§], Hae Kyung Im, PhD[1,2,§]

Departments of Medicine[1], Human Genetics[2] and Statistics[3], The University of Chicago,

Chicago, IL

^These authors contributed equally.

§Corresponding Authors:

H.K.I. haky@uchicago.edu

C.O. c-ober@genetics.uchicago.edu

Background

Childhood and adult onset asthma differ with respect to severity and co-morbidities.

Whether they also differ with respect to genetic risk factors has not been previously

investigated.

Methods

We used data from the UK Biobank to conduct genome-wide association studies

(GWASs) in 9,433 childhood onset asthma (onset before age 12) and 21,564 adult

onset asthma (onset between ages 26 and 65) cases, each compared to 318,237 non-

asthmatic controls (older than age 38), and for age of onset in 37,846 asthma cases.

Enrichment studies determined the tissues in which genes at GWAS loci were most

highly expressed, and PrediXcan, a transcriptome-wide gene-based test, was used to

identify candidate risk genes.

Findings

We detected 61 independent asthma loci: 23 were childhood onset specific, one was

adult onset specific, and 37 were shared. Nineteen loci were associated with age of

asthma onset. Genes at the childhood onset loci were most highly expressed in skin,

blood and small intestine; genes at the adult onset loci were most highly expressed in

lung, blood, small intestine and spleen. PrediXcan identified 113 unique candidate

genes at 22 of the 61 GWAS loci.

Interpretation

Genetic risk factors for adult onset asthma are largely a subset of the genetic risk for

childhood onset asthma but with overall smaller effects, suggesting a greater role for

non-genetic risk factors in adult onset asthma. In contrast, the onset of disease in

childhood is associated with additional genes with relatively large effect sizes.

Combined with gene expression and tissue enrichment patterns, we suggest that the

establishment of disease in children is driven more by allergy and epithelial barrier

dysfunction whereas the etiology of adult onset asthma is more lung-centered, with

immune mediated pathways driving disease progression in both children and adults.

<u>Funding</u>

This work was supported by the National Institutes of Health grants R01 MH107666 and

P30 DK20595 to H.K.I., R01 HL129735, R01 HL122712, P01 HL070831, and UG3

OD023282 to C.O.; N.S. was supported by T32 HL007605.

Key Words: asthma GWAS, adult onset asthma, childhood onset asthma,

transcriptome, UK Biobank

Introduction

Asthma is the most prevalent chronic respiratory disease worldwide[1]. Its diagnosis is based on the presence of reversible airflow obstruction and clinical symptoms that include wheeze, cough, and shortness of breath. Despite these shared features, asthma is likely many different conditions. In particular, childhood onset asthma and adult onset asthma differ with respect to sex ratios, triggers of exacerbation, associated co-morbidities, severity[2,3], and potentially also for genetic risk factors[4,5]. For example, the most replicated and significant genome-wide association study (GWAS) single nucleotide polymorphisms (SNPs) at the 17q12-21 locus are specific to asthma with onset of symptoms in early life[6]. Only one asthma GWAS to date was performed in adult onset cases. The GABRIEL Consortium included 1,947 cases with adult onset asthma (defined as 16 years of age or older) and 3,669 adult controls[7]. Overall, genome-wide significant variants in the combined sample of 10,365 cases and 16,110 controls revealed larger odds ratios (ORs) in the childhood onset group compared to the adult onset group, but no loci reached significance in the adult onset cases, likely due to low power. It remains unknown, therefore, whether loci other than 17q12-21 contribute specifically to childhood onset or adult onset asthma. Delineating genetic risk factors affecting age of onset from those that are shared between childhood and adult onset asthma could provide insights into the molecular mechanisms contributing to different clinical manifestations of asthma that is diagnosed at different ages.

To address this question and directly compare genetic risk architectures of adult onset and childhood onset asthma, we leveraged the U.K. Biobank (UKB), a large-scale prospective study collecting demographic, clinical, medical history, and genetic data for

nearly 500,000 participants[8]. We performed asthma GWASs in 9,433 adults with a

diagnosis of asthma before age 12 years (childhood onset cases) and 21,564 adults

with a diagnosis of asthma between age 26 and 65 years (adult onset cases), each

compared to 318,237 adults (>38 years) who did not have a diagnosis of asthma or

chronic obstructive lung disease (COPD) (controls). In addition, we conducted an age of

onset GWAS in 37,846 asthma cases. Our goals were to identify shared and distinct

genetic risk loci for childhood and adult onset asthma, and to identify genes that may

mediate the effects of associations at age of onset specific loci and those shared

between childhood and adult onset asthma.

Methods

<u>Sample composition and case definitions</u>

Data for 376,358 British white individuals from the UKB data release July 2017 were

used[8]. We extracted disease status (asthma, allergic rhinitis, atopic dermatitis (AD),

food allergy, COPD, emphysema, chronic bronchitis), age of onset of asthma, and sex

from self-reported questionnaires and hospital records (ICD10 codes) by querying our

in-house protected UKB database server. See our 'reproducible research' and other

details in the appendix (pp. 3-12).

We defined childhood and adult onset asthma using strict age of onset criteria

that would minimize the likelihood of misclassification, and considered asthma cases

with onset < 12 years of age as childhood onset cases (n=9,433) and with onset > 25

and before 66 years of age as adult onset cases (n=21,564). UKB participants without

an asthma diagnosis were included as controls (n=318,237). Individuals with COPD,

emphysema or chronic bronchitis were excluded from adult onset cases and controls.

For the age of onset GWAS, we included an additional 6,849 asthmatics with onset

between the ages of 12 and 25 years of age. Characteristics of the sample are shown in

Table 1. After genotype quality control, 10,894,596 variants were available for analyses.

See appendix (pp. 4-5) for additional details on genotype QC and phenotype definitions.

<u>Genome-wide association studies</u>

We conducted a childhood onset and adult onset GWAS by logistic regression and an

age of onset GWAS by linear regression, both using the allele dosages under an

additive genetic model, as implemented in Hail (https://github.com/hail-is/hail). In all

three GWASs, we included sex and the first 10 genetic principal components as

covariates and used a genome-wide significance threshold of 5×10^{-8}. FUMA[9], an

integrative post-GWAS annotation web-based tool, was used to define independent risk

loci and identify enriched tissues. FUMA defines independent loci using linkage

disequilibrium (LD) information from the 1000 Genomes project[10]. We specified an r^2

threshold >0.6 between genome-wide significant SNPs to represent a single locus.

Tissue enrichments were calculated in FUMA using a hypergeometric test to determine

overrepresentation of genes mapped to risk loci (by physical distance) among highly

expressed genes in each tissue in GTEx relative to all others.

Childhood onset and adult onset specific loci were defined as those that were

genome-wide significant in either the childhood onset or adult onset GWAS but were not

associated with asthma at $p<0.05$ in other group, and the 95% confidence intervals

(CIs) of the respective ORs did not overlap. All other GWAS loci that were genome-wide

signficant in at least one of the two GWASs were considered to be shared.

<u>Sensitivity analysis to adult onset misclassification</u>

We performed sensitivity analysis to assess the effects of potential poor recall of age of

onset among the adult onset asthmatics. In particular, to assure that the adult onset

cases did not include a significant proportion of childhood onset asthma in which

symptoms remitted in early life but then relapsed in adulthood. To this end, we replaced

adult onset cases with increasing proportions of randomly selected childhood onset

cases, and then tested for association at the two most significant childhood onset

specific loci. This procedure was repeated 20 times for each proportion to quantify the

sampling variability (appendix pp. 8-10).

<u>Predicted transcriptome association test</u>

We used the PrediXcan[11] framework to identify genes that may mediate associations

between genetic variants and asthma risk. PrediXcan is a software tool that estimates

tissue-specific gene expression profiles from an individual's SNP genotype profile using

prediction models trained in large references databases of genotypes and tissue-

specific gene expression profiles. Using these genotype-imputed expression profiles,

PrediXcan can perform gene-based association tests that correlate predicted

expression levels with phenotypes (e.g., asthma) to identify candidate causal genes

from GWAS data. We used a summary version of PrediXcan, which has high

concordance with the individual level version (R^2 >0.99)[12]. For predictions, we

downloaded elastic net models trained with reference transcriptome data from the GTEx

consortium[13] (http://predictdb.org) for 49 tissues (appendix Table 1).

149   PrediXcan was run separately in the childhood onset and adult onset cases,

each with the same controls. Significance was determined using a Bonferroni correction

for the 38,608 genes ($p<1.29\times10^{-6}$) that were expressed in the five tissues determined

by FUMA to be enriched (skin, lung tissue, whole blood, small intestine and spleen; see

Results). We defined childhood onset specific genes as those whose predicted

expression is significantly associated with childhood onset asthma and the variants that

predict their expression are within childhood onset specific loci. Adult onset specific

genes were similarly defined. Because SNPs at shared loci may predict the expression

of genes that are be associated only in childhood onset or adult onset cases, we also

considered genes to be age of onset specific if they were significantly associated with

asthma at $p<1.29\times10^{-6}$ in one age group and not associated with asthma at $p<0.05$ in

the other. All other genes were considered shared.

Results

<u>Genome-wide association studies of asthma</u>

We first conducted GWASs of childhood onset and adult onset asthma. These studies

revealed 61 independent loci associated with asthma, 52 were significant in the

childhood onset asthma GWAS and 19 were significant in the adult onset asthma

GWAS ($p<5\times10^{-8}$) (Figure 1). Thirty-one of the 61 loci, in 30 chromosomal regions, were

not previously reported in the GWAS catalog or in a recent large GWAS of asthma[14,15].

Among the 31 new loci, 17 were significant in the childhood onset GWAS, one was

significant in the adult onset GWAS, and 13 were significant in both (Table 1). Some of

these loci contained genes that have been associated with asthma in candidate gene

studies (e.g., *TLR10*[16] and *TLR6*[17,18] in the childhood onset GWAS; *FADS2*[19],

MUC5AC[20,21] and *TBX*21[22] in both GWASs), which provide independent validation of

our GWAS findings.

As in previous GWASs comprised largely of children, the most significant locus in

the childhood onset GWAS is at 17q12[23], with the lead SNP in the *GSDMB* gene

(Figure 1A). However, the estimated ORs for the lead SNPs at three other loci were

similar to or larger than the lead SNP at the 17q locus. The lead SNPs in *IL1RL1* at

2q12.1 and in *EMSY* at 11q13.5 had effect sizes on childhood onset asthma similar to

the lead SNP in *GSDMB* on 17q (Table 1). Both loci have been prominent in previous

asthma GWAS. The lead SNP at the 1q21.3 locus, corresponding to a nonsense

mutation (R501X) in the filaggrin (*FLG*) gene at 1q21.3, had the largest OR overall.

Variants in *FLG* have been robustly associated with food allergies[24-26] and AD[27-29], but

previous associations with asthma have been in the context of other allergic

conditions[15,30]. Whether variants in *FLG* are associated with risk for childhood onset

asthma independent of its effects on early life allergic disease is yet unknown.

To address this possibility, we repeated the childhood onset GWAS after

excluding 3,205 childhood onset cases and 5,785 controls who reported having a

history of allergic rhinitis, AD or food allergy. As expected in a smaller sample the p-

values were overall larger, but the ORs were strikingly similar (appendix Figure 1 and

Table 2). Even though the OR at the *FLG* locus on 1q21.3 decreased from 1.97 (95% CI

1.82, 2.13) to 1.61 (95% CI 1.49, 1.74), it remained both highly significant (p=2.45x10⁻

[19]) and the largest OR for childhood onset. These results suggest both a critical role for

the allergic diasthesis in the development of asthma in childhood and a shared

architecture between allergic disease and childhood onset asthma, as previously

discussed[30,31].

The most significant association in the adult onset GWAS was in the HLA region,

with independent associations at the HLA-C/B (6p21.33) and HLA-DR/DQ (6p21.32) loci

(Figure 1B). Compared to the childhood onset GWAS, effect sizes were quite small in

adult onset cases, with ORs reaching 1.1 at only five loci (2q12.1, 6p21.33, 6p21.32,

9p24.1, 10p14) (Table 2).

Among the 61 asthma loci, 23 were specific to childhood onset asthma and one

was specific to adult onset asthma (Table 2A-B). Regional association plots for the 23

loci with childhood or adult onset specific effects are shown in appendix Figure 2.

Among the remaining 38 shared loci (Table 2C), mean ORs were larger in the childhood

onset cases at all but six loci (permutation test $p<10^{-4}$; appendix p.10-11), indicating that

both more loci contribute to childhood onset asthma and even among shared loci, effect

sizes are larger in childhood onset asthma cases (Figure 2).

Finally, to directly test for loci associated with asthma age of onset, we

conducted a third GWAS including all asthma cases in UKB who met our inclusion

criteria (n=37,846). In this analysis, 19 loci were associated with age of onset ($p<5x10^{-8}$)

(appendix Figure 3 and Table 3). Age of onset loci overlapped with both the childhood

and adult onset specific and shared loci, and asthma risk alleles were nearly all

associated with earlier age of onset (Table 2; Figure 2). SNPs a the 1q21.3 locus (*FLG*)

had the largest effect on age of onset, with each copy of the asthma risk allele

(rs61816761) associated on average with 4.57 (SE 0.43) years earlier onset compared

to individuals without the risk allele (p=8.15x10^{-27}). At the 17q12 locus (rs4795399) each

copy of the risk allele was associated on average with 2.29 (SE 0.13) years earlier

onset compared to individuals without the risk allele (p=6.76x10^{-65}). Examples of

significant age of onset effects at these and other loci are shown in Figure 3. Overall,

both childhood onset specific and shared asthma risk loci were associated with younger

ages of onset, and alleles at loci associated with younger ages of onset had larger

effects compared to alleles at loci associated with later ages of onset.

<u>Tissue-specific expression of genes at associated loci</u>

Using an unbiased approach, we asked whether the tissue-specific expression of genes

that map to the 52 childhood onset loci differed from the tissue-specific expression of

genes mapped to the 19 adult onset loci. Genes at childhood onset loci (Figure 1A)

were most highly expressed in skin, whole blood, and small intestine (lower ileum)

compared to all other tissues, whereas genes at adult onset loci (Figure 1B) were most

highly expression in lung, whole blood, small intestine (lower ileum), and spleen

(enrichment for higher expression, p<10x10^{-3}) (appendix Figure 4 and Table 4). These

patterns suggest both overlapping and distinct underlying pathways and mechanisms

associated with asthma that begins in childhood and asthma with onset in adulthood.

<u>Predicted transcriptome-wide association test</u>

To better understand molecular pathways and to narrow the list of candidate causal

genes at associated loci, we focused on the five tissues that most highly expressed the

genes at childhood onset or adult onset loci: skin, lung tissue, whole blood, small

intestine, and spleen. We used PrediXcan[11] to identify genes whose expression is

predicted by variants associated with asthma in the childhood onset or adult onset

GWAS and potentially mediate the effects of associated SNPs on asthma risk.

This analysis identified 113 unique, candidate causal genes at 22 of the 61

GWAS loci ($p<1.4 \times 10^{-6}$) (Figures 4-5) (appendix Figures 5-6 and Table 5). These

included 39 genes associated with childhood onset asthma at eight of the childhood

onset specific loci and 76 genes associated with childhood and/or adult onset asthma at

13 of the shared loci. Variants at the one adult onset specific locus at 2q22.3 did not

predict the expression of any genes in the five tissues.

The predicted genes most significantly associated with childhood onset asthma

were at the 17q12 locus (Z score >10) in skin (*ORMDL3*, *ERBB2*, *PGAP3*, *GSDMA*, 2

long noncoding RNAs), lung (*ORMDL3*, *GSDMB*, *GSDMA*, *PGAP3*, *PNMT*), blood

(*ORMDL3*, *GSDMB*, *IKZF3*, *MED24*), small intestine (*GSDMA*, *GSDMB*, *PGAP3*), and

spleen (*ORMDL3*, *GSDMB*, *ZPBP2*, *MED24*). Some genes were predicted to be more

highly expressed in asthmatics (e.g., *ORMDL3*, *GSDMB*, *PGAP3*, *ERBB2*), while others

were predicted to be less expressed in asthmatics (e.g., *GSDMA*, *MED24*, *IKZF3*)

(Figure 4A).This pattern of expression reflects the broad regulatory effects of SNPs and

tissue specificifity of gene expression at this locus[23]. The childhood onset asthma locus

at 1q21.3 includes genes essential for epidermal differentiation and maintaining

essential barrier function. The predicted expression of nine genes at this locus were

associated with childhood onset asthma. Higher predicted expression of *CRNN*, *CRCT1*

and *THEM5* in skin, of *PSDM4* in lung and blood, and of *LINGO4* in skin, lung, and

blood were associated with increased asthma risk. Lower predicted expression of

SPRR2D in skin, of *S100A12* in lung and blood, of *FLG* in skin, lung and spleen, and of

TDRKH in skin, lung, blood and spleen were associated with increased asthma risk.

S100A12 has been previously implicated in asthma[32] and *FLG* variants have been

associated with AD and food allergies, and asthma in the context of other allergic

diseases[15,24-30]. Other childhood onset specific genes previously implicated in asthma

but not previously reported in asthma GWASs are *CCL20*[33] at 2q36.3 and *TLR10*[16] at

4p14 in whole blood, and *TLR6*[17,18] at 4p14, *AP5B1* at 11q13.1 and *SERPINB7*[34] at

18q21.33 in skin.

The 5q31.1 region had independent loci that were both childhood onset specific

and shared in the GWASs. Although the predicted expression of all eight asthma genes

at this extended locus were shared, five genes were more significantly associated with

childhood onset asthma: higher predicted expression of *RAD50* in skin but lower

predicted expression of *SEPT8* in skin and lung, *IL4* in skin, lung and blood, and *AFF4*

small intestine were associated with increased risk for asthma (Figure 4B). The

remaining three genes had similar associations with childhood and adult onset asthma,

with predicted lower expression of *IRF1* in skin and spleen and predicted higher

expression of *PDLIM4* in skin and *SLC22A5* in all five tissues associated with increased

asthma risk. *IL4, RAD50, SLC22A5* and *PDLIM4* have been highlighted in previous

asthma GWAS[35,36], and *KIF3A* was identified in a GWAS of the atopic march[37] and

associated with childhood onset asthma in a candidate gene study[38].

In contrast to all other loci, predicted expression of 44 genes at two independent

shared loci in the HLA region (6p21.32 and 6p21.33; referred to as the HLA region from

hereon in) were associated with childhood asthma only (n=1; in skin and lung), adult

onset asthma only (n=3; in skin only), or both (n=39; in multiple tissues). (Figure 5). The

shear number of genes in this region with predicted expression associated with asthma,

the generally broad tissue expression patterns, and three associated with asthma only

in adult onset cases are consistent with this locus being among the two most signficant

loci in nearly all asthma GWASs, and the most significant locus in a previous small

GWASs of adult onset asthma[7] and in adults with asthma[36,39].

Among the remaining 37 shared loci (Figure 2B), SNPs at 12 predicted the

expression of 23 unique genes, all of which were associated with both childhood onset

and adult onset asthma. These include *IL18R1*, *IL18RAP*, and *ILRL2* at 2q12.1 in

multiple tissues, *TSLP* at 5q22.1 in skin, *SMAD3* at 15q22.33 in skin and blood, *LRP1* at

12q13.3 in skin, *IL4R* at 16p12.1 in blood, and *CLEC16A* at 16p13.13 in lung. Loci

associated with *IL18R1*, *IL18RAP*, *IL18RAP*, *ILRL2, TSLP, SMAD3*, *LRP1*, *IL4R* and

CLEC16A were reported in previous asthma GWAS[15,35,40].

Discussion

This study is the first large GWAS of age of onset of asthma to include both childhood

and adult onset cases. These GWASs revealed 61 independent asthma loci, 23 specific

to childhood onset, one specific to adult onset, and 37 shared; with overall larger effect

sizes for childhood onset asthma at nearly all loci. Moreover, the predicted expression

of 41 of the 113 implicated genes were associated specifically with childhood onset

asthma, compared to the predicted expression of three genes associated specifically

with adult onset asthma. Our findings of more childhood onset asthma loci and

potentially causal genes and the larger effect sizes of risk alleles in childhood onset

cases are particularly striking given that there were nearly 2.5-times more adult onset

than childhood onset cases in this study. Thus, despite having substantially less power

to detect loci specific to childhood onset asthma, our analyses revealed many more

childhood onset asthma loci. Similarly, the asthma risk alleles at 19 loci identified in the

age of onset GWAS were all associated with younger age of onset. These findings are

consistent with decreased estimates of asthma heritability with increasing age of

onset[41]. Taken together, our study shows for the first time that genetic risk for adult

onset asthma is largely a subset of the genetic risk loci for childhood onset asthma, but

with overall smaller effect sizes, suggesting a larger role for environmental risk factors in

adult onset asthma.

Despite the overlap of adult onset and childhood onset loci, distinct causal

pathways contributing to each were suggested by tissue enrichments: childhood onset

loci were enriched for genes with highest expression in skin whereas adult onset loci

were enriched for genes with highest expression in lung and spleen; both were enriched

for genes highly expressed in whole blood and small intestine. The highlighting of skin

as a target tissue for childhood onset asthma supports the widely held idea that asthma

in childhood is due to impaired barrier function in the skin and other epithelial surfaces.

This model proposes that compromised epithelial barriers promote sensitization to food

and airway allergens and to wheezing illnesses in early life[31,42]. In fact, childhood onset

specific loci identified here have been associated with AD or food allergies, such as

FLG on 1q21.3 with the atopic march[37], AD[27-29] and food allergies[24-26], *KIF3A* on 5q31.1

and *AP5B1*/*OVOL1* on 11q13.1 with the the atopic march[37] and AD[43], *SERPINB7* on

18q21.33 with food allergies[34], and *CRNN* (cornulin) on 1q21.3 with AD concomitant

with asthma and reduced expression in AD-affected skin[44]. Variants at those loci were

all associated with earlier age of asthma onset. We further show that these loci are

associated with childhood onset asthma, even after exclusion of cases with a history of

allergic diseases. In contrast, the enrichment for genes highly expressed in lung and

spleen at adult onset loci suggests a more lung-centered, and potentially immune

mediated, etiology for asthma with onset later in life. The prominant role of the HLA

region in the adult onset asthma GWAS and the fact that predicted expression of three

HLA region genes was associated only with adult onset asthma only further highlights a

central role for immune processes driving asthma pathogenesis in adults. The fact that

both childhood onset and adult onset asthma loci were enriched for genes that are most

highly expressed in whole blood cells and small intestine further indicate a shared

immune etiology, as suggested from a large GWAS that included both children and

adults[35].

Combining GWAS with a transcriptome-wide association test that uses

combinations of associated SNPs to predict gene expression in different tissues

revealed significant complexity at the two most highly associated asthma loci. SNPs at

the 17q12 locus predicted expression of 18 childhood onset asthma genes and SNPs at

the HLA region predicted expression of 42 genes: three were associated with adult

onset asthma and most were not HLA genes *per se*. In this regard, it is notable that the

HLA-DRB1, *HLA-DQB1*, and *HLA-DQA1* genes, which are the most associated HLA

loci with autoimmune diseases, are predicted to have reduced expression in both

childhood onset and adult onset asthma. Instead, HLA genes with less clear functions

have increased predicted expression in asthma (Figure 5). These results strengthen the

argument that multiple genes contribute to asthma risk at the HLA and 17q12 loci and

probably account for the highly significant GWAS p-values observed at these loci in

nearly all studies. It is also likely that these genes have both tissue specific and broad

effects in epithelium, lung, and immune tissues.

The new loci identified in our study include the first adult onset asthma specific

association at 2q22.3. The lead SNP at 2q22.3 is intergenic between *TEX41* and

ACVR2A. The predicted expression of *ACVR2A* was not associated with asthma in our

study, despite it being expressed in lung, blood, small intestine and spleen. *TEX41* was

not expressed in any of the five tissues investigated. Interestingly, a GWAS also

performed in UKB subjects implicated variants near *TEX41* in heavy vs. never smoking

behavior[45]. However, even after removing adult onset cases and controls with reported

'ever smoking', the p-value for this SNP remained significant and the OR slightly

increased (OR 1.077 [95% CI 1.05, 1.11], p=2.26×10^{-8}; n=12,132 cases and 176,704

controls). Variants in or near this gene, which encodes a lincRNA, have been

associated with cardiovascular and immune mediated traits[14], making this a potentially

interesting candidate gene for adult onset asthma.

Our study had limitations. First, diagnoses of asthma and allergic disease in

study subjects were from self reported questionnaires and medical records (ICD10

codes). Thus, it is possible that diagnoses, age of onset, or both are misspecified in

some subjects. On the one hand, the large sample size and our ability to replicate

nearly all previously reported asthma loci (appendix Table 6) suggest that our analyses

were robust to any inaccuracies in the data. On the other hand, it is possible that the

adult onset asthmatics included cases with poor recall of childhood onset asthma in

which symptoms remitted and then relapsed later in life[46]. Our sensitivity analysis

suggested that if even as few as 5% of the adult onset cases were misclassified we

should have observed some signal of association at childhood onset loci, which we did

not. Second, the gene expression data used to predict candidate target genes included

heterogeneous tissues and were collected mostly from adults. As a result, our study

may have missed relevant genes whose expression is developmentally regulated or

environment specific. Our finding of candidate genes at only 22 of the 61 asthma loci

may be due in part to the importance of both in asthma pathogenesis. Moreover, all

inference based on gene expression is using imputed expression. It is possible,

therefore, that some relevant genes were more difficult to impute and not included in our

analysis, although a recent comparative study showed that PrediXcan is a more robust

method for prediction of gene expression than other related methods[47]. Third, because

of the ethnic composition of the UKB, this study was limited to individuals of European

ancestry only. As a result, we could not evaluate the genetic risk architecture or assess

the effects of age of onset specific loci in other populations. Lastly, although many of the

new loci discovered in our GWASs include genes previously implicated in asthma, the

new genetic associations reported here need to be replicated in other populations.

In the largest asthma GWAS to date, we show that genetic risk loci for adult

onset asthma is largely a subset of the loci associated with childhood onset asthma,

with overall smaller effect sizes for onset at later ages. These data suggest that

childhood onset specific loci and those associated with age of onset play a role in

disease initiation, whereas the other associated loci reflect shared pathways of disease

progression. The differences in the target tissues that most highly express the genes at

associated loci and the predicted expression of genes at age specific and shared loci

provides additional genetic and molecular evidence for both shared and distinct

pathogenic pathways in childhood onset and adult onset asthma. It is therefore likely

that the most effective treatments will also differ between these two groups, and that

strategies for precision medicine should be further personalized to account for age of

asthma onset.

Author's Contributions

All authors were involved in the conception and design of the study and in writing the

manuscript. M. P. and N.S. conducted all analyses and prepared figures and tables,

under the overall supervision of D.L.N., C.O., and H.K.I.

URLs

UK Biobank, [https://www.ukbiobank.ac.uk/]

Hail: https://github.com/hail-is/hail

Acknowledgements

References

1.   Collaborators GBDCRD. Global, regional, and national deaths, prevalence,

disability-adjusted life years, and years lived with disability for chronic obstructive

pulmonary disease and asthma, 1990-2015: a systematic analysis for the Global

Burden of Disease Study 2015. *Lancet Respir Med* 2017; **5**(9): 691-706.

2.   Bush A, Menzies-Gow A. Phenotypic differences between pediatric and adult

asthma. *Proc Am Thorac Soc* 2009; **6**(8): 712-9.

3.   Busse W, Banks-Schlegel SP, Larsen GL. Childhood- versus adult-onset

asthma. *Am J Respir Crit Care Med* 1995; **151**(5): 1635-9.

4.   Dijk FN, de Jongste JC, Postma DS, Koppelman GH. Genetics of onset of

asthma. *Curr Opin Allergy Clin Immunol* 2013; **13**(2): 193-202.

5.   Holloway JW, Arshad SH, Holgate ST. Using genetics to predict the natural

history of asthma? *J Allergy Clin Immunol* 2010; **126**(2): 200-9; quiz 10-1.

6.   Bouzigon E, Corda E, Aschard H, et al. Effect of 17q21 variants and smoking

exposure in early-onset asthma. *N Engl J Med* 2008; **359**(19): 1985-94.

7.   Moffatt MF, Gut IG, Demenais F, et al. A large-scale, consortium-based

genomewide association study of asthma. *N Engl J Med* 2010; **363**(13): 1211-21.

8.   Bycroft C, Freeman C, Petkova D, et al. Genome-wide genetic data on ~500,000

UK Biobank Participants. *bioRxiv* 2017.

9.   Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping

and annotation of genetic associations with FUMA. *Nature communications* 2017; **8**(1):

1826.

10.   Consortium TGP, Auton A, Brooks LD, et al. A global reference for human

genetic variation. *Nature* 2015; **526**(7571): 68-74.

11.   Gamazon ER, Wheeler HE, Shah KP, et al. A gene-based association method

for mapping traits using reference transcriptome data. *Nat Genet* 2015; **47**(9): 1091-8.

12.   Barbeira AN, Dickinson SP, Bonazzola R, et al. Exploring the phenotypic

consequences of tissue specific gene expression variation inferred from GWAS

summary statistics. *Nature communications* 2018; **9**(1): 1825.

13.   Battle A, Brown CD, Engelhardt BE, Montgomery SB. Genetic effects on gene

expression across human tissues. *Nature* 2017; **550**(7675): 204-13.

14.   MacArthur DG, Balasubramanian S, Frankish A, et al. A systematic survey of

loss-of-function variants in human protein-coding genes. *Science* 2012; **335**(6070): 823-

8.

15.   Zhu Z, Lee PH, Chaffin MD, et al. A genome-wide cross-trait analysis from UK

Biobank highlights the shared genetic architecture of asthma and allergic diseases. *Nat*

Genet 2018.

16.   Lazarus R, Raby BA, Lange C, et al. TOLL-like receptor 10 genetic variation is

associated with asthma in two independent samples. *Am J Respir Crit Care Med* 2004;

170(6): 594-600.

17.   Tantisira K, Klimecki WT, Lazarus R, et al. Toll-like receptor 6 gene (TLR6):

single-nucleotide polymorphism frequencies and preliminary association with the

diagnosis of asthma. *Genes Immun* 2004; **5**(5): 343-6.

18.   Lau MY, Dharmage SC, Burgess JA, et al. The interaction between farming/rural

environment and TLR2, TLR4, TLR6 and CD14 genetic polymorphisms in relation to

early- and late-onset asthma. *Sci Rep* 2017; **7**: 43681.

19.   Standl M, Sausenthaler S, Lattka E, et al. FADS gene cluster modulates the

effect of breastfeeding on asthma. Results from the GINIplus and LISAplus studies.

Allergy 2012; **67**(1): 83-90.

20.   Rousseau K, Byrne C, Griesinger G, et al. Allelic association and recombination

hotspots in the mucin gene (MUC) complex on chromosome 11p15.5. *Ann Hum Genet*

2007; **71**(Pt 5): 561-9.

21.   Johnson L, Shah I, Loh AX, et al. MUC5AC and inflammatory mediators

associated with respiratory outcomes in the British 1946 birth cohort. *Respirology* 2013;

18(6): 1003-10.

22.   Akahoshi M, Obara K, Hirota T, et al. Functional promoter polymorphism in the

TBX21 gene associated with aspirin-induced asthma. *Human genetics* 2005; **117**(1):

16-26.

23.   Stein MM, Thompson EE, Schoettler N, et al. A decade of research on the

17q12-21 asthma locus: Piecing together the puzzle. *J Allergy Clin Immunol* 2018;

142(3): 749-64 e3.

24.   Brown SJ, Asai Y, Cordell HJ, et al. Loss-of-function variants in the filaggrin gene

are a significant risk factor for peanut allergy. *J Allergy Clin Immunol* 2011; **127**(3): 661-

7.

25.   Marenholz I, Kerscher T, Bauerfeind A, et al. An interaction between filaggrin

mutations and early food sensitization improves the prediction of childhood asthma. *J*

Allergy Clin Immunol 2009; **123**(4): 911-6.

26.   Venkataraman D, Soto-Ramirez N, Kurukulaaratchy RJ, et al. Filaggrin loss-of-

function mutations are associated with food allergy in childhood and adolescence. *J*

Allergy Clin Immunol 2014; **134**(4): 876-82 e4.

27.   Bisgaard H, Halkjaer LB, Hinge R, et al. Risk analysis of early childhood eczema.

J Allergy Clin Immunol 2009; **123**(6): 1355-60 e5.

28.   Palmer CN, Irvine AD, Terron-Kwiatkowski A, et al. Common loss-of-function

variants of the epidermal barrier protein filaggrin are a major predisposing factor for

atopic dermatitis. *Nat Genet* 2006; **38**(4): 441-6.

29.   Weidinger S, Illig T, Baurecht H, et al. Loss-of-function variations within the

filaggrin gene predispose for atopic dermatitis with allergic sensitizations. *J Allergy Clin*

Immunol 2006; **118**(1): 214-9.

30.   Ferreira MA, Vonk JM, Baurecht H, et al. Shared genetic origin of asthma, hay

fever and eczema elucidates allergic disease biology. *Nat Genet* 2017; **49**(12): 1752-7.

31.   Busse WW. The atopic march: Fact or folklore? *Ann Allergy Asthma Immunol*

2018; **120**(2): 116-8.

32.   Yang Z, Yan WX, Cai H, et al. S100A12 provokes mast cell activation: a potential

amplification pathway in asthma and innate immunity. *J Allergy Clin Immunol* 2007;

119(1): 106-14.

33. Collison A, Hatchwell L, Verrills N, et al. The E3 ubiquitin ligase midline 1

promotes allergen and rhinovirus-induced asthma by inhibiting protein phosphatase 2A

activity. *Nat Med* 2013; **19**(2): 232-7.

34. Marenholz I, Grosche S, Kalb B, et al. Genome-wide association study identifies

the SERPINB gene cluster as a susceptibility locus for food allergy. *Nature*

communications 2017; **8**(1): 1056.

35. Demenais F, Margaritte-Jeannin P, Barnes KC, et al. Multiancestry association

study identifies new asthma risk loci that colocalize with immune-cell enhancer marks.

Nat Genet 2018; **50**(1): 42-53.

36. Li X, Howard TD, Zheng SL, et al. Genome-wide association study of asthma

identifies RAD50-IL13 and HLA-DR/DQ regions. *J Allergy Clin Immunol* 2010; **125**(2):

328-35 e11.

37. Marenholz I, Esparza-Gordillo J, Ruschendorf F, et al. Meta-analysis identifies

seven susceptibility loci involved in the atopic march. *Nature communications* 2015; **6**:

8804.

38. Kovacic MB, Myers JM, Wang N, et al. Identification of KIF3A as a novel

candidate gene for childhood asthma using RNA expression and population allelic

frequencies differences. *PloS one* 2011; **6**(8): e23714.

39. Lasky-Su J, Himes BE, Raby BA, et al. HLA-DQ strikes again: genome-wide

association study further confirms HLA-DQ in the diagnosis of asthma among adults.

Clin Exp Allergy 2012; **42**(12): 1724-33.

40.   Pickrell JK, Berisa T, Liu JZ, Segurel L, Tung JY, Hinds DA. Detection and

interpretation of shared genetic influences on 42 human traits. *Nat Genet* 2016; **48**(7):

709-17.

41.   Thomsen SF, Duffy DL, Kyvik KO, Backer V. Genetic influence on the age at

onset of asthma: a twin study. *J Allergy Clin Immunol* 2010; **126**(3): 626-30.

42.   Han H, Roan F, Ziegler SF. The atopic march: current insights into skin barrier

dysfunction and epithelial cell-derived cytokines. *Immunol Rev* 2017; **278**(1): 116-30.

43.   Johansson E, Biagini Myers JM, Martin LJ, et al. KIF3A genetic variation is

associated with pediatric asthma in the presence of eczema independent of allergic

rhinitis. *J Allergy Clin Immunol* 2017; **140**(2): 595-8 e5.

44.   Trzeciak M, Sakowicz-Burkiewicz M, Wesserling M, et al. Altered Expression of

Genes Encoding Cornulin and Repetin in Atopic Dermatitis. *Int Arch Allergy Immunol*

2017; **172**(1): 11-9.

45.   Wain LV, Shrine N, Miller S, et al. Novel insights into the genetics of smoking

behaviour, lung function, and chronic obstructive pulmonary disease (UK BiLEVE): a

genetic association study in UK Biobank. *Lancet Respir Med* 2015; **3**(10): 769-81.

46.   Sears MR, Greene JM, Willan AR, et al. A longitudinal, population-based, cohort

study of childhood asthma followed to adulthood. *N Engl J Med* 2003; **349**(15): 1414-22.

47.   Fryett JJ, Inshaw J, Morris AP, Cordell HJ. Comparison of methods for

transcriptome imputation through application to two common complex diseases. *Eur J*

Hum Genet 2018.

Table 1. Characteristics of the asthma cases and controls. NA, not applicable.

	Childhood Onset (N=9,433)	Adolescent or Young Adult Onset (N=6,849)	Adulthood Onset (N=21,564)	Controls (N=318,237)
Age at recruitment (yr)				
Mean (±SD)	55±8	52±8	57±8	57±8
Range	40 - 70	40 - 70	40 - 70	39 - 73
Age at asthma onset (yr)[a]				
Mean (±SD)	6±3	19±4	44±10	NA
Range	0 - 11	12 - 25	26 - 65	NA
Sex (% female)	40.7	57.4	63.6	53.5
Asthma medication use in the past 12 months, N (%)	1,383 (14.7)	1,124 (16.4)	4,081 (18.9)	NA
Current smokers N (%)	802 (8.5)	688 (10.0)	1,379 (6.4)	30,056 (9.4)
Allergic diseases ever				
Allergic rhinitis, N (%)	2,537 (26.9)	2,042 (29.8)	4,660 (21.6)	27,289 (8.6)
Atopic dermatitis, N (%)	1,126 (11.9)	473 (6.9)	927 (4.3)	7,168 (2.3)
Food allergy, N (%)	138 (1.5)	63 (0.9)	172 (0.8)	1,229 (0.4)
Any allergic disease, N (%)	3,205 (34.0)	2,321 (33.9)	5,306 (24.6)	33,808 (10.6)

Table 2. Regions with SNPs that were genome-wide significant in the childhood onset or adult onset GWAS. Information for the SNP with smallest p-value at each locus is shown.

Locus[a]	rsID	Position[b]	Nearby Genes[c]	Allele[d]	RAF[e]	Childhood Onset Asthma GWAS			Adult Onset Asthma GWAS			Age of Onset GWAS		
						OR	95% CI	p-value	OR	95% CI	p-value	Beta	SE	p-value
A. Childhood Onset Specific Loci														
1q21.3	rs61816761	1:152285861	*FLG, HRNR, FLG2*	G/A	0.024	1.970	1.823-2.129	1.88E-65	1.016	0.948-1.088	6.56E-01	-4.571	0.426	8.15E-27
1q25.1	rs7518129	1:173163568	*TNFSF4, TNSF18, PRDX6*	A/G	0.310	1.111	1.077-1.146	2.21E-11	0.997	0.977-1.019	8.17E-01	-0.849	0.145	4.89E-09
2q36.3[f]	rs10175070	2:228670575	*CCL20, SLC18A3*	A/G	0.251	1.122	1.086-1.160	4.31E-12	1.000	0.978-1.023	9.92E-01	-0.862	0.154	2.24E-08
3q28	rs12634152	3:188121019	*LPP, FLJ42393, LPP-AS1*	C/T	0.547	1.133	1.100-1.166	1.08E-16	1.007	0.987-1.027	4.98E-01	-0.968	0.136	1.21E-12
4p14	rs5743618	4:38798648	*TLR1, TLR10, TLR6*	A/C	0.774	1.249	1.203-1.295	3.19E-32	1.008	0.985-1.032	5.09E-01	-1.578	0.163	4.53E-22
5q13.2[f]	rs10036789	5:71695918	*PTCD2, ZNF366*	C/G	0.460	1.085	1.054-1.117	4.51E-08	1.016	0.996-1.036	1.12E-01	-0.511	0.137	1.95E-04
5q31.1[f]	rs2051809	5:132056874	*KIF3A, IL4, CCNI2*	C/A	0.247	1.175	1.137-1.214	3.24E-22	1.016	0.994-1.040	1.58E-01	-0.933	0.155	1.59E-09
6p25.3[f]	rs9391997	6:409119	*IRF4, DUSP22, EXOC2*	A/G	0.527	1.102	1.070-1.135	6.89E-11	0.995	0.976-1.015	6.35E-01	-0.597	0.136	1.06E-05
7p15.3[f]	rs34880821	7:2277545	*IL6, TOMM7*	G/A	0.283	1.106	1.071-1.141	4.73E-10	1.000	0.979-1.022	9.88E-01	-0.667	0.149	7.59E-06
8q24.21[f]	rs13277355	8:128777719	*MYC, TMEM75*	G/A	0.274	1.110	1.075-1.146	1.65E-10	1.013	0.991-1.036	2.46E-01	-0.660	0.151	1.20E-05
9q34.3[f]	rs117137535	9:140500443	*ARRDC1, ZMYND19, EHMT1*	G/A	0.026	1.307	1.195-1.429	4.16E-09	0.977	0.913-1.046	5.10E-01	-2.464	0.446	3.42E-08
10p15.1[f]	rs943451	10:6621773	*PRKCQ, PFKFB3, SFMBT2*	C/T	0.318	1.125	1.091-1.161	7.51E-14	1.021	1.000-1.043	5.30E-02	-0.613	0.145	2.54E-05
11q13.1[f]	rs479844	11:65551957	*AP5B1, OVOL1*	A/G	0.555	1.106	1.074-1.139	1.48E-11	1.015	0.995-1.035	1.41E-01	-0.657	0.135	1.22E-06
11q23.3[f]	rs12365699	11:118743286	*CXCR5, DDX6*	A/G	0.833	1.167	1.120-1.216	1.57E-13	1.004	0.978-1.031	7.43E-01	-1.396	0.185	4.42E-14
12q13.2[f]	rs62623446	12:55368291	*TESPA1, MUCL1, NEUROD4*	C/T	0.072	1.188	1.127-1.252	1.40E-10	1.038	0.999-1.077	5.32E-02	-1.079	0.253	1.95E-05
12q24.12[f]	rs10774625	12:111910219	*ATXN2, SH2B3, BRAP*	A/G	0.504	1.087	1.056-1.119	2.07E-08	0.997	0.978-1.017	7.92E-01	-0.713	0.135	1.43E-07
12q24.31[f]	rs1696361	12:121363823	*SPPL3, HNF1A*	C/T	0.359	1.109	1.076-1.142	1.19E-11	1.019	0.998-1.040	6.96E-02	-0.696	0.140	6.84E-07
17q12	rs4795399	17:38061439	*GSDMB, ZPBP2, ORMDL3*	C/T	0.529	1.406	1.365-1.448	1.45E-111	1.005	0.985-1.025	6.28E-01	-2.286	0.134	6.76E-65
17q21.2[f]	rs8066625	17:40390629	*STAT5B, GHDC, STAT5A*	G/A	0.100	1.151	1.098-1.206	4.90E-09	1.006	0.973-1.040	7.33E-01	-0.846	0.227	1.88E-04
17q21.32[f]	rs56308324	17:45819206	*TBX21, TBKBP1, OSBPL7*	A/T	0.132	1.132	1.086-1.180	3.47E-09	1.025	0.996-1.055	8.72E-02	-0.714	0.195	2.54E-04
18q21.33[f]	rs4574025	18:60009814	*TNFRSF11A, KIAA1468, ZCCHC2*	C/T	0.535	1.090	1.058-1.122	9.25E-09	0.986	0.967-1.006	1.69E-01	-0.871	0.136	1.61E-10
18q21.33[f]	rs12964116	18:61442619	*SERPINB7, SERPINB11, SERPINB2*	A/G	0.036	1.336	1.247-1.432	2.56E-16	1.005	0.953-1.059	8.62E-01	-1.916	0.351	4.87E-08
19p13.3[f]	rs4807630	19:1170445	*SBNO2, GPX4, STK11*	C/T	0.308	1.093	1.060-1.128	2.06E-08	1.013	0.992-1.035	2.32E-01	-0.487	0.147	9.25E-04
B. Adult Onset Specific Loci														
2q22.3[f]	rs12617922	2:146156679	*TEX41, ACVR2A*	A/G	0.518	0.983	0.955-1.012	2.36E-01	1.066	1.045-1.087	1.59E-10	0.478	0.136	4.46E-04
C. Shared Asthma Loci														
1q32.1	rs12023876	1:203093201	*ADORA1, PPFIA4*	T/G	0.668	1.102	1.068-1.137	1.11E-09	1.032	1.010-1.053	3.37E-03	-0.567	0.144	8.65E-05
2p25.1	rs13416555	2:8441735	*RNF144A, ID2*	G/C	0.705	1.130	1.094-1.167	1.94E-13	1.040	1.018-1.063	3.50E-04	-0.715	0.150	1.91E-06
2q12.1	rs72823641	2:102936159	*IL1RL1, IL1RL2, IL18R1*	A/T	0.863	1.415	1.349-1.484	3.01E-46	1.109	1.077-1.142	3.72E-12	-1.687	0.208	4.73E-16
2q37.3	rs34290285	2:242698640	*D2HGDH, ING5, GAL3ST2*	A/G	0.745	1.177	1.137-1.219	2.06E-20	1.081	1.056-1.106	2.16E-11	-0.821	0.159	2.56E-07
3p22.3[f]	rs35570272	3:33047662	*GLB1, TRIM71, TMPPE*	G/T	0.396	1.100	1.068-1.133	2.68E-10	1.046	1.025-1.067	9.84E-06	-0.454	0.138	1.03E-03
4q27	rs2069763	4:123377482	*IL2, ADAD1, IL21*	C/A	0.334	1.126	1.092-1.160	1.59E-14	1.032	1.011-1.054	2.44E-03	-0.631	0.142	8.39E-06
5p15.2[f]	rs16903574	5:14610309	*FAM105A, TRIO, OTULIN*	C/G	0.077	1.236	1.174-1.302	9.01E-16	1.052	1.013-1.091	7.75E-03	-1.055	0.250	2.51E-05
5q22.1	rs1837253	5:110401872	*SLC25A46, TSLP*	T/C	0.740	1.211	1.169-1.253	2.33E-27	1.088	1.064-1.113	2.77E-13	-0.733	0.158	3.72E-06
5q31.1	rs17622378	5:131778452	*C5orf56, SLC22A5, IRF1*	G/A	0.573	1.103	1.071-1.136	8.52E-11	1.077	1.055-1.098	3.17E-13	-0.044	0.137	7.46E-01
6p22.1[f]	rs1117490	6:30170510	*TRIM26, TRIM15, TRIM39*	T/C	0.234	1.100	1.063-1.137	2.78E-08	1.059	1.036-1.084	6.83E-07	-0.453	0.158	4.07E-03
6p21.33	rs2428494	6:31322197	*HLA-B, HLA-C, MICA*	T/A	0.477	1.157	1.124-1.191	7.94E-23	1.104	1.082-1.126	4.77E-23	-0.648	0.136	1.78E-06
6p21.32	rs28407950	6:32626348	*HLA-DQA1, HLA-DQB*	T/C	0.756	1.354	1.306-1.405	1.27E-59	1.148	1.121-1.175	7.67E-31	-1.284	0.163	4.03E-15
6q15	rs1321859	6:91011673	*BACH2, MAP3K7*	T/C	0.649	1.102	1.068-1.136	9.32E-10	1.076	1.054-1.098	5.68E-12	-0.268	0.144	6.28E-02
7p21.1[f]	rs4473914	7:20426263	*ITGB6, MACC1, ABCB5*	C/T	0.604	1.092	1.060-1.125	9.02E-09	1.035	1.014-1.056	8.44E-04	-0.338	0.139	1.51E-02
7p15.1[f]	rs917115	7:28172586	*JAZF1, TAX1BP1, CREB5*	T/C	0.208	1.112	1.074-1.152	1.73E-09	1.042	1.017-1.067	7.35E-04	-0.577	0.164	4.17E-04
8q21.13	rs4739738	8:81291645	*TPD52, ZBTB10*	A/G	0.359	1.117	1.084-1.151	3.45E-13	1.057	1.036-1.079	8.63E-08	-0.450	0.140	1.27E-03
9p24.1	rs992969	9:6209697	*RANBP6, IL33*	G/A	0.252	1.248	1.209-1.289	6.76E-42	1.100	1.076-1.125	3.14E-17	-1.026	0.152	1.35E-11
10p14[f]	rs7894791	10:8591369	*GATA3, CELF2*	A/C	0.587	1.103	1.071-1.137	9.43E-11	1.024	1.004-1.045	1.94E-02	-0.494	0.137	3.24E-04
10p14	rs1775554	10:9054340	*GATA3, CELF2*	C/A	0.577	1.117	1.084-1.150	2.85E-13	1.121	1.098-1.143	5.17E-29	-0.002	0.138	9.91E-01
11p15.5[f]	rs12788104	11:1123739	*MUC6, MUC5AC*	A/G	0.688	1.029	0.997-1.061	7.99E-02	1.064	1.041-1.087	1.41E-08	0.147	0.148	3.22E-01
11q12.2[f]	rs174621	11:61630104	*FADS2, FADS1, FADS3*	A/G	0.772	1.013	0.978-1.049	4.63E-01	1.071	1.046-1.097	1.46E-08	0.508	0.163	1.87E-03
11q13.5[g]	rs61894547	11:76248630	*EMSY, THAP12, LRRC32*	C/T	0.052	1.463	1.382-1.548	2.17E-39	1.093	1.047-1.141	5.46E-05	-2.227	0.284	4.42E-15

	rs7936312	11:76293726	*EMSY, LRRC32*	G/T	0.477	1.194	1.160-1.230	4.89E-33	1.060	1.040-1.081	4.10E-09	-0.978	0.134	3.49E-13
12q13.11	rs56389811	12:48205358	*HDAC7, SLC48A1, VDR*	T/C	0.761	1.022	0.988-1.058	2.11E-01	1.078	1.054-1.104	2.36E-10	0.485	0.161	2.55E-03
12q13.2	rs705699	12:56384804	*RAB5B, CDK2, SUOX*	G/A	0.425	1.106	1.074-1.139	1.43E-11	1.052	1.031-1.072	6.10E-07	-0.481	0.136	4.20E-04
12q13.3	rs3122929	12:57509102	*STAT6, NAB2, LRP1*	C/T	0.404	1.134	1.101-1.167	6.59E-17	1.061	1.040-1.082	5.73E-09	-0.564	0.138	4.07E-05
12q21.1 [f]	rs11178648	12:71533210	*TSPAN8, PTPRR, LGR5*	T/C	0.592	1.087	1.055-1.120	4.16E-08	1.052	1.031-1.073	6.62E-07	-0.183	0.138	1.86E-01
13q32.3	rs1887704	13:99974492	*UBAC2, DOCK9, TM9SF2*	C/G	0.681	1.124	1.089-1.161	5.48E-13	1.048	1.026-1.070	1.57E-05	-0.476	0.147	1.15E-03
14q24.1	rs1950897	14:68760141	*RAD51B, ZFYVE26, ZFF36L1*	T/C	0.287	1.096	1.062-1.131	1.38E-08	1.031	1.009-1.053	5.39E-03	-0.520	0.148	4.52E-04
15q22.2	rs11071559	15:61069988	*RORA, ANXA2, VPS13C*	T/C	0.872	1.205	1.151-1.262	2.57E-15	1.069	1.037-1.101	1.22E-05	-0.958	0.210	5.11E-06
15q22.33	rs56062135	15:67455630	*SMAD3, SMAD6, AAGAB*	C/T	0.237	1.194	1.156-1.234	2.97E-26	1.083	1.058-1.108	6.27E-12	-0.681	0.155	1.10E-05
16p13.13	rs35032408	16:11215424	*CLEC16A, CIITA, RMI2*	G/T	0.785	1.152	1.111-1.195	4.59E-14	1.073	1.048-1.100	1.10E-08	-0.568	0.169	7.91E-04
16p12.1	rs3785356	16:27349168	*IL4R, NSMCE, IL21R*	C/T	0.297	1.122	1.087-1.157	5.69E-13	1.036	1.014-1.058	1.24E-03	-0.641	0.147	1.30E-05
16q12.1	rs2066844	16:50745926	*NOD2, SNX20, CYLD*	C/T	0.048	1.194	1.121-1.272	3.52E-08	1.049	1.003-1.097	3.76E-02	-0.743	0.306	1.54E-02
17q21.33	rs28406364	17:47454507	*ZNF652, PHB*	C/T	0.377	1.108	1.076-1.142	1.39E-11	1.036	1.015-1.057	6.55E-04	-0.489	0.140	4.81E-04
19q13.11	rs10414065	19:33721455	*LRP3, CEBPA*	T/C	0.934	1.207	1.133-1.287	6.37E-09	1.108	1.064-1.155	1.01E-06	-1.078	0.291	2.12E-04
21q22.12	rs11088309	21:36464631	*RUNX1, SETD4*	C/G	0.143	1.039	0.997-1.083	6.82E-02	1.079	1.050-1.109	4.83E-08	0.063	0.189	7.38E-01

[a]Cytogenetic band. [b]SNP position, Genome Reference Consortium Build 37 (hg19). [c]The gene in which the SNP is located is indicated first, followed by the previous gene and the next gene; for intergenic SNPs, only the previous and next genes are shown. [d]Alleles are shown as non-risk/risk alleles, where the risk allele is the allele associated with increased asthma risk. [e]RAF, risk allele frequency in the UK Biobank. [f]Not reported in previous GWAS. [g]The SNP with the smallest p-value differs in the childhood onset GWAS and the adult onset GWAS at the same locus; both SNPs are shown.

Figure Legends

Figure 1. GWAS of childhood onset and adult onset asthma. Miami plot showing results for the childhood onset versus controls GWAS (blue, panel A) and adult onset versus controls GWAS (red, panel B). Each point corresponds to a SNP; the y-axes show the $-\log_{10}$p-values from the childhood onset GWAS (panel A) and adult onset GWAS (panel B). The x-axis shows the position of each SNP along the 22 autosomes.

Figure 2. Forest plot showing the odds ratios (ORs) and 95% confidence intervals from the childhood onset (blue) and adult onset (red) GWASs, and betas and standard errors (gray) from the age of onset GWAS for the 61 asthma associated loci. Left panel: Childhood onset specific loci (top) and adult onset specific locus (bottom); right panel: Shared loci. Loci within each group are sorted by OR in the childhood onset GWAS (largest to smallest).

Figure 3. Age of onset effects (ORs) for lead SNPs at three genome-wide signficant childhood onset specific loci (1q21.3, 4p14 and 17q12), and three genome-wide signficant shared loci (2q12.1, 6p21.32, and 11q13.5). The sample sizes for the age of onset bins are [0,5]: n=4,637; [6,10]: n=4,255; [11,15]: n=2,684; [16,20]: n=2,128; [21,25]: n=2,578; [26,30]: n=2,923; [31,35]: n=2,599; [36,40]: n=3,571; [41,45]: n=2,967; [46,50]: n=3,266, [51,55]: n=2,544; [56,65]: n=3,694.

Figure 4. Results of PrediXcan studies at non-HLA region loci. Genes whose predicted expression was significantly associated with asthma in either the childhood onset cases or the

adult onset cases are shown for skin (left panels), lung (middle panels) and whole blood (right panels). Values shown for skin combine both sun exposed and not sun exposed skin, showing the most significant statistic of the two. Results using gene expression in spleen and small intestine are shown in appendix Figure 5. The z-scores on the x- and y-axes are from transcriptome-wide tests of association with asthma using SNP sets that predict the expression of that gene. The diagonal dashed lines show the expected when associations are the same in the childhood onset and adult onset cases. The horizontal/vertical dashed lines correspond to z-scores ±4.84 (p=1.29x10^{-6}); the horizontal/vertical dotted lines correspond to z-scores ±1.96 (p=0.05). Colored backgrounds correspond to the chromosome location of each gene (see Key). Upper panels: Genes that are associated with childhood onset asthma (no genes were associated with adult onset asthma). Lower panels: Shared genes associated with both childhood onset and adult onset asthma. HLA region genes (6p21.32 and 6p21.33) are shown in Figure 5.

Figure 5. Results of PrediXcan studies of HLA region genes. Genes in the HLA region whose predicted expression was associated with childhood onset or adult onset asthma are shown for skin (left panel), lung tissue (middle panel) and whole blood (right panel). Results using gene expression in spleen and small intestine are shown in appendix Figure 6. The genes on darker shaded backgrounds correspond to shared genes; the four genes on lighter color backgrounds are age-specific. See Figure 4 for additional details.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5