A well-controlled BioID design for endogenous bait proteins

Giel Vandemoortele[1,2], Delphine De Sutter[1,2], Aline Moliere[1,2], Jarne Pauwels[1,2], Kris Gevaert[1,2] and

Sven Eyckerman[1,2,*]

[1]VIB Center for Medical Biotechnology, VIB, B-9000 Ghent, Belgium

[2]Department of Biochemistry, Ghent University, B-9000 Ghent, Belgium

Keywords: protein protein interaction; BioID proximity labeling; (T2A) autocleavage; genome

engineering; p53; base editor

*Correspondence to Prof. Dr. Sven Eyckerman

VIB Center for Medical Biotechnology

A. Baertsoenkaai 3, B-9000 Ghent, Belgium

Tel: +32-9-264.92.18

Fax: +32-9-264.94.90

Email: sven.eyckerman@vib-ugent.be

ABSTRACT

The CRISPR/Cas9 revolution is profoundly changing the way life sciences technologies are used. Many

assays now rely on engineered clonal cell lines to eliminate overexpression of bait proteins. Control

cell lines are typically non-engineered cells or engineered clones implying a considerable risk for

artefacts because of clonal variation. Genome engineering can also transform BioID, a proximity

labelling method that relies on fusing a bait protein to a promiscuous biotin ligase, BirA*, resulting in

the tagging of vicinal proteins. We here propose an innovative design to enable BioID for endogenous

proteins wherein we introduce a T2A-BirA* module at the C-terminus of endogenous p53 by genome

engineering, leading to bi-cistronic expression of both p53 and BirA* under control of the endogenous

promoter. By targeting a Cas9-cytidine deaminase base editor to the T2A auto-cleavage site, we can

efficiently derive an isogenic population expressing a functional p53-BirA* fusion protein. Using

quantitative proteomics we show significant benefits over classical ectopic expression of p53-BirA*,

and we provide a first well-controlled view on the proximal proteins of endogenous p53 in colon

carcinoma cells. This novel application for base editors expands the CRISPR/Cas9 toolbox and can be

a valuable addition for synthetic biology.

ABBREVIATIONS

37	AP-MS	Affinity Purification – Mass Spectrometry
38	dCas9	Catalytically inactive 'dead' Cas9
39	DDA	Data-Dependent Acquisition
40	DSB	DNA double Strand Break
41	GO	Gene Ontology
42	Indel	Insertion or deletion
43	IRES	Internal Ribosome Entry Site
44	LFQ	Label-Free Quantitation
45	nCas9	Nickase Cas9
46	PEI	Polyethylenime
47	rAAV	recombinant Adeno-Associated virus
48	UGI	Uracil DNA Glycosylase Inhibitor

INTRODUCTION

Elucidating the composition and function of protein complexes is crucial for advancing our understanding of key cellular processes, both in health as well as disease [1]. Fuelled by an ever-growing arsenal of so-called interactomics technologies [2], vast amounts of data are now available for further exploration [3, 4]. With expanding lists of interactors for every protein challenged, there is a growing consensus in the field that many reported interactions are highly contextual or spatially restricted [5]. In addition, it became apparent that approaches in which a bait protein is overexpressed lack the subtlety that is needed to further dissect protein-protein interactions and assess physiological functions of individual interactions [6]. Indeed, protein overexpression can have a detrimental effect on protein localization and folding, and distort downstream signalling [5, 7]. Protein overexpression also tends to mask complex feedback control mechanisms affecting a protein at physiological levels and generally increases the overall stress status of a cell [8, 9]. Several efforts have been made to avoid bait protein overexpression, with the use of primary antibodies to pull-down endogenous protein complexes as prime example [10, 11]. In reality, the specificity and the sensitivity of antibodies are often inadequate for studying endogenous protein complexes in a reliable manner and can therefore lead to problems with reproducibility [12]. Despite improved methodologies it remains challenging to generate specific antibodies with favourable affinities for each protein [13, 14] especially when one wants to specifically study mutant proteins bearing amino acid substitutions or truncated proteins originating from, for instance, alternative translation start sites (i.e. proteoforms [15]). For such situations, Flp-In^{TM} cell lines allowing tunable expression provide a valuable alternative [16]. However, this requires thoughtful and precise fine-tuning of expression levels to mimic as closely as possible physiological protein levels [17]. Artificial chromosomes have also been used to express transgenes at physiological levels [18].

A technology that is gaining popularity to map protein interactomes is BioID. This method relies on a promiscuous biotin ligase (BirA*) fused to a bait protein of interest [19]. BirA* is a mutagenized *E. coli*

biotin ligase (carrying the R118G mutation) that no longer displays substrate specificity. Upon

expressing a bait-BirA* fusion protein, proximity-dependent biotinylation occurs of proteins that are

near-neighbours of this bait protein in the cell. As endogenous biotinylation is a rare and highly specific

modification in mammalian cells [20, 21], biotinylated prey proteins are easily discriminated after

purification by streptavidin matrices and mass spectrometry (MS) analysis. Moreover, the need for

lysis is removed since the labelling of prey proteins occurs in intact cells before the homogenization

step. Since its initial publication in 2012, BioID has been used to study proximal and interacting

proteins (i.e. the proxisome) of bait proteins in several organisms and cell types producing datasets

complementary to affinity purification − MS (AP-MS) approaches [22]. In bait overexpression

experiments, as controls, BirA* is often expressed alone or fused to irrelevant baits (e.g. eDHFR, eGFP),

although cellular localizations and expression levels might differ between BirA*-fusion proteins and

BirA* alone, which therefore confound results [19].

Bi-cistronic expression systems can provide a way to ensure similar expression levels for a bait protein

and a detached BirA* module. 2A peptides cause ribosomal skipping during translation and

recommencement of translation immediately afterwards [23]. 2A peptides display increased

efficiency when compared to Internal Ribosome Entry Site (IRES) sequences and are only 18 amino

acids in length, which is an additional advantage when considering genome engineering approaches

[24].

Since the emergence of CRISPR/Cas9, many applications have been published with several of them

outgrowing classical genome engineering [25]. A recent application is on base editing in which Cas9 is

fused to a cytidine deaminase. This enzyme converts cytosine into uracil which has the base-pairing

properties of thymine [26] and thus introduces mutations. The base editing activity can be targeted

to confined sequence windows of approximately five nucleotides by loading Cas9 with a specific guide

RNA (gRNA). As base editing is typically performed by nickase Cas9 (nCas9) or catalytically inactive

(dead, dCas9) fusion constructs, no DNA double strand breaks (DSBs) are introduced, implying that

minimal numbers of insertion or deletion (indel) mutations are generated in this process. To avoid

base excision repair which would revert the introduced mutation, extra moieties were added to these

base editors, such as an uracil DNA glycosylase inhibitor (UGI) [27]. Up to now, base editing

applications mainly focussed on correcting disease mutations, on genetic diversification using tiled

gRNA libraries [27-29], or on introducing premature stop codons as alternative to classic knock-out

strategies [30].

p53 is arguably the most studied protein in health and disease [31, 32]. Historically dubbed the

guardian of the genome [33], the encoding gene is mutated in 50% of all reported human cancer cases

[34], making it an attractive starting point for the development of anticancer strategies [35]. Because

of the long-lasting research focus on p53, the current interactome of p53 consists of over 1000 protein

interaction partners (BioGRID, version 3.4 [3]). p53 acts as a homotetrameric, sequence-specific

transcription factor capable of inducing large gene networks comprising hundreds of target genes [36,

37]. Known pathways regulated by p53, either by transactivation or protein interactions, include DNA

damage repair, cell cycle arrest, senescence and apoptosis following exposure to a plethora of stress

conditions such as DNA damage, telomere erosion, hypoxia, replication stress and oncogene

activation. More recently, additional processes under p53 control such as autophagy, metabolism and

ferroptosis were revealed illustrating the need for continued research on this protein [38].

In this report, we use a novel knock-in strategy to enable BioID experiments for endogenous p53 in

HCT116 colon carcinoma cells as classic overexpression experiments in this cell model gave suboptimal

results. In our knock in strategy, a control cell line was first generated by introduction of a T2A-BirA*

cassette at the C-terminus of endogenous p53 using CRISPR/Cas9 genome engineering. These cells

express the BirA* protein at equal levels to p53 and thus provide a good system to assess biotinylation

background. Targeted inactivation of the T2A peptide sequence by a Cas9 cytidine deaminase fusion

(BE3) results in the generation of a cell line that expresses a functional p53-BirA* fusion protein from

the endogenous promoter allowing proximal protein mapping for endogenous p53. This reduces

genome engineering efforts considerably and eliminates artefacts due to clonal variation when a

selection step is added in the protocol. Furthermore, the concept of auto-cleavage switching in fusion

proteins can find applications in synthetic biology.

EXPERIMENTAL PROCEDURES

Cell culture

HCT116 colon carcinoma cells were purchased from ATCC (ATCC CCL-247) and cultured in McCoy's 5A

modified medium (Gibco, 22330070). Genomically characterized HEK293T embryonic kidney cells [39]

were cultured in DMEM (Gibco, 61965026) supplemented with fetal bovine serum (FBS) (Gibco,

10270106) to a final concentration of 10%, 12500 units penicillin-streptomycin (Gibco, 15070063) and

HEPES (Gibco, 15630056) to a final concentration of 10 mM. Growth medium was switched to McCoy's

5A modified medium or DMEM supplemented with 10% dialyzed FBS (Gibco, 26400044) for HCT116 2

weeks prior to seeding cells for a BioID experiment. Cells were passaged when cultures reached 70%

confluence. All cell lines were deemed clear of mycoplasma by using a mycoplasma PCR detection kit

(Minerva Biolabs, MIN-11-1100).

Plasmids and cloning procedures

The BioID biotin ligase BirA* was transferred from the pcDNA3.1 mycBioID vector (Addgene #35700)

by PCR to generate the pAav-TP53-T2A-BirA* targeting construct (available via Addgene, #115652).

All primers used in this manuscript can be found in table S1. Homology regions were generated by PCR

on genomic HCT116 DNA as described before [40, 41]. The pMet7-FLAG-PQS1-TP53-T2A-BirA*-Myc

construct was generated by standard restriction enzyme cloning procedures using the FLAG-PQS1-p53

expression vector published earlier [40]. This construct was further mutagenized by PCR to obtain a

construct bearing an inactivated T2A sequence for a BioID transient transfection experiment. A

puromycin selection cassette was incorporated in the BE3 (Addgene, plasmid #73021) vector for

endogenous mutagenesis purposes (BE3$_{puro}$) using a linear dsDNA fragment (IDT DNA Technologies,

gBlock). gRNA sequences incorporated in a U6-driven expression cassette were ordered as gBlocks

(Table S2) prior to blunt-ended ligation in a pCR-Blunt vector (Thermo Fisher Scientific, K2750).

CRISPR/Cas9 genome engineering

HCT116 *TP53*$^{+/T2A-BirA*}$ cells were generated using a combination of CRISPR/Cas9 and recombinant

Adeno-associated virus (rAAV)-mediated template delivery as described before [40, 41]. Briefly, a

TP53 C-terminal targeted gRNA (5' ACGCACACCUAUUGCAAGCA 3') was cloned in a Cas9 expression

construct (Addgene #48139) as described by Ran *et al.* [42]. Cells were seeded in a 24-well plate 24 h

before transfection with the Cas9 plasmid using FugeneHD (Promega, E2311). 48 h after transfection,

cells were infected with targeting construct packaged in rAAV. Neomycin selection (Gibco, 11811031)

was performed for two weeks starting 24 h after infection. Afterwards, cells were manually diluted

and seeded as single cells in 96-wells for further clonal expansion until on average, clones reached

70% confluence at which point they were split into a lysis plate and culture plate. Screening of clonal

populations in the lysis plate was based on junction PCR screening using GoTaq G2 Hot Start

polymerase (Promega, M7405). Input lysates for PCR were obtained using DirectPCR lysis reagent

(Viagenbiotech, 301C). Positive clones were seeded at 10^3 cells per 96-well 24 h prior to treatment

with 2.5 µM TAT-Cre (Excellgen, EG1001). 24 h after treatment, the medium was aspirated and

replaced with fresh growth medium. Four days after TAT-Cre treatment, cells were subjected to a

second round of single cell seeding and PCR screening. Clones showing aberrant growth rates or

morphology were omitted for further characterization. PCR-positive clones were subjected to

Western blot using 1/1000 diluted anti-p53 antibody (Calbiochem, OP43T) and near-infrared

fluorescent secondary antibodies (LI-COR). Additionally, Southern blot analysis was performed to

verify single genomic integration of the floxed neomycin resistance cassette. [α-^{32}P] Southern blot

probe was generated as described before.[41] For every retained clone, engineered loci were verified

using Sanger sequencing as described earlier. Primers used for screening, sequencing and generation

of the Southern blot probe can be found in table S1.

T2A mutagenesis

7.2×10^5 HEK293T cells were seeded in 6-well plates the day prior to transfection. A total amount of

800 ng DNA was mixed with 4 µl polyethylenime (PEI) for every transfection condition and incubated

for 10 min at room temperature. Meanwhile cells were placed on 1.5 ml DMEM medium

supplemented with 2% FCS. 6 h after addition of the transfection mix, the medium was removed and

replaced by DMEM with additives after washing once with Phosphate Buffered Saline (PBS) (Gibco,

14190094). 72 h post transfection cells were washed with PBS before detachment using a cell scraper

(Greiner, 541070) and lysis in 2X SDS-PAGE XT sample buffer (Bio-Rad, 1610791) and XT reducing agent

(Bio-Rad, 1610792) by heating for 10 min at 95°C. Afterwards, 100 µl MilliQ water was added to dilute

samples prior loading on a QIAshredder homogenizer spin-column (Qiagen, 79656) and running them

on a Criterion XT 4-12% BisTris gel (Bio-Rad, 3450123). After blotting, the PVDF membrane was

incubated with anti-FLAG (Sigma, F1804) and anti-Myc (in-house produced) primary antibodies and

detected with LI-COR near-infrared fluorescent secondary antibodies. For endogenous mutagenesis,

BE3$_{puro}$ vector and gRNA plasmids were transfected using FugeneHD in HCT116 $TP53^{+/T2A-BirA*}$ cells. 24

h after transfection, 2 µg/ml puromycin was added to the growth medium for 48 h to enrich for BE3$_{puro}$

transfected cells prior to expansion for proteomics experiments.

Illumina targeted sequencing

For targeted sequencing, subconfluent 25 cm^2 bottles containing HCT116 $TP53^{+/T2A-BirA*}$ cells after

BE3$_{puro}$ treatment and subsequent enrichment were pelleted by centrifugation at 500 g for 5 min and

washed once with PBS. To each pellet, 500 µl lysis buffer [20 mM Tris-HCl pH 7.5, 5 mM EDTA, 150

mM NaCl, 400 U Proteinase K (Sigma, P4850), 0.2% SDS] was added and incubated for 16 h at 37°C.

Afterwards, genomic DNA was obtained by phenol-chloroform extraction and ethanol precipitation.

Next, the T2A locus was amplified and provided with Nextera overhangs by PCR using Herculase

polymerase (Agilent, 600675). Sequencing was performed using the MiSeq Nano v2 kit (Illumina,

MS1022002) by the VIB Nucleomics core facility (http://www.nucleomics.be/). FASTQ files were

uploaded to CRISPRESSO [43] to analyse results. The CRISPRESSO allele frequency output file (Table

S3) was used as input to generate the sequence logo using the online tool WebLogo [44].

Nuclear and cytoplasmic extraction

For every cell line, cells were cultured in McCoys medium supplemented with a final concentration of

10% dialyzed FBS as described above. 1 mM doxorubicin (Sigma, D1515) and 50 µM biotin (Sigma,

B4639) was added to the growth medium 24 h prior to the start of the experiment at which 7×10^6 cells

were pelleted by centrifugation at 500 g for 5 min. After washing once with ice-cold PBS and

centrifugation at 500 g for 5 min, the supernatant was aspirated from the cell pellet. Extraction was

done using the NE-PER kit (Thermo Fisher Scientific, 78833) according to the instructions provided by

the manufacturer. In brief, 500 µl ice-cold CER 1 buffer was added to every cell pellet before

resuspension by vortexing. After 10 min incubation on ice, 27.5 µl ice-cold CER 2 buffer was added to

the mix and samples were centrifuged at 16000 g on 4°C for 5 min. Supernatant resulting from this

centrifugation step was stored as cytoplasmic extract. The pellet fraction was resolved in 250 µl ice-

cold NER and vortexed vigorously every 10 min for 15 s during an incubation period on ice for a total

of 40 min. Afterwards, nuclear extract was obtained by centrifugation at 16000 g for 10 min at 4°C and

the supernatant containing this nuclear extract was transferred to a new tube.

In parallel, 10^6 cells were washed once with PBS prior to resuspension in 150 µl ice-cold RIPA lysis

buffer [50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% NP-40, 1 mM EDTA, 1 mM EGTA, 0.1% SDS, 0.5%

sodium deoxycholate (DOC) and complete protease inhibitor cocktail (Roche, 11873580001)].

Following 10 min incubation on ice, lysates were centrifuged at 16000 g for 15 min at 4°C and the

supernatant was transferred to a new tube (= total cell lysate sample). For every fraction, the protein

concentrations as determined by a Bradford assay and 40 µg was loaded on ExpressPlus SDS-PAGE

(GenScript, M42015) and afterwards blotted on a PVDF membrane. α-tubulin was used as cytoplasmic

marker (Sigma, T5168).

BioID affinity purification

For every sample three 145 cm^2 culture dishes (Nunc) were seeded with 4.5×10^6 HCT116 *TP53*[+/T2A-BirA*]

cells cultured on McCoy's 5A modified medium supplemented with dialyzed FBS. 24 h after seeding 1

µM doxorubicin was added. To every culture dish biotin was added to a final concentration of 50 µM

at the start of the treatment. 24 h after treatment the growth medium was replenished to deprive

free biotin from the culture. 3 h later, the growth medium was aspirated and cells were washed once

with cold PBS before detachment in PBS using a cell scraper. After pelleting cells by centrifugation for

5 min at 500 g on 4°C, a second PBS washing step occurred prior to cell lysis in a 15 ml tube. 2 ml of

ice-cold RIPA lysis buffer containing 250 U of benzonase (Sigma, E1014) was added to each pellet. Lysis

was obtained by incubation with agitation for 1 h on 4°C prior to sonication (30% amplitude, 5 x 6 s

burst, 2 s interruption) on ice. The insoluble fraction of the lysate was removed by centrifugation for

15 min at 16000 g on 4°C and the supernatant was transferred to a new 15 ml tube. The total protein

concentration was determined by the Bradford protein assay to normalize input material to the

maximum shared total protein amount throughout the samples. For every sample, 90 µl Streptavidin

Sepharose High Performance bead suspension (GE Healthcare, 17-5113-01) was used for enrichment

of biotinylated proteins. Beads were pelleted by centrifugation at 500 g for 1 min and washed once in

wash buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% NP-40, 1 mM EDTA, 1 mM EGTA, 0,1% SDS)

prior to addition to the lysate. Samples were incubated for affinity purification on a rotator at 4°C for

3 h. Afterwards, beads were recovered by centrifugation of the samples at 500 g for 1 min at 4 °C and

aspirating the supernatant. Beads were washed three times in wash buffer, twice in 50 mM

ammonium bicarbonate pH 8.0 and once with 20 mM Tris-HCl pH 8.0, 2 mM $CaCl_2$ prior to

resuspension in 20 µl 20 mM Tris-HCl pH 8.0. Trypsin digest occurred overnight by addition of 1 µg

trypsin (Promega, V5111) to each sample. Digestion mixtures were depleted of beads upon

centrifugation at 500 g for 1 min, after which the supernatant was transferred to a mass spectrometry

vial. To ensure complete digestion, peptide mixtures were incubated with an additional 500 ng trypsin

for 3 h before addition of formic acid to a final concentration of 2%. All experiments were performed

in biological triplicate for downstream label-free quantitative proteome analysis.

For transient transfection BioID experiments, $8.5x10^6$ HCT116 cells were seeded in a 56 cm^2 culture

dish (Nunc) and transfected the day after with pMet7-FLAG-PQS1-TP53-T2A-BirA*-Myc, the pMet7-

FLAG-PQS1-TP53-MUTT2A-BirA*-Myc or pSV-SPORT mock vector using FugeneHD. 24 h post

transfection, cells were transferred to 3 T145 cm^2 dishes per sample. 24 h after transfer, 1 µM

doxorubicin and 50 µM biotin were added 24 h prior to harvest. Affinity purification of biotinylated

proteins was performed as described above.

LC-MS/MS instrument analysis

Peptide mixtures were analyzed by LC–MS/MS on an Ultimate 3000 RSLC nano LC (Thermo Fisher Scientific) in-line connected to a Q-Exactive mass spectrometer (Thermo Fisher Scientific). The peptides were first loaded on a trapping column (made in-house, 100 µm internal diameter (I.D.) × 20 mm, 5 µm beads C18 Reprosil-HD, Dr. Maisch, Ammerbuch-Entringen, Germany). After flushing the trapping column, peptides were loaded in solvent A (0.1% formic acid) on a reverse-phase column (made in-house, 75 µm I.D. x 250 mm, 3 µm Reprosil-Pur-basic-C18-HD beads packed in the needle, Dr. Maisch, Ammerbuch-Entringen, Germany) and eluted by an increasing concentration solvent B (0.1% formic acid in acetonitrile) using a linear gradient from 2% solvent B up to 55% solvent B in 120 min, followed by a washing step with 99% solvent B, all at a constant flow rate of 300 nl/min. The mass spectrometer was operated in data-dependent acquisition (DDA), positive ionization mode, automatically switching between MS and MS/MS acquisition for the 5 most abundant peaks in a given MS spectrum. Source voltage was set at 3.4 kV, with a capillary temperature of 275°C. One MS1 scan (m/z 400–2000, AGC target 3×10^6 ions, maximum ion injection time 80 ms), acquired at a resolution of 70000 (at 200 m/z), was followed by up to 5 tandem MS scans (resolution 17500 at 200 m/z) of the most intense ions fulfilling predefined selection criteria (AGC target 5×10^4 ions, maximum ion injection time 80 ms, isolation window 2 Da, fixed first mass 140 m/z, spectrum data type: centroid, underfill ratio 2%, intensity threshold $1.3xE^4$, exclusion of unassigned, 1, 5-8 and >8 positively charged precursors, peptide match preferred, exclude isotopes on, dynamic exclusion time 12 s). HCD collision energy was set to 25% normalized collision energy and the polydimethylcyclosiloxane background ion at 445.120025 Da was used for internal calibration (lock mass).

Mass spectrometry data processing and interpretation

Obtained Xcalibur Raw Files (.raw) were analyzed using MaxQuant and maxLFQ algorithms (MaxQuant version 1.5.8.3) [45, 46]. Spectra were searched against the human UniProt sequence database (as available on 01/09/2017). Methionine oxidation, N-terminal acetylation and lysine biotinylation were set as variable modifications, with a maximum of 5 modifications per peptide. The minimum peptide

length was set at 7 amino acids and a maximum peptide mass of 4600 Da was used. PSM, protein and

site false discovery rates were set at 0.01. The minimum Label-Free Quantitation (LFQ) ratio count was

2 and the Fast LFQ option was disabled. 20 ppm and 4.5 ppm mass accuracies were used for the first

and main search respectively. After completion of searches, LFQ intensities were loaded in Perseus

(version 1.5.5.3) [47] for further analysis.

Samples were annotated on condition for analysis. Proteins only identified by site, reverse hits and

Perseus contaminant proteins were removed from the data matrix. Retained LFQ intensities were

transformed to log2 scale and three valid values in at least one of the two sample groups were needed

in order for the protein to be retained for further analysis. Missing values were imputed from a normal

distribution of intensities (0.3 width, 2.2 downshift). A two-sided t-test (0.05 FDR, 1000

randomizations) was performed to reveal differentially enriched proteins in the volcano plot. Gene

ontology (GO) analyses were performed using the database for annotation, visualization and

integrated discovery (DAVID 6.8) [48]. The mass spectrometry proteomics data have been deposited

to the ProteomeXchange Consortium via the PRIDE [49] partner repository with the dataset identifier

PXD010771.

RESULTS

BioID by forced expression of a p53-BirA* fusion protein in colon carcinoma cells

Proximity biotinylation is an exciting development of recent years to identify vicinal proteins of a target protein which includes the associated protein complexes [19, 22]. While the downstream purification of biotinylated proteins of the *in vivo* biotinylation reaction is a powerful and sensitive procedure, great care should be taken with the design of the experiment and the levels of BirA* [50]. Most BioID studies rely on inducible expression systems based on Flp recombinase integration (Flp-In™ T-REx™ system). This powerful system allows low expression which can be tuned to match endogenous levels. Despite being around for well over 10 years, only few engineered Flp-In cell lines are available and virtually all BioID studies using this system are therefore performed in 293 cells [50-52].

We set out to explore the so-called 'proxisome' of p53 in well-known and well-characterized colon carcinoma HCT116 cells. The model system has been used extensively for the study of p53 function, in part because of the availability of many knock out cell lines [53]. We were faced with the lack of an inducible HCT116 cell line and we therefore decided to first explore an experiment with forced expression of a p53-BirA* construct. We transiently transfected HCT116 cells with a fusion construct that contains a C-terminal BirA* module cloned in frame through an inactivated T2A peptide that served as a long linker (p53-MUTT2A-BirA*). As control experiments we used either mock-transfected cells or cells transfected with a bi-cistronic p53-T2A-BirA* construct (Figure 1A and 1B). This last construct ensured equal levels of the free BirA* module in the control cells, providing a distinct advantage over expression of a free unfused BirA* protein which often requires tuning to obtain equal expression levels. We performed triplicate experiments to allow LFQ analysis using MaxQuant and Perseus workflows. Using the mock control, an extensive list of significant proteins for p53-BirA* was obtained by LFQ analysis (Figure 1C). Of these, 17.1% were known p53 interaction partners in BioGRID. Some well-known interaction partners can be discerned in the list, but despite the comprehensive list, some key interaction partners such as MDM2 are lacking. This extensive list can be expected because

the control conditions lack any BirA*. When assessing the T2A-BirA* control condition we found some

distinct differences, pointing to removal of contaminants by the presence of BirA* in this control.

However, the retrieved lists were still surprisingly long with 408 significant proteins (Figure 1D), of

which 20.3% were known BioGRID p53 interaction partners. This large number of hits is likely due to

the excessive levels of BirA*, resulting in much background biotinylation in both transient expression

approaches. Levels of the fusion protein are well beyond the endogenous levels in the cells (Figure

1B). In addition, GO analysis on 'molecular function' showed very few links to p53 function, with 'p53

binding' only retrieved at position 38 and 88 for the mock and T2A-BirA* control respectively. GO

analysis for 'cellular component' showed strong enrichment of cytoplasmic and membrane proteins

(Table S4 and S5) which contradicts the predominant nuclear function of p53. Thus the results of these

analyses hint to the presence of substantial amounts of biologically irrelevant, false positive prey

proteins.

Generation of a control cell line for BioID on endogenous p53

The long lists in the overexpression experiment are likely caused by the high expression levels of the

bait protein. While this can be solved by introducing an inducible expression cassette containing a

p53-BirA* fusion construct through lentiviral transduction [54], we decided to explore the exciting

developments in the genome editing field to engineer the BirA* module in one allele of the *TP53* gene

in HCT116 cells. We started by first creating a control T2A-BirA* cell line as we reasoned that the best

way to ensure similar levels in the control condition is to use a bi-cistronic approach (Figure 2A). Similar

BirA* control expression levels are also maintained when p53 protein levels are changed due to stress

conditions on the cells (e.g. DNA damage). We used CRISPR/Cas9 to introduce a double stranded DNA

(dsDNA) break in the last coding exon of *TP53* and provided a single strand DNA (ssDNA) repair

template containing a T2A-BirA* module and a floxed expression cassette flanked by two homology

regions. Generation of this control cell line was performed using a two-step engineering approach as

described earlier ([40], Figure S1). Clones bearing insertions were identified by PCR (Figure S2). In the

second step, the floxed selection cassette is removed by CRE recombinase, again followed by PCR

screening of single cell clones (Figure S3). Single integration for these clones was shown by Southern

blot (Figure S4A), and by Sanger sequencing of both alleles ensuring an intact second allele

(Supplementary Figure S4B and S4C). Because of current limits in knock-in efficiencies, single cell

clones were selected and evaluated for the introduction of the knock-in construct and for CRE-based

removal of the selection cassette.

Deriving an experimental cell line using a base editor approach

An experimental cell line containing an engineered *TP53* locus with a C-terminal BirA* module can be

obtained using a similar genome engineering approach as for the control cell line, but there are a

number of things to consider. Firstly, the seeding, culturing and screening of single cell clones related

to current knock-in efficiencies requires a significant effort effectively doubling genome engineering

hands-on time to perform a BioID experiment. Secondly, the generation of the experimental cell line

would also require two selection steps of single cell clones. This could introduce clonal variation and

may introduce artefacts in the data. To solve these issues, we sought to derive an isogenic population

from the control cell line wherein the T2A site would not be present. This can be obtained by another

CRISPR/Cas9 knock-in approach using a template with an alternative linker (e.g. an inactive T2A site).

However, the efficiencies currently obtained for knock-in are too low to allow such an approach and

would still require clonal selection (see higher). As an alternate way to achieve T2A inactivation, we

turned to a recently reported Cas9-cytidine deaminase base editor BE3 [27]. Targeted inactivation of

the 2A sequence would ensure restoration of bait-BirA* transgene expression (Figure 2B). 2A peptides

have conserved amino acid residues that are critical for peptidyl transferase inhibition that causes

ribosomal skipping (Figure 2C) [55, 56]. Therefore, the targeted mutagenesis of one of these key

residues should enable efficient abrogation of 2A ribosomal skipping activity. To explore this, we

generated an expression vector that expresses a fusion construct of the tumor suppressor protein p53

linked to BirA* by a T2A sequence. Appropriate codon usage of the T2A sequence was verified to

ensure the presence of several target nucleotides for BE3 (Figure 2C). Varying amounts of a FLAG-

TP53-T2A-BirA*-MYC expression construct were co-transfected with the BE3 construct and T2A-

specific gRNA. Two different gRNAs were assessed for base editing activity (Table S2). Conversion of

the auto-cleavage site was clearly detectable by the presence of a high molecular weight band

corresponding to a p53-BioID fusion construct on Western blot analysis (Figure 2D).

T2A conversion after integration by genome engineering

A puromycin selection cassette was first cloned into the original BE3 vector allowing for efficient

enrichment of transfected cells. This ensures that non-transfected cells, and thus non-edited cells, are

efficiently removed in a short puromycin selection step. These non-edited cells would generate

additional noise to downstream LFQ proteomics analysis [45]. The gRNA directly targets the conserved

amino acid residues on the T2A sequence while leaving p53 and BirA* intact (inactivated T2A is

hereafter indicated as MUTT2A). When assessing p53 translation products before or after BE3

transfection, a clear signal for p53-MUTT2A-BirA* was observed in puromycin-resistant cells (Figure

3A), a result which was obtained for multiple clones. Notably, an additional band with an apparent

molecular weight corresponding to WT p53 was also present. This extra band can be explained by the

heterozygous knock-in of T2A-BirA*. In HCT116 $TP53^{+/T2A-BirA*}$ the modified p53 allele has 22 extra

amino acids due to translation of the GSG linker and the T2A peptide. This band could still be observed

in the enriched cell population post-transfection, hinting at a portion of cells that did not undergo T2A

sequence conversion. This partial conversion can be attributed to the intrinsic activity of BE3 that is

below 100%. Furthermore, it is likely that not all editing will result in T2A inactivation as BE3

mutagenesis activity is centered around positions 4 – 8 in the protospacer sequence [27], therefore

mutations might be introduced that have no impact on T2A activity or only attenuate its potency. To

quantify the mutagenesis rate we performed next-generation sequencing on the resulting cell

population showing a conversion rate over 40% (Figure 3B). In agreement with earlier reports on base

editing, CRISPRESSO analysis showed that the rate of indel formation in the resulting population was

negligible and the mutagenesis window was restricted to the predicted nucleotides (Figure S5).

Next to efficient T2A inactivation, proper localization of p53-MUTT2A-BirA* and BirA* is clearly crucial

for the effectiveness and reliability of the system. Ideally, both the bait and the biotinylation activity

of the engineered cell lines should follow the same subcellular distribution before and after BE3

treatment to provide relevant background protein signals. To evaluate this, we derived nuclear and

cytoplasmic fractions of the engineered cells upon p53 activation by doxorubicin. We could observe a

clear nuclear enrichment for p53 in both parental and engineered cells, indicative that proper p53

localization is not impaired by knock-in of T2A-BirA* or base editor treatment (Figure 3C). As expected,

only in the BE3 treated engineered cells a strong signal for p53-T2A-BirA* could be observed in the

nuclear fraction. Furthermore, assessment of the biotinylation pattern showed an intense laddering

pattern in engineered cells when compared to the parental HCT116 cells (Figure 3C). Both before and

after BE3 treatment, biotinylation was mainly detectable in the nuclear fraction, which indicates that

the engineered cells with cleaved BirA* (p53-T2A-BirA*) provide a good control condition.

Additionally, biotinylated WT p53 can be clearly observed in BE3 treated cells, hinting that fusion to

BirA* is not affecting the oligomerization properties of p53 (Figure 3C).

Conversion of inserted T2A-BirA* as a potent tool to study the p53 proxisome

We then performed endogenous BioID experiments using the engineered and converted HCT116

$TP53^{+/MUTT2A-BirA*}$ cells. Experiments were done in triplicate to allow MaxQuant LFQ analysis [45]. Cells

were harvested and lysed after provoking a p53 response by doxorubicin. Puromycin-selected HCT116

$TP53^{+/T2A-BirA*}$ cells transfected with BE3 in absence of T2A-directed gRNA served as control condition

(Figure 4A). Following a two-sided t-test (FDR: 0.05, s0: 0.1) we observed 186 potential p53 interactors

for the mutagenized cells (Figure 4B and table S6), resulting in a high-quality proxisome dataset for

p53. String pathway analysis showed a highly interconnected network around p53 with complexes

involved in transcription and chromatin remodelling (Figure S6). Among the most differential proteins

is p53 (p-value <0.00001), supporting that T2A is mutagenized in a way that abrogates ribosomal

skipping by the peptide. In the control condition treated with doxorubicin, p53 was detected at lower

levels, hinting at rare events whereby the T2A peptide did not cause interruption of translation in

accordance with the Western blot analysis (Figure 3A). Next to p53, 63 (34.2%) known direct p53

interaction partners were identified including several hallmark interactors such as MDM2 [57],

TP53BP1 [58], EP400 [59], and Sin3B [60]. Noteworthy, we also detected proteins that affect the post-

translational (PTM) signature of p53 such as the phosphatase PPP1R10 [61]. Several proteins that

could not directly be linked to p53 have been already implied in DNA damage pathways such as

DMAP1 [62] and EPC1 [63]. For virtually all significantly enriched proteins, multiple unique peptides

were detected, enabling a robust protein inference (Table S6). GO analysis showed highly specific

enrichment for p53-related 'cellular component' and 'molecular function' terms (Table S7). 'p53

binding' can be found at position 13 with a p-value of 1.21×10^{-10}. These results support the concept

that BioID on the endogenous protein level can be exploited to further characterize dynamic p53

interactions in a physiologically relevant context.

To show the benefit of using a converted T2A design strategy, we also compared non-engineered WT

control cells (transfected with BE3 and selected on puromycin) to the p53-MUTT2A-BirA* cells as this

would mimic a simpler design strategy (Figure S7A). This analysis revealed 397 differential proteins

with a clear enrichment of RNA-binding proteins and chaperones (Figure S7B; Table S8) with 'p53

binding' found on position 17 with a p-value of 3.62×10^{-8}. When compared to the T2A-BirA* control

condition, 232 proteins were exclusively significant for the analysis using transfected HCT116 WT cells

(Figure S7C). A part of this extended protein list is also found in the BioGRID database as interactors

of p53. Strikingly, when these additional p53 BioGRID hits were assessed for 'cellular component'

terms they were mainly assigned as cytoplasmic proteins (180 proteins; p-value 9.19×10^{-14}), and their

'molecular function' was linked to functions such as RNA-binding (51 proteins; p-value 4.83×10^{-17}),

ribosomal constituents (34 proteins; p-value 1.03×10^{-17}) and poly(A) RNA or mRNA binding (142 and

15 proteins; p-values 2.02×10^{-67} and 1.14×10^{-6} respectively; Table S8). While these RNA-binding

proteins and chaperones can be *bona fide* interaction partners for p53, they likely turn up with many

different bait proteins (or directly with BirA*). Consequently, proteins with functions related to RNA

are typically enriched in contaminant repositories for affinity purification – mass spectrometry

experiments [64].

DISCUSSION

Since its initial emergence, BioID has been widely adopted by the proteomics community to detect protein-protein interactions and vicinal proteins because of its distinct advantages over classic AP-MS [22]. However, virtually all efforts rely on ectopic (over)expression of bait proteins fused to BioID [22]. To our knowledge, there is currently only one report where BioID is performed at the endogenous level [65]. The endogenous application of BioID solves two important issues in typical MS-based interactomics approaches. Firstly, the need for lysis is obviated since prey labelling occurs in intact cells before the homogenization step, and secondly, overexpression artefacts are removed by using the endogenous bait protein. This last issue is supported by our experiments with ectopic expression of p53-BirA* wherein we clearly show extensive lists of candidate proximal proteins pointing to the presence of many contaminants. Coupling expression levels of both bait-BirA* and control BirA* might be of particular relevance for proteins, such as p53, that have varying expression levels depending on the cellular state. One longstanding question in the p53 field is how p53 is able to elicit different responses to separate stimuli or stress conditions in a cell. We believe that future studies with BioID at the endogenous level may reveal some of the subtle interaction changes or altered feedback loops that are potentially overlooked in overexpression studies, thereby providing missing pieces to the p53 puzzle. Furthermore, the majority of p53 research has been based on overexpression experiments and therefore PPI databases may be riddled with interactions that are spatially and physiologically not possible in an intact cell. Although highly specific anti-p53 antibodies are commercially available for conducting endogenous immunoprecipitation MS experiments, these will always have the drawback that protein complexes need to be kept intact during purification. While crosslinking of protein complexes may improve conservation during purification, crosslinking is prone to inter-experimental variation in efficiency, suffers from ambiguous experimental data and can lead to problematic identification of crosslinked peptides by search algorithms [66].

We engineered a single cell line that can be used both as a control and an experimental cell line

without the need to go through an additional intermediary single cell stage. Our approach relies on

mutagenesis of a T2A sequence incorporated as a linker between a bait protein of interest and a

functional tag sequence, here BirA*. By mutagenizing the T2A site, ribosomal skipping is prevented

and a functional fusion construct is generated. We showed the concept of T2A mutagenesis on vectors

for overexpression and on a knock-in HCT116 $TP53^{+/T2A-BirA*}$ cell line. In addition, we show correct

localization of the BirA* enzyme in the modified cells, both before and after T2A inactivation. Based

on these results, we believe that our approach is suitable for cytoplasmic and nuclear bait proteins.

The current system does not allow to target the released BirA* to restricted compartments (e.g.

mitochondria), which may be solved by adding localization signals. Proper localization should be

carefully assessed for every bait protein, which is required for any interactome study.

As site-directed mutagenesis is accomplished by introduction of a CRISPR/nCas9-cytidine deaminase

fusion (BE3), a simple short selection step suffices to obtain a strong enrichment of cells transiently

expressing the BE3 base editor and therefore having a high chance of expressing p53-MUTT2A-BirA*.

In this way, there is no need to generate separate control cell lines, or to have a design wherein

engineered single cell knock in clones are compared to parental populations. Such a design would

inevitably lead to higher background levels as presented in this work (Figure S7, table S8). Until

genome engineering progresses to efficiencies wherein clonal screening is no longer needed, T2A

mutagenesis may be a valuable approach to obtain appropriate isogenic control cell lines. While the

devised T2A system allows rigorous control of expression levels, also when gene expression levels are

modulated by various stimuli (i.e. doxorubicin effects on p53 levels), the free BirA* cannot replicate

the exact localization of the bait protein. A similar localization would allow discrimination between

proteins in close proximity and interaction partners (both direct or indirect) [67]. As with all BioID

designs, downstream validation is still required to assess this aspect. While we opted for a classic well-

controlled knock-in approach, novel CRISPR/Cas9 strategies would also allow the selection of isogenic

populations (e.g. CRISPR co-selection [68], CRISPaint [69]).

CONCLUSIONS

Taken together, our elegant base editing approach to derive isogenic experimental cell lines from a control cell line enables BioID experiments at the endogenous level. Bringing together the advantages of proximal biotinylation in intact cells with endogenous bait expression levels and with quantitative proteomics pushes protein complex analysis to the next level and constitutes a useful addition to study hub proteins such as p53.

SUPPORTING INFORMATION

Table S1: Primers used for cloning or screening. (PDF)

Table S2: gRNA gblock sequences cloned in pCR-Blunt vector for transfection. (PDF)

Table S3: CRISPRESSO allele frequency output. (Excel)

Table S4: GO analysis of the results from an analysis of a transient transfection p53 BioID experiment using a mock vector as control. (PDF)

Table S5: GO analysis of the results from an analysis of a transient transfection p53 BioID experiment using a bi-cistronic p53-T2A-BirA* expression construct as control. (PDF)

Table S6: Perseus output endogenous T2A inactivation. (Excel)

Table S7: GO analysis of significantly enriched proteins from converted p53-MUTT2A-BirA* HCT116 cells when compared to isogenic non-converted p53-T2A-BirA* cells. (PDF)

Table S8: GO analysis of significantly enriched proteins from converted p53-MUTT2A-BirA* HCT116 cells when compared to parental HCT116 cells. (PDF)

Table S9: Perseus data matrices generated for all analyses. (Excel)

Figure S1: Targeting strategy and primer scheme used for PCR screening. (PDF)

Figure S2: PCR screening of neomycin resistant clonal populations. (PDF)

Figure S3: PCR for assessing the removal of the selection cassette upon TAT-Cre treatment. (PDF)

Figure S4: Validation of targeted T2A-BirA* insertion. (PDF)

Figure S5: CRISPRESSO analysis of the T2A region after BE3 base editing. (PDF)

Figure S6: STRING analysis of proximal proteins for endogenous p53. (PDF)

Figure S7: Evaluation of BE3 transfected non-engineered HCT116 WT cells as control for p53 BioID. (PDF)

ACKNOWLEDGEMENTS

We thank the VIB Proteomics Core for running the MS samples. We thank Elise Wyseure and Noortje

Samyn for technical assistance. pCMV-BE3 was provided by David Liu (Addgene plasmid #73021).

AUTHOR CONTRIBUTIONS

G.V. designed and performed cell line experiments with help of D.D.S, performed BioID experiments,

analyzed data and wrote the manuscript. G.V. and D.D.S performed cell line engineering and related

characterization experiments with help of A.M. J.P. operated the MS and analyzed the samples. S.E.

conceived and managed the project and wrote the manuscript. K.G. is co-promotor of G.V. and read

and corrected the manuscript.

FUNDING

G.V. is a PhD student funded by the Fund for Research - Flanders (FWO grant 3G050913N to K.G. and

S.E.) and BOF-GOA (2016000602 to S.E). K.G. acknowledges support from the Flanders Agency for

innovation by Science and Technology (IWT, SBO grant 60839).

CONFLICT OF INTEREST

The authors declare no competing interests.

REFERENCES

[1] S. Petrakis, M.A. Andrade-Navarro, Editorial: Protein Interaction Networks in Health and Disease, Front Genet 7 (2016) 111.

[2] K. Titeca, I. Lemmens, J. Tavernier, S. Eyckerman, Discovering cellular protein-protein interactions: Technological strategies and opportunities, Mass Spectrom Rev (2018) 1-33.

[3] A. Chatr-Aryamontri, R. Oughtred, L. Boucher, J. Rust, C. Chang, N.K. Kolas, L. O'Donnell, S. Oster, C. Theesfeld, A. Sellam, C. Stark, B.J. Breitkreutz, K. Dolinski, M. Tyers, The BioGRID interaction database: 2017 update, Nucleic Acids Res 45(D1) (2017) D369-D379.

[4] K. Luck, G.M. Sheynkman, I. Zhang, M. Vidal, Proteome-Scale Human Interactomics, Trends Biochem Sci 42(5) (2017) 342-354.

[5] T.J. Gibson, M. Seiler, R.A. Veitia, The transience of transient overexpression, Nat Methods 10(8) (2013) 715-21.

[6] G. Prelich, Gene overexpression: uses, mechanisms, and interpretation, Genetics 190(3) (2012) 841-54.

[7] J.E. Dumont, S. Dremier, I. Pirson, C. Maenhaut, Cross signaling, cell specificity, and physiology, Am J Physiol Cell Physiol 283(1) (2002) C2-28.

[8] Y.B. Zhang, J. Howitt, S. McCorkle, P. Lawrence, K. Springer, P. Freimuth, Protein aggregation during overexpression limited by peptide extensions with large net negative charge, Protein Expr Purif 36(2) (2004) 207-16.

[9] T.K. Kim, J.H. Eberwine, Mammalian cell transfection: the present and the future, Anal Bioanal Chem 397(8) (2010) 3173-8.

[10] A. Malovannaya, R.B. Lanz, S.Y. Jung, Y. Bulynko, N.T. Le, D.W. Chan, C. Ding, Y. Shi, N. Yucer, G. Krenciute, B.J. Kim, C. Li, R. Chen, W. Li, Y. Wang, B.W. O'Malley, J. Qin, Analysis of the human endogenous coregulator complexome, Cell 145(5) (2011) 787-99.

[11] L. Fagerberg, B.M. Hallstrom, P. Oksvold, C. Kampf, D. Djureinovic, J. Odeberg, M. Habuka, S. Tahmasebpoor, A. Danielsson, K. Edlund, A. Asplund, E. Sjostedt, E. Lundberg, C.A. Szigyarto, M. Skogs, J.O. Takanen, H. Berling, H. Tegel, J. Mulder, P. Nilsson, J.M. Schwenk, C. Lindskog, F. Danielsson, A. Mardinoglu, A. Sivertsson, K. von Feilitzen, M. Forsberg, M. Zwahlen, I. Olsson, S. Navani, M. Huss, J. Nielsen, F. Ponten, M. Uhlen, Analysis of the human tissue-specific expression by genome-wide integration of transcriptomics and antibody-based proteomics, Mol Cell Proteomics 13(2) (2014) 397-406.

[12] M.G. Weller, Quality Issues of Research Antibodies, Anal Chem Insights 11 (2016) 21-7.

[13] A. Bradbury, A. Pluckthun, Reproducibility: Standardize antibodies used in research, Nature 518(7537) (2015) 27-9.

[14] E. Marcon, H. Jain, A. Bhattacharya, H. Guo, S. Phanse, S. Pu, G. Byram, B.C. Collins, E. Dowdell, M. Fenner, X. Guo, A. Hutchinson, J.J. Kennedy, B. Krastins, B. Larsen, Z.Y. Lin, M.F. Lopez, P. Loppnau, S. Miersch, T. Nguyen, J.B. Olsen, M. Paduch, M. Ravichandran, A. Seitova, G. Vadali, M.S. Vogelsang, J.R. Whiteaker, G. Zhong, N. Zhong, L. Zhao, R. Aebersold, C.H. Arrowsmith, A. Emili, L. Frappier, A.C. Gingras, M. Gstaiger, A.G. Paulovich, S. Koide, A.A. Kossiakoff, S.S. Sidhu, S.J. Wodak, S. Graslund, J.F. Greenblatt, A.M. Edwards, Assessment of a method to characterize antibody selectivity and specificity for use in immunoprecipitation, Nat Methods 12(8) (2015) 725-31.

[15] D. Gawron, E. Ndah, K. Gevaert, P. Van Damme, Positional proteomics reveals differences in N-terminal proteoform stability, Molecular systems biology 12(2) (2016) 858.

[16] J.Z. Torres, J.J. Miller, P.K. Jackson, High-throughput generation of tagged stable cell lines for proteomic analysis, Proteomics 9(10) (2009) 2888-91.

[17] S. Senkel, C. Waldner, G.U. Ryffel, H. Thomas, Improved conditional expression systems resulting in physiological level of HNF4alpha expression confirm HNF4alpha induced apoptosis in the pancreatic beta-cell line INS-1, BMC Res Notes 2 (2009) 210.

[18] M.Y. Hein, N.C. Hubner, I. Poser, J. Cox, N. Nagaraj, Y. Toyoda, I.A. Gak, I. Weisswange, J.
Mansfeld, F. Buchholz, A.A. Hyman, M. Mann, A human interactome in three quantitative
dimensions organized by stoichiometries and abundances, Cell 163(3) (2015) 712-23.
[19] K.J. Roux, D.I. Kim, M. Raida, B. Burke, A promiscuous biotin ligase fusion protein identifies
proximal and interacting proteins in mammalian cells, J Cell Biol 196(6) (2012) 801-10.
[20] A. Chapman-Smith, J.E. Cronan, Jr., The enzymatic biotinylation of proteins: a post-translational
modification of exceptional specificity, Trends Biochem Sci 24(9) (1999) 359-63.
[21] A. Chapman-Smith, J.E. Cronan, Jr., Molecular biology of biotin attachment to proteins, J Nutr
129(2S Suppl) (1999) 477S-484S.
[22] R. Varnaite, S.A. MacNeill, Meet the neighbors: Mapping local protein interactomes by
proximity-dependent labeling with BioID, Proteomics 16(19) (2016) 2503-2518.
[23] Z. Liu, O. Chen, J.B.J. Wall, M. Zheng, Y. Zhou, L. Wang, H. Ruth Vaseghi, L. Qian, J. Liu,
Systematic comparison of 2A peptides for cloning multi-genes in a polycistronic vector, Sci Rep 7(1)
(2017) 2193.
[24] A.L. Szymczak, C.J. Workman, Y. Wang, K.M. Vignali, S. Dilioglou, E.F. Vanin, D.A. Vignali,
Correction of multi-gene deficiency in vivo using a single 'self-cleaving' 2A peptide-based retroviral
vector, Nat Biotechnol 22(5) (2004) 589-94.
[25] R. Barrangou, J.A. Doudna, Applications of CRISPR technologies in research and beyond, Nat
Biotechnol 34(9) (2016) 933-941.
[26] C. Kuscu, M. Adli, CRISPR-Cas9-AID base editor is a powerful gain-of-function screening tool, Nat
Methods 13(12) (2016) 983-984.
[27] A.C. Komor, Y.B. Kim, M.S. Packer, J.A. Zuris, D.R. Liu, Programmable editing of a target base in
genomic DNA without double-stranded DNA cleavage, Nature 533(7603) (2016) 420-4.
[28] Y. Ma, J. Zhang, W. Yin, Z. Zhang, Y. Song, X. Chang, Targeted AID-mediated mutagenesis (TAM)
enables efficient genomic diversification in mammalian cells, Nat Methods 13(12) (2016) 1029-1035.
[29] N.M. Gaudelli, A.C. Komor, H.A. Rees, M.S. Packer, A.H. Badran, D.I. Bryson, D.R. Liu,
Programmable base editing of A*T to G*C in genomic DNA without DNA cleavage, Nature 551(7681)
(2017) 464-471.
[30] C. Kuscu, M. Parlak, T. Tufan, J. Yang, K. Szlachta, X. Wei, R. Mammadov, M. Adli, CRISPR-STOP:
gene silencing through base-editing-induced nonsense mutations, Nat Methods 14(7) (2017) 710-
712.
[31] B. Vogelstein, D. Lane, A.J. Levine, Surfing the p53 network, Nature 408(6810) (2000) 307-10.
[32] J.A. Royds, B. Iacopetta, p53 and disease: when the guardian angel fails, Cell Death Differ 13(6)
(2006) 1017-26.
[33] D.P. Lane, Cancer. p53, guardian of the genome, Nature 358(6381) (1992) 15-6.
[34] A.M. Goh, C.R. Coffill, D.P. Lane, The role of mutant p53 in human cancer, J Pathol 223(2) (2011)
116-26.
[35] V.P. Almazov, D.V. Kochetkov, P.M. Chumakov, The use of p53 as a tool for human cancer
therapy, Mol Biol (Mosk) 41(6) (2007) 947-63.
[36] T. Riley, E. Sontag, P. Chen, A. Levine, Transcriptional control of human p53-regulated genes,
Nat Rev Mol Cell Biol 9(5) (2008) 402-12.
[37] M. Fischer, Census and evaluation of p53 target genes, Oncogene 36(28) (2017) 3943-3956.
[38] E.R. Kastenhuber, S.W. Lowe, Putting p53 in Context, Cell 170(6) (2017) 1062-1078.
[39] Y.C. Lin, M. Boone, L. Meuris, I. Lemmens, N. Van Roy, A. Soete, J. Reumers, M. Moisse, S.
Plaisance, R. Drmanac, J. Chen, F. Speleman, D. Lambrechts, Y. Van de Peer, J. Tavernier, N.
Callewaert, Genome dynamics of the human embryonic kidney 293 lineage in response to cell
biology manipulations, Nat Commun 5 (2014) 4767.
[40] G. Vandemoortele, A. Staes, G. Gonnelli, N. Samyn, D. De Sutter, E. Vandermarliere, E.
Timmerman, K. Gevaert, L. Martens, S. Eyckerman, An extra dimension in protein tagging by
quantifying universal proteotypic peptides using targeted proteomics, Sci Rep 6 (2016) 27220.

[41] G. Vandemoortele, D. De Sutter, S. Eyckerman, Robust Generation of Knock-in Cell Lines Using
CRISPR-Cas9 and rAAV-assisted Repair Template Delivery, Bio-protoc 7(7) (2017) e2211.
[42] F.A. Ran, P.D. Hsu, J. Wright, V. Agarwala, D.A. Scott, F. Zhang, Genome engineering using the
CRISPR-Cas9 system, Nat Protoc 8(11) (2013) 2281-2308.
[43] L. Pinello, M.C. Canver, M.D. Hoban, S.H. Orkin, D.B. Kohn, D.E. Bauer, G.C. Yuan, Analyzing
CRISPR genome-editing experiments with CRISPResso, Nature biotechnology 34(7) (2016) 695-7.
[44] G.E. Crooks, G. Hon, J.M. Chandonia, S.E. Brenner, WebLogo: a sequence logo generator,
Genome research 14(6) (2004) 1188-90.
[45] J. Cox, M.Y. Hein, C.A. Luber, I. Paron, N. Nagaraj, M. Mann, Accurate proteome-wide label-free
quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ, Mol
Cell Proteomics 13(9) (2014) 2513-26.
[46] S. Tyanova, T. Temu, J. Cox, The MaxQuant computational platform for mass spectrometry-
based shotgun proteomics, Nat Protoc 11(12) (2016) 2301-2319.
[47] S. Tyanova, T. Temu, P. Sinitcyn, A. Carlson, M.Y. Hein, T. Geiger, M. Mann, J. Cox, The Perseus
computational platform for comprehensive analysis of (prote)omics data, Nat Methods 13(9) (2016)
731-40.
[48] W. Huang da, B.T. Sherman, R.A. Lempicki, Systematic and integrative analysis of large gene lists
using DAVID bioinformatics resources, Nature protocols 4(1) (2009) 44-57.
[49] J.A. Vizcaino, A. Csordas, N. del-Toro, J.A. Dianes, J. Griss, I. Lavidas, G. Mayer, Y. Perez-Riverol,
F. Reisinger, T. Ternent, Q.W. Xu, R. Wang, H. Hermjakob, 2016 update of the PRIDE database and its
related tools, Nucleic Acids Res 44(D1) (2016) D447-56.
[50] J.P. Lambert, M. Tucholska, C. Go, J.D. Knight, A.C. Gingras, Proximity biotinylation and affinity
purification are complementary approaches for the interactome mapping of chromatin-associated
protein complexes, Journal of proteomics 118 (2015) 81-94.
[51] X. Liu, K. Salokas, F. Tamene, Y. Jiu, R.G. Weldatsadik, T. Ohman, M. Varjosalo, An AP-MS- and
BioID-compatible MAC-tag enables comprehensive mapping of protein interactions and subcellular
localizations, Nature communications 9(1) (2018) 1188.
[52] E. Coyaud, M. Mis, E.M. Laurent, W.H. Dunham, A.L. Couzens, M. Robitaille, A.C. Gingras, S.
Angers, B. Raught, BioID-based Identification of Skp Cullin F-box (SCF)beta-TrCP1/2 E3 Ligase
Substrates, Molecular & cellular proteomics : MCP 14(7) (2015) 1781-95.
[53] P. Qiu, H. Guan, P. Dong, S. Li, C.T. Ho, M.H. Pan, D.J. McClements, H. Xiao, The p53-, Bax- and
p21-dependent inhibition of colon cancer cell growth by 5-hydroxy polymethoxyflavones, Molecular
nutrition & food research 55(4) (2011) 613-22.
[54] P. Samavarchi-Tehrani, H. Abdouni, R. Samson, A.C. Gingras, A versatile lentiviral delivery toolkit
for proximity-dependent biotinylation in diverse cell types, Molecular & cellular proteomics : MCP
(2018).
[55] M.L. Donnelly, G. Luke, A. Mehrotra, X. Li, L.E. Hughes, D. Gani, M.D. Ryan, Analysis of the
aphthovirus 2A/2B polyprotein 'cleavage' mechanism indicates not a proteolytic reaction, but a
novel translational effect: a putative ribosomal 'skip', J Gen Virol 82(Pt 5) (2001) 1013-25.
[56] P.A. Radcliffe, K.A. Mitrophanous, Multiple gene products from a single vector: 'self-cleaving' 2A
peptides, Gene Therapy 11 (2004) 1673.
[57] U.M. Moll, O. Petrenko, The MDM2-p53 interaction, Molecular cancer research : MCR 1(14)
(2003) 1001-8.
[58] R. Cuella-Martin, C. Oliveira, H.E. Lockstone, S. Snellenberg, N. Grolmusova, J.R. Chapman,
53BP1 Integrates DNA Repair and p53-Dependent Cell Fate Decisions via Distinct Mechanisms, Mol
Cell 64(1) (2016) 51-64.
[59] N.L. Lill, S.R. Grossman, D. Ginsberg, J. DeCaprio, D.M. Livingston, Binding and modulation of
p53 by p300/CBP coactivators, Nature 387(6635) (1997) 823-7.
[60] N. Bansal, R. Kadamb, S. Mittal, L. Vig, R. Sharma, B.S. Dwarakanath, D. Saluja, Tumor
suppressor protein p53 recruits human Sin3B/HDAC1 complex for down-regulation of its target
promoters in response to genotoxic stress, PLoS One 6(10) (2011) e26156.

[61] S.J. Lee, C.J. Lim, J.K. Min, J.K. Lee, Y.M. Kim, J.Y. Lee, M.H. Won, Y.G. Kwon, Protein
phosphatase 1 nuclear targeting subunit is a hypoxia inducible gene: its role in post-translational
modification of p53 and MDM2, Cell Death Differ 14(6) (2007) 1106-16.
[62] T. Koizumi, M. Negishi, S. Nakamura, H. Oguro, K. Satoh, M. Ichinose, A. Iwama, Depletion of
Dnmt1-associated protein 1 triggers DNA damage and compromises the proliferative capacity of
hematopoietic stem cells, Int J Hematol 91(4) (2010) 611-9.
[63] Y. Wang, V. Alla, D. Goody, S.K. Gupta, A. Spitschak, O. Wolkenhauer, B.M. Putzer, D.
Engelmann, Epigenetic factor EPC1 is a master regulator of DNA damage response by interacting
with E2F1 to silence death and activate metastasis-related gene signatures, Nucleic Acids Res 44(1)
(2016) 117-33.
[64] D. Mellacheruvu, Z. Wright, A.L. Couzens, J.P. Lambert, N.A. St-Denis, T. Li, Y.V. Miteva, S. Hauri,
M.E. Sardiu, T.Y. Low, V.A. Halim, R.D. Bagshaw, N.C. Hubner, A. Al-Hakim, A. Bouchard, D. Faubert,
D. Fermin, W.H. Dunham, M. Goudreault, Z.Y. Lin, B.G. Badillo, T. Pawson, D. Durocher, B. Coulombe,
R. Aebersold, G. Superti-Furga, J. Colinge, A.J. Heck, H. Choi, M. Gstaiger, S. Mohammed, I.M. Cristea,
K.L. Bennett, M.P. Washburn, B. Raught, R.M. Ewing, A.C. Gingras, A.I. Nesvizhskii, The CRAPome: a
contaminant repository for affinity purification-mass spectrometry data, Nature methods 10(8)
(2013) 730-6.
[65] C.B. Mulholland, M. Smets, E. Schmidtmann, S. Leidescher, Y. Markaki, M. Hofweber, W. Qin, M.
Manzo, E. Kremmer, K. Thanisch, C. Bauer, P. Rombaut, F. Herzog, H. Leonhardt, S. Bultmann, A
modular open platform for systematic functional studies under physiological conditions, Nucleic
Acids Res 43(17) (2015) e112.
[66] J. Rappsilber, The beginning of a beautiful friendship: cross-linking/mass spectrometry and
modelling of proteins and multi-protein complexes, J Struct Biol 173(3) (2011) 530-40.
[67] B.T. Lobingier, R. Huttenhain, K. Eichel, K.B. Miller, A.Y. Ting, M. von Zastrow, N.J. Krogan, An
Approach to Spatiotemporally Resolve Protein Interaction Networks in Living Cells, Cell 169(2) (2017)
350-360 e12.
[68] D. Agudelo, A. Duringer, L. Bozoyan, C.C. Huard, S. Carter, J. Loehr, D. Synodinou, M. Drouin, J.
Salsman, G. Dellaire, J. Laganiere, Y. Doyon, Marker-free coselection for CRISPR-driven genome
editing in human cells, Nature methods 14(6) (2017) 615-620.
[69] J.L. Schmid-Burgk, K. Honing, T.S. Ebert, V. Hornung, CRISPaint allows modular base-specific
gene tagging using a ligase-4-dependent mechanism, Nature communications 7 (2016) 12338.

FIGURES

Figure 1. Transfection p53 BioID experiments. (A) Workflow used for transient expression LFQ

proteomics experiments. All experiments were conducted in presence of 1 µM doxorubicin. Top panel:

Control transfection with an empty vector. Middle panel: p53-MUTT2ABirA* transfection leading to

biotinylation of vicinal p53 proteins. Bottom panel: p53-T2A-BirA* control transfection leading to

biotinylation of background proteins. (B) Western blot for p53 in the transient expression experiment.

Different lanes depict p53 protein levels in the empty vector transfection (EV), in the fusion p53-

MUTT2ABirA* transfection (MUTT2A), and in the transfection with the bi-cistronic p53-T2A-BirA*

(T2A) control. (C) and (D) Volcano plots showing differential LFQ intensity levels (X-axis) and p-values

(Y-axis) in transient transfection BioID experiments using a mock and BirA* control transfection

respectively.

Figure 2. Targeted T2A inactivation. (A) Incorporation of a sequence coding for a T2A peptide (cyan)

between the bait (p53) and BirA* will lead to translation of two separate proteins. (B) After

transfection, the BE3 base editor will be guided to the T2A sequence by (a) specific gRNA(s) and will

convert cytidines mainly to thymidines. Dark blue: inactivated T2A (MUTT2A) (C) Layout of the T2A

region in the TP53-T2A-BirA* test construct. Conserved T2A residues are underlined. Green: p53 C-

terminus; cyan: linker and T2A sequence; red: BirA* N-terminus; grey: gRNA used; magenta: PAM-

motif. (D) Overexpression pilot experiment to assess T2A mutagenesis of a target plasmid. Cells were

lysed 72 h after transfection with a MYC-tagged p53-T2A-BirA* construct, and loaded on SDS-PAGE

for Western blotting using an anti-MYC antibody. Upon mutagenesis a band corresponding to the 82

kDa p53-MUTT2A-BirA* protein is visible. The two gRNAs targeting different residues of the T2A

sequence were tested in parallel for potency (left: gRNA 1; right: gRNA 2). In both cases, expression of

p53-BirA* could be observed at varying amounts of target concentrations. gRNA 2 was selected for

further endogenous use.

Figure 3. Endogenous T2A mutagenesis and cell line characterization (A) Endogenous T2A

mutagenesis. Cells were lysed 72 hours after BE3 transfection and loaded on SDS-PAGE for Western

blot analysis using an anti-p53 antibody. p53 or p53-BioID expression is not influenced by BE3

expression in parental HCT116 cells or in HCT116 *TP53*[+/BirA*] control cell lines. T2A inactivation could

be observed for all four HCT116 *TP53*[+/T2A-BirA*] clones tested resulting in expression of the p53-

MUTT2A-BirA* fusion protein. Clone 4 was used for further experiments. (B) Sequence logo depicting

mutation frequencies of BE3-treated cells. Paired-end sequencing was performed on the cell

population after BE3 treatment and enrichment. The relevant T2A region is depicted in the sequence

logo. C > T conversions could only be observed in the targeted residues. Red: expected BE3-induced

mutation prevalence according to the position of gRNA 2. (C) Nuclear and cytoplasmic extraction

experiments of doxorubicin-treated cell lines. p53 is mostly present in the nuclear extract of all cell

lines. Both before and after BE3 treatment, BirA* shows an identical localization in the nucleus, in

accordance with p53. In both populations, biotinylation is elevated when compared to parental

HCT116 cells, with the majority of biotinylated substrates detected in the nuclear extract. N: nuclear

extract; C: cytoplasmic extract; T: Total cell lysate.

Figure 4. Mapping the endogenous p53 proxisome using BioID. (A) Proteomics workflow used for

HCT116 *TP53*[+/T2A-BirA*]. Experiments were done in triplicate to support LFQ analysis. The puromycin

resistance cassette in the BE3$_{Puro}$ vector was used for enrichment of the transfected cells. Cells were

allowed to recover prior to doxorubicin treatment to activate p53. Zoom: graphic representation of

the promiscuous biotinylation in HCT116 *TP53*$^{+/T2A-BirA*}$ (top) and HCT116 *TP53*$^{+/MUTT2A-BirA*}$ (bottom).

Blue: p53-proximal proteins. Black: biotin. (B) Volcano plot showing the differential LFQ intensity levels

(X-axis) and the p-values (Y-axis) for HCT116 *TP53*$^{+/T2A-BirA*}$ cells before and after T2A conversion in

presence of doxorubicin. Known p53 interactors indexed in BioGRID are highlighted in blue. Significant

hits discussed in the main text are labeled.