Imaging human cortical responses to intraneural microstimulation using magnetoencephalography

George C. O'Neill[a*], Roger H. Watkins[b], Rochelle Ackerley[bc], Eleanor L. Barratt[a], Ayan Sengupta[a], Michael Asghar[a], Rosa Maria Sanchez Panchuelo[a], Matthew J. Brookes[a], Paul M. Glover[a], Johan Wessberg[b] , Susan T. Francis[a]

[a]*Sir Peter Mansfield Imaging Centre, School of Physics and Astronomy, University of Nottingham, Nottingham, UK*
[b]*Department of Physiology, University of Gothenburg, Gothenburg, Sweden*
[c]*Aix Marseille Univ, CNRS, LNSC (Laboratoire de Neurosciences Sensorielles et Cognitives – UMR 7260), Marseille, France*

10   **Abstract**

The sensation of touch in the glabrous skin of the human hand is conveyed by thousands of fast conducting mechanoreceptive afferents, which can be categorised into four distinct types. The spiking properties of these afferents in the periphery in response to varied tactile stimuli are well-characterised, but relatively little is known about the spatiotemporal properties of the neural representations of these different receptor types in the human cortex. Here, we use the novel methodological combination of single unit intraneural microstimulation (INMS) with magnetoencephalography (MEG) to localise cortical representations of individual touch afferents in humans, by measuring the extracranial magnetic fields from neural currents. We find that by assessing modulations of the beta (13-30 Hz) rhythm during single unit INMS, significant changes in oscillatory amplitude occur in the contralateral primary somatosensory cortex within and across a group of fast adapting type I mechanoreceptive afferents, which
20   corresponded well to the induced response from matched vibrotactile stimulation. Combining the spatiotemporal specificity of MEG with the selective single unit stimulation of INMS enables the interrogation of the central representations of different aspects of tactile afferent signalling within the human cortices. The fundamental finding that single unit INMS ERD responses are robust and consistent with natural somatosensory stimuli will permit us to more dynamically probe the central nervous system responses in humans, to address questions about the processing of touch from the different classes of mechanoreceptive afferents and the effects of stimulus frequency and waveform.

Keywords: magnetoencephalography, somatosensory cortex, intraneural microstimulation, single unit

*Corresponding author
Email address: george.oneill@nottingham.ac.uk

Introduction

In humans, our sense of touch is conveyed by an array of discrete low-threshold mechanoreceptive afferents distributed throughout the skin, and haptic sensations are derived from a combination of activity across stimulated receptors (Johnson, 2001; Saal and Bensmaia, 2014). The sensory afferents from these receptors project to the somatosensory cortices, via the thalamus, enabling the decoding of the nature and location of the stimulus evoking the sensation. Our somatosensory abilities support a wide range of systems in the body, from integrating with the motor cortex for force feedback and fine motor control (Ackerley and Kavounoudias, 2015), to emotional impact, providing feedback from social situations (McGlone et al., 2014).

Much of our knowledge about human mechanoreceptive afferents and their electrophysiological properties stems from microneurography studies (Vallbo and Hagbarth, 1968), where an electrode is placed in a peripheral nerve to register axonal impulses. Microneurography allows the characterisation of *single* mechanoreceptive afferent responses when the associated receptive field in the skin is excited by touch. Four distinct low-threshold mechanoreceptor classes exist in glabrous (non-hairy) skin, each with their own distinct spiking pattern in response to somatosensory stimuli and spatial representations within the skin (Vallbo et al., 1984a). In addition to registering impulses, it is possible to pass a small current (of approximately a few microamperes) down the electrode to elicit artificial impulses in the same mechanoreceptive afferent, in the absence of contact to the skin. Here, a small, corresponding simulated tactile sensation can be felt, i.e. the projected receptive field (at the same site as the physiological receptive field), which is thought to represent the sensation conveyed by these individual mechanoreceptors. This process has been termed single unit intraneural microstimulation (INMS; Torebjörk and Ochoa, 1980; Vallbo, 1981) and allows precise impulse patterns to be evoked in individual nerve fibres, something traditional transdermal or intraneural stimulation cannot provide. Single unit INMS has shown that physiologically-defined mechanoreceptor types have distinct perceptual correlates in humans (Ochoa and Torebjörk, 1983; Vallbo et al., 1984b). Fast adapting type I (FA1) and type II (FA2) afferents elicit a vibration or fluttering sensation when stimulated, whilst slowly adapting (SA) type I (SA1) give rise to a pressure or pulling sensation. Slowly-adapting type II (SA2) afferents do not typically produce a clear, identifiable sensation. The type I units, which are more numerous (Johansson and Vallbo, 1979), provide small, point-like sensations. The use of single unit INMS has the advantage over traditional tactile stimuli of being able to produce a defined and controllable pattern of firing within a single afferent and single afferent class. This allows the precise examination of how input patterns from peripheral mechanoreceptive afferents are perceived and represented in the central nervous system.

Despite knowledge of the peripheral and perceptive properties of single mechanoreceptive afferents, little is known about how their responses to stimulation manifest in the human brain. Using single unit INMS, it is possible to probe the sense of touch at the "quantal" level of individual afferents, something which cannot be achieved for any of the other senses. There have been relatively few studies in which neuroimaging has been combined with single unit INMS to investigate the neural effects of selective mechanoreceptive afferent stimulation. A study by Kelly et al., (1997) used high density electroencephalography (EEG) over the primary somatosensory cortex (S1) and showed that stimulating single FA1 or SA1 units projecting to the hand gave characteristic peaks in power at the same frequency at which the units were stimulated. Only two studies have been performed using functional magnetic resonance imaging (fMRI) to map the spatial profile of the cortical responses to single unit INMS (Sanchez Panchuelo et al., 2016; Trulsson et al., 2001). The early study, performed at 3.0 T, showed that individual type I units have corresponding neural activations in S1 and the secondary somatosensory cortex (S2), and the responses were in good spatial agreement with activations from vibrotactile stimulation (Trulsson et al., 2001). The more recent study was performed at ultra-high field (7.0 T) using high spatial resolution to show that functional activation patterns of units located on the hand digits were spatially well-localised within the expected digit regions, ascertained from maps of digit somatotopy obtained from vibrotactile stimulation to the skin of the fingertips (Sanchez Panchuelo et al., 2016). Magnetoencephalography (MEG) measures the extracranial magnetic field associated with extracellular current flow.

MEG provides unique spatiotemporal resolution, offering vastly improved temporal resolution over fMRI (which is limited by the haemodynamic lag of the BOLD effect, and it being an indirect measure of neural activity) and an improved spatial resolution compared to EEG (as magnetic fields are not distorted by the skull to the same extent as electric potentials). In this paper, we performed single unit INMS, using a recently developed system

designed for compatibility with MEG (Glover et al., 2017). We aimed to demonstrate that this approach can
100   localise the cortical response to single unit INMS and capture the spatiotemporal dynamics of responses to simulation of individual mechanoreceptors, providing the feasibility of performing future studies to probe how the dynamics of different patterns of activity in these mechanoreceptors are represented cortically.

Methods

Experimental Procedure

A total of 9 healthy participants (age 24 – 68 years, 5 female) volunteered to undergo microneurography/INMS and simultaneous MEG recording. Each participant was given detailed information about the procedure and provided their written informed consent. Ethical approval was granted by the University of Nottingham Medical School Ethics Committee, with all procedures conducted in line with the Declaration of Helsinki.

Experimental sessions consisted of 1) microneurography for the characterisation of a single mechanoreceptive
110   afferent (Vallbo and Hagbarth, 1968); 2) assessment of the sensation to single unit INMS; 3) MEG recordings during INMS. These steps were repeated and ~2-3 individual mechanoreceptive afferents were typically stimulated during MEG data collection in a single experimental session. The participant was seated comfortably with their head just below the MEG helmet, with their arm held still using a vacuum cushion. A high-impedance, insulated tungsten recording/stimulating electrode (15 mm length, shaft diameter 200 µm, tip diameter ~5 µm; FHC, Bowdoin, ME) was then inserted percutaneously into the left median nerve, approximately 30 mm proximal from the wrist crease. A second uninsulated reference electrode was placed subcutaneously 50 mm away from the first and both were connected to an INMS system specifically designed to operate without generating measurable magnetic interference (Glover et al., 2017). A diagram of the electrode setup is shown in Figure 1.

For experimental phase 1, the INMS system was in 'record mode', and a real time visual and audio readout of the
120   potential difference between the two electrodes was output to the experimenter within the magnetically shielded room. The location of mechanoreceptive afferents was determined by gentle touch across the hand, delivered by the experimenter, until a unitary recording was detected. The precise location and force threshold of the unit was then identified using von Frey monofilaments and marked with ink. Once the mechanoreceptive afferent was identified, the response pattern and spatial extent on the skin was examined to identify the receptor type as SA1, FA1, SA2 or FA2 (Vallbo et al., 1984a). The INMS system was then switched to 'stimulate mode', to assess the sensation to INMS (experimental phase 2). A train of constant current pulses (pulse width 200 µs) were delivered at 60 Hz for 1 s. The pulses were delivered repetitively and the current was increased gradually (resolution 0.1 µA), until the participant felt a sensation in their hand, which was time locked to the stimulation. The participant was asked to describe the nature and location of the sensation; if there was an exact correspondence with the
130   physiological properties of the unit (e.g. location, receptive field size), the investigation proceeded to the MEG data collection (experimental phase 3). The paradigm during the MEG recording consisted of 80 trials of a 1 s long train of pulses (as described above), with a base inter-trial-interval of 10 ± 1 s to jitter the stimulus. After every 10 trials, the quality and intensity of the sensation was confirmed verbally with the participant, as small movements of the electrode can diminish the intensity of the sensation, presumably reflecting failure to generate impulses with each stimulation pulse. In the case where a participant no longer felt the sensation, the pulse amplitude was increased until a similar quality of sensation was again felt again in the same location (Torebjörk et al., 1987). If a comparable sensation was achieved by increasing the stimulation intensity, the MEG protocol was resumed, but if comparable sensation intensity was not recovered, or there was evidence of a change in the unit stimulated (e.g. its quality), the MEG recording was terminated.

140   Immediately after INMS, the participant underwent a short experiment, where mechanical vibrotactile stimulation was applied to the skin at each microstimulated unit's receptive field location to determine the amplitude at which the vibrotactile stimulation matched the INMS sensation. A small piezoelectric vibrotactile stimulator (Dancer Design, St. Helens, UK) controlled by an Arduino microcontroller (http://arduino.cc) was used to deliver a vibrotactile stimulation at 60 Hz. The amplitude was adjusted until the sensation felt similar in intensity to that from the INMS. These amplitudes were noted and used in a subsequent MEG experiment in which vibrotactile stimulation at each microstimulated unit's receptive field was carried out with identical timings to the INMS paradigm. This vibrotactile/MEG follow up session was performed at a later date to allow for the maximum number of units to be stimulated within each INMS experimental session.

Figure 1 – Experimental setup for microneurography and single unit microsimulation. A) A simplified diagram showing how a single mechanoreceptive afferent is interrogated. A recording/stimulating electrode is placed in the median nerve to register activity from a single nerve fibre. If mechanical stimulation is applied to the afferent's receptive field, the electrode in 'record mode' will register the spiking pattern being sent to the central nervous system. In 'stimulate mode', passing a small current down the electrode will stimulate the mechanoreceptive afferent, eliciting a tactile sensation at the location of the receptive field in the skin. B) The experimental setup on a participant in the study.

Data Acquisition

MEG Data were recorded using a 275 channel MEG system (CTF; Coquitlam, BC) in synthetic 3rd order gradiometer configuration, with a sampling rate of 1200 Hz. An analogue trace of the INMS stimulation pulses was recorded concurrently to allow for offline synchronisation of the neuromagnetic recordings. All participants wore a bespoke flexible headcast (details of the manufacturing procedure can be found in Liuzzi et al., 2017; Meyer et al., 2017), designed to minimise head movement within the MEG system. The headcasts include cavities designed to fit three head position indicator (HPI) coils, which were periodically energised during the experiment, to determine their position within the MEG sensor array. The HPI coil locations relative to the brain were defined during headcast manufacture, meaning accurate coregistration between the sensors and the anatomy was possible. For a single participant, the headcast failed to fit and so an alternative method to coregister the brain to the sensors was employed. In this case the HPI indicators were attached to the head at the nasion and bilateral preauricular locations. A measurement of coil locations relative to the scalp was recorded using a 3D digitiser (Polhemus; Colchester, VT). Coregistration was achieved by matching the digitised surface with a surface extracted for the anatomical image. Anatomical images were acquired using a 3.0 T MR System (Philips Healthcare; Best, Netherlands) using an MPRAGE sequence (1 mm^3 isotropic resolution).

Data Analysis

MEG data were visually inspected for artefacts. If an epoch contained excessive artefacts (for example generated by muscle or eye movement, or from SQUID resets), it was omitted from further analysis. The sensor data were

then band-pass filtered to the beta band (13-30 Hz), as this frequency is known to be robustly modulated during somatosensory stimulation (Hari and Salmelin, 1997; Cheyne, 2013). MR images were segmented and cortical surfaces were reconstructed using FreeSurfer 5.3 (Dale et al., 1999; Fischl et al., 1999). Further segmentation of the scalp, inner and outer skull surfaces were generated using modified routines from FieldTrip (Oostenveld et al., 2011). The surface which defined the grey/white matter boundary was decimated to 12500 vertices per hemisphere and each vertex was used as a location for source reconstruction. Source reconstruction at each vertex was achieved with a Linearly Constrained Minimum Variance (LCMV) beamformer, with the data covariance matrix generated from the entire duration of the experiment (Brookes et al., 2008). Dipole approximation for forward modelling was calculated using a 3-shell boundary element model (BEM; Stenroos and Nummenmaa, 2016), with the optimal dipole orientation for maximal signal variance derived using an eigenvalue decomposition approach (Sekihara et al., 2004). With beamformer weights for a given vertex, w, generated, they are employed to quantify changes in power across the cortex. For every location, an activation index, A, was generated to quantify changes in source power during the stimulation (0-1 s after stimulus onset) compared to a baseline period (8.9 – 9.9 s after stimulus onset). For the i^{th} trial, A was calculated at each location using Equation 1,

$$A(i) = \frac{wC_s(i)w^T - wC_b(i)w^T}{wC_s(i)w^T + wC_b(i)w^T},$$
(1)

where $C_s(i)$ and $C_b(i)$ are the single-trial sensor-level covariance matrices which correspond to the stimulation and baseline epochs respectively. Activation indices were averaged across trials to generate a final activation image for each experiment. The activation index was scaled between -1 and 1, with 0 denoting no modulation in power between the two epochs. To allow for a group level analysis, all images were transformed to a standard cortical surface derived from the MNI-305 template brain.

Statistical Inference of Significant Power Changes

To identify which locations corresponded to a significant change in power during stimulation (INMS or vibrotactile), we employed non-parametric permutation testing (Maris and Oostenveld, 2007; Nichols and Holmes, 2002; Pantazis et al., 2005; Singh et al., 2003) at both the individual subject and group levels. At the individual subject level, our null hypothesis was that the oscillatory power measured in the stimulation and the baseline epochs within a trial were identical. Therefore, if we were to randomly shuffle the labelling of the covariance matrices within trials, prior to generating A in Equation 1, then averaging across trials to form a null averaged activation index, $\tilde{A}$, would not give a significantly different result to the genuine averaged activation index, $\bar{A}$. 2000 possible permutations of $\tilde{A}$ were generated to create a null distribution for each vertex location. To correct for multiple comparisons, the threshold free cluster enhancement score (TFCE; Smith and Nichols, 2009) was calculated for every vertex on each iteration of $\tilde{A}$. For a given location r, the TFCE score was calculated as

$$\text{TFCE}\left(\tilde{A}(r)\right) = \int_0^{h_{max}} e\left(\tilde{A}(r), h\right)^{0.5} h^2 \, dh$$
(2)

where $e(\tilde{A}(r), h)$ is a function which returns the spatial extent of a cluster at r, should $\tilde{A}(r)$ exceed a threshold value h, and 0 otherwise. Positive and negative clusters were treated as non-contiguous, and the maximum absolute value of TFCE on each iteration was used to generate a global null. Any vertex where the absolute TFCE of $\bar{A}$ exceeded the 95th percentile of this null was considered significant.

At the group level, our null hypothesis was that the activation maps simply represented zero-mean noise, so randomly multiplying any of the experiments' trial averaged activation images by -1 and then group averaging to generate a null activation map, $\tilde{A}$, would not give a significantly different result to the group averaged image $\bar{A}$. All $n!$ possible permutations were calculated and similarly to the group level analysis, multiple comparisons correction was handled by finding the critical TFCE threshold.

Results

Electrophysiology

Across the cohort of participants, 39 single unit recordings were identified. 19 of these provided percepts that matched the unit physiology, indicating successful single unit INMS (Torebjörk et al., 1987) and cortical responses to stimulation were recorded in the MEG. An excerpt from a single unit recording from an FA1 can be seen in Figure 2A. Here, we observed high spiking rates (occasionally exceeding 200 impulses.s^{-1}) when a von Frey monofilament was pressed into the receptive field location in the skin. Data from 3 units were discarded due to too few MEG trials being recorded before losing the unit sensation. The properties of each of the 16 units with full MEG recording protocols can be found in Table 1, and their locations on the hand in Figure 2B. 10 units were identified as mechanoreceptor type FA1, 5 as SA1, and 1 as FA2. In the results presented here, we primarily focus on the 10 FA1 units, given that these formed the largest group, and their perceived sensation – vibration/flutter – is similar to that of vibrotactile stimulation. For completeness, the localisation results of the other 6 units can be found in the supplementary information.

Figure 2 – A) Excerpt from a single unit peripheral neurve recording from a fast adapting type 1 (FA1) afferent (unit 6a), showing the unitary responses to mechanical stimulation of the receptive field in the skin whilst in record mode. Upon single unit INMS, the subject felt a 3 mm spot defined to have a 'prickle' sensation. The location of the receptive field can be found in panel B. (B) Receptive field locations in all 16 units in the right hand which were microstimulated whilst in the MEG scanner, with adjacent numbers corresponding to the Unit ID in Table 1. The shapes correspond to the receptor type; dark blue circles represent FA1 afferents (vibration sensation in a very small area); blue squares show SA1 afferents (pressure or pulling in a small area), and the cyan triangle represents an FA2 afferent (diffuse vibration sensation).

Participant ID	Unit ID	Type	Location	Sensation	Perceptive Field Size (mm)
1	1a	FA1	Medial D3	Vibration	Not recorded
	1b	SA1	Proximal D4	Pulling	0.1
2	2a*	FA1	Distal D3	Tingle	2
3	3a	SA1	Palm	Pulling	0.1
	3b	SA1	Palm	Pulling	1
	3c	FA1	Palm	Vibration	1
	3d	FA1	Distal D3	Flutter	1
	3e	FA1	Dorsal Medial D3	Flutter	1
4	4a	FA1	Distal D4	Vibration	1
	4b	FA1	Distal D4	Vibration	2
	4c	FA2	Distal D4	Tapping	7
	4d	SA1	Medial D4	Point-like pressure	0.1
	4e	SA1	Medial D4	Pressure	3
	4f	FA1	Distal D4	Vibration	3
5	5a	FA1	Distal D1	Tingle	2
6	6a	FA1	Palm	Prickle	3

240   *Unit acquired without headcast

Table 1 – Metadata on the 16 microstimulated units recorded in the MEG scanner. Unit identifiers are based on the order of collection. Note all locations are on the palmar side of the hand unless otherwise noted (D denotes digit number).

Figure 3 –Results of the INMS from the group of 10 FA1 units studied overlaid onto inflated cortical surfaces, gyri are represented as light grey surfaces sulci are dark grey. A) Results of the individual unit analysis, represented as a conjunction map, showing the number of FA1 units that exhibit significant changes in beta band (13-30 Hz) power for a given location. B) Group averaged activation image of the 10 FA1 units at a statistical threshold (p < 0.05), showing microsimulation elicits a reduction in beta oscillatory power over the contralateral sensorimotor area. C) Zoomed portion of panel B, with the Brodmann areas as defined by the Glasser atlas (Glasser et al., 2016) overlaid. The peak reduction of beta power (represented as the magenta dot) is located in postcentral gyrus. D) A time frequency spectrogram of the peak location from panel C, showing a distinct event related behaviour in the beta band which extends beyond the initial 1 s of stimulation.

MEG

Figure 3 shows the results of localising changes in beta power during INMS across the cohort of the 10 FA1 units. Figure 3A shows a conjunction map showing the number of individual units which have a significant ($p_{corrected} <$ 0.05) change in power. We observed a large degree of overlap across the units in the contralateral primary somatosensory and motor (M1) cortices, with 8/10 units in spatial agreement; the other 2 units showed no significant changes in power at any spatial location in the brain (see Supplementary material). In the ipsilateral

260 S1, we observed overlap of activations in 2/10 units. Figure 3B shows the map of group-averaged activation index ($\bar{A}_{INMS}$), thresholded to show only the significant changes in beta power in response to microstimulation at the group level. This map shows that there was a strong decrease in oscillatory power which was (much like the high overlap region at the individual level) situated primarily over the contralateral S1 and extended into M1. Figure 3C is a zoomed representation of the findings in Figure 3B, overlaid with the Brodmann areas of the sensorimotor

system, as defined by the Glasser atlas (Glasser et al., 2016). The location of peak reduction in beta power is represented by the magenta dot, and is situated on the borders of Brodmann areas 1 and 2, confirming that the peak modulation of power occurred in S1, rather than in M1. Figure 3D shows a group averaged time frequency spectrogram of INMS, at the peak location from Figure 3C. The spectrogram shows that at this location, it is the beta oscillatory band which has the largest fractional change in power during stimulation, where we observed event related desynchronisation (ERD). After stimulation ceased (t = 1), post-event synchronisation (ERS) occurred, with an overshoot in beta power at t = 2-3 s, before returning to baseline. Performing the same localisation analysis on beta power in the ERS epoch showed this to be significant at the group level, but individual testing showed that this was only significant in three units, all from the same participant (see Supplementary material). We also found similar temporal properties in the t = 0-1 s and t = 2-3 s epochs in the alpha band, but few units showed individual level significance and the group level significance was limited (see Supplementary material).

The equivalent statistical maps from the vibrotactile experiments are shown in Figure 4. In Figure 4A, we observed similar results at the individual level, as depicted in Figure 3B, with spatial overlap between the vibrotactile stimuli applied at the receptive field location for each unit and INMS in contralateral S1 in 9/10 corresponding vibrotactile stimuli. In the ipsilateral hemisphere we saw a greater number of overlapping responses between vibrotactile stimulation and INMS (3/10) with the area of overlap more parietal than in contralateral S1. Figure 4B shows the group-averaged activation index ($\bar{A}_{\text{Vibrotactile}}$), again with a statistical threshold. A significant decrease in beta band oscillatory power was found over the contralateral postcentral gyrus during vibrotactile stimulation, agreeing with the findings of INMS. Figure 4C shows the peak location of the group level activation index, which was located at the same vertex as the INMS experiments. The time-frequency plot of the vibrotactile data is shown in Figure 4D, where we observed a similar behaviour in the alpha and beta bands. Analyses for the alpha band during the stimulation and resynchronisation epochs and the beta in the resynchronisation epoch can be found in the Supplementary material, but in short, none of these other conditions showed a high likelihood of significance in individual subjects.

Figure 4 –Results of vibrotactile stimulation at the perceived receptive field of the 10 FA1 units studied overlaid onto inflated cortical surfaces, gyri are represented as light grey surfaces sulci are dark grey. A) Results of the individual vibrotactile analysis, represented as a conjunction map, showing how many exhibited significant changes in beta band (13-30 Hz) power for a given location. B) The group averaged activation image of the 10 vibrotactile images, with a statistical (p < 0.05) threshold applied, showing vibrotaction elicited a reduction in beta oscillatory power over the contralateral sensorimotor area. C) A zoomed in portion of panel B, with the Brodmann areas as defined by the Glasser atlas overlaid. The peak reduction of beta power (represented as the magenta dot) is located in postcentral gyrus. D) Time frequency spectrogram of the peak location from panel C, showing a distinct event related behaviour in the beta band which extends beyond the initial 1 s of stimulation.

Figure 5 shows the spatiotemporal similarities of responses to single unit INMS and vibrotactile stimulation. Figure 5A provides a conjunction map representing the number of unit locations for which there was a significant change in beta power in both the INMS *and* corresponding vibrotactile experiments. Here, we saw that for any given location there was a maximum overlap in 8 subjects over the contralateral S1, whilst in the ipsilateral hemisphere only activation in a single subject showed co-significance. Further, at the group level, permutation testing found no significant differences between $\bar{A}_{\text{INMS}}$ and $\bar{A}_{\text{Vibrotactile}}$ in the beta band during stimulation at any location within the brain. Figure 5B shows the conjunction map within the right S1 and M1. The map shows three features; locations where $\bar{A}_{\text{INMS}}$ was within 90% of the most extreme value (magenta), $\bar{A}_{\text{Vibrotactile}}$ within 90% of the most extreme value (cyan), and areas from Figure 5A where $n = 8$ (yellow). Overlaying these highlights two regions in S1 where all three conditions are satisfied (marked in black), one within Brodmann Area 3b, and

one which straddles the borders of Brodmann areas 2 and 1, the latter of which agreed with the peak locations of $\bar{A}_{\text{INMS}}$ and $\bar{A}_{\text{Vibrotactile}}$. Figure 5C shows the group averaged oscillatory power in just the beta band, and compares the INMS beta timecourse to that of the vibrotactile experiments in the same location. The shaded areas represent the standard error across subjects. Non-parametric randomisation testing of the timecourses revealed no significant differences at any time point between the INMS and vibrotactile experiments.

Figure 5 – Comparison of the spatiotemporal properties between INMS and the vibrotactile experiments. A) A conjunction map of where both INMS and vibrotactile experiments showed significant changes in beta power for a given mechanoreceptive afferent, and how many experiments spatially overlapped for a given area. B) A zoomed map of the right hemisphere's central gyri showing the largest group averaged changes in power for INMS (magenta) and vibrotactile (cyan) and where they overlap with the locations of the largest spatial agreement as depicted in panel A (yellow). Any location where all three conditions held are represented in black. Brodmann areas, as defined using the Glasser atlas, have been overlaid for reference. C) The group averaged beta power timecourses for INMS (blue) and vibrotactile (red) responses, showing similar response timecourses in the contralateral (right) S1.

Discussion

Here, we aimed to show the feasibility of combining single unit INMS (using recently-developed stimulation hardware; Glover et al., 2017) with MEG, to probe whether we could reliably capture the spatial, spectral, and temporal properties of the cortical projections of individual single unit mechanoreceptive afferents in human glabrous hand skin.

Our results show that single unit microstimulation of these quantal elements of touch yields a robust reduction of oscillatory power in the beta (13-30 Hz) band which can be spatially localised using MEG, with 80% of individual

FA1 units showing significant oscillatory power reduction in contralateral S1. Furthermore, we compared our results with oscillatory responses to vibrotactile stimulation of the same receptive field location, and showed that our ability to localise the ERD of individual mechanoreceptive afferent stimulation using INMS was as reliable as localising sources derived from vibrotactile stimulation and that the group results from single FA1 mechanoreceptive afferent stimulation revealed similar spatiotemporal patterns to those from vibrotactile stimuli. The fundamental finding that single unit INMS ERD responses are robust and consistent with natural somatosensory stimuli will permit us to more dynamically probe central nervous system responses in humans. This will allow us to address a number of questions about the processing of touch from the different classes of mechanoreceptive afferent to a range of input patterns, including how varying the stimulus frequency and waveform may modify cortical responses.

In the current study we chose to assess beta power, which was motivated by a number of studies observing modulations in beta power to tactile stimulation using both intracranial (Brovelli et al., 2004; Crone et al., 1998; Witham and Baker, 2007) and extracranial modalities (examples include, but are not limited to Bauer et al., 2006; Cheyne et al., 2003; Gaetz and Cheyne, 2006; Pfurtscheller et al., 2002; Salenius et al., 1997; van Ede et al., 2010). Bauer and colleagues used a spatial-selective tactile task to investigate how attention affected oscillatory power during tactile stimulation, they found the reduction of beta power during stimulation was not significantly affected by attentional modulation, suggesting this was likely an automatic response to afferent stimulation, independent of processing demand (Bauer et al., 2006). The same study also noted that the amount of post-stimulation resynchronisation in the somatosensory cortex was reduced if participants were instructed to attend to, compared to those who were asked to ignore, the stimulus. The sensation from microsimulation is subtle and required subjects to attend to it, thus we expected a high level of attention. It is thus interesting to note that we saw only a few units showing significant rebounds of beta power, and these all originated from the same participant, a participant well accustomed to single unit INMS and so more familiar to the sensation, thus potentially requiring less attentional load. A study has shown that anticipation may modulate event related reductions in beta, in research employing a delay between a cue and the delivery of a stimulus, where there was contralateral ERD of the beta band, with the extent of the reduction monotonically increasing with delay time, implying an anticipatory element of somatosensory beta (van Ede et al., 2011). However, in the present study we used no explicit cues and therefore have confidence that our observations in beta power reduction are entirely due to stimulation.

Recently, there has been an emerging shift in how we interpret beta band oscillations in the somatosensory and motor cortices (van Ede et al., 2018). Rather than treating beta as a sustained oscillation, which is power modulated based on processing demand, it may be better to discuss beta in terms of transient (~50 ms) events, or bursts. These punctate events have been shown to be robust across multiple species (Sherman et al., 2016; Shin et al., 2017), and their occurrence in human experiments seems to imply these are inhibitory processes (Sacchet et al., 2015; Shin et al., 2017). Furthermore, the probability of the occurrence of a burst follows the trial-averaged beta power in the sensorimotor cortex; a rarefication of bursting events occurs during the ERD period and maximal likelihood coincides with the ERS. Furthermore, there are some early indications that these beta events form long-range transient synchronisations across the brain (Sacchet et al., 2015), which could make bursts a potential mechanism of functional connections in the brain. To be able to fully understand the mechanisms of these bursts, we require experimental validations for predictions which mathematical models of this bursting mechanism (Jones et al., 2009) make. Given the precise nature of single unit INMS, it a suitable experimental candidate to explore such mechanisms.

In a previous study by Kelly and colleagues (1997), INMS was combined with EEG, where they identified sustained evoked potentials that followed modulations in stimulation frequency. They found that both single unit INMS and vibrotactile stimulation could evoke clear frequency following responses (FFRs) and these were matched to the stimulation frequency (33 Hz) (Kelly et al., 1997). In our current study, we did not observe any such FFRs for neither INMS nor vibrotactile stimulation, although we used a stimulus frequency of 60 Hz. We can identify two potential reasons why this was the case, which are both linked to the experimental design. The first can be attributed to the number of trials per experiment. In the Kelly et al. study they used 200 trials of 1 s stimulation, whereas here we used 80 trials. Our choice of 80 was to optimise the experiment for capturing modulations in beta power, which in the sensorimotor system takes several seconds to return to baseline after an event (Barratt et al., 2017; Fry et al., 2016; Pakenham et al., 2018; Robson et al., 2016), and so to gain an adequate inter-trial interval, the total number of trials is less. Assuming zero trials rejected, the Kelly et al. study has a 58% signal to noise ratio (SNR) improvement over our paradigm and this may be crucial to see such effects. The second reason, which may be more important than number of trials, is the stimulation frequency. Our choice of 60 Hz stimulation was driven by this providing a clearly perceivable constant sensation for subjects when stimulating both SA1 and FA1 afferents (Johnson, 2001). However, it is likely that this frequency of stimulation is too high

to generate a (detectible) FFR within the somatosensory cortex with MEG. Previous literature on the non-invasive imaging mechanoreceptive FFRs (see Vlaar et al., 2015 for a brief review) implies that the bulk of studies have been performed at a stimulation frequency of less than 40 Hz, and those which use higher carrier frequencies (such as Tobimatsu et al., 1999) used amplitude modulation below 40 Hz. Studies characterising the optimal frequencies show that the optimal frequency for an FFR is around 20-26 Hz (Jamali and Ross, 2013; Snyder, 1992; Tobimatsu et al., 1999). There has been a single report into the detection of subharmonics (Langdon et al., 2011), but their carrier frequencies were in this optimal spectral window.

In our current study, we used bespoke flexible headcasts to improve the reliability of the spatial localisation of the oscillatory sources and comparison of INMS and vibrotactile responses between MEG scan sessions. The use of such headcasts has been shown to minimise head movement and the inter-session errors which arise from sensor-to-anatomy co-registration (Liuzzi et al., 2017). Since it was unknown how spatiotemporal responses to single unit INMS would manifest in the human cortex, we wanted to ensure that we maximized the robustness of localisation of the sources between INMS and vibrotactile sessions using optimal methods. However, given that one unit (Unit 2a) was captured without a headcast and was localised to somatosensory cortex (see Supplementary material), with a source amplitude similar to that of the vibrotactile stimulation, future studies may not necessarily require a headcast, as long as sufficient practices in reducing head motion and co-registration errors are adhered to. Further the use of the headcast required care when inserting the participant in into the MEG sensor array once a unit was found, which has the potential of causing small electrode movements, which may lead to the loss of the single unit nature of the stimulation. We originally intended to perform the search for mechanoreceptive afferents whilst the participant was within the MEG sensor array, however due to an unacceptable high level of capacitively-coupled mains-borne interference related to the MEG system earth in our installation this interference obscured the neural responses, making effective identification of single unit recordings near impossible whilst the head was inside the MEG sensor array. The reference and recording electrodes used for INMS have very different impedances which makes the Common Mode Rejection Ratio performance for the pre-amplifier quite poor, but usually in screened room environments this is not an issue. We ameliorated this problem by lowering the participant out of the MEG sensor array and cutting the power to the MEG gantry during the search, with the former of these solutions necessitating careful entry and exit into the sensor array before and after data collection. It should be noted that cryogen-free on-scalp MEG solutions are starting to come into use (Boto et al., 2018, 2017), and these will potentially provide flexibility for combined single unit INMS without compromising on spatial specify (Boto et al., 2016; Iivanainen et al., 2017).

In conclusion, combining the spatiotemporal specificity of MEG with the single unit INMS enables the interrogation of the central representations of different aspects of tactile afferent signalling within the human cortices. Our initial findings imply that beta power in the somatosensory cortex is significantly modulated from the stimulation of single afferent mechanoreceptors, and given the precise nature of the stimulation method, INMS may serve as a controlled means to explore the mechanisms behind beta bursting events.

Acknowledgements

This work was supported by the Medical Research Council [grant number MR/M022722/1, to STF], the Swedish Research Council [grant 2017-01717, to JW], and the Knut and Alice Wallenberg Foundation [project NeuroSQUID, to JW] We also acknowledge the Medical Research Council MEG UK partnership [grant number MR/K005464/1].

References

Barratt, E.L., Tewarie, P.K., Clarke, M.A., Hall, E.L., Gowland, P.A., Morris, P.G., Francis, S.T., Evangelou, N., Brookes, M.J., 2017. Abnormal task driven neural oscillations in multiple sclerosis: A visuomotor MEG study. Hum. Brain Mapp. 38, 2441–2453. https://doi.org/10.1002/hbm.23531

Bauer, M., Oostenveld, R., Peeters, M., Fries, P., 2006. Tactile Spatial Attention Enhances Gamma-Band Activity in Somatosensory Cortex and Reduces Low-Frequency Activity in Parieto-Occipital Areas. J. Neurosci. 26, 490–501. https://doi.org/10.1523/JNEUROSCI.5228-04.2006

Boto, E., Bowtell, R., Krüger, P., Fromhold, T.M., Morris, P.G., Meyer, S.S., Barnes, G.R., Brookes, M.J., 2016. On the potential of a new generation of magnetometers for MEG: A beamformer simulation study. PLoS One 11. https://doi.org/10.1371/journal.pone.0157655

440 Boto, E., Holmes, N., Leggett, J., Roberts, G., Shah, V., Meyer, S.S., Muñoz, L.D., Mullinger, K.J., Tierney, T.M., Bestmann, S., Barnes, G.R., Bowtell, R., Brookes, M.J., 2018. Moving magnetoencephalography towards real-world applications with a wearable system. Nature 555, 657–661. https://doi.org/10.1038/nature26147

Boto, E., Meyer, S.S., Shah, V., Alem, O., Knappe, S., Kruger, P., Fromhold, T.M., Lim, M., Glover, P.M., Morris, P.G., Bowtell, R., Barnes, G.R., Brookes, M.J., 2017. A new generation of magnetoencephalography: Room temperature measurements using optically-pumped magnetometers. Neuroimage 149, 404–414. https://doi.org/10.1016/j.neuroimage.2017.01.034

Brookes, M.J., Vrba, J., Robinson, S.E., Stevenson, C.M., Peters, A.M., Barnes, G.R., Hillebrand, A., Morris, P.G., 2008. Optimising experimental design for MEG beamformer imaging. Neuroimage 39, 1788–1802.

Brovelli, A., Ding, M., Ledberg, A., Chen, Y., Nakamura, R., Bressler, S.L., 2004. Beta oscillations in a large-scale sensorimotor cortical network: Directional influences revealed by Granger causality. Proc. Natl. Acad. Sci. 101, 9849–9854. https://doi.org/10.1073/pnas.0308538101

450 Cheyne, D., Gaetz, W., Garnero, L., Lachaux, J.P., Ducorps, A., Schwartz, D., Varela, F.J., 2003. Neuromagnetic imaging of cortical oscillations accompanying tactile stimulation. Cogn. Brain Res. 17, 599–611. https://doi.org/10.1016/S0926-6410(03)00173-3

Cheyne, D.O., 2013. MEG studies of sensorimotor rhythms: A review. Exp. Neurol. https://doi.org/10.1016/j.expneurol.2012.08.030

Crone, N.E., Miglioretti, D.L., Gordon, B., Sieracki, J.M., Wilson, M.T., Uematsu, S., Lesser, R.P., 1998. Functional mapping of human sensorimotor cortex with electrocorticographic spectral analysis. I. Alpha and beta event-related desynchronization. Brain 121, 2271.

Dale, A.M., Fischl, B., Sereno, M.I., 1999. Cortical surface-based analysis: I. Segmentation and surface reconstruction. Neuroimage 9, 179–194. https://doi.org/10.1006/nimg.1998.0395

Fischl, B., Sereno, M.I., Dale, A.M., 1999. Cortical surface-based analysis: II. Inflation, flattening, and a surface-based coordinate system. Neuroimage. https://doi.org/10.1006/nimg.1998.0396

460 Fry, A., Mullinger, K.J., O'Neill, G.C., Barratt, E.L., Morris, P.G., Bauer, M., Folland, J.P., Brookes, M.J., 2016. Modulation of post-movement beta rebound by contraction force and rate of force development. Hum. Brain Mapp. 37, 2493–2511. https://doi.org/10.1002/hbm.23189

Gaetz, W., Cheyne, D., 2006. Localization of sensorimotor cortical rhythms induced by tactile stimulation using spatially filtered MEG. Neuroimage 30, 899–908. https://doi.org/10.1016/j.neuroimage.2005.10.009

Glasser, M.F., Coalson, T.S., Robinson, E.C., Hacker, C.D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C.F., Jenkinson, M., Smith, S.M., Van Essen, D.C., 2016. A multi-modal parcellation of human cerebral cortex. Nature 536, 171–178. https://doi.org/10.1038/nature18933

Glover, P.M., Watkins, R.H., O'Neill, G.C., Ackerley, R., Sanchez-Panchuelo, R., McGlone, F., Brookes, M.J., Wessberg, J., Francis, S.T., 2017. An intra-neural microstimulation system for ultra-high field magnetic resonance imaging and magnetoencephalography. J. Neurosci. Methods 290, 69–78. https://doi.org/http://dx.doi.org/10.1016/j.jneumeth.2017.07.016

470 Hari, R., Salmelin, R., 1997. Human cortical oscillation: a neuromagnetic view through the skull. Trends Neurosci. 20, 44–49. https://doi.org/10.1016/S0166-2236(96)10065-5

Iivanainen, J., Stenroos, M., Parkkonen, L., 2017. Measuring MEG closer to the brain: Performance of on-scalp sensor arrays. Neuroimage 147, 542–553. https://doi.org/10.1016/j.neuroimage.2016.12.048

Jamali, S., Ross, B., 2013. Somatotopic finger mapping using MEG: Toward an optimal stimulation paradigm. Clin. Neurophysiol. 124, 1659–1670. https://doi.org/10.1016/j.clinph.2013.01.027

Johansson, R.S., Vallbo, A.B., 1979. Tactile sensibility in the human hand: relative and absolute densities of four types of mechanoreceptive units in glabrous skin. J. Physiol. 286, 283–300. https://doi.org/10.1113/jphysiol.1979.sp012619

Johnson, K.O., 2001. The roles and functions of cutaneous mechanoreceptors. Curr. Opin. Neurobiol. https://doi.org/10.1016/S0959-4388(00)00234-8

480 Jones, S.R., Pritchett, D.L., Sikora, M.A., Stufflebeam, S.M., Hamalainen, M., Moore, C.I., 2009. Quantitative Analysis and Biophysically

Realistic Neural Modeling of the MEG Mu Rhythm: Rhythmogenesis and Modulation of Sensory-Evoked Responses. J. Neurophysiol. 102, 3554–3572. https://doi.org/10.1152/jn.00535.2009

Kelly, E.F., Trulsson, M., Folger, S.E., 1997. Periodic microstimulation of single mechanoreceptive afferents produces frequency-following responses in human EEG. J. Neurophysiol. 77, 137–144.

Langdon, A.J., Boonstra, T.W., Breakspear, M., 2011. Multi-frequency phase locking in human somatosensory cortex. Prog. Biophys. Mol. Biol. 105, 58–66. https://doi.org/10.1016/j.pbiomolbio.2010.09.015

Liuzzi, L., Gascoyne, L.E., Tewarie, P.K., Barratt, E.L., Boto, E., Brookes, M.J., 2017. Optimising experimental design for MEG resting state functional connectivity measurement. Neuroimage 155, 565–576. https://doi.org/10.1016/j.neuroimage.2016.11.064

Maris, E., Oostenveld, R., 2007. Nonparametric statistical testing of EEG- and MEG-data. J. Neurosci. Methods 164, 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Meyer, S.S., Bonaiuto, J., Lim, M., Rossiter, H., Waters, S., Bradbury, D., Bestmann, S., Brookes, M., Callaghan, M.F., Weiskopf, N., Barnes, G.R., 2017. Flexible head-casts for high spatial precision {MEG}. J. Neurosci. Methods 276, 38–45. https://doi.org/https://doi.org/10.1016/j.jneumeth.2016.11.009

Nichols, T.E., Holmes, A.P., 2002. Nonparametric permutation tests for functional neuroimaging: A primer with examples. Hum. Brain Mapp. 15, 1–25. https://doi.org/10.1002/hbm.1058

Ochoa, J., Torebjörk, E., 1983. Sensations evoked by intraneural microstimulation of single mechanoreceptor units innervating the human hand. J. Physiol. 342, 633–654. https://doi.org/10.1113/jphysiol.1983.sp014873

Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M., 2011. FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput. Intell. Neurosci. 2011, 1.

Pakenham, D.O., Fry, A., Brookes, M.J., Francis, S.T., Mullinger, K.J., 2018. Post-stimulus beta responses are modulated by task duration. Hum. Brain Mapp. Under Revi.

Pantazis, D., Nichols, T.E., Baillet, S., Leahy, R.M., 2005. A comparison of random field theory and permutation methods for the statistical analysis of MEG data. Neuroimage 25, 383–394. https://doi.org/10.1016/j.neuroimage.2004.09.040

Pfurtscheller, G., Woertz, M., Müller, G., Wriessnegger, S., Pfurtscheller, K., 2002. Contrasting behavior of beta event-related synchronization and somatosensory evoked potential after median nerve stimulation during finger manipulation in man. Neurosci. Lett. 323, 113–116. https://doi.org/10.1016/S0304-3940(02)00119-2

Robson, S.E., Brookes, M.J., Hall, E.L., Palaniyappan, L., Kumar, J., Skelton, M., Christodoulou, N.G., Qureshi, A., Jan, F., Katshu, M.Z., Liddle, E.B., Liddle, P.F., Morris, P.G., 2016. Abnormal visuomotor processing in schizophrenia. NeuroImage Clin. 12, 869–878. https://doi.org/10.1016/j.nicl.2015.08.005

Saal, H.P., Bensmaia, S.J., 2014. Touch is a team effort: Interplay of submodalities in cutaneous sensibility. Trends Neurosci. https://doi.org/10.1016/j.tins.2014.08.012

Sacchet, M.D., LaPlante, R.A., Wan, Q., Pritchett, D.L., Lee, A.K.C., Hamalainen, M., Moore, C.I., Kerr, C.E., Jones, S.R., 2015. Attention Drives Synchronization of Alpha and Beta Rhythms between Right Inferior Frontal and Primary Sensory Neocortex. J. Neurosci. 35, 2074–2082. https://doi.org/10.1523/JNEUROSCI.1292-14.2015

Salenius, S., Schnitzler, A., Salmelin, R., Jousmäki, V., Hari, R., 1997. Modulation of human cortical rolandic rhythms during natural sensorimotor tasks. Neuroimage 5, 221–228. https://doi.org/10.1006/nimg.1997.0261

Sanchez Panchuelo, R.M., Ackerley, R., Glover, P.M., Bowtell, R.W., Wessberg, J., Francis, S.T., McGlone, F., 2016. Mapping quantal touch using 7 Tesla functional magnetic resonance imaging and single-unit intraneural microstimulation. Elife 5, e12812.

Sekihara, K., Nagarajan, S.S., Poeppel, D., Marantz, A., 2004. Asymptotic SNR of scalar and vector minimum-variance beamformers for neuromagnetic source reconstruction. IEEE Trans. Biomed. Eng. 51, 1726–1734. https://doi.org/10.1109/TBME.2004.827926

Sherman, M.A., Lee, S., Law, R., Haegens, S., Thorn, C.A., Hämäläinen, M.S., Moore, C.I., Jones, S.R., 2016. Neural mechanisms of transient neocortical beta rhythms: Converging evidence from humans, computational modeling, monkeys, and mice. Proc. Natl.

Acad. Sci. 113, E4885–E4894. https://doi.org/10.1073/pnas.1604135113

Shin, H., Law, R., Tsutsui, S., Moore, C.I., Jones, S.R., 2017. The rate of transient beta frequency events predicts behavior across tasks and species. Elife 6. https://doi.org/10.7554/eLife.29086

Singh, K.D., Barnes, G.R., Hillebrand, A., 2003. Group imaging of task-related changes in cortical synchronisation using nonparametric permutation testing. Neuroimage 19, 1589–1601. https://doi.org/10.1016/S1053-8119(03)00249-0

Smith, S.M., Nichols, T.E., 2009. Threshold-free cluster enhancement: Addressing problems of smoothing, threshold dependence and localisation in cluster inference. Neuroimage 44, 83–98. https://doi.org/10.1016/j.neuroimage.2008.03.061

530   Snyder, A.Z., 1992. Steady-state vibration evoked potentials: description of technique and characterization of responses. Electroencephalogr. Clin. Neurophysiol. Evoked Potentials 84, 257–268. https://doi.org/10.1016/0168-5597(92)90007-X

Stenroos, M., Nummenmaa, A., 2016. Incorporating and Compensating Cerebrospinal Fluid in Surface-Based Forward Models of Magneto- and Electroencephalography. PLoS One 11, e0159595. https://doi.org/10.1371/journal.pone.0159595

Tobimatsu, S., Zhang, Y.M., Kato, M., 1999. Steady-state vibration somatosensory evoked potentials: Physiological characteristics and tuning function. Clin. Neurophysiol. 110, 1953–1958. https://doi.org/10.1016/S1388-2457(99)00146-7

Torebjörk, H.E., Ochoa, J.L., 1980. Specific sensations evoked by activity in single identified sensory units in man. Acta Physiol. Scand. 110, 445–447. https://doi.org/10.1111/j.1748-1716.1980.tb06695.x

Torebjörk, H.E., Vallbo, Å.B., Ochoa, J.L., 1987. Intraneural microstimulation in man: Its relation to specificity of tactile sensations. Brain 110, 1509–1529.

540   Trulsson, M., Francis, S.T., Kelly, E.F., Westling, G., Bowtell, R., McGlone, F., 2001. Cortical responses to single mechanoreceptive afferent microstimulation revealed with fMRI. Neuroimage 13, 613–622.

Vallbo, Å.B., 1981. Sensations evoked from the glabrous skin of the human hand by electrical stimulation of unitary mechanosensitive afferents. Brain Res. 215, 359–363. https://doi.org/10.1016/0006-8993(81)90517-5

Vallbo, Å.B., Hagbarth, K.-E., 1968. Activity from skin mechanoreceptors recorded percutaneously in awake human subjects. Exp. Neurol. 21, 270–289.

Vallbo, Å.B., Johansson, R.S., others, 1984a. Properties of cutaneous mechanoreceptors in the human hand related to touch sensation. Hum Neurobiol 3, 3–14.

Vallbo, Å.B., Olsson, K.Å., Westberg, K.G., Clark, F.J., 1984b. Microstimulation of single tactile afferents from the human hand. Sensory attributes related to unit type and properties of receptive fields. Brain 107 (Pt 3, 727–749. https://doi.org/10.1093/brain/107.3.727

550   van Ede, F., de Lange, F., Jensen, O., Maris, E., 2011. Orienting Attention to an Upcoming Tactile Event Involves a Spatially and Temporally Specific Modulation of Sensorimotor Alpha- and Beta-Band Oscillations. J. Neurosci. 31, 2016–2024. https://doi.org/10.1523/JNEUROSCI.5630-10.2011

van Ede, F., Jensen, O., Maris, E., 2010. Tactile expectation modulates pre-stimulus A-band oscillations in human sensorimotor cortex. Neuroimage 51, 867–876. https://doi.org/10.1016/j.neuroimage.2010.02.053

van Ede, F., Quinn, A.J., Woolrich, M.W., Nobre, A.C., 2018. Neural Oscillations: Sustained Rhythms or Transient Burst-Events? Trends Neurosci. In Press.

Vlaar, M.P., van der Helm, F.C.T., Schouten, A.C., 2015. Frequency Domain Characterization of the Somatosensory Steady State Response in Electroencephalography. IFAC-PapersOnLine 48, 1391–1396. https://doi.org/10.1016/j.ifacol.2015.12.327

Witham, C.L., Baker, S.N., 2007. Network oscillations and intrinsic spiking rhythmicity do not covary in monkey sensorimotor areas. J.
560   Physiol. 580, 801–814. https://doi.org/10.1113/jphysiol.2006.124503