Full title: The nuclear RNAi factor, NRDE2, prevents the accumulation of DNA damage during mitosis in stressful growth conditions

Short Title: Heterochromatin associated proteins and DNA damage in mitosis

Aarati Asundi[1, 2], Srivats Venkataramanan[1], Atsushi Suzuki[3], Stephen N. Floor[1], Andrei Goga[1, 4, 5], Noelle L'Etoile[1]

1. Department of Cell and Tissue Biology, University of California, San Francisco, CA, USA
2. Biomedical Sciences Graduate Program, University of California, San Francisco, CA, USA.
3. Department of Oncology, Astellas, 2-5-1, Nihonbashi-Honcho, Chuo-ku, Tokyo 103-8411, Japan
4. Department of Medicine, University of California, San Francisco, CA, USA.
5. Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco, CA, USA.

Abbreviations

PGC: Proliferative Germ Cell

NRDE-2: Nuclear RNAi Defective-2

AuBK: Aurora B Kinase

Mrt: Mortal germ line

Abstract

Organisms have evolved multiple mechanisms to prevent and repair DNA damage to

protect the integrity of the genome, particularly under stressful conditions. Unrepaired

DNA damage leads to genomic instability, aneuploidy, and an increased risk for cancer.

Before the cell can divide, it must repair damaged DNA and it is thought that this

process requires global silencing of most transcription. In *C. elegans*, NRDE-2, in complex

with other nuclear factors and guided by small RNA, directs heterochromatin formation

and transcriptional silencing of targeted genes. Additionally, when *C. elegans* are

cultivated at high temperatures, NRDE-2 is required to maintain germ line immortality.

However, the role of NRDE-2 in maintaining the physical integrity of the genome is not

understood. We show here that loss of NRDE2 in either nematode or human cells

induces the accumulation of DNA damage specifically under conditions of stress, such as

cultivation at a high temperature in *C. elegans* or Aurora B Kinase oncogenic

overexpression in the MCF10A epithelial breast cell line. In addition, we found that

NRDE2 interacts with β-actin in unstressed mammalian cells. This interaction is

dramatically reduced upon DNA damage. The oligomerization state of nuclear actin

alters its association with targets, which in turn, regulates their function. Monomeric

nuclear actin binds to heterochromatin remodeling factors and transcriptional activators

while filamentous actin has been implicated in DNA repair processes. Here we show that

NRDE2 associates with actin only when DNA is intact and the bulk of nuclear actin is

monomeric. Thus, NRDE2 may dissociate from actin when it becomes filamentous as a

result of DNA damage. This implies that, NRDE2, in its role as a heterochromatin factor,

binds to monomeric actin, protecting the genome from DNA damage in stressful

conditions. In this way, heterochromatin factors may associate with the actin dependent

DNA repair process to allow appropriate mitotic progression and maintain genomic

integrity.

Introduction

Organisms have developed multiple ways to protect genomic integrity during mitotic

cell division. Maintenance of the physical and informational integrity of the genome

ensures that cells accurately transmit genetic information from one generation to the

next, thereby safeguarding against diseases associated with genomic instability, such as

cancer. Genomic instability is caused primarily through DNA damage. DNA damage can

be incurred in a number of ways, including naturally during DNA replication, meiotic

recombination, improper chromosome segregation, and transposable elements. DNA

damage can also be caused by external sources such as UV irradiation or drugs. If the

cell attempts to divide before the DNA is repaired, the unrepaired DNA damage can

result in mitotic delay, aneuploidy, or apoptosis, which in turn can lead to disorders in

the organism including cancer and infertility.

To protect against genomic instability, cells must fulfill a checkpoint at the Gap2 to

Mitosis (G2/M) transition [46] at which the cell repairs any DNA damage sustained

during the previous cell cycle phases. If the requirements of the G2/M checkpoint are

not satisfied, the cell will express a number of cell cycle inhibitors, leading to an arrest in

G2. [30]

DNA repair is especially important in the germ line where strong evolutionary pressure

to maintain the integrity of the genome of germ cells selects against harmful mutations.

Thus, the germ line provides a window into how the environment affects the cell cycle.

The nematode *Caenorhabditis elegans* has a well-characterized germ line and

reproductive cycle, which can be used to study the effects of improper mitosis on cell

fate and fertility. The adult *C. elegans* gonad contains a stem cell niche, which

mitotically proliferates to self-renew. These proliferating germ cells (PGCs) constitute

the only pool of mitotically dividing cells in the adult worm. As the PGCs divide, they

travel proximally along the gonad, where they transition from mitotic proliferation to

meiotic division and ultimately differentiate into spermatozoa or oocytes. Therefore,

maintaining mitotic integrity in the PGC pool is essential for ensuring the reproductive

integrity of the animal.

Previous studies show that the number of PGCs is susceptible to internal molecular

changes as well as external stresses, which can impact the worm's fertility. In addition to

canonical signaling, such as the GLP-1/Notch, [31-32] DAF-7/TGFb [33] and IGF-1/insulin

[34] pathways, many small RNA pathway proteins have also been extensively studied in

the context of maintaining the PGC pool and fertility. For example, mutations in CSR-1

(argonaute that binds endogenous 22G siRNAs)[36], EGO-1 (putative RdRP required for

small RNA biosynthesis) [37, 38], DCR-1 (cleaves dsRNA) [39, 40] and EKL-1 (Tudor

domain protein required in 22G RNA biosynthesis) [36, 41] have all been shown to have

significant mitotic and meiotic germ cell defects. Furthermore, external stresses such as

starvation or changes in cultivation temperature can directly impact the PGC pool and

subsequently affect fertility. [34, 35]

In particular, the nuclear RNAi protein, NRDE-2 is required to maintain reproductive

integrity of *C. elegans* in response to environmental stress. *Nrde-2* null mutants remain

fertile indefinitely at when grown under normal conditions at 20°C, however, when

propagated at 25°C they produce fewer progeny each successive generation until

becoming completely sterile by the fourth generation. [1, 16] This progressive sterility is

termed a mortal germ line phenotype (Mrt). *Nrde-2* null mutant worms grown at 25°C

also express high levels of repetitive RNA as compared to wild type worms propagated

under the same conditions. This is consistent with the hypothesis that NRDE-2 acts to

silence transposons and repress repetitive sequences, [16, 47] which are naturally

occurring DNA damaging agents and can affect the informational and physical integrity

of the genome.

We and others have hypothesized that the Mrt phenotype seen in *C. elegans* mutants

lacking heterochromatin factors, piRNA, or nuclear RNAi factors, including NRDE-2,

might result from accumulated DNA damage in the germ line. [47] The function of

NRDE-2 in preventing DNA damage accumulation seems to be conserved across species

but had not been directly studied in the *C. elegans* germ line. In the fission yeast, *S.*

pombe, loss of the NRDE-2 homolog (known as nrl1) results in increased DNA damage,

which was posited to arise from unresolved R-loops. [17] R-loops are RNA-DNA hybrids

that arise from transcription of repetitive elements in which the RNA message invades

the double stranded DNA helix. The DNA-RNA hybrid is more stable than the DNA-DNA

hybrid and can cause DNA damage when the replication fork collides with it.

Transcriptional silencing of repetitive elements by heterochromatin factors such as

NRDE2 was thought to prevent the possibility of R-loop formation. [48] A recent study in

HEK293 (human embryonic kidney) cells, however, showed the human homolog of

NRDE-2 (known as NRDE2 or C14orf102) plays a role in DNA damage response and

preventing the accumulation of double stranded breaks, independently of R-loop

formation and resolution. [22] Thus, NRDE2 may act in its capacity as a heterochromatin

factor to repress transcription of repetitive elements to limit both R-loop formation and

transposition, which are two events that pose threats to the integrity of the genome.

Recently, it has been increasingly appreciated that heterochromatic regions of the

genome (nuclear speckles) are associated with monomeric actin and that

heterochromatin remodelers such as SWI/SNF , SWR1 and INO80 are tightly bound to

actin monomers and this association is important for their ability to bind to chromatin.

[45] Thus, stressors such as high temperature and DNA damage have been shown to tilt

the balance of nuclear actin from monomeric to filamentous, which has been shown to

be required for efficient DNA damage repair. [25] How heterochromatin associated

factors and nuclear actin polymerization status are linked, however, remains unclear.

In this paper, we show evidence that NRDE2 prevents the accumulation of DNA damage

particularly under conditions of stress in proliferating cell populations. In the *C. elegans*

germ line, NRDE-2 loss resulted in the accumulation of DNA damage only when the

worms were propagated at a high temperature. This correlated with a decrease in the

number of PGCs and was consistent with the previously reported Mrt phenotype.

Moreover, in human cells, we found that NRDE2 loss did not significantly impact the

proliferation of normal epithelial breast cells MCF10A unless they overexpress the

oncogene Aurora B Kinase (AuBK). In these cells, we observed a delay in mitotic

prophase as well as an increase in DNA damage. This indicates that NRDE2 may have a

role in repair of DNA damage. Finally, we observed a strong protein-protein interaction

between NRDE2 and β-actin in normally proliferating MCF10A cells, which was

weakened upon the introduction of DNA damaging agents. Thus, we suggest a novel link

between this heterochromatin factor, actin, and DNA repair.

Results

Loss of NRDE-2 coupled with propagation at high temperature incurs DNA damage in

the mitotic zone of the *C. elegans* gonad

Previous studies in *C. elegans* have shown that loss of NRDE-2 does not affect germ line

immortality when the worms are propagated at 20°C. We have observed that although

nrde-2 (gg91), null mutants remain fertile indefinitely when cultivated at 20°C, these

worms have a lower brood size as compared to wild type (Fig S1, p=0.0148, n=4). In

contrast, if the worms are cultivated at 25°C, a more stressful temperature, *nrde-2*

(gg91) mutants exhibit a Mrt phenotype that increases in severity with each successive

generation.[1] We asked whether the decrease in brood size of *nrde-2 (gg91)* mutant

worms was a result of physiological defects in the gonad. We observed a significant

decrease in the number of proliferating germ cells (PGCs) in the mitotic zone in *nrde-2*

(gg91) worms propagated at 20°C as compared to wild type (Fig 1A and B, left,

p=0.0115).   When we propagated the worms at the stressful temperature of 25°C for 2

generations (G2), we found that the number of PGCs was significantly decreased as

compared to wild type G2 worms grown at 25°C (Fig 1A and B, left, p<0.0001).

Since the Mrt phenotype in *nrde-2 (gg91)* mutants depends on cultivation at 25°C, we

asked whether the cells accumulated DNA damage at this temperature, which would

ultimately cause sterility. We used immunofluorescence to stain for RAD-51, a marker of

DNA damage repair, and found that *nrde-2 (gg91)* G2 worms grown at 25°C had

significantly increased RAD-51 foci in the mitotic zone as compared to wild type G2

worms grown at the same temperature (Fig 1A and B, right, p<0.0001). We then asked

whether the aberrant accumulation of DNA damage in the mitotic PGCs resulted in

defects as the cells entered meiosis in the proximal gonad and progressed towards

oocyte formation. We expressed the apoptotic cell corpse marker, CED-1::GFP, which

labels apoptotic cell corpses in the proximal loop of the gonad, in *nrde-2 (gg91)* mutants

and wild type worms. We found that when the worms were propagated at 20°C, *nrde-2*

(gg91) mutants showed a similar number of CED-1::GFP labeled germ cell corpses as

wild type worms. However, *nrde-2 (gg91)* G1 and G2 worms showed increased numbers

of CED-1::GFP labeled germ cell corpses as compared to wild type G1 and G2 worms,

respectively (Figure 1C and D, p<0.0001 for both).

We conclude that when worms are cultivated at a normal, non-stressful temperature,

loss of NRDE-2 has minor endurable impacts on fertility and integrity of the *C. elegans*

germ line. However, when NRDE-2 loss is coupled with an external stress, such as a high

cultivation temperature, the DNA damage and subsequent cell death in the gonad

compounds over the generations, ultimately resulting in sterility. Therefore, we

propose the NRDE-2 may have a role in preventing DNA damage accumulation in the

mitotic PGCs in order to maintain integrity of the *C. elegans* germ line.

Loss of NRDE2 coupled with Aurora B Kinase overexpression in MCF10A cells results in

DNA damage accumulation

To gain a biochemical and cell biological understanding of the role of NRDE-2, we

decided to utilize the actively dividing MCF10A cell line. MCF10A cell are a normal

human epithelial cell line derived from reduction mammoplasty. [4]

We used a lentiviral vector to stably express an eGFP-tagged version of human NRDE2 in

the MCF10A cells. The eGFP::NRDE2 localized to puncta within the nucleus (Fig 2Ai).

These puncta were reminiscent of staining of heterochromatin and heterochromatin

associated factors. [49] Once the cell entered prophase, as characterized by chromatin

condensation, the eGFP signal became diffuse and remained so throughout mitosis (Fig

2Aii-v) until cytokinesis, when eGFP::NRDE2 seemed to re-localize to puncta in the

nuclei of the daughter cells (Fig 2Avi). The dynamics of this pattern of staining is similar

to that of human RNA pol II transcription machinery proteins TBP (TATA box-binding

protein) and TAF12 (TATA box-binding protein associated factor 12) in HeLa cells. [29]

These data are therefore consistent with a conserved role for NRDE2 as a transcriptional

regulator in human cells.

To further test the hypothesis that NRDE-2 regulates DNA damage and mitosis under

stressful conditions, we stably overexpressed an HA-tagged version of the oncogene

Aurora B kinase (AuBK OE) in eGFP::NRDE-2 expressing MCF10A cells. AuBK regulates a

number of mitotic events, including chromatin condensation, [5] microtubule-

kinetochore attachment, [6], [7] and abscission. [8] Loss of this regulation has been

linked to genetic instability, which results in aneuploidy, a key hallmark of cancer.

Accordingly, AuBK over-expression has been linked to many cancer types and correlates

with poor prognosis for patients. [11-12] The localization of eGFP::NRDE2 in AuBK OE

MCF10A cell lines throughout the cell cycle was identical to that of eGFP::NRDE2 in

control cells (Fig 2A vii and xii).

Consistent with AuBK's role in mediating chromatin condensation at the G2/M

transition, AuBK OE MCF10A cells showed increased levels of phosphorylated Histone

H3 (Fig 2B, middle, p=0.0337, n=4), a known target of Aurora B Kinase. We also

observed a significant increase in γH2AX (H2AX phosphorylation on S139), a common

marker of DNA damage (Fig 2B, right, p=0.015, n=5). Therefore, we utilized AuBK

overexpression as a tool to study the effects of NRDE2 loss on DNA damaged cells.

Next, we assessed the efficacy of using NRDE2 targeted siRNA as a tool to knockdown

NRDE2 in MCF10A cells. We observed that cells treated with NRDE2 siRNA showed >90%

decrease in eGFP::NRDE2 expression as compared to cells treated with a non-targeting

siRNA control (Fig 2C). We then asked whether the amount of DNA damage in control or

AuBK OE cells was altered upon NRDE2 knockdown. Interestingly, we found when

control cells were treated with NRDE2 siRNA, there was no increase in γH2AX as

compared to control cells treated with non-targeting siRNA (Fig S3, p=0.11, n=5).

However, loss of NRDE2 in AuBK OE cells resulted in increased DNA damage as

compared to AuBK OE cells treated with non-targeting siRNA (Fig 2D, p=0.048, n=5). We

conclude that loss of NRDE2 has a minimal effect on DNA damage in normally

proliferating cells. However, in AuBK OE cells, which show an increased basal level of

DNA damage, loss of NRDE2 further increases the accumulation of DNA damage.

Loss of NRDE2 coupled with Aurora B Kinase overexpression in MCF10A cells results in

mitotic delay

We asked whether the increase in DNA damage observed in the AuBK OE cells upon

NRDE2 loss would result in an observable change in mitotic cell proliferation. We

observed that when AUBK OE cells were treated with NRDE2 siRNA, there was a

decrease in overall metabolic cell activity in the culture (Fig S2) and a significant increase

in the number of rounded cells (Fig 3A). Cells commonly round up due to a stall in

mitosis or when the cell is about to undergo apoptosis, both of which could explain the

observed decrease in metabolic cell activity. To differentiate between these two

possibilities, we used live cell imaging to record cell growth after NRDE2 knockdown. We

found that AuBK OE cells treated with non-targeting siRNA took a longer time to

complete mitosis than control cells. Furthermore, when AuBK OE cells were treated with

NRDE2 siRNA, there was a significant increase in the average time through mitosis (Fig

3B and C). However, we also noticed that a small proportion of AUBK-overexpressing

cells required a significantly longer time to complete mitosis upon NRDE2 siRNA

treatment (Fig 3B and C). The rounded morphology of the cells, the compaction of the

chromatin and the concentration of the DNA at the cell center, indicated that this stall

may occur during the early stages of mitosis.

We propose that the rounded cells observed upon NRDE2 knockdown in AUBK-

overexpressing cells is due to a delay in mitosis. To further characterize this delay, we

calculated the mitotic index of these cells. Consistent with the time lapse data, we found

that AuBK OE cells had a higher mitotic index than control cells. However, we found no

significant difference between the mitotic index of non-targeting siRNA treated cells and

the NRDE2 siRNA treated cells in either the control or AuBK OE (Fig S3). We surmised

that this could be due to the fact that the percentage of cells with a significant delay in

mitosis is so low that these cells were not sufficient to affect the mitotic index.

We next asked whether we could identify the stage at which mitosis was delayed in

AUBK-OE cells. We quantified the number of cells in each stage of mitosis in a fixed

population of cells and found more cells in prophase upon NRDE2 knock down (Fig 3E p=

0.025, n=4). This is also consistent with the time lapse analysis of AuBK OE cells delayed

in early mitosis (Fig 3C) leading us to conclude that the delay seen in NRDE2 knockdown

AUBK-overexpressing cells may be occurring at prophase.

Mammalian NRDE2 interacts with β-actin in healthy, normally proliferating cells, and

this interaction is weakened upon DNA damage

Our finding that NRDE2 specifically mitigates DNA damage accumulation in sensitized or

stressed cell populations led us to ask whether NRDE2 interacts with different partners

under these conditions. We decided to probe this question using co-

immunoprecipitation experiments, using an anti-GFP antibody to detect eGFP::NRDE2 in

MCF10A lysates (Fig 4A and Fig S5).

In *C. elegans*, NRDE-2 interacts with other nuclear RNAi factors NRDE-1 and NRDE-4, as

well as the worm specific argonaute NRDE-3. [11] In contrast, NRDE2 does not interact

with other RNAi factors in S. pombe yeast cells or in human cells. [22, 43, 44] Consistent

with these findings, we saw no interaction between eGFP::NRDE2 and any of the known

human Argonautes AGO-1/2/3/ or -4 (Fig S6).

Interestingly, we observed a strong interaction between NRDE2 and β-actin in control

cells (Fig 4A). Furthermore, this interaction seemed to be weakened in AuBK OE cells

(Fig 4A and B, p=0.036, n=5). Although it is important to note that co-IP experiments do

not necessary reflect a direct interaction between two proteins, these results may

suggest an interesting function for NRDE2 in DNA damaged cells. Nuclear actin forms

filaments upon DNA damage to aid in repair. [25] Since AuBK overexpression increases

DNA damage in MCF10A cells, we hypothesized that the interaction between NRDE2

and β-actin may be weakened in these cells due to the nuclear actin forming filaments

in response to DNA damage. To test whether DNA damage altered the interaction

between NRDE2 and β-actin, we treated the cells with cisplatin, a DNA damaging

reagent. In both control and AuBK overexpressing MCF10A cells, the interaction

between NRDE2 and β-actin was reduced as compared to untreated cells (Fig 4A and B,

p=0.045, n=3). Therefore, we propose that NRDE2 interacts with β-actin only when the

DNA is intact. However, when DNA damage in induced by stressors such as AuBK over-

expression or the presence of cisplatin, the interaction between NRDE2 and β-actin is

disrupted, perhaps as a result of the nuclear actin forming filaments to aid in DNA

repair.

Discussion

This study describes a previously uncharacterized role for NRDE2, a putative nuclear

RNA-binding protein, in protecting against DNA damage in stressed cells. Our study is

the first to study the effect of NRDE2 loss in normal, unstressed cells as compared to in

stressed cells. We have also shown that this protective role for NRDE2 in stressed cells is

conserved in *C. elegans* and mammalian mitotic cells.

Previously, NRDE2 was described to protect against the accumulation of DNA damage,

as assessed by an increase in γH2AX levels upon NRDE2 loss in HEK293 cells. [22] We

believe these results are consistent with these results as HEK293 cells are not a normally

proliferating cell line. The HEK293 line was originally obtained by transforming human

embryonic kidney cells with human adenovirus type 5, resulting in a 4.5kb insertion of a

viral fragment in chromosome 19 which interferes with cell death and cell cycle control

pathways. [23] HEK293 cells are also hypotriploid, [24] which may further affect mitosis.

Consistent with this point, we found that NRDE2 knockdown resulted in an increase of

γH2AX levels only in AuBK-overexpressing MCF10A cells, not in the normal, unstressed

control cells. Since AuBK overexpressing causes an increase in basal levels of DNA

damage, we propose that NRDE2 in mammalian cells results in an accumulation of DNA

damage only in cells with increases mitotic stress, similar to the results seen in *C.*

elegans.

In *C. elegans*, loss of NRDE-2 alone does not seem to severely compromise the mitotic

PGCs or brood size, as evidenced by the worms remaining indefinitely fertile and

exhibiting relatively normal development. Upon the addition of a stressful 25°C

cultivation temperature, however, there is a drastic reduction in the number of PGCs

and brood size. Under these stressful conditions, DNA damage accumulates in the germ

cells, presumably resulting in the previously observed Mrt phenotype. This trans-

generational Mrt phenotype is similar to fertility defects seen in the RNAi spreading

defective mutants rsd-2 and rsd-6. RSD-2 and RSD-6, along with NRDE-2 were proposed

to induce genome wide epigenetic silencing to maintain germ cell immortality and

fertility. RSD-2 and RSD-6 are also thought to act in a different pathway than the 22G-

RNA associating argonautes CSR-1 and ALG3/4, which display Mrt phenotypes due to

defects in spermatogenesis. [16] The effect of NRDE-2 loss on spermatogenesis remains

unclear.

The defects seen in *nrde-2 (gg91)* mutants may also be due to improper chromatin

condensation and chromosome segregation. It has been previously reported that CSR-1

is required for the localization of CDE-1, a nucleotidyltransfease protein that is

responsible for the uridylation of siRNAs bound by CSR-1, to mitotic chromosomes in

embryos. [19] Loss of CDE-1 results in an accumulation of siRNAs and improper gene

silencing, as well as mitotic and meiotic chromosome segregation defects. NRDE-2, by

contrast, associates with other NRDE pathway proteins to recruit methlytransferases

MES-2 and SET-25, which direct the deposition of methylation marks to silence loci

targeted by siRNAs. [20-21] Therefore, NRDE-2 loss may also result in improper gene

silencing in the *C. elegans* germ line, resulting in mitotic defects, which are exacerbated

by the stress of high temperature.

Interestingly, we observed a strong association between NRDE2 and β-actin in MCF10A

cells that was diminished upon DNA damage causing agents. Since NRDE2 localizes to

puncta in the interphase nucleus, we hypothesize that NRDE2 likely interacts with

nuclear actin. One potential model for this interaction is that NRDE2 binds to

monomeric nuclear actin in the unstressed cell. Monomeric nuclear actin associates

with chromatin remodeling complexes, and might be involved in chromatin binding. [45]

However, upon DNA damage, nuclear actin assembles into filaments and promotes DNA

damage repair. [25] NRDE2 may be displaced or unable to bind to filamentous actin, and

as a result, show a weakened interaction on β-actin upon DNA damage. It is still unclear

whether NRDE-2 functions in preventing the formation of DNA damage or as a

chromatin associated sensor of DNA damage in the β-actin dependent DNA repair

pathway. Further study is needed to characterize the function of NRDE2, and possibly

other heterochromatin factors, in preventing DNA damage accumulation and

maintaining genomic stability. The interaction between NRDE2 and β-actin is especially

exciting as it may indicate a new mechanistic role for NRDE2 in the nucleus beyond gene

silencing and epigenetic inheritance.

Materials and Methods

C. elegans strains

All strains were cultured on Nematode Growth Medium (NGM) plates seeded with E.

coli OP50. Bristol N2 was the wild type strain used. Loss of NRDE-2 was studied using the

mutant allele *nrde-2 (gg91).* Worms were crossed into a strain expressing ced-1p::GFP+

lin-15 (+) to study apoptotic cells.

Gonad Staining

Gonads were dissected from young adults (24 hours post-L4 stage) in 1xEGG buffer

(25mM HEPES, pH7.3; 118mM NaCl, 48mM KCl, 2mM CaCl2, 2mM MgCl2) and fixed in

2% (final concentration) formaldehyde. The following primary antibodies were used:

rabbit anti-RAD-51 and chicken anti-HTP-3. Fixed gonads were imaged with exposure

times of 100ms with DAPI, GFP, TRITC, and CY5 filter cubes and a mercury arc lamp on a

Zeiss AxioPlan2 epifluorescence microscope (operated by MicroManager 1.4.13) with a

60x 1.3 DIC oil objective and a QIClick camera (QImaging). Images were recorded with

a Z-optical spacing of 2 μm and analyzed using the Image J software.

<u>Progeny Assay</u>

Sixteen L4-staged worms were individually placed on a 5cm NGM plate with op50. Every

24 hours, the worm was moved to a fresh plate for 10 consecutive days. 48 hours after

the original worm was removed from the plate, the number of living progeny was

counted.

<u>Cell Culture</u>

MCF10A cells lines expressed a vector conferring puromycin/ blastocydin antibiotic

resistance that was empty (control) or that drove the overexpression of Aurora B Kinase

(AuBK OE). MCF10A cells were maintained in Dulbecco's modified Eagle's medium/F-12,

phenol red free medium supplemented with 5% heat inactivated horse serum, 20 ng/ml

recombinant EGF (Invitrogen), 0.5 μg/ml hydrocortisone (Sigma), 100 ng/ml cholera

toxin (Sigma), and 10 μg/ml insulin (Sigma). U20S cells were maintained using

Dulbecco's modified Eagle's medium with phenol red supplemented with 10% FBS.

<u>siRNA knockdown</u>

We used the following Thermo Scientific siRNAs: siGENOME Non-Targeting siRNA

Pool#2 (D-001206-14-20), siGENOME Human NRDE2 (55051) siRNA- SMARTpool (M-

013794-00-0005). siRNAs were added to the cell culture at a final concentration of 5nM

in OPTIMEM Reduced Serum Media with Lipofectamine 2000 Reagent (1ug/ml final

concentration) (Invitrogen) for 24 hours. Cells were then washed 1x with PBS and fresh

MCF10A media was added. 48 hours after changing media, the cells were harvested for

analysis.

<u>Western Blot</u>

Cultured cells were homogenized in Laemmli Buffer (60 mM Tris–HCl pH 7.6, 2% SDS,

1mM DTT) containing COMPLETE protease inhibitor cocktail (Roche #118361700001)

and phosphatase inhibitors (Santa Cruz Biotechnology)). Protein concentrations were

determined by performing DC Protein Assay (Bio-Rad) using BSA as standard. Protein

extracts were resolved using Bolt 4%-12% Bis-Tris PAGE gels (Invitrogen) with 1x MES

SDS Running Buffer (Life Technologies). Transfer to nitrocellulose membranes (Life

Technologies) was performed on an iBlot apparatus (Invitrogen). Membranes were

probed with primary antibodies overnight on a 4°C shaker and then incubated with

horseradish peroxidase (HRP)-conjugated secondary antibodies, and signals were

visualized with ECL (Bio-Rad) or Visualizer Western Blot Detection Kit, rabbit (Millipore).

The following primary antibodies were used: GFP (Rockland, 600-101-215), phospho-

Histone H3 (Ser10) (Cell Signaling Technologies, #9701), phospho-Histone H2A.X (Ser

139) (Cell Signaling Technologies, #9718), HA-tag (Cell Signaling Technologies #2367),

PARP (Cell Signaling Technologies, #9542), β-actin HRP (Santa Cruz Biotechnology, sc-

47778)

Cell Immunofluorescence

Cells were grown on a sterilized glass coverslip (Fisherbrand). After siRNA treatment, the

cells were fixed with ice cold 100% methanol for 3 minutes at -20°C and rehydrated with

TBST (1xTBS, 0.1% Tween 20). Antibodies were made up in blocking solution (2gBSA

powder, 0.1g NaN3, 100mL TBST). The following primary antibodies were used: human

anti-centromere (CREST, 1:50, Antibodies Incorporated 15-234) mouse anti-a-tubulin

(DM1a, 1:1000, Sigma T6199), rabbit anti-g-tubulin (1:1000, Sigma T3559). Cells were

mounted with Vectashield Mounting Medium with DAPI (Vector Laboratories H-1200).

Time lapse Imaging

MCF10A cells expressing H2B::mCherry were imaged at 37°C with a 100x NA 1.49

objective lens (CFI APO TIRF; Nikon) on an inverted microscope system (TE2000 Perfect

Focus System; Nikon) equipped with a Borealis modified spinning disk confocal unit

(CSU10; Yokogawa) with 200-mW, 405 nm, 488 nm, 561 nm and 643 nm solid-state

lasers (LMM5; Spectral Applied Research), electronic shutters, a Clara cooled scientific-

grade interline CCD camera (Andor), and controlled by NIS-Elements software (Nikon).

Co-Immunoprecipitation

Cells were collected and lysed in cold lysis buffer (150mM NaCl/ 1.0% IGEPAL/ 50mM

Tris-Cl (pH 7.4)) containing Complete protease inhibitor cocktail (Roche, 118361700001)

by nutating at 4°C for 30 minutes. The lysate was spun at 14K G, and supernatant was

transferred to a new tube, and cleared using 50ul of Dynabeads Protein A (Invitrogen,

10001D). Pre-cleared lysate was removed from the beads using magnetic separation

(input sample was collected at this stage), added to 5 ul of anti-GFP antibody (Rockland,

600-101-215), and nutated at 4°C for 2 hours. Antibody-protein complexes were

isolated by the addition of Dynabeads Protein A (Invitrogen, 10001D), overnight

nutation and magnetic separation. The beads were washed with the lysis buffer

containing protease inhibitors five times. IP western blot bands were normalized to 5%

sample input.

Acknowledgements

Thanks to Dr. Scott Kennedy for providing the nrde-2 (gg91) strain and to Dr. Needhi

Bhalla (UC Santa Cruz) for providing the CED-1::GFP strain. We would like to thank Dr.

Gina Caldas (Dernburg lab at UC Berkeley) for reagents and advice about *C. elegans*

gonad immunofluorescence staining. We extend a special thanks to Christina Hueschen

(Dumont lab at UCSF) for creating a nuclear GFP quantification MatLab script. We would

also like to thank Dr. Sophie Dumont and Dr. Torsten Wittmann (UCSF) for use of their

microscopes and advice regarding imaging and the cell cycle. Srivats Venkataramanan

and Stephen N. Floor are supported by funds from the California Tobacco-Related

Disease Research Grants Program Office of the University of California, 27KT-0003 and

the Program for Breakthrough Biomedical Research, which is partially funded by the

Sandler Foundation

References

[1] Buckley BA, Burkhart KB, Gu SG, Spracklin G, Kershner A, Fritz H, Kimble J, Fire A,

Kennedy S. A nuclear Argonaute promotes multigenerational epigenetic inheritance and

germline immortality. Nature. 2012 Sep 20;489(7416):447-51. doi:

10.1038/nature11352.

[2] Burton NO, Burkhart KB, Kennedy S. Nuclear RNAi maintains heritable gene silencing

in *Caenorhabditis elegans*. Proc Natl Acad Sci U S A. 2011 Dec 6;108(49):19683-8. doi:

10.1073/pnas.1113310108.

[3] Zhou Z, Hartwieg E, Horvitz HR. CED-1 is a transmembrane receptor that mediates

cell corpse engulfment in *C. elegans*. Cell. 2001 Jan 12;104(1):43-56.

[4] Soule HD, Maloney TM, Wolman SR, Peterson WD Jr, Brenz R, McGrath CM, Russo J,

Pauley RJ, Jones RF, Brooks SC. Isolation and characterization of a spontaneously

immortalized human breast epithelial cell line, MCF-10. Cancer Res. 1990 Sep

15;50(18):6075-86.

[5] Goto H, Yasui Y, Nigg EA, Inagaki M. Aurora-B phosphorylates Histone H3 at serine28

with regard to the mitotic chromosome condensation. Genes Cells. 2002 Jan;7(1):11-7.

[6] Lampson MA, Grishchuk EL. Mechanisms to Avoid and Correct Erroneous

Kinetochore-Microtubule Attachments. Biology (Basel). 2017 Jan 5;6(1). pii: E1. doi:

10.3390/biology6010001.

[7] Welburn JP, Vleugel M, Liu D, Yates JR 3rd, Lampson MA, Fukagawa T, Cheeseman

IM. Aurora B phosphorylates spatially distinct targets to differentially regulate the

kinetochore-microtubule interface. Mol Cell. 2010 May 14;38(3):383-92. doi:

10.1016/j.molcel.2010.02.034.

[8] Steigemann P, Wurzenberger C, Schmitz MH, Held M, Guizetti J, Maar S, Gerlich DW.

Aurora B-mediated abscission checkpoint protects against tetraploidization. Cell. 2009

Feb 6;136(3):473-84. doi: 10.1016/j.cell.2008.12.020.

[9] Fell VL, Walden EA, Hoffer SM, Rogers SR, Aitken AS, Salemi LM, Schild-Poulter C.

Ku70 Serine 155 mediates Aurora B inhibition and activation of the DNA damage

response. Sci Rep. 2016 Nov 16;6:37194. doi: 10.1038/srep37194.

[10] Monaco L, Kolthur-Seetharam U, Loury R, Murcia JM, de Murcia G, Sassone-Corsi P.

Inhibition of Aurora-B kinase activity by poly(ADP-ribosyl)ation in response to DNA

damage. Proc Natl Acad Sci U S A. 2005 Oct 4;102(40):14244-8.

[11] Guang S, Bochner AF, Burkhart KB, Burton N, Pavelec DM, Kennedy S. Small

regulatory RNAs inhibit RNA polymerase II during the elongation phase of transcription.

Nature. 2010 Jun 24;465(7301):1097-101. doi: 10.1038/nature09095.

[12] Chieffi P. Aurora B: A new promising therapeutic target in cancer. Intractable Rare

Dis Res. 2018 May;7(2):141-144. doi: 10.5582/irdr.2018.01018.

[13] Tang A, Gao K, Chu L, Zhang R, Yang J, Zheng J. Aurora kinases: novel therapy

targets in cancers. Oncotarget. 2017 Apr 4;8(14):23937-23954. doi:

10.18632/oncotarget.14893.

[14] Zuazua-Villar P, Rodriguez R, Gagou ME, Eyers PA, Meuth M. DNA replication stress

in CHK1-depleted tumour cells triggers premature (S-phase) mitosis through

inappropriate activation of Aurora kinase B. Cell Death Dis. 2014 May 22;5:e1253. doi:

10.1038/cddis.2014.231.

[15] Kabeche L, Nguyen HD, Buisson R, Zou L. A mitosis-specific and R loop-driven ATR

pathway promotes faithful chromosome segregation. Science. 2018 Jan

5;359(6371):108-114. doi: 10.1126/science.aan6490.

[16] Sakaguchi A, Sarkies P, Simon M, Doebley AL, Goldstein LD, Hedges A, Ikegami K,

Alvares SM, Yang L, LaRocque JR, Hall J, Miska EA, Ahmed S. *Caenorhabditis elegans*

RSD-2 and RSD-6 promote germ cell immortality by maintaining small interfering RNA

populations. Proc Natl Acad Sci U S A. 2014 Oct 14;111(41):E4323-31. doi:

10.1073/pnas.1406131111.

[17] Aronica L, Kasparek T, Ruchman D, Marquez Y, Cipak L, Cipakova I, Anrather D,

Mikolaskova B, Radtke M, Sarkar S, Pai CC, Blaikley E, Walker C, Shen KF, Schroeder R,

Barta A, Forsburg SL, Humphrey TC. The spliceosome-associated protein Nrl1 suppresses

homologous recombination-dependent R-loop formation in fission yeast. Nucleic Acids

Res. 2016 Feb 29;44(4):1703-17. doi: 10.1093/nar/gkv1473.

[18] Conine CC, Moresco JJ, Gu W, Shirayama M, Conte D Jr, Yates JR 3rd, Mello CC.

Argonautes promote male fertility and provide a paternal memory of germline gene

expression in *C. elegans*. Cell. 2013 Dec 19;155(7):1532-44. doi:

10.1016/j.cell.2013.11.032.

[19] van Wolfswinkel JC, Claycomb JM, Batista PJ, Mello CC, Berezikov E, Ketting RF.

CDE-1 affects chromosome segregation through uridylation of CSR-1-bound siRNAs. Cell.

2009 Oct 2;139(1):135-48. doi: 10.1016/j.cell.2009.09.012.

[20] Mao H, Zhu C, Zong D, Weng C, Yang X, Huang H, Liu D, Feng X, Guang S. The Nrde

Pathway Mediates Small-RNA-Directed Histone H3 Lysine 27 Trimethylation in

Caenorhabditis elegans. Curr Biol. 2015 Sep 21;25(18):2398-403. doi:

10.1016/j.cub.2015.07.051.

[21] Burkhart KB, Guang S, Buckley BA, Wong L, Bochner AF, Kennedy S. A pre-mRNA-

associating factor links endogenous siRNAs to chromatin regulation. PLoS Genet. 2011

Aug;7(8):e1002249. doi: 10.1371/journal.pgen.1002249.

[22] Richard P, Ogami K, Chen Y, Feng S, Moresco JJ, Yates JR 3rd, Manley JL. NRDE-2,

the human homolog of fission yeast Nrl1, prevents DNA damage accumulation in human

cells. RNA Biol. 2018 Aug 2:1-9. doi: 10.1080/15476286.2018.1467180.

[23] Lin YC, Boone M, Meuris L, Lemmens I, Van Roy N, Soete A, Reumers J, Moisse M,

Plaisance S, Drmanac R, Chen J, Speleman F, Lambrechts D, Van de Peer Y, Tavernier J,

Callewaert N. Genome dynamics of the human embryonic kidney 293 lineage in

response to cell biology manipulations. Nat Commun. 2014 Sep 3;5:4767. doi:

10.1038/ncomms5767.

[24] Bylund L, Kytölä S, Lui WO, Larsson C, Weber G. Analysis of the cytogenetic stability

of the human embryonal kidney cell line 293 by cytogenetic and STR profiling

approaches. Cytogenet Genome Res. 2004;106(1):28-32.

[25] Belin BJ, Lee T, Mullins RD. DNA damage induces nuclear actin filament assembly by

Formin -2 and Spire-½ that promotes efficient DNA repair. [corrected]. Elife. 2015 Aug

19;4:e07735. doi: 10.7554/eLife.07735. Erratum in: Elife. 2015;4. doi:

10.7554/eLife.11935.

[26] Caridi CP, D'Agostino C, Ryu T, Zapotoczny G, Delabaere L, Li X, Khodaverdian VY,

Amaral N, Lin E, Rau AR, Chiolo I. Nuclear F-actin and myosins drive relocalization of

heterochromatic breaks. Nature. 2018 Jul;559(7712):54-60. doi: 10.1038/s41586-018-

0242-8.

[27] Kim asutis KM, Kozminski KG. Cell cycle checkpoint regulators reach a zillion. Cell

Cycle. 2013 May 15;12(10):1501-9. doi: 10.4161/cc.24637.

[28] Hamperl S, Cimprich KA. Conflict Resolution in the Genome: How Transcription and

Replication Make It Work. Cell. 2016 Dec 1;167(6):1455-1467. doi:

10.1016/j.cell.2016.09.053.

[29] Segil N, Guermah M, Hoffmann A, Roeder RG, Heintz N. Mitotic regulation of TFIID:

inhibition of activator-dependent transcription and changes in subcellular localization.

Genes Dev. 1996 Oct 1;10(19):2389-400.

[30] Taylor WR, Stark GR. Regulation of the G2/M transition by p53. Oncogene. 2001

Apr 5;20(15):1803-15.

[31] Kimble J, Simpson P. The LIN-12/Notch signaling pathway and its regulation. Annu

Rev Cell Dev Biol. 1997;13:333-61.

[32] Fox PM, Schedl T. Analysis of Germline Stem Cell Differentiation Following Loss of

GLP-1 Notch Activity in *Caenorhabditis elegans*. Genetics. 2015 Sep;201(1):167-84. doi:

10.1534/genetics.115.178061.

[33] Dalfó D, Michaelson D, Hubbard EJ. Sensory regulation of the *C. elegans* germline

through TGF-β-dependent signaling in the niche. Curr Biol. 2012 Apr 24;22(8):712-9. doi:

10.1016/j.cub.2012.02.064.

[34] Michaelson D, Korta DZ, Capua Y, Hubbard EJ. Insulin signaling promotes germline

proliferation in *C. elegans*. Development. 2010 Feb;137(4):671-80. doi:

10.1242/dev.042523. Erratum in: Development. 2014 Jan;141(1):237.

[35] Begasse ML, Leaver M, Vazquez F, Grill SW, Hyman AA. Temperature Dependence

of Cell Division Timing Accounts for a Shift in the Thermal Limits of *C. elegans* and

C. briggsae. Cell Rep. 2015 Feb 4. pii: S2211-1247(15)00007-8. doi:

10.1016/j.celrep.2015.01.006.

[36] Claycomb JM, Batista PJ, Pang KM, Gu W, Vasale JJ, van Wolfswinkel JC, Chaves DA,

Shirayama M, Mitani S, Ketting RF, Conte D Jr, Mello CC. The Argonaute CSR-1 and its

22G-RNA cofactors are required for holocentric chromosome segregation. Cell. 2009 Oct

2;139(1):123-34. doi: 10.1016/j.cell.2009.09.014.

[37] Vought VE, Ohmachi M, Lee MH, Maine EM. EGO-1, a putative RNA-directed RNA

polymerase, promotes germ line proliferation in parallel with GLP-1/notch signaling and

regulates the spatial organization of nuclear pore complexes and germ line P granules in

Caenorhabditis elegans. Genetics. 2005 Jul;170(3):1121-32.

[38] Maine EM, Hauth J, Ratliff T, Vought VE, She X, Kelly WG. EGO-1, a putative RNA-

dependent RNA polymerase, is required for heterochromatin assembly on unpaired dna

during *C. elegans* meiosis. Curr Biol. 2005 Nov 8;15(21):1972-8.

[39] Duchaine TF, Wohlschlegel JA, Kennedy S, Bei Y, Conte D Jr, Pang K, Brownell DR,

Harding S, Mitani S, Ruvkun G, Yates JR 3rd, Mello CC. Functional proteomics reveals

the biochemical niche of *C. elegans* DCR-1 in multiple small-RNA-mediated pathways.

Cell. 2006 Jan 27;124(2):343-54.

[40] Sy Bukhari SI, Vasquez-Rifo A, Gagné D, Paquet ER, Zetka M, Robert C, Masson JY,

Simard MJ. The microRNA pathway controls germ cell proliferation and differentiation in

C. elegans. Cell Res. 2012 Jun;22(6):1034-45. doi: 10.1038/cr.2012.31.

[41] Rocheleau CE, Cullison K, Huang K, Bernstein Y, Spilker AC, Sundaram MV. The

Caenorhabditis elegans ekl (enhancer of ksr-1 lethality) genes include putative

components of a germline small RNA pathway. Genetics. 2008 Mar;178(3):1431-43. doi:

10.1534/genetics.107.084608.

[42] Menck CF, Munford V. DNA repair diseases: What do they tell us about cancer and

aging? Genet Mol Biol. 2014 Mar;37(1 Suppl):220-33.

[43] Lubas M, Christensen MS, Kristiansen MS, Domanski M, Falkenby LG, Lykke-

Andersen S, Andersen JS, Dziembowski A, Jensen TH. Interaction profiling identifies the

human nuclear exosome targeting complex. Mol Cell. 2011 Aug 19;43(4):624-37. doi:

10.1016/j.molcel.2011.06.028.

[44] Ogami K, Richard P, Chen Y, Hoque M, Li W, Moresco JJ, Yates JR 3rd, Tian B,

Manley JL. An Mtr4/ZFC3H1 complex facilitates turnover of unstable nuclear RNAs to

prevent their cytoplasmic transport and global translational repression. Genes Dev.

2017 Jun 15;31(12):1257-1271. doi: 10.1101/gad.302604.117.

[45] Virtanen JA, Vartiainen MK. Diverse functions for different forms of nuclear actin.

Curr Opin Cell Biol. 2017 Jun;46:33-38. doi: 10.1016/j.ceb.2016.12.004.

[46] Barnum KJ, O'Connell MJ. Cell cycle regulation by checkpoints. Methods Mol Biol.

2014;1170:29-40. doi: 10.1007/978-1-4939-0888-2_2.

[47] McMurchy AN, Stempor P, Gaarenstroom T, et al. A team of heterochromatin

factors collaborates with small RNA pathways to combat repetitive elements and

germline stress. Heard E, ed. eLife. 2017;6:e21666. doi:10.7554/eLife.21666.

[48] Hamperl S, Cimprich KA. Conflict Resolution in the Genome: How Transcription and

Replication Make It Work. Cell. 2016 Dec 1;167(6):1455-1467. doi:

10.1016/j.cell.2016.09.053.

[49] Larson AG, Elnatan D, Keenen MM, Trnka MJ, Johnston JB, Burlingame AL, Agard

DA, Redding S, Narlikar GJ. Liquid droplet formation by HP1α suggests a role for phase

separation in heterochromatin. Nature. 2017 Jul 13;547(7662):236-240. doi:

10.1038/nature22822.

Figure 1A: Loss of NRDE-2 significantly decreases the number of mitotic germ cells and increases the accumulation of RAD-51 foci in these cells in worms grown for two generations at 25°C. Maximum intensity projections of DAPI stained gonads with RAD-51 foci visualized in the green channel using an antibody against RAD-51. (Left panels) The red line indicates mitotic proliferating germ cell zone, the green line indicates transition zone where DAPI labeled chromosomes exhibit a "crescent-like" morphology, and the blue line indicates meiotic zone. (Right panels) Zoomed in image of one focal plane of mitotic PGCs. The white arrow indicates a single RAD-51puncta. **Figure 1B: (Left Panel) Loss of NRDE-2 at 20°C decreases the number of PGCs and this is exacerbated when the worms are grown for two generations at 25°C.** Mean number $\pm$ SEM of PCGs in gonads of wild type propagated at 20°C (209.6 ± 7.518, n=14) and *nrde-2(gg91)* mutants propagated at 20°C (181.6 ± 6.830, n=22) are different (*p=0.0115). PGCs of G2 wild type worms propagated at 25°C (207.9 ± 5.542, n=16) and G2 *nrde-2(gg91)* mutants propagated at 25°C (101.0 ± 5.418, n=21) is different (****p<0.0001). **(Right panel) Loss of NRDE2 at 25°C increases the number of RAD-51 foci per PGC nucleus.** The focal plane with the greatest number of RAD-51 foci was chosen for each germ cell nucleus. The number of RAD-51 foci per nuclear focal plane cell were averaged across 50 mitotic PGCs (10 PGCs closest to the distal tip cell in 5 gonads per condition). Mean number $\pm$ SEM of RAD-51 foci in gonads of wild type propagated at 20°C (0.3400 ± 0.07346, n=50) and *nrde-2(gg91)* mutants propagated at 20°C (0.3800 ± 0.1026, n=50) are not different. RAD-51 foci of G2 wild type worms propagated at 25°C (0.6000 ± 0.1245, n=50) have more RAD-51 foci than G2 *nrde-2(gg91)* mutants propagated at 25°C (9.200 ± 0.2665, n=50, ****p<0.0001). All data represents 3 experimental days and the p-values were determined using unpaired Student's t test. **Figure 1C: Loss of NRDE-2 increases the number of CED-1::GFP-labeled cell corpses in worms grown for 1 or 2 generations at 25°C.** CED-1::GFP expressed in *nrde-2(gg91)* and wild type *C. elegans* gonads (white arrows indicate apoptotic cell corpses). **Figure 1D: Loss of NRDE-2 increases the number of apoptotic corpses in the gonads of worms grown for 1 or 2 generations at 25°C.** Quantification of CED-1::GFP labeled germ cells. Wild type worms propagated at 20°C (6.071 ± 0.4409, n=42) have the same number of corpses as *nrde-2(gg91) mutants propogated* at 20°C (7.302 ± 0.4897, n=43). Wild type worms (6.263 ± 0.4869, n=38) have fewer cell corpses than *nrde-2(gg91)* mutants when grown for 1 generation at 25°C (10.05 ± 0.5379, n=19, ****p<0.0001). Wild type worms (7.510 ± 0.4278, n=51) have fewer cell corpses than *nrde-2 (gg91)* mutants when grown for 2 generations at 25°C (15.86 ± 0.9987, n=44, ****p<0.0001). Cell corpses identified as described in Figure 1D. Data represents 3 experimental days and the p-values were determined using unpaired Student's t test.

Figure 2A: In MCF10A human breast cells with endogenous or overexpressed AuBK levels, eGFP-NRDE2 localizes to puncta within the nucleus when chromatin is de-condensed during interphase and cytokinesis. In each circumstance, eGFP-NRDE2 is diffuse throughout all other stages of mitosis. Epifluorescent images of cells expressing eGFP-tagged NRDE2 as visualized in the green channel, chromatin as visualized by DAPI staining in the blue and the cytoskeleton as visualized by an antibody against α-tubulin in the Cy5 (far red) channel. (Top row) MCF10A cells expressing control vector. (Bottom row) MCF10A cells over-expressing AuBK. **Figure 2B: AuBK overexpression in MCF10A cells as a means to stress cells by increasing basal levels of DNA damage.** Western blot band intensities (see figure 2D for representative blot) were measured and normalized to β-actin band intensity. HA-tag band intensity in HA-tagged AUBK overexpressing cells is higher than in control cells (n=4; p=0.0046). Phosphorylated Histone H3 band intensity in AUBK overexpressing cells is higher than in control cells (n=4; p=0.0337). Phosphorylated H2AX band intensity in AUBK overexpressing cells is higher than in control cells (n=5; p=0.0146). All p-values determined by paired Student's t test. **Figure 2C: Average GFP intensity in MCF10A cells expressing eGFP-tagged NRDE2 decreases specifically in cells treated with NRDE2 targeted siRNA.** (Left) Epifluorescent images of MCF10A cells expressing eGFP-tagged NRDE2 as visualized in the green channel, the cytoskeleton as visualized by an antibody against α-tubulin in the far red channel and DNA as visualized by DAPI staining in the blue channel. Control or AuBK overexpressing MCF10A cells were treated with non-targeting siRNA or NRDE2 siRNA. (Top right) Quantification of epifluorescent GFP intensity of control and AuBK overexpressing cells treated with NRDE2 siRNA and normalized to the respective non-targeting siRNA treated cells. (Bottom right) Representative western blot using anti-GFP to detect eGFP-tagged NRDE2 inMCF10A cells treated with non-targeting or NRDE2 targeted siRNA. All cells treated with siRNA for 72 hours before imaging. Data represents 4 experimental days. **Figure 2D: MCF10A cells that overexpress HA-tagged AuBK show an increase in phosphorylated Histone H3 and an increase in the DNA damage marker γH2AX.** Representative blot: Anti-HA-tag used to detect AuBK in cells transformed with AuBK-HAtag in MCF10A cells. AuBK overexpressing cells have increased levels of phosphorylated Histone H3 and increased levels of γH2AX expression as compared to control cells. Western blot band intensities were measured and normalized to β-actin band intensity. See Fig S3 for quantification and statistics.

Figure 3A: Loss of NRDE2 in MCF10A cells overexpressing AuBK results in an increase in the percentage of cells with a rounded morphology. (Top) Brightfield images of cells 72 hours after siRNA treatment taken with 4x objective and 20x objective (inset). (Bottom) Quantitation of cell morphology. (Left paired points) Control or AuBK OE MCF10A cells show no difference in the percentage of round cells in cultures treated with non-targeting siRNA. (Middle paired points) MCF10A control cells show no difference in the percentage of round cells upon NRDE2 siRNA treatment as compared to non-targeting siRNA treatment. (Right paired points) MCF10A AuBK OE cells show a significant increase (*p=0.0110) in the percentage of round cells upon NRDE2 siRNA treatment as compared to non-targeting siRNA treatment. Data represents 6 experimental days. All p-values determined by paired Student's t test. **Figure 3B: Loss of NRDE2 in MCF10A cells overexpressing AuBK delays time through mitosis.** Time required for individual MCF10A cells to through mitosis. Control cells treated with NRDE2 siRNA (24.87 ± 0.3902, n=318) do not take significantly longer than control cells treated with non-targeting siRNA (23.79 ± 0.2442, n=492) to progrss through mitosis (p=not significant, Mann-Whitney test). AuBK over-expressing cells treated with NRDE2 siRNA (31.40 ± 0.7220, n=472) take significantly longer than cell trated with non-targeting siRNA (26.72 ± 0.4120, n=451) to progress through mitosis (****p<0.0001, Mann-Whitney test) **Figure 3C: Loss of NRDE delays time through mitosis in a subset of MCF10A cells overexpressing AuBK** of live MCF10A cells expressing H2B::mCherry visualized in the red channel to visualize chromatin compaction and chromosome separation. **Figure 3D: Loss of NRDE2 in AuBK overexpressing MCF10A cells significantly increases the proportion of cells in prophase.** Quantification of the percentage of cells in each phase of mitosis shows that AuBK overexpressing MCF10A cells have significantly more cells in prophase after treatment with NRDE2 siRNA as compared to treatment with non-targeting siRNA (p= 0.0258, determined by student's t test). Data represents 4 experimental days.

Figure 4A: eGFP::NRDE2 interacts with β-actin in control cells. This interaction is weakened in AuBK OE cells and when cells are treated with cisplatin. Representative western blots of MCF10A cell lysates. MCF10A cells were treated with 60uM cisplatin for 16 hours. Co-IP was performed using goat Anti-GFP antibody to pull down eGFP::NRDE2 from MCF10A cell lysates. Western blots were developed using rabbit anti-GFP polyclonal antibody to detect eGFP::NRDE2 and soluble GFP, anti-β-actin HRP antibody and mouse anti-β-tubulin antibody (negative control). **Figure 4B: The amount of β-actin per unit of IP'd GFP::NRDE2 decreases when DNA damage is induced in cells by AuBK OE or by cisplatin drug treatment.** (Left) The amount of β-actin per unit of co-IP'd eGFP::NRDE2 decreases when MCF10A cells overexpress AuBK as compared to MCF10A control cells (*Permutation test exact p-value =0.036). (Right) The amount of β-actin per unit of co-IP'd eGFP::NRDE2 decreases when MCF10A control cells are treated with 60uM cisplatin as compared to untreated control MCF10A cells. (*p-value =0.045 as assessed by paired student's t test). Note: Blue data points represent quantification of the same samples.

Figure S1: Loss of NRDE-2 significantly decreases brood size as compared to wild type worms when propagated at 20°C. Each data point represents one biological replicate of 16 worms per experiment (n=4; *p=0.0148, paired Student's t test).

Figure S2: NRDE2 knockdown decreases cell viability in AuBK OE cells but not in control cells. The Cell Titer Glo assay measures the number of metabolically active cells by quantifying the amount of ATP present in the cell culture. The luminescent signal was normalized to non-targeting siRNA treated cells in both control cells and AuBK OE cells. KIF11 siRNA was used as a positive control. Control cells treated with NRDE2 siRNA (100.0 ± 2.890, n=4) had no change in luminescence as compared to control cells treated with non-targeting siRNA. AuBK OE cells treated with NRDE2 siRNA (85.55 ± 4.167, n=4) had a lower amount of luminescence as compared to AuBK OE cells treated with non-targeting siRNA (*p=0.0133). P-values determined using unpaired Student's t test.

Figure S3 (Top) Loss of NRDE2 in control cells increases pHH3 levels as compared to control cells treated with non-targeting siRNA (n=4, p=0.0068). In contrast, loss of NRDE2 in AuBK OE cells does not change pHH3 levels (p=???). (Bottom) Loss of NRDE2 in control cells does not alter levels of γH2AX (p=????), however in AuBK OE cells, loss of NRDE2 increases γH2AX levels. (n=5, *p= 0.0483) All p-values determined using paired student's t test.

Figure S4: Loss of NRDE2 in MCF10A cells does not significantly change the mitotic index. Mitotic Index calculated by dividing total number of cells in mitosis by total number of cells in the population. AuBK overexpressing cells have a significantly higher basal mitotic index than control cells (*p= 0.0315, student's paired t test).

Figure S5: Representative western blot of MCF10A cell lysates. Co-IP was performed using goat Anti-GFP antibody to pull down eGFP::NRDE2 from MCF10A cell lysates. Western blots were developed using rabbit anti-GFP polyclonal antibody to detect eGFP::NRDE2 and soluble GFP.

Figure S6: NRDE-2 does not interact with the mammlian argonautes Ago 1,2,3 or 4. Goat anti-GFP used to pull down eGFP::NRDE2 in MCF10A lysates in co-IP assay. Western blots against human Ago1-4 show no interaction with eGFP::NRDE2. Arrows point to 97kD, the expected size of each Ago protein.

Figure S7: Control and AuBK OE MCF10A cells treated with nocodozole show rounded morphology consistant with stall in G2/M. Cells treated with cisplatin show rounded, fragmented morphology, consistant with apoptosis. Representative brightfield images of cells taken 16 hours after 0.1ug/ml nocodozole or 60uM cisplatin drug treatment with 4x objective and 20x objective (inset).