Title: **Divergent neuronal DNA methylation patterns across human cortical development:**

Critical periods and a unique role of CpH methylation

Authors

Price AJ[1,2]*, Collado-Torres L[1,3]*, Ivanov NA[1], Xia W[1], Burke EE[1], Shin JH[1], Tao R[1], Ma L[1], Jia

Y[1], Hyde TM[1,4,5], Kleinman JE[1,5], Weinberger DR[1,2,4,5,6], Jaffe AE[1,2,3,7,8,+]

Affiliations

1. Lieber Institute for Brain Development, Johns Hopkins Medical Campus, Baltimore, MD, USA

2. McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of

Medicine, Baltimore, MD, USA

3. Center for Computational Biology, Johns Hopkins University, Baltimore, MD, USA

4. Department of Neurology, Johns Hopkins School of Medicine, Baltimore, MD, USA

5. Department of Psychiatry, Johns Hopkins School of Medicine, Baltimore, MD, USA

6. Department of Neuroscience, Johns Hopkins School of Medicine, Baltimore, MD, USA

7. Department of Mental Health, Johns Hopkins Bloomberg School of Public Health, Baltimore,

MD, USA

8. Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore,

MD, USA

*equally contributing authors

Corresponding Author: Andrew E Jaffe, 855 N Wolfe St, Ste 300; Baltimore MD 21205.

Phone: 1-443-287-6864; Email: andrew.jaffe@libd.org

Abstract

We have characterized the landscape of DNA methylation (DNAm) across the first two decades of human neocortical development in NeuN+ neurons using whole-genome bisulfite sequencing and compared them to non-neurons (primarily glia) and prenatal homogenate cortex. We show that DNAm changes more dramatically during the first five years of postnatal life than during the entire remaining period. We further refined global patterns of increasingly divergent neuronal CpG and CpH methylation (mCpG and mCpH) into six developmental trajectories and found that in contrast to genome-wide patterns, neighboring mCpG and mCpH levels within these regions were highly correlated. We then integrated paired RNA-seq data and identified direct regulation of hundreds of transcripts and their splicing events exclusively by mCpH levels, independently from mCpG levels, across this period. We finally explored the relationship between DNAm patterns and development of brain-related phenotypes and found enriched heritability for many phenotypes within DNAm features we identify.

Introduction

Neurons are unique cells that persist throughout the lifespan, accumulating programmed developmental changes and environmental experience that fine tune neural circuitry in the brain. During development and maturation, neurons undergo precisely coordinated cascades of genetic regulation that combine with experience to shape the cellular output via progressive changes to the epigenome. DNA methylation (DNAm) is an integral facet of the epigenome that plays a role in establishing cell identity and developmental trajectories as well as adapting to experience via regulation of gene expression. Previous large-scale studies of DNAm across human brain development have been limited to homogenate tissue[1] or have used microarray technologies[2], creating ambiguity about the extent of cell type-specific developmental DNAm changes and effects on transcript isoforms across the genome[3].

To better characterize the DNAm landscape across human cortical development, we performed whole-genome bisulfite sequencing (WGBS, see Methods) on homogenate tissue and on a neuron-enriched population isolated from 24 human dorsolateral prefrontal cortex (DLPFC) samples aged 0-23 years using NeuN-based fluorescence-activated nuclear sorting (FANS, **Figure S1A**). To complement these data, we sequenced eight FANS-derived NeuN- postnatal samples and 20 homogenate prenatal cortical samples, for a total of 75 samples after quality control (**Table S1**). We fully characterized the landscape of DNAm at both CpG and non-CpG (CpH) dinucleotides in these samples, allowing for a finer dissection of differential DNAm functional specificity. We also sequenced matched transcriptomes of homogenate cortical samples from these donors and a subset of three nuclear transcriptomes each from NeuN+ and NeuN- samples to assess the functional consequences of epigenomic remodeling (53 total transcriptomes, **Table S2**). By exploring DNAm patterns in neurons across prenatal and postnatal human brain development, we show that the first five years of postnatal life are a

critical period in epigenetic plasticity, and we identify developmental shifts in neuronal DNAm in

both the CpG and CpH context. We also clarify the relationship between CpG but particularly

CpH methylation (mCpG and mCpH, respectively) and gene expression and splicing in neuronal

development, and explore the ramifications of these insights for neuropsychiatric disease.

Results

After data processing, quality control and filtering, we analyzed 18.7 million cytosines in

the CpG context at an average coverage of 15x (see methods). Comparable to previous

reports[1,4,5], CpGs were overall highly methylated (71-76% CpGs with β>80%, **Table S3**).

While NeuN antibody labels most mature neuronal subtypes in human cortex, some

neurons will not be labeled and will be captured in the NeuN- fraction amidst a diverse array of

non-neuronal cell types, including oligodendrocytes, astrocytes, microglia, and epithelial cells.

Gene expression differences between fractions confirmed, however, that NeuN+ and NeuN-

samples are enriched for neuronal and glial-lineage cells, respectively (**Figure S1B-D**).

Therefore in this work we refer to NeuN+ and NeuN- samples as "neurons" and "glia" although

we acknowledge that these samples do not perfectly reflect these identities and mask more

granular differences between subcellular identities contained within.

Developmental DNAm changes identified in homogenate cortex were strongly

confounded by shifting cell type proportions (OR=7.5, $p<10^{-100}$, **Figure S2A**)[2]. While

homogenate measurements were positively correlated with developmental changes that

occurred in both neuronal and glial cell types (ρ=0.79, $p<10^{-100}$), cell type-specific developmental

changes were less consistently observed in homogenate preparations (ρ=-0.26, $p<10^{-100}$,

Figure S2B-D). Overall, ~40% of cell type-specific developmental DNAm changes could not be

detected at all in homogenate cortex (**Figure S2E**), and many of the cell type-specific effects

could not be accurately identified in homogenate tissue. These results highlight the importance

of measuring DNAm in the appropriate cellular context for improved resolution to detect true

developmental changes.

DNAm as a map of putative functional genomic states

Local CpG methylation (mCpG) patterns are known to distinguish genomic states of

DNA and chromatin. For instance, unmethylated regions (UMRs) are associated with promoters,

with a subset of longer UMRs (DNAm valleys, DMVs) that overlap developmental genes often

encoding transcription factors (TFs)[6,7]; low-methylated regions (LMRs) often signify enhancer

sequence[8]; and partially methylated domains (PMDs) are associated with heterochromatin and

late replicating DNA[9–11]. To better resolve the developing regulatory landscape in postnatal

neurons and glia and in bulk prenatal cortex, we assessed the temporal dynamics of these

selected DNAm patterns in the CpG context. Compared to prenatal homogenate cortex and

postnatal glial cells, postnatal neurons showed a general accumulation of mCpG, at a rate 50%

faster than the other cells. This was evident in the LMR and to a lesser extent the UMR

landscape, since fewer and smaller LMRs were identified as neuronal development progressed

(**Figure S3A-B**). As expected, UMRs and LMRs were highly enriched for transcription start sites

(TSSs) and enhancers in DLPFC chromatin state data from the Roadmap Epigenomics

Consortium[12] (**Figure S3C**). Interestingly, LMRs were similarly enriched in these states in both

adult and fetal brain; this correspondence may reflect a shared regulatory landscape established

early in development.

While PMDs are a common feature of most cell types, they have not been conclusively

identified in neurons. Here we identified a range of 245 to 404 PMDs per neuronal sample

(**Figure S4A**). PMDs were especially enriched for heterochromatin and, interestingly, enhancers

in our postnatal neuronal samples (**Figure S4B**). 65.4% of PMD base pairs were also identified

as PMD in an independent WGBS dataset of NeuN-sorted human neurons (**Figure S4C**).

40.3-61.0% of PMD bases per neuronal sample were identified as common PMD sequence, and

9.3-15.0% bases were additionally identified as PMD in at least one sample in a recent study

profiling PMDs in multiple cell types and tissues[13] (**Figure S4D**). These data suggest that

although the neuronal genome was overall highly methylated, a small but consistent portion

displayed the characteristics of PMDs.

We further identified significant neuronal DMV changes through accumulation of mCpG

that revealed regulators of cell identity and development and their temporal windows of

expression change. Compared to bulk prenatal cortex, postnatal neurons and glia showed

marked reduction in the size of DMVs (**Figure S5A**). Methylation shifts within DMVs led to

inclusion and exclusion of transcription factor genes in an age-dependent manner, and on

average transcription factor genes were higher expressed in the age group in which the gene

was escaping the DMV state by accumulating DNAm (**Figure S5B-C**). These results underscore

the substantial DNAm landscape alterations that neurons and glia undergo during development

in defined mCpG patterns, including previously unobserved PMDs.

Developmental shifts in neuronal mCpG highlight synaptic remodeling during first five years of

postnatal life

We next quantified more localized changing mCpG levels by exploiting the correlation

between neighboring mCpG levels to identify genomic regions with changing mCpG. We

identified 11,179 differentially methylated regions (DMRs, FWER<5%, see Methods) in the CpG

context between cell types (covering 31.1 Mb) that replicated in independent WGBS data[1]

(98.4% concordant, ρ=0.925, **Figure S6A**). Many of these DMRs overlapped genes involved in

neuronal or glial-specific processes (**Figure S6B**). We found fewer DMRs for developmental

mCpG changes compared with cell type differences, the majority being within rather than across

cell types (2,178 versus 129 DMRs, at ~5% change in DNAm per decade of life, FWER<5%).

Among the 2,178 cell type-specific developmental DMRs (cdDMRs, 3 Mb, **Table S4**), neuronal

mCpG patterns seemed to diverge from an immature landscape shared by glia and prenatal

cortex (**Figure 1A**), with the largest changes occuring in the first five years of life. Indeed, the

magnitude of DNAm changes in neurons and glia in samples five years and younger was double

that of older samples (**Figure S6C-D**). These results provide epigenetic correlates to the known

developmental processes occuring in the cortex in the first five postnatal years, including prolific

synaptogenesis and gliogenesis.

We further parsed these cdDMRs using k-means clustering to partition the cdDMRs into

six groups with unique DNAm characteristics (**Figure 1B**). 71.1% of cdDMRs were in groups

characterized by increasing neuronal and/or decreasing glial DNAm over postnatal development

(**Figure 1B;** Groups 1, 2 & 6). A varying proportion of each cdDMR group corresponded to

sequence differentially methylated by neuronal subtype from publicly available data[14] depending

on the trajectory of neuronal methylation patterns in the group, suggesting that assorted

neuronal subclasses contribute to these developmental patterns (**Figure S7A-B**). Gene

ontology enrichment in the six groups suggested that these groups are associated with a

continuum of biological roles, many relating to functions specific to the cell type with decreasing

methylation (**Figure 1C**). For example, **Figure 1D** shows a Group 3 cdDMR within *SNAP25*, a

presynaptic neuronal gene, in which neurons uniquely and progressively lost DNAm over

development. This pattern suggests increased repression of a neuronal fate in maturing glia not

mirrored in neurons over postnatal development in Group 3 cdDMRs. Likewise, the opposite

pattern was observed in a Group 6 cdDMR within *MBP*, an oligodendrocyte gene encoding a

component of the myelin sheath, in which glia but not neurons progressively lost DNAm (**Figure 1E**).

We lastly compared these cdDMR groups to a list of putative enhancers active in human brain development curated by evolutionary age[15] and found strong enrichment for these sequences across all six groups (**Figure S8A**). Human accelerated regions, or conserved sequences that have experienced rapid mutation in the human lineage[16], were also enriched for dynamic DNAm remodeling (**Figure S8B**), suggesting that our CpG-based cdDMRs may be enriched for sequences related to higher cognitive functions associated with the human DLPFC.

Overlapping cdDMRs with the mCpG features identified above provided additional insight to the potential functional genomic states underlying these regions. For instance, cdDMRs scarcely overlapped heterochromatic PMDs; cdDMRs losing neuronal mCpG were positively correlated with increasing LMR overlap, potentially reflecting enhancer element activation during cortical maturation in these groups (Groups 3 & 5 cdDMRs; both with t>3.8, ρ>0.63, and FDR<2.7e-03). Curiously, a high proportion of cdDMRs gaining DNAm in glia but not in neurons (Group 4 cdDMRs) overlapped DMVs early in development in glia but steadily lost DMV status over time (t=-4.3, ρ=-0.87, FDR=1.3e-02, **Figure S9A**). Assessing chromatin state from the homogenate Roadmap Epigenomics brain maps, in contrast, lacked the resolution to provide this nuance: all six cdDMR groups were similarly enriched for transcriptional (particularly TSS-flanking) and enhancer chromatin states, and depleted for heterochromatin and quiescent states (**Figure S9B**). These results confirm the role of dynamic DNAm in helping establish epigenomic states that guide cell lineage differentiation and emphasize the utility of creating genome-wide DNAm maps to better parse the functional diversity of cell type-specific developmental DNAm remodeling in the human cortex, a process that is particularly critical during the first five years of postnatal development.

Abundant neuronal CpH methylation is highly correlated with neighboring CpG methylation

Unlike in most other somatic tissues and cell types, mCpH is an abundant, conserved feature of the neuronal epigenome[1,4]. We therefore analyzed 58.1 million cytosines in CpH contexts (H=A,T, or C) that had evidence of methylation across the samples (coverage ≥5, at least 5 samples with β>0, see Methods). As shown previously[1], mCpH sites were predominantly lowly methylated (92-99% CpHs with β<20%, **Table S3**). While mCpH was distributed throughout the genome (**Figure S10A**), it was greater in neurons than glia (98.9% of 7,682,075 differentially methylated CpHs were in neurons, at FDR<5%) and mostly accumulated across postnatal development (99.3% of 3,194,618 CpHs, at FDR<5%; **Table S5**). Most mCpH accumulated primarily in either the CAG or CAC context over the first five years of postnatal life—similarly to mCpG—followed by a tapered global increase into adulthood (**Figure S10B**).

While the majority of mCpH in embryonic stem cells (ESCs) occurs in the CAG context, previous work has shown that ESCs undergo loss of mCAG during neuronal differentiation followed by preferential accumulation of mCAC[17]. Here we further refined these patterns and found a cell type-specific relationship with trinucleotide context: overall, total mCAG increased 40% faster than mCAC in neurons, while in glia, mCAG accumulated 50% slower than mCAC (**Figure S10C**). Taking into account relative genome-wide proportions of CAG and CAC though, neuronal mCAG accumulated 30% slower than mCAC (**Figure 2A**). mCH that was greater in glia than neurons, or in younger than older neurons, was more likely to be in the CAG than CAC context (OR>4.13, p<2.2e-16). Interestingly, the 3,286 and 1,744 genes that contained significantly increasing and decreasing mCAC versus mCAG over development, respectively, were associated with different biological processes related to neuronal function and activity, particularly involving the synapse (**Figure S10D**). These results reinforce that trinucleotide

context may be regulated by distinct mechanisms playing non-redundant biological roles in

human brain development.

We next examined the relationship between neighboring mC levels by measuring

autocorrelation, or how correlated the methylation level of a cytosine is with that of cytosines

progressively further away. Unlike in the CpG context, where neighboring mCpG levels were

highly correlated as previously described[18], neighboring CpH DNAm levels across the genome

were not autocorrelated. Within the cdDMRs, however, while mCpH levels separately remained

uncorrelated, together all methylated cytosines (i.e., mCpH+mCpG) showed similar

autocorrelation as mCpG levels alone (**Figure 2B**). This was especially surprising given that

there were about two times as many CpHs than CpGs within these regions and that the CpG

and CpH were relatively interspersed, suggesting potential functional convergence in the

developmentally regulated patterns identified by mCpG in these regions. Indeed, unsupervised

hierarchical clustering of CpH within the cdDMRs showed infant neuronal mCpH levels were

even more similar to glia compared to older neurons than mCpG (**Figure 2C**). Examining the

mean mCpH compared to mCpG within the k-means cdDMR clusters showed that the groups

gaining mCpG were the most correlated with mCpH trajectories within the cdDMRs (ρ=0.97,

t=17.6, p=2.0e-14), and that although mCpG (unlike mCpH) is present at high levels prenatally,

both mCpG and mCpH accumulate at similar rates over postnatal development in these groups,

once again especially in the first five years of postnatal life where the majority of the methylation

change takes place (t=-0.091, p=0.94; **Figure S11**). These results emphasize the potential

regulatory importance of cdDMRs and putative functional agreement between both contexts of

DNAm in these regions.

mCpG and mCpH levels influence transcript isoform use

Previous studies show that both mCpG and mCpH in gene bodies but particularly in the

promoter and first 2 kb of the gene are negatively associated with gene expression, and that

genic mCpH is the most discriminating predictor of gene expression[1,5]. To anchor our DNAm

patterns in transcriptional activity, we compared our WGBS data with NeuN-sorted nuclear

RNA-seq data (see Methods). We took the average DNAm levels across six groups—infant

(ages 0-1), child (ages 1-10), and teen (age 10+) within both cell types (neuronal and

glial)—and calculated associations between DNAm and expression. Gene expression was

negatively correlated with mCpG regardless of age and cell type in both promoter sequence and

gene bodies ($-0.42 < \rho < -0.22$, p~0; 57,332 genes, $p < 10^{-100}$; **Figure S12A**). Interestingly, mCpG

in exons was significantly but weakly positively correlated with exon expression in infancy,

particularly in glial samples ($\rho = 0.094$, $p < 10^{-100}$), which may relate to the previously-identified

positive relationship between mCpG and expression and higher methylation in exons than

introns[9].

Across promoters, gene bodies, and exons, only neurons showed a negative correlation

between gene expression and mCpH, a relationship that became stronger over development

(**Figure S12B**). This pattern was consistent with the preferential accumulation of mCpH in

neurons as the brain matures. mCAC and mCAG showed similar patterns of increasingly strong

negative correlation preferentially in neurons between methylation and expression across these

features (**Figure S12C-D**). Both mCpG and mCpH surrounding exon-exon splice junctions were

weakly negatively correlated with expression of the junction in neurons (**Figure S12E**).

Because mCpG has previously been associated with alternative splicing[19] and mCpH is

15-20% greater in exons than in introns[9], we hypothesized that accumulating mCpH may

contribute to the diversity of alternative splicing characteristic of the brain particularly during

development. Leveraging our single-base resolution data, we were able to identify genome-wide

functional correlates of mCpH, independent of nearby mCpG, by associating DNAm with nearby

expression in the same cortical samples. Specifically, we tested whether methylation levels

directly associated with gene or exon expression levels as well as the "percent spliced in" (PSI)

of alternative splicing events using the 22 neuronal samples with matching homogenate polyA+

RNA-seq data (see Methods). We found 40,940 CpG and 40,303 CpH associations that explain

changes in these three expression summarizations at FDR<5% with a genome-wide $p<5 \times 10^{-4}$.

We further identified 220,622 marginal (p<0.01) CpG associations with expression within 1 kb

around the associated CpH. While an independent association of mCpH at the gene and PSI

summarizations was rare, there were substantially more exons exclusively regulated by local

mCpH, largely in the CHH context, in developing postnatal neurons (**Figure 3A**). Three

examples of methylation-associated isoform changes are shown in **Figure 3B**.

      Regardless of context specificity, these expression-associated cytosines were depleted

in gene promoters and instead enriched in gene bodies and flanking regions (**Table 1**, see

Methods). Both contexts were enriched for the high-GC 3' and 5' canonical splice site

sequences (FDR<1.1e-04), although the associated cytosine could be either inside or outside

the corresponding expression feature. Only 3.5-13.7% of expression-associated cytosines

overlapped DMR sequence after stratifying by expressed feature and dinucleotide context,

indicating that these associations may arise from a more individualized mC effect than the

DMRs.

      Although the majority of these DNAm-expression associations were independent of

development despite being identified in developing neurons, the mCpH changes at these sites

were independently associated with age and expression. The genes including PSI events

regulated by local DNAm levels in both CpG and CpH contexts were consistently enriched for

neuronal components (**Figure S13**), while genes containing methylation-associated alternative

exons were enriched for synaptic signaling and neurotransmitter transport (**Figure 3C** and

Figure S14), suggesting that we are detecting true neuronal mC-expression associations

despite measuring splicing in homogenate RNA-seq. Many of these genes were also

differentially expressed between neuronal and glial nuclear RNA (FDR<0.05, **Table 1**). Most, but

not all, expression-associated cytosines at the gene- and exon-level showed significant

decreases in expression as methylation levels increased.

The associations between these putatively regulatory cytosines and nearby expression

levels can be explored in a webtool (https://jhubiostatistics.shinyapps.io/wgbsExprs/). Results

can be interactively summarized such as in **Table 1** for user-selected subsets and visualized as

in **Figure 3B** or via the UCSC genome browser (**Figure S15**). By integrating neuronal mCpG

and mCpH levels with accompanying RNA-seq data in the same brains, we have identified, for

the first time, direct regulation of hundreds of transcripts and their splicing events exclusively by

mCpH, independent of mCpG levels, across the first two decades of human cortical

development.

DNAm patterns shed light on the active cell type and timing of neuropsychiatric phenotype

development

Previous work has attributed a high proportion of neuropsychiatric trait heritability to

neuron-specific DNA methylation patterns[20]. Given the role of dynamic DNAm in marking DNA

sequence function over development, we examined the relationship between our methylation

features with heritability for 30 human behavioral-cognitive traits, psychiatric and neurological

disorders, and non-brain-related traits[21] (**Table S6**), hypothesizing that DNAm patterns may

illuminate not only the active cell type but potential critical timeframes for genomic activity in

these complex phenotypes. We used stratified linkage disequilibrium score regression (LDSC)[22]

to estimate the proportion of heritability measured in GWAS summary statistics for each

phenotype that could be attributed to each of 16 genomic features, including 10 sets of DMRs,

LMRs identified in the prenatal, glial or neuronal methylome, human brain regulatory sequence

annotated by chromHMM or the LDSC package, or non-differential CpG clusters (**Figure 4A**). In

agreement with previous findings[20,23], human brain annotated regulatory sequence was broadly

enriched for heritability of brain-specific traits (14 of 26 brain-associated phenotypes enriched in

chromHMM or CNS (LDSC) regions at FDR≤0.05), as were neuronal features (10 of 26 brain

phenotypes enriched in neuronal hypomethylated regions, FDR≤0.05; **Table S7**). Significantly

differentially hypomethylated neuronal regions had on average 1.85 times higher enrichment

scores than non-differential neuronal LMRs, meaning the DMRs explained 1.85x more

heritability over regions containing a similar number of SNPs than the LMRs. Interestingly, body

mass index (BMI) heritability was enriched in general brain regulatory sequence and

hypomethylated neuronal DMRs (FDR≤0.05), consistent with previous evidence linking this

metabolic phenotype to regulatory sequence active in cells of the human central nervous

system[22].

In terms of developmental DNAm patterns, heritability of BMI, IQ, neuroticism, and major

depressive disorder was enriched in both postnatal neuronal and prenatal LMRs, suggesting

early action of genetic influence on the development of these phenotypes (FDR≤0.05). Few

developmental differential groups captured a significant proportion of heritability for the 30 traits

tested, perhaps because of their small size compared to the cell type-specific or non-differential

groups (cdDMRs covered ~31, 240 and 838 times less sequence and included ~32, 273 and

904 times fewer SNPs than cell type DMRs, LMRs or general brain features, respectively).

However, despite covering only 607 kilobases, group 3 cdDMRs (ie, static glial and decreasing

neuronal DNAm) were significantly enriched for heritability of schizophrenia (coefficient

z-score=2.74, FDR=0.039). Group 4 cdDMRs (111.7 kilobases; increasing glial, static neuronal

DNAm) were also enriched for heritability of PTSD (coefficient z-score=3.21, FDR=0.01).

Given the enrichment for psychiatric disease heritability measured in common SNPs in

these cdDMR groups, we then expanded our analysis to include seven curated gene sets

containing *de novo* and rare inherited variation--including rare copy number variants (CNVs) and

syndromic variants--associated with psychiatric, neurodevelopmental and neurodegenerative

disorders[24,25]. We again found significant enrichment of hypomethylated neuronal DMRs in

genes implicated in psychiatric and neurodevelopmental disorders (i.e., schizophrenia, autism

spectrum disorder (ASD), syndromal neurodevelopmental disorders, and intellectual disability;

all with OR>2.04 and FDR<1.9e-02; **Table S8**). In this analysis, we also found enrichment for

hypermethylated neuronal DMRs in ASD genes from the SFARI Gene database and

schizophrenia genes containing *de novo* mutations (both with OR>1.92 and FDR<5.0e-03).

These results confirmed a prominent role of neuronal functioning in most of the

neurodevelopmental disorders using an orthogonal measurement approach as done

previously[23]. Over postnatal development, Group 5 cdDMRs (increasing glial, decreasing

neuronal DNAm) were enriched in ASD genes from the SFARI Gene database (OR=5.7,

FDR=4.1e-03), while Group 6 cdDMRs (decreasing glial, static neuronal DNAm) were enriched

in ASD, syndromal neurodevelopmental disorder, and intellectual disability genes (all with

OR>3.1 and FDR<1.9e-02). In contrast, a curated set of neurodegenerative disorder genes

showed no enrichment for cdDMRs, perhaps reflecting lesser relevance of the first two decades

of postnatal epigenomic remodeling to the etiology of those disorders.

In the non-CpG context, we found significant enrichment of both increasing and

decreasing mCpH levels in genes associated with schizophrenia, ASD, and syndromal

neurodevelopmental disorders (all with OR>2.1 and FDR<2.0e-02). CpH hypomethylation in

neurons was also enriched in the neurodegenerative disease gene set (OR=2.6, FDR=3.7e-03).

Finally, significantly increasing mCpH was depleted in genes associated with intellectual

disability (OR=0.34, FDR=3.7e-06). While enrichment for conflicting mCpH patterns are at first

curious given the overall negative association between mCpH and gene expression, outside of

the context of DMRs, individual mCpH could be associated both positively and negatively with

expression. Indeed, many genes, exons, and PSI events whose expression both positively and

negatively associated with both mCpH and mCpG were also enriched in genes associated with

schizophrenia, ASD, and syndromal neurodevelopmental disorders (all with OR>2.1 and

FDR<2.5e-02; Table S9). Overall these results suggest that these examples of dynamic

methylation and associated isoform switching may play a role in the development of higher

cognitive functions during brain maturation associated with these diseases.

Discussion

Here we have created a single-base resolution map of the dynamic DNAm landscape

across the first two decades of postnatal human brain development in two cell type-enriched

populations. Using FANS-derived samples, we were able to identify 40% more

developmentally-regulated regions of changing DNAm than were identified in homogenate

DNAm cortical data. We profiled specific features of the DNAm landscape including LMRs,

UMRs, PMDs and DMVs and found that across features, neurons were typified by a general

accumulation of mCpG. In the absence of complementary cell type-specific chromatin data,

characterizing known DNAm features provided a more granular view of the potential functional

genomic state in these regions than the available predictions derived from a few homogenate

cortical samples. Particularly in studies using human postmortem brain, where tissue is often

subjected to long postmortem intervals and low pH that degrades less stable epigenetic

signatures, DNAm is a robust and durable marker that can be used to map the functional

genomic terrain. These DNAm maps complement recently available epigenomic maps of

different modalities generated on FANS-derived samples in the psychENCODE Consortium[27].

We further parsed the general accumulation of neuronal DNAm into six trajectories of

cell type-specific developmental patterns and found that neuronal mCpG progressively diverged

from a shared landscape with glia and bulk prenatal cortex as the brain matured. Importantly,

these diverging patterns were most striking during infancy through the first five years of

postnatal life. The human brain experiences an explosion of synaptic connections during this

time period, to nearly double the number found in the mature adult brain[28]. Although previous

work has underscored this timeframe in terms of rapid DNAm accumulation[1], this is the first

work to refine DNAm patterns to reflect cell type-specific gain and loss of mCpG and mCpH

within this critical window. By parsing these neuronal and glial DNAm patterns, we have

highlighted epigenetically dynamic regions that may be contributing to the developmental

processes such as synaptogenesis occuring during this timeframe that establish the foundation

for fine-tuning connections throughout the remainder of brain maturation. These results provide

a finely resolved depiction of epigenetic plasticity being greatest during this period of life, and

support other evidence that environmental experience during these years may have an

especially enduring impact on brain function[29].

mCpH is unusually abundant in neurons compared to other cell types and appears to

undergo trinucleotide-specific reprogramming during differentiation from ECSs[17]. While most

mCpH in ESCs occurs in the CAG context, neuronal mCpH predominantly accumulates in the

CAC context[17]. Here we elaborate on this relationship, showing that while both mCAG and

mCAC aggregate in neurons as they mature and mCAG is gained faster than mCAC overall,

mCAC accumulates proportionally faster in both neurons and glia over time. Interestingly,

although neurons and glia contained mCpH in both trinucleotide contexts, mCAG was more

likely to have higher levels in glia than neurons or be decreasing over development; indeed,

genes containing decreasing mCAG but not mCAC were strongly associated with neuronal

biological processes. mCpH trinucleotide context, therefore, may have as yet not well

understood ramifications in brain development.

In terms of the relationship between mCpH and mCpG, we found that while neighboring

mC (i.e., mCpG+mCpH) was not correlated genome-wide, mC was highly correlated within the

cdDMRs despite local mCpH not being correlated. In other words, there was a convergence of

levels of all contexts of methylation within the cdDMRs that was not detected genome-wide.

mCpH also recapitulated the pattern seen in mCpG of diverging from a shared DNAm

landscape with glia. Given that mCpH and mCpG have previously been shown to work in

concert to recruit MECP2 binding to fine-tune gene expression[30], it is sensible that levels of both

contexts would perhaps reflect a shared functional role within putatively regulatory cdDMR

sequence, since cdDMRs were also enriched for gene bodies and brain enhancer sequence.

This work quantifies this correlation for the first time, a DNAm relationship unique to only a

selection of cell types that includes neurons.

The identification of widespread association of mCpG and mCpH with expression and

specific splicing events, particularly in neuronal genes enriched for neuropsychiatric diseases,

highlights a potential novel role of mCpH and further expands the role of mCpG in the regulation

of gene expression in neurons. Splicing is predominantly a co-transcriptional process influenced

by changes in chromatin modifications and RNA binding proteins; the effects of DNAm on

splicing decisions is not yet well studied[31]. Here we found thousands of associations between

mC levels and gene, exon and PSI expression in developing postnatal neurons, particularly

featuring many exons that are exclusively associated with mCpH. Although it is not possible to

establish a causal role for mC in these data, these analyses, which are summarized in the provided website, can empower other researchers to explore the connection between DNAm and alternative isoform use, a phenomenon particularly prevalent in the developing brain that is often associated with disease[32].

We also explored the relationship between DNAm patterns and genetic associations with various phenotypes and found both expected and surprising associations. We confirmed enrichment for heritability of brain traits generally in neurons, and heritability for schizophrenia, a disorder with strong neurodevelopmental underpinnings, specifically in genomic regions losing DNAm preferentially in neurons over early postnatal development (ie, Group 3 cdDMRs). This result emphasizes the critical nature of neuronal development and maturation in early establishment of pathological connectivity and function for this adult-onset disorder, as most DNAm loss--generally associated with increased activity of a gene or regulatory element--occurred within the first five postnatal years. Indeed, Group 3 cdDMRs were present in genes such as *GRIN1*, *SYN1* and *CAMK2A*, and others involved in establishing synapse organization and function, a hallmark of early postnatal brain development, implicating abnormal genetic regulation of neuronal connectivity in schizophrenia development. Interestingly, heritability for PTSD was significantly enriched in regions preferentially gaining DNAm in the non-neurons over development (Group 4 cdDMRs), regions associated with neural precursor cell proliferation and cell-cell connective properties (Figure 1C). This association, particularly given the small amount of sequence covered by Group 4 cdDMRs, could lead to fruitful insights into susceptibility for PTSD and warrants further study.

We also found enrichment of genes associated with rare variants implicated in psychiatric and neurodevelopmental disease in a variety of cell types and developmental trajectories, highlighting the the genomic boundaries, developmental timing and cellular context

of epigenomic remodelling of regulatory elements or expressed features associated with known

risk genes. Two examples of this are *HDAC4* and *CACNA1B,* genes associated with ASD in the

SFARI Gene database. We identified a 566 bp Group 3 (decreasing neuronal, static glial DNAm)

cdDMR within an intron of *HDAC4*, a calcium-sensitive transcriptional repressor, and a 3.6 kb

Group 5 (decreasing neuronal, increasing glial DNAm) cdDMR within *CACNA1B*, a gene

encoding a voltage-gated calcium channel subunit (**Figure 4B-C)**. Even though both genes are

implicated in ASD and both cdDMRs are hypomethylated in neurons, the timing of loss of mCpG

suggests that *CACNA1B* activity occurs earlier in postnatal development than *HDAC4*. Given

that ASD onset is typically in early childhood, these risk genes may therefore have differing

implications in the etiology of ASD. Another example is a 646 bp Group 6 (decreasing glial,

static neuronal DNAm) cdDMR that overlaps the last intron and exon of *AKT3*, a

serine/threonine-protein kinase gene. Although the *AKT3*/1q44 locus has been associated with

schizophrenia risk, the mechanisms are not yet known given that *AKT3* is involved in many

biological functions[25]. Interestingly, this cdDMR selectively lost mCpG in glia beginning in

infancy, suggesting that *AKT3* activity in human DLPFC may be localized to glia beginning in

infancy or earlier (**Figure 4D**). This work provides the first *ex vivo* look at the DNAm dynamics

within human neurons and glia and thus allows for the first examination of those parameters

within the relevant organ, the brain.

Despite these insights, our data invoke several caveats. While NeuN based FANS

greatly improves identifying developmental DNAm changes over homogenate data, designating

NeuN+ and NeuN- samples as "neurons" and "glia" is not completely accurate in that NeuN-

samples will include signal from unlabeled neurons and mask non-neuronal diversity. However,

recent work assessing brain regional DNAm differences between NeuN+ and NeuN- found that

NeuN- contributed comparatively marginal variability in DNAm compared to NeuN+, suggesting

that neuronal methylomes are much more dynamic than non-neurons[20]. Likewise, while a

percentage of the bases in the DMRs identified in this work have also been previously shown to

be differentially methylated by neuronal subtypes whose unique methylomes are masked using

NeuN-based FANS[5,14,33], the proportion of these subtypes should be stable over postnatal

development[34]. Future epigenomics studies however can improve on the resolution of our study

by isolating more specific neuronal subpopulations to refine the cellular specificity of these

neuronal methylation changes largely occurring in the first few years of life.

Another caveat is that WGBS does not allow for the discrimination between mC and

hydroxymethyl-cytosines (hmC), an intermediary in the demethylation pathway. Previous work

has shown[5] that only a fraction of CpGs have measurable levels of hmC, suggesting that our

results are not confounded. In the cited study[5], hmCpG signal from homogenate cortex

represented 10% of the hypermethylation found in excitatory neuron mCpG, suggesting that

most of the mCpG signal in our neuronal data likely is true mCpG. The level of hmC in the

non-CpG context remains controversial, with some studies not identifying hmCpH[5] and others

detecting low amounts (1% in FANS-derived human glutamatergic and 0.47% in GABAergic

neurons)[35]. Further, FANS-derived human oligodendrocytes showed little hmC in the same

study[35]. Overlap of our cdDMRs with DMRs between neurons and oligodendrocytes in a study of

mC in FANS-derived human PFC samples[32] showed that cell type differences primarily reflected

true mC rather than hmC contamination in WGBS, while hmC DMRs between neuronal

subtypes primarily were reflected in hypomethylated neuronal cdDMR groups and not the

hypermethylated neuronal groups that would potentially include hmC signal contamination

(**Figure S7B**). Future work, however, should more closely examine the contribution of hmC to

the global hypermethylation seen during neurodevelopment.

By mapping the changing DNAm landscape over human postnatal neuronal and glial

development, we have identified unique trajectories of DNAm change particularly dynamic

during the first five years of life that show convergence between mCpG and mCpH, as well as

associations between single mCpG and mCpH and alternative splicing. These patterns may

also help illuminate the mechanisms through which psychological, neurological and psychiatric

traits are developed by placing known genetic contribution in an epigenomic context.

Acknowledgements

Author Contributions

● A.J.P.: Conceptualization, Formal Analysis, Investigation, Visualization, Writing – Original

Draft Preparation, Writing – Review & Editing.

● L.C.-T.: Conceptualization, Formal Analysis, Visualization, Software, Writing – Original

Draft Preparation, Writing – Review & Editing.

● N.A.I.: Software, Data Curation.

● W.X.: Investigation.

● E.E.B.: Formal Analysis, Writing – Review & Editing.

● J.H.S.: Supervision.

● R.T.: Investigation.

● L.M.: Investigation.

● Y.J.: Investigation, Supervision.

● T.M.H.: Data Curation, Resources, Writing – Review & Editing.

● J.E.K.: Data Curation, Resources, Writing – Review & Editing.

● D.R.W.: Conceptualization, Funding Acquisition, Supervision, Writing – Review &

Editing.

● A.E.J.: Conceptualization, Formal Analysis, Funding Acquisition, Supervision, Writing –

Original Draft Preparation, Writing – Review & Editing.

Funding

This project was supported by The Lieber Institute for Brain Development and by NIH grants

R21MH102791, R21MH105853, and R01MH112751. Data were generated as part of the

PsychENCODE Consortium, supported by: U01MH103392, U01MH103365, U01MH103346,

U01MH103340, U01MH103339, R21MH109956, R21MH105881, R21MH105853,

R21MH103877, R21MH102791, R01MH111721, R01MH110928, R01MH110927,

R01MH110926, R01MH110921, R01MH110920, R01MH110905, R01MH109715,

R01MH109677, R01MH105898, R01MH105898, R01MH094714, P50MH106934 awarded to:

Schahram Akbarian (Icahn School of Medicine at Mount Sinai), Gregory Crawford (Duke

University), Stella Dracheva (Icahn School of Medicine at Mount Sinai), Peggy Farnham

(University of Southern California), Mark Gerstein (Yale University), Daniel Geschwind

(University of California, Los Angeles), Fernando Goes (Johns Hopkins University), Thomas M.

Hyde (Lieber Institute for Brain Development), Andrew E. Jaffe (Lieber Institute for Brain

Development), James A. Knowles (University of Southern California), Chunyu Liu (SUNY

Upstate Medical University), Dalila Pinto (Icahn School of Medicine at Mount Sinai), Panos

Roussos (Icahn School of Medicine at Mount Sinai), Stephan Sanders (University of California,

San Francisco), Nenad Sestan (Yale University), Pamela Sklar (Icahn School of Medicine at

Mount Sinai), Matthew State (University of California, San Francisco), Patrick Sullivan

(University of North Carolina), Flora Vaccarino (Yale University), Daniel R. Weinberger (Lieber

Institute for Brain Development), Sherman Weissman (Yale University), Kevin White (University

of Chicago), Jeremy Willsey (University of California, San Francisco), and Peter Zandi (Johns

Hopkins University).

Competing Interest

The funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript. Conflict of Interest: none declared.

Data availability

Raw and processed sequence data that support the findings of this study have been deposited

at www.Synapse.org with the accession code: syn5842535.

Code availability

Code is available through GitHub at: https://github.com/LieberInstitute/brain-epigenomics.

Methods

Postmortem brain tissue acquisition and processing

As previously described in Jaffe *et al.* (2016)[2], post-mortem human brain tissue was obtained by

autopsy primarily from the Offices of the Chief Medical Examiner of the District of Columbia, and

of the Commonwealth of Virginia, Northern District, all with informed consent from the legal next

of kin (protocol 90-M-0142 approved by the NIMH/NIH Institutional Review Board). Additional

post-mortem prenatal, infant, child and adolescent brain tissue samples were provided by the

National Institute of Child Health and Human Development Brain and Tissue Bank for

Developmental Disorders (http://www.BTBank.org) under contracts NO1-HD-4-3368 and

546 NO1-HD-4-3383. Postmortem human brain tissue was also provided by donation with informed

547 consent of next of kin from the Office of the Chief Medical Examiner for the State of Maryland

548 (under Protocol No. 12-24 from the State of Maryland Department of Health and Mental

549 Hygiene) and from the Office of the Medical Examiner, Department of Pathology, Homer Stryker,

550 M.D. School of Medicine (under Protocol No. 20111080 from the Western Institute Review

551 Board). The Institutional Review Board of the University of Maryland at Baltimore and the State

552 of Maryland approved the protocol, and the tissue was donated to the Lieber Institute for Brain

553 Development under the terms of a Material Transfer Agreement. Clinical characterization,

554 diagnoses, and macro- and microscopic neuropathological examinations were performed on all

555 samples using a standardized paradigm, and subjects with evidence of macro- or microscopic

556 neuropathology were excluded, as were all subjects with any psychiatric diagnoses. Details of

557 tissue acquisition, handling, processing, dissection, clinical characterization, diagnoses,

558 neuropathological examinations, and quality control measures have also been further described

559 previously[36]. Homogenate postmortem tissue of the prefrontal cortex (dorsolateral prefrontal

560 cortex, DLPFC, BA46/9) was obtained from all subjects.

561 *Fluorescence activated nuclei sorting*

562 Nuclei were isolated from 100-300mg of pulverized DLPFC tissue using dounce homogenization

563 followed by ultracentrifugation over a sucrose density gradient. Homogenization was performed

564 on ice in 5mL lysis buffer [0.32M sucrose, 3mM magnesium acetate, 5mM calcium chloride,

565 5mM EDTA (ph 8.0), 10mM Tris-HCl (pH 8.0), 0.1% Triton X-100] and the resulting homogenate

566 was layered over 38mL sucrose buffer [1.8M sucrose, 3mM magnesium acetate, 10mM Tris-HCl

567 (pH 8.0)] and centrifuged at 139,800 x g for 2 hours at 4°C. Cellular debris and lysis and sucrose

568 buffers were removed and the pelleted nuclei were resuspended in 500uL PBS. Nuclei were

then labeled in a solution of anti-NeuN antibody conjugated to Alexa Fluor 488 (A60, Millipore,

1/1000) and 0.1% BSA, rocking for 30 minutes at 4°C, followed by the addition of DAPI. Nuclei

sorting was performed at the Johns Hopkins School of Public Health Flow Cytometry Core with

a MoFlo Legacy (Beckman Coulter) using Summit (version 4.3) software. The purity of sorted

populations was determined to be >99% based on resorting NeuN+ and NeuN- populations

through the same gates.

The identity of the NeuN+ and NeuN- populations as neuron-enriched and glia-enriched

was confirmed by sequencing nuclear RNA from each population and determining that neuronal

and glial biological processes were enriched in genes differentially expressed between the two

groups (FDR<0.05; **Figure S1B**). Likewise, cell type marker gene expression patterns also

corroborated the neuronal- and glial-enriched identities of NeuN+ and NeuN- samples (**Figure**

S1C). The estimated proportion of neurons in homogenate DNA methylation data based on

deconvolution using differentially methylated sites between NeuN+ and NeuN- samples was

highly correlated with the empirical proportion of neurons (**Figure S1D**). Raw sorting data is

shown in **Figure S16**.

Nucleic acid extraction and RNA-seq library preparations

RNA was extracted from homogenate and sorted samples using TRIzol LS Reagent (Thermo

Fisher Scientific) followed by the RNeasy MinElute Cleanup Kit (Qiagen). Genomic DNA

extraction was performed using the DNeasy Blood and Tissue Kit (Qiagen). Bisulfite conversion

of 600 ng genomic DNA was performed with the EZ DNA methylation kit (Zymo Research). RNA

sequencing libraries were made with the TruSeq RNA Library Prep Kit (Illumina) and the

RiboGone Low-Input Ribosomal RNA Removal Kit (Clontech). Library concentrations were

measured using a Qubit 2.0 and library fragment sizes were measured using a Caliper Life

Sciences LabChip GX. One hundred base-pair paired-end sequencing was run on an Illumina

HiSeq 2000.

Whole genome bisulfite library preparations and sequencing

Sequencing libraries were made with Illumina TruSeq DNA Methylation library preparation kits.

Lambda (λ) DNA sequence was spiked in at 1% concentration to assess bisulfite conversion

efficiency. Library concentrations were measured using a NanoDrop and library fragment sizes

were measured using an Agilent Bioanalyzer 2100. Libraries were spiked in with 10% PhiX to

improve base calibration calls and subsequently sequenced on an Illumina X-Ten Platform with

paired-end reads (2x150bp), targeting 30x coverage and Q30 > 70% read quality.

Data processing/alignment

We sought to align the paired-end reads for each sample to the in-silico bisulfite-treated hg19

genome, which we created using the Bismark v0.15.0[37] `bismark_genome_preparation`

program. For each library of paired-end reads (one per sample), the following processing was

performed (**Figure S17**):

· FastQC v0.11.4, to assess read quality, presence of adapter sequence, and

overrepresented sequences.

· Trimmomatic v0.35[38] to trim low quality and adapter-containing portions of the reads, with

the following parameters: `PE -threads 12 -phred33`

`ILLUMINACLIP:/Trimmomatic-0.35/adapters/TruSeq3-PE.fa:2:30:10:1`

`LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:75`. This resulted in three

sub-libraries of reads per sample: one paired-end sub-library, and then two single-end

sub-libraries where the other corresponding paired read was trimmed to a length below the

defined threshold.

·   FastQC v0.11.4, on each of the three sub-libraries, to assess the improvement in read

quality and adapter content following trimming.

·   FLASh v1.2.11[39], to merge the paired-end sub-library reads into longer single-end reads,

as reads that overlapped around CpGs and CpHs might bias or at the least double-count

the DNAm estimates. Furthermore Bismark[37] could only be run on single- or paired-end

reads and not a combination of both. This therefore further split the paired-end sub-library

into three sub-libraries: the subset of paired-end reads that were merged into longer

single-end reads, and then left and right single-end reads that could not be merged.

·   Bismark v0.15.0[37], to align each of the five now-single-end sub-libraries (left-trimmed,

right-trimmed, FLASh-merged, FLASh-left-unmerged, and FLASh-right-unmerged) to the

bisulfite-converted hg19 genome using bowtie2[40] and the --non-directional argument.

·   Resulting alignment (BAM) files across five sub-libraries were merged, sorted, and

indexed using samtools v1.3[41] to produce one large/merged BAM file per sample.

·   Alignments with evidence of duplication were removed using the MarkDuplicates program

in Picard tools v1.141, which systematically appeared to be localized to low complexity

DNA sequence near centromeres.

·   The Bismark[37] `bismark_methylation_extractor` program was run on each

post-duplicate-removed BAM file per sample to extract CpG and CpH DNAm levels.

We additionally aligned reads from each sample to the PhiX and Lambda (λ) genomes to

compute quality control metrics related to sequencing and bisulfite conversion quality. The

average percentage of reads mapping back to the λ genome was 1.32% and the average

bisulfite conversion efficiency was 98.64%. The average bisulfite conversion efficiency was not

associated (p>0.05) with cell type, age, cell type (adjusting for age), age (adjusting for cell type),

and the interaction of them in the NeuN- (glia) and NeuN+ (neuron) samples as well as for age

in the homogenate samples. The genome coverage decreased from an initial average of 43x to

10x across the processing stages as shown in **Figure S18 (coverage)**. For each of the

processing stages, there was no significant difference between cell types (adjusting for age),

age (adjusting for cell type) as well as the interaction between age and cell type (p

Bonferroni>0.05). The genome coverage was extracted from the `FASTQC` reports and by using

`bamcount` (https://github.com/BenLangmead/bamcount) v0.2.6.

Identifying methylation features

We identified PMDs, UMRs and LMRs using the bioconductor package MethylSeekR (version

1.20.0). We obtained coverage information from the cleaned set of ~18 million CpGs by

extracting coverage and methylation using the `getCoverage()` function from the bsseq

bioconductor package[42] v1.10.0. PMDs were called using `segmentPMDs()` and were visually

inspected using `plotPMDSegmentation()`. To create a more stringent cutoff for PMDs, we

filtered PMDs to those longer than 100 Kbp. We calculated the FDRs using

`calculateFDRs()`, while masking the >100 Kbp PMDs, setting the m parameter to 0.5 and

the FDR cutoff to 10. PMDs were further filtered to exclude overlaps with the UCSC "gap"

database table from hg19 except for the gaps labeled heterochromatin. We calculated UMRs

and LMRs using `segmentUMRsLMRs()`. DMVs were defined as UMRs in which pmeth was less

than or equal to 0.15 and the width was greater than or equal to 5 Kbp.

Identifying CpG differentially methylated regions (DMRs)

Using the bsseq[42] Bioconductor package v1.10.0 we loaded the Bismark[37] report files and

filtered the CpG data to keep only the bases where all samples had a minimum coverage of 3

(18,664,892 number passed the filter). We smoothed the methylation values of the remaining

CpGs using the `BSmooth()` function from bsseq with the `parallelBy="sample"` option. To

identify the age and cell type DMRs we used a model that adjusted for both covariates while for

the interaction model we included an additional interaction covariate. We identified the DMRs

using the bumphunter[43] Bioconductor package v1.14.0 using the `maxGap=1000, B=250,`

`nullMethod="permutation", smooth=FALSE` options, which tends to be conservative in

DMR identification[43]. For the `cutoff` option we used 0.1 for the cell type DMRs adjusting for

age, 0.005 for the age DMRs adjusting for cell type, and 0.009 for the age and cell type

interaction DMRs. These first two parameter cutoffs correspond to 10% minimum DNAm

differences between neurons and glia and 5% change in DNAm per decade of life across cell

types, and were chosen based on functionally-relevant change in DNAm. The cutoff for the

interaction model (cdDMRs) was based on selecting the equivalent percentile of change from

the overall age model (86th percentile) - this percentile-based cutoff was in line with

recommendations for selecting cutoffs for statistical models with less clear biological

interpretations[43]. We used a family wise error rate (FWER) threshold of 5% to determine the

DMRs: `fwer` output from the `bumphunter()` function. A small subset of DMRs involved a

single CpG, which arises from having a more significant area (length times effect size) than any

DMRs identified in null permuted data. All DMRs showed less than 10% median percent

absolute bias to technical and biological covariates as shown in **Figure S19 (sensitivity)**.

cdDMR processing

For the "interaction" DMRs with FWER<5% (cdDMRs) we extracted the methylation values from

the glial and neuronal samples using bsseq[42] v1.13.9 and then computed a mean methylation

value per DMR for each cell type. We also calculated the mean interaction coefficient for each

DMR across all the cytosines in the DMR by cell type. Using the mean coefficients by cell type

we clustered the interaction DMRs using the `kmeans()` function with `centers=6`,

`nstart=100` options. We chose `centers=6` based on biological interpretability of the results

and because k=6 results in an optimal AIC for clusters computed with mean centered and

scaled data. For each cytosine in the cdDMRs we calculated the t-statistic and coefficient for

age explaining differences in methylation adjusted for cell type: `~ age + cell type`. For

each DMR we computed the mean age coefficient by cell type and then calculated the median

absolute coefficient across all cdDMRs. The neuronal / glia ratio is 1.5 for such median absolute

age effects across the cdDMRs.

Comparing homogenate vs. cell type-specific WGBS

We first filtered CpGs to those with coverage in all 55 postnatal samples. For homogenate

samples, we used `lmFit()` and `ebayes()` from the limma[44] Bioconductor package v3.34.5 to

assess age-associated changes to DNAm levels with the linear model `~ Age`. For the cell

type-specific samples, we used a linear model `~ Age * Cell Type` to assess cell type,

overall age, and age in a cell type changes to DNAm levels. We subset CpGs to those that were

significantly differentially methylated by age in homogenate samples ($p<1\times10^{-4}$) and plotted the

coefficients for each CpG in **Figure S2A-D**. The relationship between each variable was

quantified using Fisher's exact test.

Roadmap Epigenome enrichments

We computed the relative enrichments of different genomic regions using Epigenome Roadmap

data[12] by computing the proportion of bases in each of the 15 ChromHMM states for each of the

cells and tissues provided by the Consortium. We compared the proportion of bases in each

state within each candidate region set to the overall genome and computed the corresponding

log2 enrichments between the regions and this genomic background. We compared DMR and

mCpG-based methylation feature regions to all profiled cell types in the Consortium for these

analyses.

Assessing the contribution of neuronal subtypes

The percent of neuronal subtype-specific bases were calculated by reducing the total

subtype-specific CpG-DMRs from Luo *et al.*[14], reducing the bases in each group of cdDMRs and

calculating the percent of cdDMR bases that intersected the merged subtype-specific CpG-DMR

bases.

Visualizing DMRs

Plots for the DMRs were made using bsseq[42] v1.14.0, EnsDb.Hsapiens.v75 v2.99.0, and

RColorBrewer v1.1-2. Genes within 20 kb of a DMR were retained for the plots. Genes and

exons were included using the `annoTrack` argument and we used extend=`2000` for making the

plots.

CpH processing

Using Bismark v0.16.3[37] we created report files using the methylation extractor program with the

`CX_context` and `split_by_chromosome` options for the hg19 human genome in order to

extract the methylation values for the CpHs. Then for each chromosome, using the bsseq[42]

Bioconductor package v1.10.0 we loaded the Bismark[37] report files and added the c_context

and trinucleotide_context information from Bismark using custom R code based on the bsseq

internal code that uses the data.table package v1.10.4. After combining the results for each

chromosome, we filtered the CpHs to keep only those where all samples had a minimum

coverage of 5 (58,109,566 number passed the filter).

Lister, et al. (2013) data processing

We downloaded the WGBS data from Lister *et al.* data[1] with SRA accession SRP026048. We

then processed and aligned the data following the same steps we used for our data. Using

bsseq[42] as in the *CpH processing* section, we extracted the methylation values from the

Bismark[37] report files and added the `c_context` and `trinucleotide_context` information

per chromosome. We then merged the results for all the chromosomes retaining only the CpG

and CpH positions we observed with a minimum coverage of 3 and 5 in our data, respectively.

To assess the replication of our cell type DMR results we computed mean methylation

differences across the CpG positions comparing neuron and non-neuron samples in the Lister *et*

al. data[1] using the `rowttests()` function from the genefilter package version 1.56.0. We then

computed the mean difference for each of the DMRs and compared this mean difference

against the DMR mean methylation difference derived from our data to derive the concordance

and correlation between them. To assess the replication of our age DMR results we modeled

age as a continuous variable and calculated the mean methylation difference per year for every

CpG contained in the age DMRs using `lmFit()` function from limma. Finally, we compared the

mean methylation difference in the Lister *et al.* data[1] against the observed mean methylation

difference for the age DMRs we derived.

Homogenate data processing

We processed the prenatal and postnatal homogenate brain samples using the same procedure

described in the *Data processing/alignment* section to produce Bismark[37] report files. Then

using bsseq[42] we extracted the methylation values for the CpG positions observed in our

postnatal samples in order to make them comparable to each other.

Identification of differentially methylated positions

With the set of CpGs and CpHs with a minimum coverage 3 and 5 in all samples, respectively,

and the same models for identifying the DMRs, we identified the differentially methylated

positions (DMPs), keeping the CpGs and CpHs separate. For the CpHs we further filtered to

keep only those where at least 5 samples had a methylation value greater than 0 (40,818,742,

70.2%). We used the limma Bioconductor package v3.30.13 for determining the DMPs by

running functions `lmFit()` and `eBayes()` with default parameters and FDR<5%.

 Enrichments for HARs and enhancers

We calculated the enrichment of genomic segments overlapping cdDMRs and human

accelerated regions (HARs) and enhancers[15] using Fisher's exact test. We calculated the

overlap of methylation features (DMRs, mCpH, expression-associated cytosines) and HARs or

enhancers with the entire set of CpG clusters used to identify the DMRs as background. We

corrected for multiple testing using the false discovery rate (FDR).

Stratified linkage disequilibrium score regression

GWAS summary statistics for 30 phenotypes (described in Anttila et al. and Rizzardi et

al.)[20,21] were downloaded from the sources listed in Table S6. We used LDSC (LD SCore) v1.0.0

to estimate the proportion of heritability captured in 16 sets of genomic DNAm features for each

GWAS phenotype, including the DMRs and LMRs defined above in this work, non-differentially

methylated CpG clusters (called above before the DMR analysis, excluding DMR sequence),

central nervous system annotations included in the LDSC package (referred as CNS (LDSC)),

and regions annotated as putatively regulatory in human brain using chromHMM (i.e., the union

of regions annotated as "Bivalent Enhancer", "Bivalent/Poised TSS", "Genic enhancers",

"Flanking Active TSS", "Active TSS", "Strong transcription", and "Enhancers" in the following

tracks accessed using the AnnotationHub Bioconductor package (v2.14.5)[45]: AH46920,

AH46921, AH46922, AH46923, AH46924, AH46925, AH46926, AH46927, AH46934, and

AH46935).

We first converted the GWAS summary statistics into the `.sumstats` format using

`munge_sumstats.py` and keeping only HapMap 3 SNPs (downloaded from

https://data.broadinstitute.org/alkesgroup/LDSCORE/w_hm3.snplist.bz2) as described in the

Partitioned Heritability LDSC tutorial. We made `.annot` files for each custom feature set based

on the list of SNPs in the CNS cell type annotations provided in the LDSC package and

estimated partitioned LD scores for each feature using 1000 Genomes plink files (downloaded

from https://data.broadinstitute.org/alkesgroup/LDSCORE/1000G_Phase3_plinkfiles.tgz) using

`ldsc.py`. We finally estimated the partitioned heritability for each feature-phenotype

combination by adding each feature individually to the "baseline model" including 53 baseline

annotations described in Finucane et al.[22]

Enrichments for genes associated with brain disorders

We calculated the enrichment of genes overlapping different methylation features in gene sets

described by Birnbaum *et al*.[24] We measured what fraction of the genes in each set overlapped

methylation features (ie, DMRs, mCpH, and expression-associated cytosines) with Fisher's

exact test using all expressed genes with Entrez IDs as background. We corrected for testing

multiple disorder gene sets using the false discovery rate (FDR).

RNA-seq data processing

Raw sequencing reads were mapped to the hg19/GRCh37 human reference genome with

splice-aware aligner HISAT2 v2.0.4[46]. Feature-level quantification based on GENCODE release

25 (GRCh38.p7) annotation on hg19 coordinates was run on aligned reads using featureCounts

(subread v1.5.0-p3)[47]. Using custom R code we processed the different feature counts and

created RangedSummarizedExperiment objects using the SummarizedExperiment

Bioconductor package v1.4.0.

Percent spliced in (PSI) calculation

We calculated the percent spliced in using the SGSeq[48] Bioconductor package v1.12.0 and the

Gencode v25 annotation for the GRCh37 human reference genome

(ftp.sanger.ac.uk/pub/gencode/Gencode_human/release_25/GRCh37_mapping/gencode.v25lift

37.annotation.gtf.gz) from the BAM files generated by HISAT2. We used default arguments

except for the function `analyzeVariants()` where we used a `min_denominator=10`.

Differential expression between cell types

We combined the gene counts for the polyA+ and RiboZero sequencing protocols for the

cell-sorted RNA-seq data: 3 NeuN+ and 3 NeuN- samples for a total of 12 RNA-seq sequencing

runs. We calculated the library size normalization factors using `calcNormFactors()` from

edgeR[49] (v3.22.3) and identified the differentially expressed genes using `voom()`, `lmFit()` and

`eBayes()` from limma[44,50]. We repeated this procedure for the exon counts.

Gene ontology analyses

Gene ontology enrichment analyses were performed using clusterProfiler[51] v3.6.0 using the

options `pAdjustMethod="BH"`, `pvalueCutoff=0.1`, and `qvalueCutoff=0.05` on the

Entrez IDs for each expression feature for the BP, CC and MF GO ontologies. Only cytosines or

DMRs overlapping genes were included.

Methylation vs expression associations

With the sorted RNA-seq data, we computed the average gene expression per age group

(infant, child, teen) and cell type (six groups), and correlated these values to average DNAm

levels in the same groups in both the CpG and CpH contexts at the gene promoter and body,

exon (500bp window), and splice junction (50bp into each intron) levels. With the

RangedSummarizedExperiment objects with the RNA-seq polyA homogenate data and the

bsseq objects with the CpG and CpH data we determined which CpG and CpH positions

explained changes in expression (RPKM) at the gene or exon level as well as in percent spliced

in (PSI). We retained only the expression and PSI data from the postnatal samples and matched

them by brain identifier to the neuronal methylation data with a final sample size of 22. We

filtered lowly expressed features using the `expression_cutoff()` function from the jaffelab

package v0.99.18: mean RPKM>0.22 mean for genes, 0.26 for exons. For genes and exons we

transformed the expression values to log2(RPKM + 1) and extracted the raw PSI values. Using

MatrixEQTL[52] v2.2 (GitHub b9a9f01 patch) we then identified the methylation quantitative trait

loci (QTL) for the CpG and CpH methylation data separately using the function

`Matrix_eQTL_main()` function with options `pvOutputThreshold=0`,

`pvOutputThreshold.cis=5e-4`, `useModel=modelLINEAR`, `cisDist=1000`. We identified

marginal CpG associations near CpH associations by running MatrixEQTL again for the CpG in

a +- 1kb window around the CpH positions with an association with expression at FDR <5% for

each expression feature type using the same parameters as above except for

`pvOutputThreshold.cis=0.01`. We filtered the associations to retain only those having at

least 11 samples with non-zero methylation and 11 samples with non-one methylation values to

remove extreme cases. We further restricted the results to protein coding genes and dropped

any with infinite t-statistics. To assess whether age confounds the relationship between

methylation and expression, we used a multiple linear regression model adjusting for age and

checked if the methylation coefficient was still FDR<5%. Venn diagrams in **Figure 3** were made

with the VennDiagram package v1.6.18. For the gene and PSI associations we used the gene

ID to check if it was present in the 3,473 differentially expressed genes from the sorted RNA-seq

data (described above) with higher expression in neurons and the top 5,000 DE genes with

higher expression in glia at FDR<5%; similarly we did so for exons and the top 5,000 DE exons

(FDR<5%) with higher expression in each cell type. The LIBD WBGS Expression explorer at

https://jhubiostatistics.shinyapps.io/wgbsExprs/ was made using the bsseq[42] v1.14.0, DT v0.4,

SGSeq[48] v1.12.0 and shiny v1.0.5 R packages.

Global autocorrelation

Using CpG and CpH positions with a minimum coverage of 3 and 5 respectively for all samples

we calculated the autocorrelation for the methylation levels for the CpGs, the CpHs, the CpHs

with a CHG trinucleotide context, or the CpHs with a CHH trinucleotide context. For each of the

sets, we grouped the positions using derfinder v1.12.0 into groups by a maximum distance of 1

kb. Only those groups with at least 5 Cs were further considered. For each sample we then

calculated the autocorrelation using the `acf()` function with `lag.max=4` in parallel for each

chromosome using BiocParallel v1.12.0. For each cluster of cytosines we calculated the mean

across the neuronal (NeuN+) and the glia (NeuN-) samples at each autocorrelation lag. After

combining and tidying the results, we visualized the global auto correlation using ggplot2 v2.2.1.

We repeated this same analysis for the Lister *et al.* data[1].

Autocorrelation within DMRs

Similar to the global autocorrelation, we extracted the methylation values at CpGs with a

minimum coverage of 3 and the CpHs with a minimum coverage of 5 that were within each of

the sets of DMRs (age, cell type or interaction). We then computed the autocorrelation for DMRs

with a least 5 different cytosines using the acf() function with a lag.max=4 and calculated the

mean auto-correlation among the neuronal and glial samples. The lag is proportional to the

genomic distance as shown in **Figure S20 (lag and distance)**.

	Table 1: Summary of methylation-expression associations. Mean values are shown for the

	number of samples with β>0, β<1, and expression change variables (either ΔPSI or

	Δlog2(RPKM + 1) for genes and exons). Columns from "Age Confounded" are proportions. See

	Table S10 for a full description of the variables.

Feature	Meth. Type	Direction	N	N Unique C	N Unique Features	Mean N Meth>0	Mean N Meth<1	Mean Expr Change	Prop. Age Confounded FDR >=5%	Prop. In Promoter	Prop. In Gene Body	Prop. In Gene Flanking	Prop. CHG	Prop. CHH	Prop. DE Glia	Prop. DE Neuron	Prop No Diff
gene	CpG	down	18458	18353	2074	21.9	21.9	1.39	0.0932	0.0671	0.966	0.906	0	0	0.126	0.329	0.546
		up	5217	5021	1743	21.8	21.8	1.69	0.0845	0.0571	0.979	0.92	0	0	0.304	0.193	0.503
	CpH	down	25812	25712	1611	21.1	21.1	1.4	0.115	0.0161	0.519	0.75	0.22	0.781	0.0912	0.413	0.496
		up	3842	3840	846	20.1	20.1	1.66	0.2	0.0219	0.532	0.807	0.26	0.744	0.123	0.27	0.607
exon	CpG	down	9505	4536	4168	21.9	21.9	1.23	0.104	0.19	0.934	0.991	0	0	0.0097	0.0256	0.965
		up	1328	638	687	21	21	1.93	0.191	0.175	0.928	0.988	0	0	0.0369	0.0136	0.95
	CpH	down	7874	4082	3458	21.1	21.1	1.72	0.074	0.0328	0.437	0.927	0.19	0.814	0.0086	0.0451	0.946
		up	1288	733	962	18.7	18.7	1.94	0.154	0.0435	0.481	0.918	0.28	0.722	0.0031	0.0311	0.966
psi	CpG	down	4286	2477	1315	21.7	21.7	0.124	0.0272	0.0434	0.994	0.936	0	0	0.242	0.212	0.547
		up	5782	3915	1646	21.8	21.8	0.121	0.0195	0.0398	0.991	0.919	0	0	0.261	0.222	0.517
	CpH	down	1296	945	489	16.8	16.8	0.447	0.0201	0.0274	0.537	0.676	0.36	0.644	0.187	0.286	0.526
		up	875	536	391	16.5	16.5	0.169	0.032	0.0279	0.559	0.775	0.36	0.637	0.194	0.33	0.477

Figure 1: Regional cell type-specific developmental mCpG trajectories. (A) Euclidean

distances between samples within cdDMRs shows that older neuronal samples cluster

separately from infant neuronal samples, glia regardless of age, and bulk prenatal cortex. **(B)**

Decomposing cdDMRs patterns into 6 clusters using *k-means* based on glia and neuron mean

mCpG changes per year of life. **(C)** The top five most enriched gene ontology terms for each of

the six groups in **(B)** highlights diverse biological processes amongst the groups. No terms were

enriched for Group 5. **(D)** Example Group 3 cdDMR within *SNAP25.* **(E)** Example Group 6

cdDMR within *MBP.* Gray shading indicates the boundaries of the cdDMR, and black tick marks

on the x-axis indicate the position of CpGs. Key: Neuron (NeuN+); Glia (NeuN-); Infant (0-1

years); Child (1-10 years), Teen (11-17 years) and Adult (18+ years).

Figure 2: CpH methylation patterns across brain development. (A) The proportion of CAC and CAG sites that are greater than 10% methylated in neurons and non-neurons (glia) across brain development. (B) Autocorrelation levels for different cytosine contexts in neurons. Autocorrelation levels were similar for mCpG and all cytosines, with uncorrelated levels in the CpH context. (C) Euclidean distances between samples based on mCpH within cdDMRs again cluster infant neurons dark red) with glia of all ages (light colors) rather than with older neurons.

Figure 3: Methylation associations with expression. **(A)** Venn diagrams of the methylation

associations by unique feature for the gene, exon and PSI. The sets are determined by if the

association is FDR<5% genome-wide for CpG and CpH or if it is a CpG marginally significant

within +/- 1 kb window of a CpH association. **(B)** Example associations between methylation

and expression at the gene level colored by age: red - infant, orange - child, green - teen, blue -

adult. *GUCY1A3* contains one of the top CpH differentially expressed between neurons and glia.

Expression of an exon of *TTN*, an autism-associated gene, is negatively associated with mCpH.

DOCK1 PSI of an alternative end site is negatively associated with mCpH. **(C)** Enriched

molecular function ontology terms for methylation-associated exons by the venn diagram groups

from (A).

Figure 4: DNAm patterns and brain trait heritability. **(A)** Results assessing enrichment for heritability of 30 phenotypes within 16 groups of DNAm features using Stratified Linkage Disequilibrium Score Regression (SLDSR). Each dot represents results for a single phenotype:DNAm feature pair. The color indicates the DNAm feature, and the phenotypes are stratified by column into psychiatric phenotypes, other brain-related phenotypes (ie, neurological or behavioral-cognitive), or non-brain-related traits. The upper row shows the coefficient Z score for each tested phenotype:DNAm pair, or the amount of additional heritability explained by the DNAm feature over 53 baseline features in the SLDSR model. The lower row shows the enrichment score, or the proportion of heritability attributed to the feature divided by the proportion of SNPs in the feature. For clarity enrichment scores of only the significant feature-trait combination are depicted. Filled in circles indicate significantly enriched heritability for a phenotype in a feature (coefficient p-value corrected using Holms method≤0.05). **(B)** A cdDMR overlapping *HDAC4*, a gene associated with autism spectrum disorder (ASD), shows the Group 3 pattern of decreasing neuronal and static glial DNAm. **(C)** A cdDMR overlapping *CACNA1B*, a gene associated with ASD, shows the Group 5 pattern of decreasing neuronal and increasing glial DNAm. **(D)** A cdDMR overlapping *AKT3*, a gene associated with schizophrenia,

shows the Group 6 pattern of decreasing glial and static neuronal DNAm. Gray shading

indicates the boundaries of the cdDMR, and black tick marks on the x-axis indicate the position

of CpGs.

Figure S1: Confirmation of neuronal- and glial-enriched identity of NeuN+ and NeuN-

samples. (A) Representative gating strategy from fluorescence-activated nuclear sorting by

NeuN antibody. Debris is first reduced by selecting events based on forward scatter and side

scatter, then aggregates are reduced by measuring pulse width. Autofluorescent events are

discarded by measuring true Alexa fluor-488 signal compared to signal in the FL2 channel.

Finally, singlets are determined by DAPI staining, and Alexa fluor-488 positive and negative

events are collected. **(B)** Top biological processes enriched in genes significantly differentially

expressed in nuclear RNA from NeuN+ (labeled "Neurons") and NeuN- (labeled "Glia") at

FDR<0.05. **(C)** T statistics for marker genes for neuronal and non-neuronal identity show that

the genes are differentially expressed in NeuN+ and NeuN- nuclear RNA (FDR<0.05). **(D)** The

estimated proportion of neurons in homogenate WGBS samples based on statistical

deconvolution using differentially methylated cytosines between NeuN+ and NeuN- samples is

highly correlated with the empirically-derived proportion of NeuN+ events.

Figure S2: Detecting developmental changes in homogenate vs. cell type-specific DNAm

data. (A) Developmental age effect coefficients of individual CpGs as measured in homogenate

samples compared against cell type effect coefficients in cell type-specific samples after

adjusting for age. **(B)** Developmental age effect coefficients of individual CpGs as measured in

homogenate samples compared against the developmental effect adjusting for cell type in cell

type-specific samples. **(C)** Developmental age effect coefficients as measured in homogenate

samples compared against the cumulative age and cell type interaction effect coefficients in cell

type-specific samples at the CpG level. **(D)** Developmental age effect coefficients as measured

in homogenate samples compared against the estimated cell type-specific age effects for cell

type-specific samples at the CpG level. **(E)** Overall age mean t-statistic for the cdDMRs (x-axis)

against the mean interaction t-statistic (y-axis) with the 2.5% quantile from the y-axis shown on

the x-axis. Colors are the same as those from **Figure 1B**: teal=Group 1 (decreasing glial

methylation, increasing neuronal methylation), orange=Group 2 (static glial methylation,

increasing neuronal methylation), purple=Group 3 (static glial methylation, decreasing neuronal

methylation), pink=Group 4 (increasing glial methylation, static neuronal methylation),

green=Group 5 (increasing glial methylation, decreasing neuronal methylation), gold=Group 6

(decreasing glial methylation, static neuronal methylation).

Figure S3: Unmethylated Regions (UMRs) and Low-methylated regions (LMRs). (A)
Number of LMRs and **(B) p**ercent of the genome covered by LMRs by age stratified by cell type.
Shading indicates the standard error of the linear model. **(C)** Roadmap Epigenomics Consortium
predicted chromatin-enriched states for LMRs and UMRs as defined in adult DPLFC and fetal
brain. log2(Coverage Ratio) represents the enrichment of the proportion of bases within LMRs
or UMRs in a chromatin state compared to the rest of the genome.

Figure S4: Partially methylated domains (PMDs). (A) Example PMDs on chromosomes 8, 12,

and 3. PMDs are highlighted in green. **(B)** Roadmap Epigenomics Consortium chromatin state

enrichment for PMDs. log2(Coverage Ratio) represents the enrichment of the proportion of

bases within PMDs in a state compared to the rest of the genome. **(C)** Percent of PMD

base-pairs that are replicated in the Lister *et al.* (2013) FANS samples[1]. Rows represent Lister

et al. (2013) glia and neurons, and columns represent the glia, neurons and prenatal samples

from this paper (LIBD). Each bar represents the percent of PMD bases shared between that

quadrant's cell types, using either total LIBD or total Lister PMD bases as the denominator. **(D)**

Percent of PMD base pairs per sample that are replicated in either common or unique PMDs

and high methylated domains (HMDs) identified in Zhou *et al.*[13] Samples are colored based on

whether the sample was postnatal neuron, postnatal glia, or bulk prenatal cortex.

Figure S5: DNA methylation valleys (DMVs). (A) Percent of the genome covered by DMVs in

postnatal glia and neurons and bulk prenatal cortex. **(B)** Overlap of transcription factor (TF)

genes within DMVs by age in neurons. **(C)** Expression enrichment between age groups in TF

genes excluded from DMVs in one group but not the other in neurons and glia. A negative

T-statistic signifies greater expression in the age group in which the gene is not in a DMV.

I=Infant (0-1 years); C=Child (1-10 years), T=Teen (11-17 years) and A=Adult (18+ years).

Figure S6: Differentially methylated regions (DMRs). (A) Replication of cell type DNAm differences at the CpG level between neurons and glia (NeuN+ and NeuN- samples) in our and Lister *et al.* data[1]. X and Y axes are linear model coefficients. (B) Enriched biological process ontology terms in the DMRs by cell type. The number of genes within each ontology group that overlaps a cell type DMR is listed. (C) Coefficients of linear regression on the mean mCpG level per cdDMR in samples younger than 6 years and older. (D) Coefficients of linear regression on the mean mCpG level per cdDMR in samples younger than 6 years and older.

 Figure S7: Cell type-specific developmental DMRs and neuronal subtype methylation and

 hydroxymethylation. (A) Percent of bases within each group of cdDMRs as defined in **Figure**

 1B that are differentially methylated by neuronal subtypes in Luo *et al.*[14]. **(B)** Percent of cdDMRs

within each of the 6 groups as defined in **Figure 1B** that overlap DMRs of methylcytosine (mC

captured using oxidative bisulfite sequencing), hydroxymethylcytosine (hmC), or total cytosine

methylation (tmC or mC+hmC, captured using standard bisulfite sequencing) as defined by

Kozlenkov *et al.*[50]. The plot is stratified by whether the Kozlenkov *et al.* (2018) DMRs are more

highly methylated in GABAergic or glutamatergic neurons, or in overall neurons compared to

oligodendrocytes. The number of overlapping cdDMRs are listed above each bar.

A. Enrichment for Enhancers

B. Enrichment for HARs

Figure S8: cdDMR overlap of human brain developmental enhancers and Human Accelerated Regions (HARs). (A) Enrichment for enhancers in the six clusters of cdDMRs from **Figure 1B** at the CpG-level. **(B)** Enrichment for human accelerated regions (HARs) in the six clusters of cdDMRs from **Figure 1B** at the CpG-level. Filled in circles indicate FDR<0.05, and error bars are the 95% confidence interval for the log(odds ratio).

 Figure S9: Cell type-specific, developmentally dynamic DMRs (cdDMRs) and epigenetic

states. (A) Percent of cdDMRs by the k-mean groups from **Figure 1B** overlapped by UMR,

LMR, DMV and PMD sequence across age. Lines show linear regression and shading indicates

the standard error. **(B)** Roadmap Epigenomics Consortium enriched chromatin states for the six

clusters of cdDMRs from **Figure 1B** at the CpG-level. log2(Coverage Ratio) represents the

enrichment of the proportion of bases within each cdDMR group in a chromatin state compared

to the rest of the genome.

Figure S10: CpH methylation distribution, levels and context-specific biological process

ontology. (A) Number of differentially methylated CpHs by cell type (FDR<5%) falling in

different genomic annotations across the genome. Annotation was prioritized CDS > 5'UTR >

3'UTR > Intron > Promoter > Intergenic. **(B)** Breakdown of measured CpH by trinucleotide

context. Differentially methylated mCpH by age in neurons (FDR<0.05) are colored in dark gray.

(C) Accumulation of mCpH by trinucleotide context over development, stratified by cell type. **(D)**

The top 20 biological process ontology terms enriched for genes exclusively overlapping CAG

and CAC sites with significantly increasing or decreasing methylation levels in neurons over

postnatal development (FDR<0.05).

Figure S11: Trajectories of methylation accumulation in cdDMR groups. Mean mCpH and

mCpG within the cdDMRs by cluster from **Figure 1B** across development. Loess line and

standard error shading are depicted.

1014   log2([mean junction-overlapping reads per 10 million mapped reads]+1) for the junctions on the

1015   y-axis. 10,000 pairs were chosen to reduce overplotting. Linear regression with shaded standard

1016   error and the rho for each correlation is listed for each plot. **(E)** Correlation of feature expression

1017   and methylation stratified by cytosine context (columns) and feature type (rows).

Figure S13: Cellular compartment ontology. Genes containing splicing events as measured

by "percent spliced in" (PSI) that are associated with changing CpG and CpH methylation are

enriched for cellular compartment gene ontology terms relating to neuronal features.

Figure S14: Molecular function ontology. Genes containing alternative exons that are associated with changing CpG and CpH methylation are enriched for molecular function ontology terms relating to neurons.

Figure S15: Web meQTL browser display. Interactive display of the CpHs and CpGs

associated with expression at FDR<5% as shown at

https://jhubiostatistics.shinyapps.io/wgbsExprs/. This screenshot shows the top nonCpG

(mCpH) meQTL association at the gene expression level. Information about the gene is shown

under *expression feature*, and information of the methylated C is shown under *cytosine*.

Figure S16: Raw sort data. Raw data collected from the MoFlo Legacy (Beckman Coulter)

using Summit (version 4.3) software. The density plots show DAPI signal on the y-axis and

Alexa Fluor 488-NeuN signal on the x-axis for events filtered via gates 1-3 as depicted in Figure

S1A. The histograms show the distribution of Alexa Fluor 488-NeuN signal after including the

gate for singlets based on DAPI signal (gate R14 in the raw data).

Figure S17: Data processing/alignment pipeline. Overview of the processing steps taken to

prepare the WGBS data.

Figure S18: Genome coverage across processing stages. (A) Genome coverage across the

four main processing stages: initial FASTQ files, after trimming, after mapping to the genome,

and after removing duplicated reads. **(B)** Same as (A) but with samples separated by cell type.

(C) Genome coverage across cell types and four age categories: infant, child, teen and adult.

(D) Genome coverage trajectories for each sample separated by age category and colored by

cell type.

Figure S19: DMR sensitivity analyses. This figure shows the percent absolute bias for the

three coefficients of interest used to define the DMRs: age (adjusting for cell type), cell type

(adjusting for age), and the interaction between age and cell type. For each DMR type we

computed the coefficients after adjusting for each of the covariates in the X-axis (β_{adj}) and

computed the percent absolute bias: $100 * |\beta_{adj} - \beta_{original}| / |\beta_{original}|$. All covariates considered

show less than 10% absolute bias. The covariates considered are: race, sex, PMI, pH, average

coverage, duplication percent, alignment efficiency, total number of trimmed reads, total number

of untrimmed reads, ratio of trimmed reads, and average conversion efficiency (lambda).

Figure S20: genome distance versus autocorrelation lag. For each of the cytosines

considered for the autocorrelation analysis in **Figure 2B**, we computed the genome base-pair

distance and averaged it for each group of cytosines. **(A)** Average genomic distance for each of

the three DMR models and for various contexts of cytosines. **(B)** Average genome distance for

the interaction DMRs (cdDMRs) for all cytosines, CpG and CpH. The genome distance is

proportional to the lag. This panel is the exact complement to **Figure 2B**.

Supplementary Tables

Table S1: Phenotype and sequencing metrics for the WGBS samples.

See sheet 1 of Supplementary_tables.xlsx.

Table S2: Phenotype and sequencing metrics for the RNA-seq samples.

See sheet 2 of Supplementary_tables.xlsx.

Table S3: Number of cytosines measured and distribution of methylation by cytosine context. Sample sizes for each sample type, number of CpGs and CpHs and distribution of CpG and CpHs across low, partial and high levels of methylation.

	N	CpG Context				CpH Context			
		# CpGs	Low (<20%)	Partial (20-80%)	High (>80%)	# CpHs	Low (<20%)	Partial (20-80%)	High (>80%)
Neuron	24	18.7×10^6	10.10%	13.90%	76.00%	58.1×10^6	91.60%	8.00%	0.40%
Glia	8	18.7×10^6	11.10%	17.80%	71.10%	58.1×10^6	99.40%	0.50%	0.10%
Fetal	20	18.7×10^6	12.10%	17.30%	70.60%	58.1×10^6	99.75%	0.17%	0.07%
Homogenate	23	28.2×10^6	8.50%	16.90%	73.70%	58.1×10^6	NA	NA	NA

Table S4: Cell type-specific, developmental differentially methylated regions (cdDMRs).

See sheet 4 of Supplementary_tables.xlsx.

1066 **Table S5: mC association with cell type and age in postnatal cell type-specific samples.**

1067 Number of differentially methylated CpGs and CpHs across each of the statistical models: cell

1068 type effects adjusting for age, age effects adjust for cell type, cell type and age interaction

1069 effects, as well as age in neuron samples only.

Model	CpG Context		CpH Context	
	FDR ≤ 0.05	% Significant	FDR ≤ 0.05	% Significant
Cell type	4824804	25.80%	7682075	18.80%
Age	536164	2.90%	3194618	7.80%
Cell type:Age	90227	0.50%	76	0%
Age in Neurons	NA	NA	4020371	9.80%

Table S6: GWAS traits assessed using LDSC.

See sheet 6 of Supplementary_tables.xlsx.

Table S7: Stratified linkage disequilibrium score regression results.

See sheet 7 of Supplementary_tables.xlsx.

Table S8: Enrichment of DMRs and mCpH for disease-associated gene sets.

See sheet 8 of Supplementary_tables.xlsx.

Table S9: Enrichment for disease gene sets in DNAm-splicing association features.

See sheet 9 of Supplementary_tables.xlsx.

 Table S10: Variable dictionary for Table 1. Variables used in **Table 1** with their full definition.

Variable	Description
Feature	Either gene, exon or event affect the percent spliced in (PSI).
Methylation	Methylation type. Either CpG or CpH.
Direction	down.
N	Number of associations at FDR 5%.
N Unique C	Number of unique cytosines in these associations.
N Unique Features	Number of unique features in these associations.
N Meth>0	Mean number of samples that have non-zero methylation values. Total 22.
N Meth<1	Mean number of samples that have non-one methylation values.
Expr Change	Mean change in expression (ignoring NAs).
Age Confounded	Proportion of associations that are confounded by age, that is, proportion of associations where the methylation coefficient in a multiple linear regression adjusting for age has a FDR >= 5%.
Promoter	Proportion of cytosines in these associations that overlap gene promoters. A single cytosine can overlap multiple gene regions for different genes.
Gene Body	Proportion of cytosines in these associations that overlap gene bodies.
Gene Flanking	Proportion of cytosines in these associations that overlap gene flanking regions.
CHG	Proportion of cytosines in these associations that have a CHG context.
CHH	Proportion of cytosines in these associations that have a CHH context.
Glia	Proportion of features in these associations where the feature is differentially expressed between neurons and glia, with higher expression in glia. For genes and PSI, the gene list was used. Up to the top 5,000 features DE in each direction were used.
Neuron	Proportion of features in these associations where the feature is differentially expressed between neurons and glia, with higher expression in neurons.
No Diff	Proportion of features in outside the top differentially expressed genes between glia and neuron.

Bibliography

1. Lister, R. *et al.* Global epigenomic reconfiguration during mammalian brain development. *Science* **341,** 1237905 (2013).

2. Jaffe, A. E. *et al.* Mapping DNA methylation across development, genotype and schizophrenia in the human frontal cortex. *Nat. Neurosci.* **19,** 40–47 (2016).

3. Jaffe, A. E. & Irizarry, R. A. Accounting for cellular heterogeneity is critical in epigenome-wide association studies. *Genome Biol.* **15,** R31 (2014).

4. Schultz, M. D. *et al.* Human body epigenome maps reveal noncanonical DNA methylation variation. *Nature* **523,** 212–216 (2015).

5. Mo, A. *et al.* Epigenomic signatures of neuronal diversity in the mammalian brain. *Neuron* **86,** 1369–1384 (2015).

6. Jeong, M. *et al.* Large conserved domains of low DNA methylation maintained by Dnmt3a. *Nat. Genet.* **46,** 17–23 (2014).

7. Xie, W. *et al.* Epigenomic analysis of multilineage differentiation of human embryonic stem cells. *Cell* **153,** 1134–1148 (2013).

8. Stadler, M. B. *et al.* DNA-binding factors shape the mouse methylome at distal regulatory regions. *Nature* **480,** 490–495 (2011).

9. Lister, R. *et al.* Human DNA methylomes at base resolution show widespread epigenomic differences. *Nature* **462,** 315–322 (2009).

10. Salhab, A. *et al.* Partially methylated domains are hallmarks of a cell specific epigenome topology. *BioRxiv* (2018). doi:10.1101/249334

11. Hovestadt, V. *et al.* Decoding the regulatory landscape of medulloblastoma using DNA methylation sequencing. *Nature* **510,** 537–541 (2014).

12. Roadmap Epigenomics Consortium *et al.* Integrative analysis of 111 reference human

epigenomes. *Nature* **518,** 317–330 (2015).

13. Zhou, W. *et al.* DNA methylation loss in late-replicating domains is linked to mitotic cell

division. *Nat. Genet.* **50,** 591–602 (2018).

14. Luo, C. *et al.* Single-cell methylomes identify neuronal subtypes and regulatory elements in

mammalian cortex. *Science* **357,** 600–604 (2017).

15. Emera, D., Yin, J., Reilly, S. K., Gockley, J. & Noonan, J. P. Origin and evolution of

developmental enhancers in the mammalian neocortex. *Proc Natl Acad Sci USA* **113,**

E2617-26 (2016).

16. Doan, R. N. *et al.* Mutations in human accelerated regions disrupt cognition and social

behavior. *Cell* **167,** 341–354.e12 (2016).

17. He, Y. & Ecker, J. R. Non-CG Methylation in the Human Genome. *Annu. Rev. Genomics*

Hum. Genet. **16,** 55–77 (2015).

18. Eckhardt, F. *et al.* DNA methylation profiling of human chromosomes 6, 20 and 22. *Nat.*

Genet. **38,** 1378–1385 (2006).

19. Shukla, S. *et al.* CTCF-promoted RNA polymerase II pausing links DNA methylation to

splicing. *Nature* **479,** 74–79 (2011).

20. Rizzardi, L. F. *et al.* Neuronal brain-region-specific DNA methylation and chromatin

accessibility are associated with neuropsychiatric trait heritability. *Nat. Neurosci.* **22,**

307–316 (2019).

21. Brainstorm Consortium *et al.* Analysis of shared heritability in common disorders of the

brain. *Science* **360,** (2018).

22. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide

association summary statistics. *Nat. Genet.* **47,** 1228–1235 (2015).

23. Skene, N. G. *et al.* Genetic identification of brain cell types underlying schizophrenia. *Nat.*

Genet. **50,** 825–833 (2018).

24. Birnbaum, R., Jaffe, A. E., Hyde, T. M., Kleinman, J. E. & Weinberger, D. R. Prenatal expression patterns of genes associated with neuropsychiatric disorders. *Am. J. Psychiatry* **171,** 758–767 (2014).

25. Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511,** 421–427 (2014).

26. Howell, K. R., Floyd, K. & Law, A. J. PKBγ/AKT3 loss-of-function causes learning and memory deficits and deregulation of AKT/mTORC2 signaling: Relevance for schizophrenia. *PLoS ONE* **12,** e0175993 (2017).

27. Wang, D. *et al.* Comprehensive functional genomic resource and integrative model for the human brain. *Science* **362,** (2018).

28. Stiles, J. & Jernigan, T. L. The basics of brain development. *Neuropsychol. Rev.* **20,** 327–348 (2010).

29. Tierney, A. L. & Nelson, C. A. Brain development and the role of experience in the early years. *Zero Three* **30,** 9–13 (2009).

30. Stroud, H. *et al.* Early-Life Gene Expression in Neurons Modulates Lasting Epigenetic States. *Cell* **171,** 1151–1164.e16 (2017).

31. Lev Maor, G., Yearim, A. & Ast, G. The alternative role of DNA methylation in splicing regulation. *Trends Genet.* **31,** 274–280 (2015).

32. Jaffe, A. E. *et al.* Developmental and genetic regulation of the human cortex transcriptome illuminate schizophrenia pathogenesis. *Nat. Neurosci.* **21,** 1117–1125 (2018).

33. Kozlenkov, A. *et al.* Substantial DNA methylation differences between two major neuronal subtypes in human brain. *Nucleic Acids Res.* **44,** 2593–2612 (2016).

34. Sahara, S., Yanagawa, Y., O'Leary, D. D. M. & Stevens, C. F. The fraction of cortical

GABAergic neurons is constant from near the start of cortical neurogenesis to adulthood. *J. Neurosci.* **32,** 4755–4761 (2012).

35. Kozlenkov, A. *et al.* A unique role for DNA (hydroxy)methylation in epigenetic regulation of human inhibitory neurons. *Sci. Adv.* **4,** eaau6190 (2018).

36. Lipska, B. K. *et al.* Critical factors in gene expression in postmortem human brain: Focus on studies in schizophrenia. *Biol. Psychiatry* **60,** 650–658 (2006).

37. Krueger, F. & Andrews, S. R. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics* **27,** 1571–1572 (2011).

38. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30,** 2114–2120 (2014).

39. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* **27,** 2957–2963 (2011).

40. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9,** 357–359 (2012).

41. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,** 2078–2079 (2009).

42. Hansen, K. D., Langmead, B. & Irizarry, R. A. BSmooth: from whole genome bisulfite sequencing reads to differentially methylated regions. *Genome Biol.* **13,** R83 (2012).

43. Jaffe, A. E. *et al.* Bump hunting to identify differentially methylated regions in epigenetic epidemiology studies. *Int. J. Epidemiol.* **41,** 200–209 (2012).

44. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* **43,** e47 (2015).

45. Morgan, M. *AnnotationHub: Client to access AnnotationHub resources*. (Bioconductor, 2019). at <http://bioconductor.org/packages/release/bioc/html/AnnotationHub.html>

46. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory

requirements. *Nat. Methods* **12,** 357–360 (2015).

47. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. *Bioinformatics* **30,** 923–930 (2014).

48. Goldstein, L. D. *et al.* Prediction and Quantification of Splice Events from RNA-Seq Data.

PLoS ONE **11,** e0156132 (2016).

49. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26,** 139–140

(2010).

50. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock linear model

analysis tools for RNA-seq read counts. *Genome Biol.* **15,** R29 (2014).

51. Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R package for comparing

biological themes among gene clusters. *OMICS* **16,** 284–287 (2012).

52. Shabalin, A. A. Matrix eQTL: ultra fast eQTL analysis via large matrix operations.

Bioinformatics **28,** 1353–1358 (2012).