MISSED OPPORTUNITIES OF CHLAMYDIA AND GONORRHEA DETECTION WHEN NOT USING EXTRA GENITAL SCREENING AMONG MALES

Chelsea Schafer, BS, CHES, Belinda Prado, MPH, Nora Barin, MPH, and Leslie Gama, BS.

Contact: The City of Long Beach, Department of Health and Human Services, STD/HIV

Surveillance; 2525 Grand Avenue, Long Beach, CA 90815

Phone: (310) 527-2062

E-mail: Belinda.Prado@longbeach.gov; Chelsea.A.Schafer@gmail.com

Statement of Disclosure:

There is no known conflict of interest that exists for purposes of this study.

Sources of Support:

We would like to acknowledge the California Department of Public Health, Centers for Disease

Control and Prevention, and the City of Long Beach with special thanks to the Health and

Human Services Department.

Abstract

Objective

Through extra genital screening methods, Health Departments and clinics can increase detection of sexually transmitted diseases (STDs) when used with urethral testing.

Methods

Retrospective medical review of 1,571 patient health records, who received CT/GC testing, based on at least 1-visit to the Long Beach Department of Health & Human Services, STD Clinic, between 2012 to 2015. All male patients ages 18 years or older with positive CT/GC results (242 cases) for any of the three sites (e.g. urethral, rectal, and pharyngeal); regardless of their sexual behaviors, were included in the study. Females, those under the age of 18, and patients who tested negative for all three anatomical sites were excluded (1,412 controls).

Results

At time of collection, study participants had a mean age of 37 years. Reported ethnicity indicated 56% Caucasian, 21% Hispanic or Latino, 9% Asian or Pacific Islander, 7% Other, 5% Black, and 2% More than one race. The use of extra genital screening detected 15% (242) of the 1,571 patients tested positive for at least 1-type of CT/GC infection. Our findings demonstrated that if urine was the only specimen collected, then over 29.7% of CT and 46.8% of GC cases would have been missed.

Conclusion

Testing of all three anatomical sites should continue to be performed for CT/GC detection. Cases of CT/GC are underreported if performing urethral screening alone. It is a tool for prevention and control to reduce the potential for missed cases of CT and GC. These results highlight the need for clinicians to perform extra genital screening among male patients to avoid missing a

45   diagnosis, mitigate transmission, and report the number of cases that would have gone

46   undiagnosed.

47   *Keywords: Chlamydia, Gonorrhea, extra genital screening, Sexually Transmitted*

48   *Diseases, Men's health*

Introduction

Sexually transmitted diseases (STDs) have been an ongoing public health concern as communicable disease rates have steadily increased since 2011. In 2016, Long Beach, California experienced the second highest rate of Chlamydia (CT) and third highest rate of Gonorrhea (GC) reported for the State of California [4]. People can become infected through sexual intercourse, by having contact with the bacterium *Chlamydia trachomatis* and/or *Neisseria gonorrhoeae* [5]. Once infected, individuals may experience pain during urination, discharge from penis, and swollen testes. Infection can be detected in the genitals, rectum, and throat. However, it is possible to show no signs or symptoms of infection, being asymptomatic [6-7]. Testing of these STDs is critical to determine a person's infection status and can be detected in the genitals, rectum, and throat [5-7]. Based on 2017 STD & HIV Screening Recommendations from the CDC, annual CT/GC screenings should be performed among all sexually active males, regardless of condom use [8]. Screening is recommended every 3-6 months if the individual has multiple or anonymous sex partners. Despite CDC recommendations, there is still a lack of performing extra genital screening. According to the CDC, one in three clinicians have not heard of extra genital screening [9]. Previous research shows most patients do not report their sexual behaviors and have infections which are asymptomatic [9-11]. Men who have sex with men (MSM) are disproportionately affected for contracting STDs; increasing their risk for co-infections such as human immunodeficiency virus (HIV) [12]. Young males are at greater risk for infection if residing in areas with a high prevalence of CT [8, 13]. This STD screening measure, also known as triple-site testing, has been the method of choice for CT/GC detection, as it examines pharyngeal, rectal, and urethral sites with first-void urine samples using nucleic acid amplification tests (NAATs). Extra genital screening is considered the most sensitive in regard to

specificity to cell culture and recognition of bacterial sample assays [11, 14]. There is currently no

standard recommendation on screenings for GC among heterosexual males [8].

The City of Long Beach faces one of the highest morbidity rates of CT and GC in

California [4]. From 2012-2016, CT increased by 51% (530 to 801 per 100,000), while rates of

GC have increased by 214% (98 to 309 per 100,000) [4]. In 2017, the California Department of

Public Health reported rates of CT to be 552.2 per 100,000 population, with Los Angeles having

a rate of 560 per 100,000 population. That same year, rates of GC were 190.3 per 100,000

population in California [6, 15]. Communicable disease rates have been on the rise, calling for

improved methods for early detection and STD testing. Screening programs for these harmful

infections are made widely available to reduce the burden of disease in the population. Extra

genital screening tests patients in their genital, urethral, and pharyngeal sites. Although urine is

most widely used to test for CT/GC infection, it is still recommended to test all three sites as an

individual's sexual practices can place them at greater risk for infection, depending on the type of

sex they are having [16, 18].

African American males in Long Beach are reported to have the highest rates for CT

(727.1 per 100,000 population in 2016) and GC (667.6 per 100,000 population in 2016) [4].

African American MSM have significantly higher prevalence of HIV than compared to White

males [16-17]. Information on the underlying disparities regarding prevalence and incidence of

disease is necessary for bridging the gap in services. These issues could be related to differences

in cultural backgrounds and risky sexual behaviors such as lack of condom use or high number

of sex partners [16-17]. In a previous study conducted by Kent, Chaw et al. [11] at a health clinic in

San Francisco, researchers noticed that if patients were only screened at urethral sites, 53% of

CT and 64% of GC infections would have been missed. In the same study, among MSM

population, more than 70% of CT infections would have been missed if they only tested for GC

infection. This would have been based on a false assumption that among those who tested

positive for GC infection, that patients were also tested for CT infection [11]. Other researchers

have reported similar findings stating that if a clinic does not perform extra screening measures,

then roughly 70% of CT/GC cases, would have gone unreported [19-20].

Due to high rates of STDs in the City of Long Beach, there is increased need for

improved detection, patient education, and surveillance of CT/GC among residents. Extra genital

screening of CT/GC detects more cases of infections, when using nucleic acid amplification tests

(NAATs), enhancing sensitivity and specificity [16, 21]. STD testing is performed

opportunistically, regardless if patients show any signs or symptoms, since many of the cases

seen are asymptomatic [6]. Examining high-risk populations, based on their sexual behaviors and

infection status, is most cost effective and has demonstrated a reduction in the incidence of

CT/GC infections [6]. Peters, et al. [6] and his research team followed screening protocols for

MSM, collecting urine samples and urethral swabs. Their reasoning and methodology were

based upon national testing guidelines obtained from the United Kingdom, Norway, Australian,

and the United States of America, Centers for Disease Control and Prevention [6].

The purpose of this study was to determine the number of CT/GC cases that would have

been missed and left untreated if not performing extra genital screening among male patients

who were tested at the LBDHHS, STD Clinic. Patient and provider education are needed to

increase awareness, while reducing stigma associated among those seeking STD testing and

treatment. The transmission and susceptibility of CT/GC infection is reported to be enhanced

among patients with HIV [22]. Performing urethral screening alone creates the opportunity for

missed infections in other anatomical sites. Health Departments and STD Clinics should

continue to perform extra genital screening of all three anatomical sites as there is limited data at

the national level, underestimation of disease prevalence, and risk for HIV [19-20]. STD screening

guidelines should continue to be followed to reduce incidence of new cases.

Materials and Methods

Retrospective medical review of 242 cases identified as male and tested CT/GC positive

after extra genital screening. A total of 1,571 patients were tested for CT/GC at the LBDHHS,

STD Clinic during a 4-year period, 2012-2015. Study participants were chosen based on whether

they received STD testing during at least 1-visit to the on-site clinic at LBDHHS. Females, those

under the age of 18, and patients who tested negative for all three anatomical sites were excluded

(1,412 controls). All male patients ages 18 years or older with positive CT/GC results (242

cases) for any of the three sites, including urethral, rectal, and pharyngeal; regardless of their

sexual behaviors, were included in the study. All health records were collected during patient

intake with no attempt to re-contact. Patients' medical charts were pulled from NextGen v5.6 to

assess data on STD test results and demographics. The LBDHHS lab performed internal

validation of nucleic acid amplification testing (NAAT) to screen extra genital sites, as well as

urine testing to test for CT/GC positivity. NAATs are the most sensitive assays with a specificity

similar to cell culture and are considered the method of choice for CT/GC detection. In

particular, sensitivity and specificity of STD testing is reported to be significantly enhanced by

application of molecular techniques with NAATs [21]. All patients were evaluated in a clinical

setting and diagnosed in accordance to California Department of Public Health (CDPH)

guidelines, with serological confirmation [13]. Test positivity was defined as the proportion of

males who had positive extra genital and/or urogenital CT/GC test [3, 18, 23]. Patients are offered

treatment once diagnosed, following California STD treatment guidelines from 2015 [13].

Recommended regimens for adults who tested positive for CT genital, rectal, and pharyngeal

infections include Azithromycin (1 g po) or Doxycycline (100 mg po bid x 7 days). Adults

positive for GC, are recommended dual therapy with Ceftriaxone 250 mg IM and Azithromycin

1 g po [13].

To determine the amount of missed CT/GC infections, analysis was done using Pearson's

Chi-Square test for association to generate a cross-tabulation and frequency tables using SPSS

Statistics 24 [3, 18, 23]. To determine the number of missed CT/GC infections: cases were assessed

using those who tested positive per disease at site of infection, including those positive for both

pharyngeal and rectal, yet negative urethral sample [3, 18, 23]. Recent sexual contact determined

symptoms of CT and/or GC. Convenience sampling methodology was used for patient selection,

with persons who received STD testing at the on-site Clinic. Sexual behaviors were recorded into

categorical variables, with patients grouped as either men who have sex with men (MSM), men

who have sex with women (MSW), or men who have sex with men and women (MSMW). For

operationalization measurement in our health record review, we chose variables that had

exhaustive and mutually exclusive categories. To increase the likelihood of the successful

achievement of our study objective, application of sample size selection using threshold

probability of a (two-tailed) α test =.05, β =.20, and r=.20, indicated a minimum sample size of

194 cases for statistical significance [24].

Results

At time of collection, age of patients ranged from 15-75, with a mean age of 37 years.

52.5 % identified as "*Not Hispanic or Latino(a)*," 42.6% "*Hispanic or Latino(a),*" and 5.0%

"*Unknown or Not Reported*" for ethnicity. Patients reported race indicated, 56% White, 21%

African American or Black, 9% Asian or Pacific Islander, 5% Hispanic or Latino, and 2% More

than one race. Our results indicated that an estimated 29.7% of CT and 46.7% of GC infections would have been missed if only performing urethral testing. There were 5.9% (10 cases) pharyngeal, 20.8% (35 cases) rectal, and 2.9% (5 cases) pharyngeal-rectal cases who tested positive for CT and were negative by urine (Table 1).

Table 1. Chlamydia (CT) by Extra Genital Site among Male patients, Long Beach, 2012-2015 (n=242)	
Extra Genital Screening Test Results	# Cases
*Pharyngeal (negative), Rectal (negative), and Urethral (negative)	118
Pharyngeal (positive) and Urethral (negative)	10
Pharyngeal (positive), Rectal (positive), and Urethral (negative)	5
Rectal (positive) and Urethral (negative)	35
Urethral (positive)	74
Total:	**242**
% Missed CT	**29.7% (50/168)**

This table includes all cases who tested positive for a single site, those who tested positive for two sites (Rectal & Pharyngeal) and those who tested negative for all three sites.
*Those who were negative by Urethral, and negative/not done for Pharyngeal and/or Rectal test.

In addition, there were 16.0% (25 cases) of pharyngeal, 16.7% (26 cases) rectal, and 14.1% (22 cases) pharyngeal-rectal cases who tested positive for GC and were negative by urine (Table 2).

Table 2. Gonorrhea (GC) by Extra Genital Site among Male patients, Long Beach, 2012-2015 (n=242)	
Extra Genital Screening Test Results	# Cases
*Pharyngeal (negative), Rectal (negative), and Urethral (negative)	83
Pharyngeal (positive) and Urethral (negative)	25
Pharyngeal (positive), Rectal (positive), and Urethral (negative)	22
Rectal (positive) and Urethral (negative)	26
Urethral (positive)	86
Total:	**242**
% Missed GC	**46.8% (73/156)**

This table includes all cases who tested positive for a single site, those who tested positive for two sites (Rectal & Pharyngeal) and those who tested negative for all three sites.
*Those who were negative by Urethral, and negative/not done for Pharyngeal and/or Rectal test.

To determine if there was a difference in the amount of cases reported from 2012 to

2016, a percent change calculation was performed. This demonstrated an increase of CT rates by

51% (530 to 801 per 100,000) and GC rates by 214% (98 to 309 per 100,000).

Discussion

These findings provide data on the incidence and prevalence of CT/GC cases in The City

of Long Beach. Results from our analysis show the need for implementing extra genital

screening, in addition to urine testing, among male patients in clinical settings to prevent cases

from going undiagnosed. The use of extra genital screening with NAATs may improve detection

of asymptomatic cases or those that would otherwise be undiagnosed if a patient does not report

their sexual health history. Testing of all three anatomical sites should be performed routinely at

health clinics to detect for infection early since patients can be asymptomatic. Fewer cases of

CT/GC are to be reported among sexually active males. It is recommended to continue these

practices across clinical settings to control the spread of infection and ensure the appropriate

treatment is being offered. Due to lack of data at the national level, and provider education on

extra genital screening, there is underestimation of CT/GC prevalence [9, 19]. About one in three

clinicians have not heard of extra genital screening, which has resulted in the lack of performing

these testing procedures [9].

Study limitations included selection bias through the use of convenience sampling

methodology. Distinguishing temporal precedence was determined to be a consequence of using

retrospective health records for assessment. This study contributes to the field of epidemiology

by providing information on CT/GC prevalence among males. Early detection and treatment of

these bacterial infections can help in the prevention and control of other STIs, such as HIV and

HPV. The Health Department is one of the most well-suited entities for having a surveillance

program, as they can reach the most vulnerable populations through targeted interventions.

However, there is still a lack of performing extra genital screenings for CT/GC, even if indicated.

It is recommended by the Centers for Disease Control and Prevention (CDC) that annual

screenings be done based on those at greatest risk for infection [22].

Evaluation of CT/GC is important for ensuring that the appropriate medical tests are

being done, patients receive adequate treatment, and a reduction STD rates. Extra genital

screening needs increased support as a recommended STD testing procedure. This is especially

true among MSM, due to sexual practices that put them at increased risk for infection such as

anal intercourse. Despite national organizational recommendations, more support is still needed

if wanting to include this STD testing and screening method for other populations such as

females. Surveillance data might not be well representative of CT testing and reported rates.

Findings from researchers Kustec, et al. [25] indicate underreporting of CT cases among females

to the National Institute of Public Health (NIPH) [25]. Health programs should bring increased

awareness, while removing stigma associated among those seeking STD testing and treatment.

This can be done by providing routine extra genital screening at Health Department Clinics for

prevention and control. Testing of all three anatomical sites can reduce opportunities for missed

infections, as fewer cases of CT/GC are more likely to be reported if urethral testing only.

Further research is still needed on determining the amount of cases who receive treatment and

follow through on CT/GC infection.

References

(1)   Centers for Disease Control and Prevention. 2015 Sexually Transmitted Disease Treatment

Guidelines: Screening Recommendations and Considerations Referenced in Treatment

Guidelines and Original Sources, https://www.cdc.gov/std/tg2015/screening-

recommendations.htm (2016, accessed 7 September 2018).

(2)   Peters RP, Verweij SP, Morré SA, et al. Evaluation of sexual history-based screening of

anatomic sites for chlamydia trachomatis and neisseria gonorrhoeae infection in men having

sex with men in routine practice. BCM Infec Dis. 2011, 11: 203.

(3)   McLaughlin J and Castrodale L. *Gonococcal Infection Update ─ Alaska, 2015.* Report for

State of Epidemiology, Department of Health and Social Services. Bulletin No. 15, 26 May,

2016.

(4)   Long Beach Department of Health and Human Services. STD/HIV Monitoring Report:

2016 Annual Report. Report for the Long Beach Department of Health and Human Services.

2016.

(5)   Centers for Disease Control and Prevention. Sexually Transmitted Diseases: What are

STDs, https://www.cdc.gov/std/ (2018, accessed 7 September 2018).

(6)   California Department of Public Health. Sexually Transmitted Diseases Control Branch –

Gonorrhea, https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/Gonorrhea.aspx (2018,

accessed 7 September 2018)

(7)   California Department of Public Health. Sexually Transmitted Diseases Control Branch –

Chlamydia, https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/Chlamydia.aspx (2018,

accessed 7 September 2017).

(8)   Centers for Disease Control and Prevention. STD & HIV Screening Recommendations,

https://www.cdc.gov/std/prevention/screeningreccs.htm (2017, accessed 7 September 2018).

(9)   National Coalition of STD Directors. MSM Sexual Health Standards of Care: Addressing

the Sexual Health Crisis among Gay, Bisexual, and Other Men who have Sex with Men

(MSM). Report for the National Coalition of STD Directors. 2017.

(10)   Callander D, Bourne C, Pell C, et al. Recording the sexual orientation of male patients

attending general practice. *Family Practice* 2015, 32:1, 35–40.

(11)   Kent CK, Chaw JK, Wong W, Liska S, Gibson S, Hubbard G, Klausner JD. Prevalence

of Rectal, Urethral, and Pharyngeal Chlamydia and Gonorrhea Detected in 2 Clinical Settings

Among Men who have Sex with Men. Clinical Infectious Disease 2005, 41:1, 67-74.

(12)   Centers for Disease Control and Prevention. STDs in Men Who Have Sex with Men.

Public Health Impact, https://www.cdc.gov/std/stats16/msm.htm (2017, accessed 7

September 2018).

(13)   California Department of Public Health. California STD Treatment Guidelines Table for

Adults & Adolescents 2015,

https://www.cdph.ca.gov/Programs/CID/DCDC/CDPH%20Document%20Library/STD-

Treatment-Guidelines-Color.pdf (2015, accessed 7 September 2018).

(14)   Centers for Disease Control and Prevention. STD Screening,

https://www.cdc.gov/std/treatment/2010/qanda/screening.htm (2012, accessed 7 September

2018).

(15)   California Department of Public Health. Sexually Transmitted Diseases in California:

2017 Snapshot,

https://www.cdph.ca.gov/Programs/CID/DCDC/CDPH%20Document%20Library/STDs-CA-2017Snapshot.pdf (2017, accessed 7 September 2018).

(16)	Sultan B, White JA, Fish R, Carrick G, Brima N, Copas A, Benn P, et al. The "3 in 1" Study: Pooling Self-Taken Pharyngeal, Urethral, and Rectal Samples into a Single Sample for Analysis for Detection of Neisseria gonorrhoeae and Chlamydia trachomatis in Men Who Have Sex with Men. *Journal of Clinical Microbiology* 2016, *54*:3, 650–656.

(17)	Sullvian PS, Peterson J, Sanchez TH, et al. Understanding Racial HIV/STI Disparities in Black and White Men Who Have Sex with Men: A Multilevel Approach. PLoS ONE 2014, 9:3, e90514.

(18)	Trebach JD, Chaulk CP, Page KR, Tuddenham S, and Ghanem KG. *Neisseria gonorrhoeae* and *Chlamydia trachomatis* among Women Reporting Extragenital Exposures. *Sexually Transmitted Diseases* 2015, *42*:5, 233–239.

(19)	Chan PA, Robinette A, Montgomery M, Almonte A, Cu-Uvin S, Lonks JR, Hardy EJ, et al. Extragenital Infections Caused by *Chlamydia trachomatis* and *Neisseria gonorrhoeae*: A Review of the Literature. *Infectious Diseases in Obstetrics and Gynecology 2016*, 17.

(20)	Sandkovsky U, O'Neill J, Sayles H, et al. High Rates of Asymptomatic Chlamydia and Gonorrhea Infection Among HIV Infected MSM. In: *ID Week conference on HIV Prevention, Early Diagnosis, STDs*, San Diego: California, 2015 October 8, Abstract No. 120.

(21)	Centers for Disease Control and Prevention. Screening Tests To Detect *Chlamydia trachomatis* and *Neisseria gonorrhoeae* Infections - 2002. *Morbidity and Mortality Weekly Report*, https://www.cdc.gov/mmwr/preview/mmwrhtml/rr5115a1.htm (2002, accessed 7 September 2018).

(22)   Bernstein K, Marcus J, Nieri G, et al. Rectal Gonorrhea and Chlamydia Reinfection Is

Associated with Increased Risk of HIV Seroconversion. JAIDS Journal of Acquired Immune

Deficiency Syndromes 2009; 53(4):537-43.

(23)   Patton ME, Kidd S, Llata E, Stenger M, Braxton J, Asbel L, Weinstock H, et al.

Extragenital Gonorrhea and Chlamydia Testing and Infection Among Men Who Have Sex

with Men—STD Surveillance Network, United States, 2010–2012. *Clinical Infectious*

Diseases 2014, 58:11, 1564–1570.

(24)   Hulley SB, Cummings SR, Browner WS, Grady D, Newman TB. Designing Clinical

Research: An Epidemiologic Approach. 4th ed. Philadelphia, PA: Lippincott Williams &

Wilkins, 2013, Appendix 6A, pg. 73.

(25)   Kustec T, Keše D, and Klavs I. Under-reporting of sexually transmitted infection with

chlamydia trachomatis - a revision of surveillance system is required. *Slovenian Journal of*

Public Health 2016, *55*:3, 174–178.

331