Social relationships, social isolation, and the human gut microbiota

Short title: Social relationships and the gut microbiota

Kimberly A. Dill-McFarland[1,2†#], Zheng-Zheng Tang[3,4#], Julia H. Kemis[1], Robert L. Kerby[1],

Guanhua Chen[3], Alberto Palloni[5], Thomas Sorenson[1], Federico E. Rey[1*], Pamela Herd[2,5*]

[1]Department of Bacteriology, U. of Wisconsin-Madison, 1550 Linden Drive, Madison, WI, USA

53706;

[2]Center for the Demography of Health and Aging, 1180 Observatory Drive, Madison, WI, USA,

53706;

[3]Department of Biostatistics and Medical Informatics, U. of Wisconsin-Madison, 600 Highland

Avenue, Madison, WI, USA, 53792

[4]Wisconsin Institute for Discovery, 330 N Orchard St, Madison, WI, USA, 53715

[5]Department of Sociology, U. of Wisconsin-Madison, 1180 Observatory Drive, Madison, WI,

USA, 53706

[†]Current address: Department of Microbiology and Immunology, U. of British Columbia, 2350

Health Sciences Mall, Vancouver, BC, Canada, V6T 1Z3

# Equal contribution

*F. E. Rey

5157 Microbial Sciences, 1550 Linden Drive, Madison, WI, USA 53706

24 608-890-2046

25 ferey@wisc.edu

26 *P. Herd

27 3454 Social Science, 1180 Observatory Drive, Madison, WI, USA, 53706

28 608-262-9451

29 pherd@lafollette.wisc.edu

ABSTRACT

Social relationships shape human health and mortality via behavioral, psychosocial, and

physiological mechanisms, including inflammatory and immune responses. Though not tested in

human studies, recent primate studies indicate that the gut microbiome may also be a biological

mechanism linking relationships to health. Integrating microbiota data into the 60-year-old

Wisconsin Longitudinal Study, we found that socialness with family and friends is associated with

differences in the human fecal microbiota. Analysis of spouse (N = 94) and sibling pairs (N = 83)

further revealed that spouses have more similar microbiota and more bacterial taxa in common

than siblings, with no observed differences between sibling and unrelated pairs. These differences

held even after accounting for dietary factors. The differences between unrelated individuals and

married couples was driven entirely by couples who reported close relationships; there were no

differences in similarity between couples reporting somewhat close relationships and unrelated

individuals. Moreover, the microbiota of married individuals, compared to those living alone, has

greater diversity and richness, with the greatest diversity among couples reporting close

relationships, which is notable given decades of research documenting the health benefits of

marriage. These results suggest that human interactions, especially sustained, close marital

relationships, influence the gut microbiota.

INTRODUCTION

Social relationships exert a sustained influence on human health and mortality with social isolation having strong negative consequences and high levels of social integration far exceeding the protective effects on mortality of individual level behaviors such as smoking cessation or maintaining a normal weight [1,2]. Research in the social sciences has shown that individuals who cohabitate in marriage and marital like relationships have better health than do unpartnered adults[3]. For both social relationships generally, and marriage specifically, health benefits are largely achieved in the context of high-quality relationships. The robust links between these relationships and health are related to stress, behaviors, and psychosocial resources, among other factors [2]. In part, social support may impact one's health by reinforcing healthy habits, reducing the impacts of stress, and preventing the use of unhealthy "self-medications" like smoking and drinking [2]. Additional research points to stress-related biological processes that may also contribute to the positive impacts of social relationships through changes in inflammatory processes, metabolic syndrome, and neurological functioning [4,5].

Recent work in the field of microbiology points to another possible biological mechanism linking human relationships and health: the microbiome. The microbial communities that inhabit mammals have profound effects on biology and health [6]. Gastrointestinal (GI) microbial communities impact host health by modulating the epigenome [7], brain function [8], and metabolism of drugs and nutrients [9] as well as impacting immune system function [10] and development [11]. While the microbiota reaches an adult-like configuration by three to five years of age [12], considerable variation exists between adults [13], and differences are mediated by a number of factors. Most notable among these are diet [14] and host genetics [15], which also correlate with health. An individual's microbiota structure (*i.e.* relative abundance) and composition (*i.e.* who's there) can

change rapidly in response to inputs like diet [16] and antibiotics [17]. Nonetheless, there is evidence

that an individual's microbiota remains relatively stable over many years [18–20], perhaps in part

because a person's behaviors also tend to be consistent over many years.

While a number of factors like diet are known to impact both the microbiota and health [21],

less is known regarding social relationships. Most existing research has focused on animal models,

which has produced compelling evidence that social interactions, via a range of different types of

physical contact, influences the gut microbiota through microbial sharing between individuals [22–

26]. Additionally, states of isolation, such as maternal neglect, influence the gut microbial

composition in animal models [27] at least in part through stress [28,29]. Thus, the gut microbiota may

play a role in some of the long-term health effects of social relationships.

But despite this tantalizing evidence, studies in human populations remain relatively small

in number [30]. There are a few studies exploring how mother-infant interactions influence the

development of the infant's gut microbiome and even how broader social interactions influence

the milk microbiome [31,32]. In terms of adults, there is evidence regarding the influence of

cohabitation, may influence the gut microbiome. A few recent studies have found that individuals

living together had more similar gut [33] and skin [33,34] microbiota. Interestingly, however, another

study found that married cohabitating couples had no more similarity in the composition of their

gut microbiota than did unrelated individuals [35].

Thus, while it does appear that living together may influence the gut microbiome, human

studies have not investigated how adult relationships, rather than just simply living in the same

space, may influence the gut microbiome. The quality of the relationship may matter. Closer

relationships likely lead to even closer shared environments, via mechanisms such as time spent

physically together. Indeed, one recent study of wild baboons found that close partners within

social groups had more similar gut microbiotas [36]. Studies have also have not more generally

compared how living alone versus living with an intimate partner influences the gut microbiome;

individuals living alone are on average, de facto, more socially isolated than those living with

someone, and animal studies have generally shown that social isolation leads to decreased

microbial diversity [22,37–39]. Though causality is not certain, decreased microbial diversity is

associated with obesity, cardiac disease, and type 2 diabetes, and a range of other inflammatory

disorders [40–47]. More broadly, there is extensive evidence that cohabitating couples in later life

have substantially improved physical and psychological well-being compared to single adults [48–

50]. Thus, similar mechanisms might explain some of the variance in findings in humans.

An important hindrance to research examining social relationships and the GI microbiota

is the availability of human samples with sufficiently well-characterized life course measures of

broader social environments and conditions. Thus, most microbiological research in this field is

based on animal models [22–25]. However, there are now a wide array of well-characterized

longitudinal studies in the social sciences that have generated decades of research documenting

relationships between broader social environments and mortality [5,51–54]. These data can provide a

platform for studies of the human microbiota to advance knowledge for both social scientists and

microbiologists, including whether social conditions influence the gut microbiota and whether the

gut microbiota is a mediating biological mechanism explaining how social conditions influence

health.

Here, we leverage a multidisciplinary collaboration to investigate the links between human

interaction, the microbiota, and human health. We utilized data in the nearly 60-year Wisconsin

Longitudinal Study (WLS) [54], which constitutes a random sample of 1 in 3 1957 Wisconsin high

school graduates (N = 10,317), as well as selected spouses and siblings surveyed periodically

during their adult life. We correlate the fecal microbiota of 408 older individuals (58 – 91 yo) from

WLS with extensive health and behavioral data, as well as compare spouse and sibling pairs within

the cohort. Overall, this project demonstrates the promise of joint participation between social

scientists and microbiologists in efforts to more fully understand the gut microbiota and its impacts

on human health.

RESULTS and DISCUSSION

We employed 16S rRNA gene sequencing to characterize the fecal microbiota of 408 individuals,

including Wisconsin Longitudinal Study (WLS) graduates (N = 179, 76 ± 0.5 years old), siblings

of graduates (134, 74 ± 6.4), spouses of graduates (63, 76 ± 3.7), and spouses of siblings (32, 73 ±

6.1). We then correlated these communities to longitudinal survey data collected from 1957 to

2015 as part of WLS [54]. For more details on this data collection, see [55]. A total of 24.5 million

high-quality sequences were obtained for 408 fecal samples (60,000 ± 19,000 SD sequences per

sample) after quality filtering in mothur. All samples achieved sufficient coverage as determined

by Good's coverage > 99% (Dataset S1).

In the WLS graduate cohort, we identified several factors correlated with gastrointestinal

(GI) microbiota including sex, antibiotics, dietary protein, high blood sugar, and heart disease (Fig.

1, Fig. S1, Table S1). These factors were reported in the previous literature [57,58] with diet playing

a particularly strong role [14,16,56]. Thus, we assessed diet across a number of measures including

habitual intake of protein, vegetables, and fruits (Text S1) during the year prior to the fecal sample

collection (for details, see METHODS, Statistical analysis for graduates). While overall dietary

dissimilarity (Bray-Curtis and Jaccard) across these three categories correlated with gut microbiota

dissimilarity, only the total frequency of dietary protein consumption was robustly associated with

microbial composition using either univariate or multivariate analyses (Table S1). Thus, we note

that all analyses have adjusted for potential confounders including age, sex, antibiotics, dietary

protein, and chronic conditions (diabetes and heart disease) unless stated otherwise. In some

analyses –that are noted below—we do also include vegetable and fruit dietary data.

Social interactions and the human fecal microbiota Human interactions were also associated

with differences in gut microbiota and diversity. Specifically, we found that individuals that were

cohabitating with a spouse or partner had more similar microbiota composition with their

cohabitating spouse/partner as well as higher diversity and richness than unmarried, non-

cohabitating individuals (unweighted UniFrac $P = 0.029$ Shannon $P = 0.005$, Chao $P = 0.011$, Fig.

2). Since all cohabitating pairs were male-female and sex was a strong determinant of the

microbiota in this study ($P < 0.001$, Table S2), increased diversity may be partially due to sustained

exchange of microorganisms between the sexes, though we were not able to test this given that

there were no same sex couples in these data. Increases in diversity seen here are consistent with

a previous cohabitation study in pigs [59] and may have implications for human health, as previous

work indicates that increased gut microbial diversity is associated with lower risks of irritable

bowel syndrome (IBS), Crohn's disease, ulcerative colitis, and other GI afflictions [60].

Social interactions with relatives and friends were stronger predictors of gut microbial

diversity in non-cohabiting individuals than cohabiting spouses/partners (unweighted UniFrac $P =$

0.0030, Shannon $P = 0.042$, Chao $P = 0.063$, Fig. S2) (Table S2). Here, social interactions were

defined as the sum of "How many times during the past four weeks have you gotten together with

relatives/friends?" The associations may have been weaker for cohabitating spouses due to their

higher microbial diversity; ecological theory supports that diverse communities are more resilient

and resistant to invasion by new species [61]. Thus, one explanation for these differential associations

is that the more diverse microbiotas of individuals already cohabiting with a spouse may not

have been as strongly influenced by increasing social interactions while the less diverse

microbiotas of those living alone were more strongly influenced by invasion of new species

through social exposures. It is also possible that cohabiting couples share the same friends and

socialize together with these friends. However, factors contributing to the resilience of the human

gut microbiota require further exploration to confirm this hypothesis.

Spouses have more similar microbes than siblings and unrelated individuals Previous studies

have established that the GI microbiota reaches an adult-like configuration by 3 to 5 years of age

[18,62,63] and that during adulthood, communities are stable on the time scale of years [19,20]. Thus,

microbial communities established in early life may persist and, aside from extreme perturbation,

remain stable across one's adult lifetime. However, our analyses comparing sibling, couple, and

unrelated pairs challenge the assumption that microbial communities established in early life will

be largely unperturbed in later life (for details, see METHODS, Statistical Analysis for spouse and

siblings). In fact, we find no evidence for a remaining influence of early life on the composition

of the gut microbiota among older adults. In this older cohort, spouses were more similar than

unrelated subjects (unweighted UniFrac P = 3.2E-5) or sibling pairs (unweighted UniFrac P =

0.033, Fig. 3). Further, the length of the cohabiting marital relationship was positively correlated

with similarity (unweighted UniFrac, P = 0.031)   In contrast, siblings were no more similar than

unrelated pairs by any beta-diversity metric (P > 0.3, Fig. 3A, Fig. S3A, D, G) (Table S2). We also

found no evidence that the physical proximity of siblings—as measured by physical distance

between siblings—influenced gut microbial similarity. Thus, adult factors like marriage with

cohabitation (spouses) appear to have a greater influence on the adult gut microbiota than early-

life environment or genetics (siblings).

This is further supported by our findings that childhood farm status was not associated with

microbial richness (Chao $P = 0.342$) while working on a farm as an adult correlated with higher

richness (Chao $P = 0.005$). Farm-driven differences in the microbiota are of particular interest,

because adolescents that grew up on a farm have more diverse microbial communities [64] and

reduced risk of asthma and other atopic diseases both during childhood [65] and as adults [66]. Given

the results here, it appears that the microbially-driven protective effects of early farm exposures

are not due to the persistence of protective microorganisms acquired in early-life. Protection may,

instead, be conferred by immune development and training by early-life microbes as suggested

previously [67].

Our results are also in contrast with previous work showing that genetically related

individuals harbor more similar microbial communities than unrelated individuals, regardless of

current cohabitation [35,68–70]. However, these previous studies investigated children [68], young adults

[35,69], or a wide age range [70], and therefore, cumulative changes across a lifetime may not have

reached a level sufficient to overcome early-life factors impacting the microbiota. Additionally,

sibling pairs in other studies were twins [35,68–70], and many focused on monozygotic twins (same

sex and age) [35,69,70] as opposed to this study where siblings were often of opposite sexes (43%) and

ranged from less than a year to 18 years apart in age. Also, the unrelated group in this study may

have exhibited higher homogeneity than unrelated groups in other studies, because most grew up

in and/or currently live in the state of Wisconsin. Thus, compared to previous studies, siblings

were likely less similar and unrelated pairs more similar across our cohort. Furthermore, genetic

effects on the microbiota are often small [70] and detection may require a larger human cohort than

used here. Taken together, these factors may have contributed to the lack of significant differences

observed between sibling and unrelated groups even though average sibling beta-diversity was

intermediate between spouses and unrelated individuals.

Increased microbial similarity, diversity, and richness in closer relationships For both spouse

and sibling relationships, microbiota similarity was associated with self-reported relationship

closeness (unweighted UniFrac $P = 0.0079$). Closeness was measured by participant responses to

"How close are you and your current spouse/sibling?" on a scale of not at all (1) to very (4). Due

to the small sample sizes in the categories "Not very" (N=13) and "Not at all" (N=4), we combined

these two groups into "Not" close. Across spouses and siblings, individuals in very close

relationships harbored gut microbial communities more similar to their close social partners than

those in not very close relationships (Fig. 3B), though this relationship was not significant within

the spousal and sibling pair groups separately (Fig. 3C). Moreover, differences between spouses

and unrelated individuals, in terms of closeness (Fig. 2), as well as the enhanced diversity and

richness in cohabitating couples versus individuals living alone (Fig. 2) were driven by spouses

reporting very close relationships. This was in contrast to couples reporting only somewhat close

relationships as these pairs did not have higher gut microbiota similarity than unrelated pairs

(Table S3) nor did they display microbial diversity or richness different from non-cohabitating

individuals (Table S3). Importantly, the apparent impacts of relationship closeness do not appear

to be mediated by similarities in diet since overall dietary dissimilarity (Bray-Curtis and Jaccard)

did not significantly differ according to relationship closeness (ANOVA $P > 0.5$; Table S4). We

note that these included sensitivity tests that modeled diet based on the protein consumption, but

also overall diet that captured vegetable and fruit consumption.

While diet is often correlated with the GI microbiota [56], closeness points to the less well-

understood contributions of human interactions and shared behaviors. Close proximity and

frequent physical contact were correlated with microbiota similarity among primates with direct

microbial sharing between individuals contributing to similarity [22,23]. In this study, relationship

closeness may represent a summative measure of time spend together, physical affection, and other

human interactions with the potential to result in microbial sharing. Indeed, there is evidence that

the salivary microbiome influences the gut microbiome and the salivary microbiome may be

influenced by kissing [71,72]. In these data, this is supported by the fact that spouses had more

operational taxonomic units (OTUs, a proxy for microbial species) in common ($30.4 \pm 7.32\%$)

than siblings ($26.4 \pm 7.47\%$, t-test $P = 4.39E\text{-}04$) (Dataset S2). Also, when comparing the spouse

and sibling pair within a family represented in this dataset, a person tended to have more OTUs in

common with his or her very close spouse ($25.4 \pm 7.9\%$) than his or her very close sibling ($22.2 \pm$

6.4%, N = 12 families, $P = 0.074$, Fig. 3D). This is also true when comparing very close spouses

($22.9 \pm 5.8\%$) and somewhat close siblings within a family ($20.6 \pm 5.5\%$, N = 17 families, $P =$

0.027, Fig. 3E).

Shared taxa with close human relationships. In general, highly abundant genera and OTUs were

shared between many spouse and sibling pairs while less abundant shared taxa were specific to

one pair type and shared by a small number of pairs within that type (Dataset S3). OTUs that were

commonly found among spouses or siblings ($> 50\%$ of pairs) but rare in the unrelated dataset ($<$

70% individuals, $< 49\%$ unrelated pairs) may represent bacterial species easily shared by close

human interaction. These OTUs were predominately from the phylum Firmicutes (16 of 22 OTUs)

with representatives of families Lachnospiraceae and Ruminococcaceae (Dataset S4).

Interestingly, most of these potentially shared OTUs were from strictly anaerobic taxa, indicating

that persisting in an oxygen-rich environment in-between hosts may not be a limiting factor in very

close human relationships. Transmission, in these cases, could be mediated by direct contact

similar to mechanisms of vertical transmission from mother to child [73].

Taxa commonly associated with reduced disease incidence or severity like *Akkermansia*

muciniphila [74], *Bifidobacterium* spp. [75,76], *Collinsella aerofaciens* [76], and *Ruminococcus bromii* [77]

as well as potentially harmful taxa like *Clostridium spiroforme* [78,79] were often present in both

persons in a spouse or sibling pair. Several of these potentially shared OTUs were associated with

disease incidence in the larger dataset. In particular, *Ruminococcus bromii*, *Lachnospira* spp. and

unclassified Ruminococcaceae and Lachnospiraceae OTUs were less abundant in those with high

blood sugar (Fig. 4, Dataset S4). These results are in contrast to previous reports of more abundant

Ruminococcaceae/*Ruminococcus* [80,81] and Lachnospiraceae [80] associated with diabetes in humans

and may point to important differences in the impacts of the microbiota on metabolic health in

older populations. Overall, though, this indicates that GI microbial species with the potential to

impact host health may be shared by close human interactions. However, it cannot be discounted

that these apparent health associations may be mediated by diet as those with high blood sugar

often consume specific diets to manage disease.

Overall, our findings indicate that in order to understand environmental influences on the

gut microbioata, we must now consider the many microbiotas with which this individual interacts.

Socialness with family and friends is associated with differences in the fecal microbiota. These

differences held even after accounting for dietary factors, though given this is the first study of its

kind, it will be critical for future work to validate this finding. Thus, it is possible that relationships

with others may influence the gut microbiota and consequent health outcomes, either through

direct microbial transfer or reinforcement of healthy microbiota behaviors. We further found not

only that married couples had more similar gut microbiota but also that the microbiota of married

individuals, compared to those living alone, has greater diversity and richness. Key to both of these

findings, however, was that they were driven by individuals reporting that they were very close to

their spouse as opposed to somewhat close. Close marriage relationships had a stronger influence

than the shared genetic factors and early life environments among siblings. This finding is

interesting, in part, because it parallels an extensive body of evidence demonstrating robust links

between high quality marriages and morbidity and mortality. Future work could attempt to

disentangle the mechanisms linking close relationships to microbial composition. For example,

while we did not find evidence that shared diet was primarily responsible for these findings, we

could not test precise frequencies of physical contact and intimacy as an alternative explanatory

mechanism. Importantly, the types of physical contact and intimacy change over the life course,

with sexual intimacy becoming far less frequent in later life, but other kinds of intimate physical

contact remaining important. Regardless of the mediating mechanism, from a social and

population health science perspective, decades of evidence that social relationships, especially

close ones like marriage, influence morbidity and mortality make the central finding of significant

interest. For example, even if future work finds a greater role for shared diets, it is still the social

relationships that drive that shared diet. Overall, these results provide support for the gut

microbiome as a possible mediating pathway between social relationships, especially marriage,

and health and mortality. These findings, in the context of the robust body of evidence linking

social relationships to human morbidity and mortality, provide fodder for further work examining

the role of the gut microbiome as a possible biological mediator in these relationships [32]. Further

microbiota work across time in a more diverse population should be undertaken with the many

longitudinal social science studies currently underway in an effort to increase our understanding

of the complex interactions between human behavior, the microbiota, and health.

METHODS

Wisconsin Longitudinal Study (WLS) WLS is based on a one-third sample of all 1957

Wisconsin high school graduates (N = 10,317) as well as selected siblings and spouses [54].

Graduates originally enrolled with an in-person questionnaire upon graduating high school in

1957, which was followed by data collection in 1964, 1975, 1992, 2004, and 2011. Siblings were

surveyed in 1977, 1994, 2005, and 2011; spouses were surveyed in 2004 or 2006. The content of

WLS surveys changed to reflect the participants' life course with an education focus in the initial

data collection, familial and career outcomes in young adulthood / midlife, and health, cognitive

functioning, psychological well-being, non-work activities, caregiving, bereavement, social

support, and end-of-life preparations in later rounds. WLS data collection was approved by the

Institutional Review Board (IRB) at the University of Wisconsin-Madison (2014-1066, 2015-

0955). Informed consent, the content and procedures of which were included in the IRB approval,

was obtained from participants. All methods were performed in accordance with relevant

guidelines and regulations.

Study design A total of 500 individuals were randomly drawn from the full WLS dataset

constrained based on the following: 1) participated in the 2011 interviews; 2) lived in one of 10

counties in Wisconsin that included both northern rural counties and southern more urban counties;

and 3) were part of a sibling pair. Individuals were removed from the study if they did not give

consent, their sample did not arrive for processing chilled, but not frozen, within 48 hrs of

collection, or their sample did not yield at least 10,000 sequences for analysis. This resulted in 408

individuals being included in this study.

An additional survey was administered at the time of fecal sampling, which detailed dietary

data from the prior three days, prescription/antibiotic use, current living situation, and additional

health information. This as well as selected data from the larger WLS study focused on health,

spouse/sibling relationships, and social interactions were used in this study (Text S1). Data,

documentation, and other materials are accessible at http://www.ssc.wisc.edu/wlsresearch/.

Access to the full dataset can be obtained through wls@ssc.wisc.edu.

Sample collection Stool samples were collected by participants in November 2014, January 2015,

or April 2015 following provided instructions (Text S2). Participants stored samples at ~4 °C in

their refrigerator or in a NanoCool box (Albuquerque, NM) with cooling cartridge and customized

foam insert, supplemented with a single ice pack. Interviewers picked-up samples from

participants within 24 hours of collection and shipped samples in fresh NanoCool boxes for arrival

at UW-Madison within 48 hours of collection. Upon arrival, an aliquot of feces was collected for

DNA extraction and immediately stored at -80°C until further processing. The use of WLS and

fecal microbiota data were approved by the Institutional Review Board at the University of

Wisconsin-Madison (2017-0600).

DNA extraction Genomic DNA was extracted from fecal aliquots using a bead-beating protocol

[45]. Briefly, feces (~100 mg) were re-suspended in a solution containing 500 μl of extraction buffer

[200 mM Tris (pH 8.0), 200 mM NaCL, 20 mM EDTA], 210 μl of 20% SDS, 500 μl

phenol:chloroform:isoamyl alcohol (pH 7.9, 25:24:1) and 500 μl of 0.1-mm diameter

zirconia/silica beads. Samples were mechanically disrupted using a bead beater (BioSpec Products,

Barlesville, OK; maximum setting for 3 min at room temperature), followed by centrifugation,

recovery of the aqueous phase, and precipitation with isopropanol. QIAquick 96-well PCR

Purification Kit (Qiagen, Germantown, MD) was used to remove contaminants. Isolated DNA was

eluted in 5 mM Tris/HCl (pH 8.5) and was stored at -80 °C until further use. We also note that we

used negative controls.

Sequencing PCR was performed using universal primers flanking the variable 4 (V4) region of

the bacterial 16S rRNA gene [82]. We used negative controls for each PCR reaction. PCR reactions

where the negative control yielded a product were not sequenced until the problem was solved.

Samples were processed all together, not in batches, in a random order (i.e., not clustered by

family). Additionally, unlike other specimens (e.g., saliva, skin), DNA contamination from

reagents is in general not a problem for fecal samples given the high DNA content of the sample

(10^{12} microbes/g of feces). In one reaction per sample, 10 - 50 ng DNA, 10 μM each primer, 12.5

μl 2X HotStart ReadyMix (KAPA Biosystems, Wilmington, MA, USA), and water to 25 μl were

used. Cycling conditions were initial denaturation of 95 °C for 3 min followed by 25 cycles of 95

°C for 30 s, 55 °C for 30 s, and 72 °C for 30 s, with a final extension of 72 °C for 5 min. PCR

products were purified with the QIAquick 96-well PCR Purification Kit (Qiagen, Germantown,

MD, USA). Samples were quantified by Qubit Fluorometer (Invitrogen, Carlsbad, CA, USA) and

equimolar pooled. The pool plus 5% PhiX control DNA was sequenced through the U. of

Wisconsin-Madison Biotechnology Center with the MiSeq 2x250 v2 kit (Illumina, San Diego, CA,

USA) using custom sequencing primers [82]. All DNA sequences are available upon institutional

review board (IRB) or other ethics board approval through wls@ssc.wisc.edu.

Sequence clean-up All sequences were demultiplexed on the Illumina MiSeq. Sequence clean-

up and processing was performed with mothur v.1.36.1 [83] following a protocol similar to [82].

Briefly, paired-end sequences were combined into contigs with default parameters (match bonus

= 1, mismatch penalty = -1, gap penalty = -2, gap extend penalty = -1, insert quality ≥ 20,

mismatch quality difference ≥ 6). Poor-quality sequences, including those with ambiguous base

pairs, homopolymers greater than 8, or outside 200 – 500 bp in length, were discarded.

Sequences were then aligned to the SILVA 16S rRNA gene reference alignment database [84] and

trimmed to the V4 region. To reduce sequencing error, sequences with 2 or fewer differences

were pre-clustered. Chimera detection and removal were performed using UCHIME [85]. Final

sequences were then classified to the GreenGenes database [86]. Singletons were removed to

facilitate downstream analyses. All sequences were grouped into 98% operational taxonomic

units (OTUs) by uncorrected pairwise distances and average neighbor clustering in mothur.

Clustering performed on uncorrected pairwise distances revealed no differences in clusters at 97

vs 98% similarity. Therefore, the stricter cutoff was reported Coverage was assessed by Good's

coverage, and then samples were normalized to whole number counts by percent

relative abundance to approximately 10,000 sequences per sample (9,914 - 10,061 after

rounding.

Statistical analysis for graduates Graduates were assessed separately from siblings and spouses

to avoid potential interactions, and the graduate subset was not significantly different from other

groups (PERMANOVA *P* Bray-Curtis *P* = 0.56, Jaccard *P* = 0.57, weighted UniFrac *P* = 0.33,

unweighted UniFrac *P* = 0.24). Alpha-diversity was assessed with Shannon's diversity and Chao's

richness calculated in mothur. Differences in alpha-metrics were assessed in R v3.3.2 [87] by linear

regression with the Benjamini-Hochberg correction for multiple comparisons across each metric.

Microbial beta-diversity was assessed for Bray-Curtis, Jaccard, weighted, and unweighted

UniFrac metrics with results shown for unweighted UniFrac unless otherwise noted. Dietary beta-

diversity was assessed for Bray-Curtis and Jaccard metrics as well as corresponding nMDS axes

calculated from habitual intake of specific sources of protein (N = 4), vegetables (N = 76), and

fruits (N = 24) expressed as times consumed per week (protein), proportions of total types (all),

and presence/absence of individual types (all). Differences in beta-diversity were tested with

permutational analysis of variance (PERMANOVA, adonis) in the vegan package [88] with the

Benjamini-Hochberg correction for multiple comparisons across each metric and a maximum of

5000 permutations. All variables were modeled using independent, univariate tests and dietary

variables were additionally modeled using multivariate tests of all components (protein,

vegetables, fruits). Co-variance of microbial and dietary beta metrics was measured using Mantel's

test. The factors that associated with the microbiome in univariate models (*i.e.* age, sex, antibiotics,

dietary protein, high blood sugar, and heart disease [57]) were adjusted for in regression models as

potential confounders. Beta-diversity was visualized by non-metric multidimensional scaling

(nMDS) plots with arrows from significant variables (PERMANOVA) fitted to the ordination

using maximum correlation (envfit, vegan). All tests were assessed at significance $P < 0.05$ and

trends $0.05 < P < 0.1$.

Statistical analysis for spouses and siblings

For the spouse and sibling similarity analysis, the unit of the observation is the pair (*i.e.* spouse,

sibling, or unrelated pair defined below) and the variables used in the analysis are distance in

individual measurements between the two members of the pair. Specifically, beta-diversity metrics

were used to quantify the distance in microbial and overall diet whereas absolute difference were

calculated to quantify the distance in all the other variables (*e.g.* age, sex, dietary protein). We

sampled unrelated pairs from the data in order to compare the spouse or sibling pair with unrelated

pairs. In particular, the unrelated individuals cannot be siblings, spouses, or in-laws, and each

unrelated pair will match the corresponding spouse or sibling pair in sex and antibiotics usage.

Beta-diversity distances were compared among spouse, sibling, and unrelated pairs using linear

regression while adjusting for the distance in age, sex, dietary protein, health conditions (if

available). P-values were averaged across 1000 rounds of unrelated pair sampling. For closeness

analysis, we removed age and sex from the model because the two variables are highly correlated

with pair type (*i.e.* sibling/spouse pair can be accurately classified using the difference of the age

or sex between the two members of the pair). For comparing OTU sharing among spouse and

sibling within a family, we used mixed-effect models to account for family clustering. All tests

were assessed at significance $P < 0.05$ and trends $0.05 < P < 0.1$.

REFERENCES

1.   House, J. S., Landis, K. R. & Umberson, D. Social relationships and health. *Science (80-.).* **241,** 540–545 (1988).
2.   Umberson, D., Crosnoe, R. & Reczek, C. Social relationships and health behavior across life course. *Annu. Rev. Sociol.* **36,** 139–157 (2010).
3.   Robles, T. F. & Kiecolt-Glaser, J. K. The physiology of marriage: pathways to health. *Physiol. Behav.* **79,** 409–416 (2003).
4.   McEwen, B. S. Brain on stress: How the social environment gets under the skin. *Proc. Natl. Acad. Sci. U. S. A.* **109,** 17180–17185 (2012).
5.   Yang, Y. C. *et al.* Social relationships and physiological determinants of longevity across the human life span. *Proc. Natl. Acad. Sci. U. S. A.* **113,** 578–583 (2016).
6.   Grice, E. A. & Segre, J. A. The human microbiome: Our second genome. *Annu. Rev. Genomics Hum. Genet.* **13,** 151–170 (2012).
7.   Krautkramer, K. A. *et al.* Diet-microbiota interactions mediate global epigenetic programming in multiple host tissues. *Mol. Cell* **64,** 982–992 (2016).

8.   Rieder, R., Wisniewski, P. J., Alderman, B. L. & Campbell, S. C. Microbes and mental
health: A review. *Brain. Behav. Immun.* (2017). doi:10.1016/j.bbi.2017.01.016

9.   Flint, H. J., Bayer, E. A., Rincon, M. T., Lamed, R. & White, B. A. Polysaccharide
utilization by gut bacteria: potential for new insights from genomic analysis. *Nat Rev*
Micro **6,** 121–131 (2008).

10.   Lin, L. & Zhang, J. Role of intestinal microbiota and metabolites on gut homeostasis and
human diseases. *BMC Immunol.* **18,** 2 (2017).

11.   Gensollen, T., Iyer, S. S., Kasper, D. L. & Blumberg, R. S. How colonization by
microbiota in early life shapes the immune system. *Science* **352,** 539–544 (2016).

12.   Rodríguez, J. M. *et al.* The composition of the gut microbiota throughout life, with an
emphasis on early life. *Microb. Ecol. Health Dis.* **26,** 10.3402/mehd.v26.26050 (2015).

13.   Arumugam, M. *et al.* Enterotypes of the human gut microbiome. *Nature* **473,** 174–180
(2011).

14.   Wu, G. D. *et al.* Linking long-term dietary patterns with gut microbial enterotypes.
Science **334,** 105–108 (2011).

15.   Bonder, M. J. *et al.* The effect of host genetics on the gut microbiome. *Nat Genet* **48,**
1407–1412 (2016).

16.   David, L. A. *et al.* Diet rapidly and reproducibly alters the human gut microbiome. *Nature*
505, 559–563 (2014).

17.   Panda, S. *et al.* Short-term effect of antibiotics on human gut microbiota. *PLoS One* **9,**
e95476 (2014).

18.   Yatsunenko, T. *et al.* Human gut microbiome viewed across age and geography. *Nature*
486, 222–227 (2012).

19.   Faith, J. J. *et al.* The long-term stability of the human gut microbiota. *Science* **341,**
1237439 (2013).

20.   Costello, E. K. *et al.* Bacterial community variation in human body habitats across space
and time. *Science* **326,** 1694–1697 (2009).

21.   Flint, H. J., Scott, K. P., Louis, P. & Duncan, S. H. The role of the gut microbiota in
nutrition and health. *Nat Rev Gastroenterol Hepatol* **9,** 577–589 (2012).

22.   Moeller, A. H. *et al.* Social behavior shapes the chimpanzee pan-microbiome. *Sci. Adv.* **2,**
(2016).

23.   Amato, K. R. *et al.* Patterns in gut microbiota similarity associated with degree of sociality
among sex classes of a neotropical primate. *Microb. Ecol.* 1–9 (2017).
doi:10.1007/s00248-017-0938-6

24.   Amaral, W. Z. *et al.* Social influences on Prevotella and the gut microbiome of young
monkeys. *Psychosom. Med.* **Epub ahead,** (2017).

25.   Tung, J. *et al.* Social networks predict gut microbiome composition in wild baboons. *Elife*
4, e05224 (2015).

26.   Amaral, W. Z., Lubach, G. R. & Coe, C. L. 114. Social influences on the gut microbiome
of developing monkeys. *Brain. Behav. Immun.* **40,** (2014).

27.   Bailey, M. T. & Coe, C. L. Maternal separation disrupts the integrity of the intestinal
microflora in infant rhesus monkeys. *Dev. Psychobiol.* **35,** 146–155 (1999).

28.   O'Mahony, S. M. *et al.* Early life stress alters behavior, immunity, and microbiota in rats:
implications for irritable bowel syndrome and psychiatric illnesses. *Biol. Psychiatry* **65,**
263–267 (2009).

29.   Bailey, M. T. *et al.* Exposure to a social stressor alters the structure of the intestinal

microbiota: implications for stressor-induced immunomodulation. *Brain. Behav. Immun.*
25, 397–407 (2011).

30. Archie, E. A. & Tung, J. Social behavior and the microbiome. *Curr. Opin. Behav. Sci.* **6,**
28–34 (2015).

31. Meehan, C. L. *et al.* Social networks, cooperative breeding, and the human milk
microbiome. *Am. J. Hum. Biol.* e23131 (2018).

32. Münger, E., Montiel-Castro, A. J., Langhans, W. & Pacheco-López, G. Reciprocal
interactions between gut microbiota and host social behavior. *Front. Integr. Neurosci.* **12,**
21 (2018).

33. Song, S. J. *et al.* Cohabiting family members share microbiota with one another and with
their dogs. *Elife* **2,** e00458 (2013).

34. Ross, A. A., Doxey, A. C. & Neufeld, J. D. The skin microbiome of cohabiting couples.
mSystems **2,** (2017).

35. Erwin G. Zoetendal Wilma M. Akkermans-van Vliet, J. Arjan G. M. de Visser, Willem M.
de Vos, A. D. L. A. The host genotype affects the bacterial community in the human
gastronintestinal tract. *Microb. Ecol. Health Dis.* **13,** 129–134 (2001).

36. Grieneisen, L. E., Livermore, J., Alberts, S., Tung, J. & Archie, E. A. Group living and
male dispersal predict the core gut microbiome in wild baboons. *Integr. Comp. Biol.* **57,**
770–785 (2017).

37. Levin, I. I. *et al.* Stress response, gut microbial diversity and sexual signals correlate with
social interactions. *Biol. Lett.* **12,** 20160352 (2016).

38. Lombardo, M. P. Access to mutualistic endosymbiotic microbes: an underappreciated
benefit of group living. *Behav. Ecol. Sociobiol.* **62,** 479–497 (2008).

39. Li, H. *et al.* Pika population density is associated with the composition and diversity of gut
microbiota. *Front. Microbiol.* **7,** 758 (2016).

40. Manichanh, C. *et al.* Reduced diversity of faecal microbiota in Crohn's disease revealed
by a metagenomic approach. *Gut* **55,** 205–211 (2006).

41. Scher, J. U. *et al.* Decreased Bacterial Diversity Characterizes an Altered Gut Microbiota
in Psoriatic Arthritis and Resembles Dysbiosis of Inflammatory Bowel Disease. *Arthritis*
Rheumatol. (Hoboken, N.J.) **67,** 128–139 (2015).

42. de Goffau, M. C. *et al.* Fecal Microbiota Composition Differs Between Children With β-
Cell Autoimmunity and Those Without. *Diabetes* **62,** 1238–1244 (2013).

43. Wang, M. *et al.* Reduced diversity in the early fecal microbiota of infants with atopic
eczema. *J. Allergy Clin. Immunol.* **121,** 129–134 (2008).

44. Schippa, S. *et al.* A distinctive 'microbial signature' in celiac pediatric patients. *BMC*
Microbiol. **10,** 175 (2010).

45. Turnbaugh, P. J. *et al.* A core gut microbiome in obese and lean twins. *Nature* **457,** 480
(2008).

46. Lambeth, S. M. *et al.* Composition, Diversity and Abundance of Gut Microbiome in
Prediabetes and Type 2 Diabetes. *J. diabetes Obes.* **2,** 1–7 (2015).

47. Menni, C. *et al.* Gut microbial diversity is associated with lower arterial stiffness in
women. *Eur. Heart J.* **39,** 2390–2397 (2018).

48. Kim, J. & Waite, L. J. Relationship quality and shared activity in marital and cohabiting
dyads in the National Social Life, Health, and Aging Project, Wave 2. *Journals Gerontol.*
Ser. B Psychol. Sci. Soc. Sci. **69,** S64–S74 (2014).

49. Lewin, A. C. Health and Relationship Quality Later in Life: A Comparison of Living

Apart Together (LAT), First Marriages, Remarriages, and Cohabitation. *J. Fam. Issues* **38,**
1754–1774 (2016).
50. Wong, J. S. & Waite, L. J. Marriage, social networks, and health at older ages. *J. Popul.*
Ageing **8,** 7–25 (2015).
51. Levine, M. E., Cole, S. W., Weir, D. R. & Crimmins, E. M. Childhood and later life
stressors and increased Inflammatory gene expression at older ages. *Soc. Sci. Med.* **130,**
16–22 (2015).
52. Seeman, M., Merkin, S., Karlamangla, A., Koretz, B. & Seeman, T. Social status and
biological dysregulation: The "status syndrome" and allostatic load. *Soc. Sci. Med.* **118,**
143–151 (2014).
53. Das, A. Psychosocial distress and inflammation: Which way does causality flow? *Soc. Sci.*
Med. **170,** 1–8 (2016).
54. Herd, P., Carr, D. & Roan, C. Cohort profile: Wisconsin longitudinal study (WLS). *Int. J.*
Epidemiol. **43,** 34–41 (2014).
55. Herd, P. *et al.* The influence of social conditions across the life course on the human gut
microbiota: a pilot project with the Wisconsin Longitudinal Study. *J Gerontol B Psychol*
Sci Soc Sci (2017). doi:10.1093/geronb/gbx029
56. Scott, K. P., Gratz, S. W., Sheridan, P. O., Flint, H. J. & Duncan, S. H. The influence of
diet on the gut microbiota. *Pharmacol. Res.* **69,** 52–60 (2013).
57. Borgo, F. *et al.* Body Mass Index and sex affect diverse microbial niches within the gut.
Front. Microbiol. **9,** 213 (2018).
58. Knight, R. *et al.* The microbiome and human biology. *Annu. Rev. Genomics Hum. Genet.*
18, 65–86 (2017).
59. Pajarillo, E. A. B., Chae, J. P., Kim, H. B., Kim, I. H. & Kang, D.-K. Barcoded
pyrosequencing-based metagenomic analysis of the faecal microbiome of three purebred
pig lines after cohabitation. *Appl. Microbiol. Biotechnol.* **99,** 5647–5656 (2015).
60. Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K. & Knight, R. Diversity,
stability and resilience of the human gut microbiota. *Nature* **489,** 220–230 (2012).
61. Levine, J. M. & D'Antonio, C. M. Elton revisited: a review of evidence linking diversity
and invasibility. *Oikos* **87,** 15–26 (1999).
62. Koenig, J. E. *et al.* Succession of microbial consortia in the developing infant gut
microbiome. *Proc. Natl. Acad. Sci. U. S. A.* **108,** 4578–4585 (2011).
63. Palmer, C., Bik, E. M., DiGiulio, D. B., Relman, D. A. & Brown, P. O. Development of
the human infant intestinal microbiota. *PLoS Biol* **5,** e177 (2007).
64. Dicksved, J. *et al.* Molecular fingerprinting of the fecal microbiota of children raised
according to different lifestyles. *Appl. Environ. Microbiol.* **73,** 2284–2289 (2007).
65. Ege, M. J. *et al.* Exposure to environmental microorganisms and childhood asthma. *N.*
Engl. J. Med. **364,** 701–709 (2011).
66. House, J. S. *et al.* Early-life farm exposures and adult asthma and atopy in the Agricultural
Lung Health Study. *J. Allergy Clin. Immunol.* (2017). doi:10.1016/j.jaci.2016.09.036
67. Wlasiuk, G. & Vercelli, D. The farm effect, or: when, what and how a farming
environment protects from asthma and allergic disease. *Curr. Opin. Allergy Clin.*
Immunol. **12,** 461–466 (2012).
68. Stewart, J. A., Chadwick, V. S. & Murray, A. Investigations into the influence of host
genetics on the predominant eubacteria in the faecal microflora of children. *J. Med.*
Microbiol. **54,** 1239–1242 (2005).

69. Tims, S. *et al.* Microbiota conservation and BMI signatures in adult monozygotic twins.
ISME J. **7,** 707–717 (2013).
70. Goodrich, J. K. *et al.* Human genetics shape the gut microbiome. *Cell* **159,** 789–799
(2014).
71. Cao, X. Intestinal inflammation induced by oral bacteria. *Science (80-.).* **358,** 308 LP-309
(2017).
72. Kort, R. *et al.* Shaping the oral microbiota through intimate kissing. *Microbiome* **2,** 41
(2014).
73. Ley, R. E., Peterson, D. A. & Gordon, J. I. Ecological and evolutionary forces shaping
microbial diversity in the human intestine. *Cell* **124,** 837–848 (2017).
74. Everard, A. *et al.* Cross-talk between Akkermansia muciniphila and intestinal epithelium
controls diet-induced obesity. *Proc. Natl. Acad. Sci.* **110,** 9066–9071 (2013).
75. Arboleya, S., Watkins, C., Stanton, C. & Ross, R. P. Gut Bifidobacteria populations in
human health and aging. *Front. Microbiol.* **7,** 1204 (2016).
76. Malinen, E. *et al.* Association of symptoms with gastrointestinal microbiota in irritable
bowel syndrome. *World J. Gastroenterol.* **16,** 4532–4540 (2010).
77. Kang, S. *et al.* Dysbiosis of fecal microbiota in Crohn's disease patients as revealed by a
custom phylogenetic microarray. *Inflamm. Bowel Dis.* **16,** 2034–2042 (2010).
78. Babudieri, S. *et al.* Diarrhoea associated with toxigenic Clostridium spiroforme. *J. Infect.*
12, 278–279 (1986).
79. Borriello, S. P. Clostridial disease of the gut. *Clin. Infect. Dis.* **20,** S242–S250 (1995).
80. Zhang, X. *et al.* Human gut microbiota changes reveal the progression of glucose
intolerance. *PLoS One* **8,** e71108 (2013).
81. Ciubotaru, I., Green, S. J., Kukreja, S. & Barengolts, E. Significant differences in fecal
microbiota are associated with various stages of glucose tolerance in African American
male veterans. *Transl. Res.* **166,** 401–411 (2015).
82. Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D.
Development of a dual-index sequencing strategy and curation pipeline for analyzing
amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl. Environ.*
Microbiol. (2013). doi:10.1128/aem.01043-13
83. Schloss, P. D. *et al.* Introducing mothur: open-source, platform-Independent, community-
supported software for describing and comparing microbial communities. *Appl. Environ.*
Microbiol. **75,** 7537–7541 (2009).
84. Pruesse, E. *et al.* SILVA: a comprehensive online resource for quality checked and
aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res.* **35,**
7188–7196 (2007).
85. Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C. & Knight, R. UCHIME improves
sensitivity and speed of chimera detection. *Bioinformatics* **27,** 2194–2200 (2011).
86. DeSantis, T. Z. *et al.* Greengenes, a chimera-checked 16S rRNA gene database and
workbench compatible with ARB. *Appl. Environ. Microbiol.* **72,** 5069–5072 (2006).
87. R Core Team. R: A language and environment for statistical computing. (2017).
88. Oksanen, J. *et al.* vegan: community ecology package. (2015).
625

AUTHOR CONTRIBUTIONS

FER and PH designed research. JHK, RLK, and TS completed sample processing and sequencing. AP contributed to the conceptual and analytic approach to conducting the analyses. KADM, ZZT, and GC analyzed the data. KADM, ZZT, GC, FER, and PH contributed to the final manuscript.

Alberto Palloni[5]

ACKNOWLEDGMENTS

The authors wish to thank members of the Rey laboratory for their insights and support. This work was supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture (2016-67017-24416, FER), the National Institutes for Aging, National Institutes for Health (AG041868, PH), the Center for the Demography of Health and Aging (AG017266, PH), and the Clinical and Translational Science Award (CTSA), National Institutes for Health National Center for Advancing Translational Sciences (NCATS) (UL1TR000427, ZZT). Additional support was provided by the Vice Chancellor for Research and Graduate Education (VCGRE) at the U. of Wisconsin-Madison. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

The authors declare that we have no competing interests as defined by Nature Research, or other interests that might be perceived to influence the results and/or discussion reported in this paper.

FIGURES

Figure 1. Factors associated with the overall fecal microbiota. Non-metric multidimensional

scaling (nMDS) of unweighted UniFrac for all graduates (N = 179). Variables found to be

significant (PERMANOVA P < 0.05, red) and trends (0.05 < P < 0.1, black) are shown as fitted

arrows. Arrows point toward increasing values (dietary protein), toward affirmative responses

(high blood sugar, antibiotics, heart disease), or from male to female (sex).

658   **Figure 2.** Cohabitation is associated with increased alpha-diversity. Boxplots of (A) Shannon's

659   diversity and (B) Chao's richness of graduates that are (blue) or are not (red) cohabitating with a

660   spouse or partner. All spouses/partners were cohabitating while all non-cohabitating individuals

661   were unmarried. **P < 0.01, *P < 0.05

Figure 3. Microbial sharing in spouse and sibling relationships. Unweighted UniFrac distances of

(A) spouse, sibling, and unrelated pairs, (B) spouses and siblings grouped by relationship

closeness, and (C) spouses and siblings separated by relationship closeness. Groups (A) were

compared in linear regression model adjusting for potential confounders (*e.g.* age, sex, diet, health

conditions). P-values were averaged across 1000 rounds of unrelated pair sampling. Closeness

groups (B,C) were compared in linear regression models adjusting for potential confounders. (D,

E) Average percentages of shared OTUs within family groups including a related spouse and

sibling pair. Families included those with both very close spouses and siblings (D, N = 12) and

those with very close spouses and somewhat close siblings (E, N = 17). Percentages are of the total

number of OTUs across all three individuals, and circle sizes are proportional to total percentages

represented. ***P < 0.001, **P < 0.01, *P < 0.05, •P< 0.1

Figure 4. Percent relative abundance of OTUs that are commonly shared between spouses and

that differed between those with (grey) and without (white) high blood sugar. (A) Low abundance

and (B) more highly abundant OTUs. Means with standard error bars are shown. Kruskal-Wallis

FDR *$P < 0.05$, •$P < 0.1$