Prioritize diversity or declining species? Trade-offs and synergies in spatial planning for the conservation of migratory birds

Wilson, S[a,b]., Schuster R.[b,c], Rodewald, A. D.[d,e], Bennett, J.R.[b], Smith, A.C[b,f]. and Arcese P.[g],

[a] Wildlife Research Division, Environment and Climate Change Canada, National Wildlife Research Centre, 1125 Colonel By Drive, Ottawa, Ontario, Canada, K1S 5B6

[b] Department of Biology, 1125 Colonel By Drive, Carleton University, Ottawa ON, K1S 5B6 Canada

[c] Ecosystem Science and Management Program, 3333 University Way, University of Northern British Columbia, Prince George BC, V2N 4Z9 Canada

[d] Cornell Lab of Ornithology, 159 Sapsucker Woods Rd., Ithaca, NY 14850, USA

[e] Department of Natural Resources, Cornell University, Fernow Hall, #111, Ithaca, NY 14853, USA

[f] Canadian Wildlife Service, Environment and Climate Change Canada, National Wildlife Research Centre, 1125 Colonel By Drive, Ottawa, Ontario, Canada, K1S 5B6

[g] Department of Forest and Conservation Sciences, 2424 Main Mall, University of British Columbia, Vancouver BC, V6T 1Z4 Canada

Corresponding Author: S. Wilson, Wildlife Research Division, Environment and Climate Change Canada, National Wildlife Research Centre, 1125 Colonel By Drive, Ottawa, Ontario, Canada, K1S 5B6, email:scott.wilson@canada.ca

Abstract

Stemming biodiversity loss requires strategic conservation guided by well-articulated and achievable targets, whether they be proactive (e.g., protect diverse places) or reactive (e.g., protect threatened or declining species). Both types of targets can be effective, but there are trade-offs, especially for broadly-distributed ecosystems or taxa, such as migratory species, a group for which conservation has been challenged by limited knowledge of distributions throughout the annual cycle. We combined fine scale abundance-based distribution models using crowd-sourced data with population trend analyses to evaluate how conservation targets emphasizing proactive (high species diversity) versus reactive (declining species richness and average severity of decline) approaches influenced spatial priorities for the conservation of Neotropical migratory birds (n=112 species), during the non-breeding period in the Western Hemisphere. We also examined how priority landscapes for each target were affected by the magnitude and changes in human footprint, forest cover, and land protection. Spatial priorities were strikingly different when targets emphasized areas of high species diversity (southern Mexico, northern Central America) versus areas with more severe average declines across species (Andean cordilleras of South America). Targeting richness of declining species was most congruent with high diversity. Only a fraction of non-breeding landscapes (0.03%), mostly in southern Central America and northern Colombia, were selected across all targets. Priority areas for the most severe average declines across the community were subject to rapid increases in human footprint, unlike areas selected for high diversity or declining species richness. Our results highlight how even different reactive conservation approaches may emphasize different regions with varying threats and opportunities for conservation. Overlap between areas prioritized for the conservation of high species diversity and declining species richness in the

northern Neotropics indicate a potential for conservation to support declining species without

compromising conservation of diverse communities of Neotropical migrants. However, our study

provides new evidence that rapid land conversion in the northern Andes may drive declines of

Neotropical migrants overwintering in the region; conserving these communities of Neotropical

migrants will require an emphasis on protecting and restoring habitat in northwest South

America.

Keywords: citizen science, conservation prioritization, eBird, human footprint, Neotropical

migrant, population decline, protected area

Introduction

Stemming the unprecedented rates of current biodiversity loss (Pimm et al., 2014; Ceballos,

Ehrlich, & Dirzo, 2017) will require strategic and sustained investment in conservation but

socio-economic limits on such investments make difficult choices inevitable (Wilson, McBride,

Bode & Possingham, 2006; Carwardine et al., 2012; Martin et al., 2018). Although sophisticated

decision-support tools can help inform such choices, the selection of a conservation target is a

key initial step as different targets will often lead to different outcomes (e.g. Orme et al., 2005;

Grenyer et al., 2006; Klein et al., 2009; Somveille, Manica, Butchart, & Rodrigues, 2013).

Potential targets include areas of high species diversity (Buchanan, Donald & Butchart, 2011),

critical habitat for threatened species (ESA, 1973; SARA, 2002), endemism (Myers, Mittermeier,

Mittermeier, da Fonseca, & Kent, 2000) and areas of intact wilderness (Watson et al., 2009).

Targets may also reflect contrasting conservation paradigms. For example, prioritizing high

species diversity may be viewed as a proactive approach to conserving areas of high biodiversity

value, whereas targeting areas with a smaller number of species in rapid decline, or a large

number of species in modest decline, could be viewed as a more reactive response to species in

need of conservation effort (*cf.* Myers et al., 2000; Buchanan et al., 2011; Martin et al., 2018). In

this paper, we illustrate the consequences of proactive versus reactive approaches for area-based

conservation strategies of > 100 Neotropical migratory birds during the non-breeding season in

the Western Hemisphere.

Migratory species are declining globally (Wilcove & Wikelski, 2008; Harris, Thurgood,

Hopcraft, Cromsight, & Berger et al., 2009) and present unique challenges in conservation

planning (Martin et al., 2007; Runge, Martin, Possingham, Willis, & Fuller, 2014). Because

migrants traverse vast distances throughout the annual cycle we often do not know what

locations and periods of the annual cycle most limit population growth. While threats on both the

breeding and non-breeding grounds likely combine to influence abundance in most cases, the

relative influence of threats at different stages of the annual cycle may vary among species

within a community and among different populations (Rushing, Ryder, & Marra, 2016;

Morrison, Robinson, Butler, Clark, & Gill, 2016; Taylor & Stutchbury, 2016; Wilson et al.

2018a; Wilson, Smith, & Naujokaitis-Lewis, 2018b). This issue creates uncertainty for multi-

species conservation efforts because it is unclear which species would benefit from management

actions within a particular region. A proactive strategy could still emphasize areas with high

species diversity to protect against future threats in the region while simultaneously benefiting

declining species in the community. However, this may be an inefficient use of scarce

conservation resources if the recovery of threatened species is the priority. In that case our

approach could focus on areas showing the strongest average declines across species. A region

with steep community-wide declines provides evidence that threats within those regions are

negatively impacting populations.

Neotropical migrant songbirds have recently received much attention (e.g. NABCI, 2012;

NABCI, 2016) in large part due to long-term population declines of several species

(Environment Canada, 2017; Sauer et al., 2017). The majority of conservation investment for

this group is focused on the breeding period of the annual cycle in North America and primarily

through reactive single-species conservation measures for those species undergoing the steepest

declines (ESA, 1973; SARA, 2002). Yet, there is a growing concern and recognition that rapid

land-use change on the non-breeding grounds is a key driver of declines for some threatened

Neotropical migrants (Taylor & Stutchbury, 2016; Kramer et al., 2018; Wilson et al., 2018a).

Thus, we may be able to more effectively conserve these species if we focus conservation

resources on this period of the annual cycle. Historically, there has been far less information on

the non-breeding season distribution and ecology of migrants in Neotropical regions, which may

have in part contributed to the greater conservation focus on the breeding grounds. However,

advances in crowd-sourced data and their application to species distribution models now allow

us to generate weekly estimates of the abundance and distribution for most Neotropical

migratory passerines over their entire annual cycles (Fink et al., 2010; Sullivan et al., 2014; La

Sorte et al., 2017; Fink et al., 2018). Precise estimates of abundance are critical for predicting

fine-scale species habitat use and response to landscape change that will allow for conservation

planning at local scales (Possingham & Wilson, 2005).

Our first objective was to evaluate how the geographic regions and ecosystems targeted

for conservation would differ depending on a proactive approach towards areas of high species

diversity versus a reactive approach towards areas with the strongest average declines across

species and the largest number of declining species. To accomplish this, we used distribution

models based on crowd-sourced data for 112 Neotropical migrants during their stationary non-

breeding period (hereafter 'non-breeding period'). The abundance-based 8.34×8.34km spatial

grain we used in this analysis is one of the finest spatial resolutions to date for assessing the

location of conservation targets at a hemispheric scale. We were also interested in identifying

complementarity where conservation efforts might enhance protection of diversity hotspots and

species in need of conservation attention. Our second objective was to examine the extent of

recent land-use change and protected area status for priority areas based on proactive versus

reactive conservation targets. We had two goals within this objective. First, by examining human

land-use change and degree of protection within priority areas, we provide key information for

the urgency of conservation and types of conservation actions that might be most suitable.

Second, we identified the extent to which hotspots of community-wide decline and declining

species richness coincided with rapid and recent land-use change within those regions. The

identification of these regions would provide evidence for the location of threats during the non-

breeding season that are contributing to population declines across the community.

Methods

Species selection

We initially identified 117 species of Neotropical migratory passerines for which weekly

distributions were available based on eBird spatio-temporal exploratory models (Fink et al.,

2010; Fink et al., 2018; Supporting Information (SI) Table S1). We set the non-breeding period

as November 14 – March 14 to exclude spring and fall migration and retained all species with

weekly distribution maps throughout that period (n = 112 species). Species fell into two broad

groups based on their breeding and stationary non-breeding ranges: 1) species that are solely

migratory, wherein all individuals breed in North America north of the US-Mexico border and

migrate south of the Tropic of Cancer during the non-breeding period (n = 95 species); 2) species

with migratory and resident populations in the Neotropics, wherein individuals from migratory

populations north of the US-Mexico border migrate south of the Tropic of Cancer during the

non-breeding period (n = 17 species). STEM maps were not available for all Neotropical

migrants passerines and thus a small number were not included in our analysis.

Spatiotemporal exploratory models

We used the eBird citizen-science database (Sullivan et al., 2014) to estimate the weekly

abundance of all species in each eBird cell (hereafter 'landscape', Fig. 1) during the non-

breeding season. Estimates of abundance were derived previously using spatiotemporal

exploratory models (STEM) and detailed descriptions of the development and application of

STEM are provided elsewhere (Fink et al., 2010; Fink et al., 2018). Briefly, the eBird data used

in STEM includes species counts from complete checklists collected under the "traveling",

"stationary", and "areal" protocols from January 1, 2004 to December 31, 2016 within the spatial

extent bounded by 180° to 30° W Longitude (as well as Alaska between 150° E and 180° E).

This resulted in a dataset consisting of 14 million checklists collected at 1.7 million unique

locations. The STEM developed from this database are an ensemble of local regression models

generated from a spatiotemporal block subsampling design. Abundance of each species and in

each week are assumed to be stationary. Zero-inflated boosted regression trees are used to

predict the observed counts (abundance) of species based on spatial covariates (e.g. relationships

with land cover types), temporal covariates to account for trends, and predictors that describe the

observation/detection process to incorporate variation in detectability. Estimates of relative

abundance are rendered at weekly temporal resolution and at an 8.34×8.34 km spatial

resolution. The quantity estimated per eBird landscape was the expected number of birds of a

given species by a typical eBird participant on a search starting from the center of the pixel from

7:00 to 8:00 AM while traveling 1 km. The final set of models contained estimates of abundance

for all weeks throughout the annual cycle; for this analysis, we restricted the time window to the

non-breeding period as described above.

Human Footprint

We used the global human footprint index (Sanderson et al., 2002; Venter et al., 2016) to identify

the extent and trends in human pressures for all eBird landscapes. The human footprint index has

been shown to be a strong predictor of ecological and conservation metrics (Yackulic, Sanderson,

& Uriarte, 2011; Hand, Cushman, Landguth, & Lucotch, 2014). The index is a composite

measure of human impact derived from eight separate measures: 1) built environments, 2) crop

land, 3) pasture land, 4) human population density, 5) night-time lights, 6) railways, 7) roads,

and 8) navigable waterways. These eight measures are individually weighted based on their

relative levels of human pressure and summed to create a single standardized estimate. The index

varies from 0 (no footprint) to 50 (very high footprint) and is estimated at a 1km resolution

across all global terrestrial lands except Antarctica. The index was first measured in 1993 and

again in 2009 and therefore, for all eBird landscapes in our analysis we obtained an average

footprint in these two years as well as the change over the 16-yr period. With the $1km^2$ resolution

there were approximately 70 estimates of footprint in each landscape and these were averaged to

create a single measure.

Land Cover

We used the 2015 data set of the global land cover map (ESA Climate Change Initiative, 2017)

at a 300m resolution to examine the extent of forest cover and diversity of land covers among the

priority landscapes for each target as well as for the areas where priority landscapes of different

targets overlap. The land cover data layer was clipped to the study region with 32 land cover

types that comprised at least one pixel. With the 300m resolution there were approximately 773

land cover pixels in each eBird landscape and for each landscape, we calculated the proportion of

pixels that comprised forest cover and a Shannon index of land cover diversity using the Vegan

package (Oksanen et al., 2017) in R Version 3.4.2 (http://www.r-project.org/).

Protected area coverage

We used the World Database on Protected Areas (WDPA, UNEP-WCMC, 2018) to estimate the

extent of protected area coverage for the three conservation targets. The WDPA includes seven

categories of protected area coverage: (Ia) strict nature reserve, (Ib) wilderness area, (II) national

park, (III) national monument, (IV) habitat/species management, (V) protected

landscape/seascape, (VI) managed resource protected area. Following La Sorte et al. (2017) we

defined these seven classes into three representing high (Ia, Ib), medium (II, III) and low (IV, V, VI)

protection status. We then intersected the WDPA layer with the eBird landscape layer. For all

targets we calculated the proportion of landscapes that contained each of the 7 protected area

types and the high, medium and low protected area categories.

Analyses

We used the eBird abundance information to calculate Shannon diversity, median long-term

BBS trends (hereafter referred to as "average severity of decline") and number of declining

species for all eBird landscapes as defined above. A few species in our analysis also over-winter

in regions well outside of the Neotropics (e.g. Yellow-rumped Warbler). Therefore, we defined

the non-breeding region based on all landscapes containing five or more species. This definition

allowed us to include the distributions from all species while excluding large, peripheral areas

that are well outside of the main non-breeding range for the vast majority of Neotropical

migrants. Long-term estimates of population trend in Neotropical migratory birds do not exist for

the non-breeding period; thus, we used continent-wide trends estimated for each species from

1966-2015 via the North American Breeding Bird Survey (BBS, Environment Canada, 2017;

Sauer et al., 2017). In this case, for each species present in each landscape we used the long-term

continental trend to estimate the median trend across all species detected in that landscape. To

estimate the richness of declining species in each landscape we summed the number of species

present with a long-term continental trend below 0. Population trends were unavailable for three

species (Table S1).

For each target, we defined the top 10% and 5% (see Appendix 1) of landscapes as a

priority (i.e. based on highest diversity, average severity of decline and declining species

richness). This approach of selecting a top percentage of values in each target category is similar

to that used in earlier studies (e.g. Orme et al. 2005; Hof, Araújo, Jetz, & Rahbek, 2011). We

then identified the geographic regions of these priority landscapes for each target individually as

well as the regions where the priority landscapes of different targets overlapped. To quantify the

spatial overlap among targets, we used a Bray-Curtis measure of dissimilarity between the

priority landscapes. Dissimilarity calculations were performed using the Vegan package

(Oksanen et al. 2017) in R Version 3.4.4.

Results

Geographic variation in priority landscapes

Our results highlighted a strong north-south contrast between regions containing the highest

diversity and those with the most severe average declines across the community (Fig. 2). This

contrast resulted in very different regions where landscapes were prioritized for the two targets.

Priority landscapes for high species diversity were concentrated along the Pacific slope of

Mexico, the Yucatan peninsula, western Cuba and southern Mexico through Nicaragua (Fig. 2,

Supporting Information (SI) Fig. S1). In contrast, priority landscapes based on average severity

of decline were mostly in Andean regions of Colombia through Bolivia (Fig. 2, S1). The regional

concentration of priority landscapes for declining species richness was also in the northern

Neotropics, similar to diversity, although with fewer priority landscapes on Mexico's Pacific

slope (Fig. 2).

Variation in focal regions for the proactive versus reactive conservation targets were even

more apparent when we examined the extent to which the targets overlapped. Of 96,079

landscapes in the non-breeding region, only 26 were in the top 10% for scenarios based on all

three targets (Table 1). Overlapping landscapes were mainly in two regions; the Cordillera de

Talamanca of Costa Rica and west Panama, and the Cordillera Occidental of Colombia (Fig. S2).

The overlap was also low between landscapes for average severity of decline and high diversity

targets (n=35 landscapes, Table 1, Fig. S3) and between average severity of decline and number

of declining species (n=155 landscapes, Table 1, Fig. S4). In contrast, there was a strong,

positive correlation (r=0.79) in the target areas for high diversity and number of declining

species, particularly in southern Mexico, the Yucatan peninsula and Honduras (Table 1, Fig. S5).

Human footprint and land cover among target landscapes

The magnitude and trend in the human footprint index differed among the three targets. Priority

landscapes for diversity and average severity of decline had a smaller average human footprint

than the non-breeding region overall, whereas sites prioritized based on the number of declining

species had footprints slightly above the region average (Table 2). From 1993 to 2009, the

human footprint index declined by 6.4% in priority landscapes based on high diversity and

number of declining species, but increased by 11% for priority landscapes for average severity of

decline. This pattern was also reflected in a higher proportion of landscapes where the footprint

increased between 1993 and 2009 for priority average severity of decline landscapes (67.2%)

than for priority diversity (43.0%) or declining species richness (42.5%) landscapes. The three

targets had similar land cover patterns with relatively greater forest cover and lower land cover

diversity compared to the regional average (Table 2).

Differences in protected area coverage among target landscapes

Across the entire non-breeding region, 15.9% of all landscapes had some form of protected area

status (Table 3). Priority landscapes for high diversity and number of declining species targets

were similar in their protected area coverage, with 19.44% and 18.86% respectively having some

form of protection and the majority of those with low protection status (69% and 66%

respectively), mostly in the form of managed resource protected areas (Table 3, A1). Priority

landscapes for average severity of decline had a similar percentage of landscapes with some form

of protection (18.12%) but in contrast to the other two targets, the majority of these (64%) had

medium protection status and were primarily National Parks (Table S2). Thirteen of the 26

landscapes (50%) that met all three targets had protected area status with medium status in 7 (all

National Parks) and low status in 7 (all managed resource protection areas) with one landscape

containing both types (Table 3, S2).

Discussion

To our knowledge this is the first study combining fine-scale abundance-based distribution data

with population trend analyses to examine the complementarity of proactive versus reactive

conservation approaches for migratory species. Our results highlight the importance of target

selection when prioritizing landscapes for conservation (Possingham & Wilson 2005; Klein et

al., 2009). We observed marked differences in landscapes prioritized for conservation using a

proactive target that emphasized high species diversity versus a reactive target focusing on areas

with more severe average declines across the community. When the target was high species

diversity priority landscapes were distributed mainly in the northern portion of the non-breeding

region in southern Mexico, the Yucatan Peninsula and northern Central America, whereas

targeting areas with more severe average declines across the community emphasized the Andean

cordilleras of South America. Interestingly, the priority landscapes for an alternative reactive

approach that emphasized the total number of declining species, irrespective of the extent of that

decline, coincided to a much greater extent with those selected for high diversity than those

selected for average severity of decline.

Global analyses of resident taxa offer mixed results on the extent to which areas

containing threatened species are congruent with areas of high species diversity. In amphibians,

strong spatial overlap exists between the total number of species and the number at risk of

extinction (Hof et al., 2011), but in birds and mammals there is less congruence at global scales

(Orme et al., 2005; Schipper et al., 2008; Jenkins, Pimm, & Jopa, 2013; Pimm et al., 2014). To

our knowledge, only one previous study has addressed this topic for migratory species and found

little congruence between areas of high richness, endemism and threatened status at global scales

(Somveille et al., 2013). Whereas some regional studies show high spatial overlap in species

diversity and endangerment (e.g. Chilean mammals, Cofré & Marquet, 1999), it is not evident in

other cases (e.g. British fauna, Prendergast, Quinn, Lawton, Eversham, & Gibbons, 1993). The

extent of congruence between these two targets often depends on the degree to which threatening

anthropogenic processes overlap areas rich in resident taxa (Orme et al., 2005; Hof et al., 2011)

as well as the timeframe examined (Bennett & Arcese, 2013). For example, habitat conversion

by humans might initially lead to spatial correlations between diversity, population decline and

human influence at regional scales, but eventually leads to negative correlations between rare

species occurrence and human influence as declining species are extirpated from human-

dominated landscapes (Bennett & Arcese, 2013).

Our results on the extent of congruence among landscapes supporting diversity versus

declining species were complex and depended on how we used the information on population

trend to define a species in need of conservation attention. When our reactive approach

emphasized areas with species showing more severe average declines, the results were consistent

with those described above with little congruence between diversity and species in need of

conservation attention when threatening processes and diversity do not overlap. Landscapes in

more southern regions where average species declines were greater showed a recent increase in

the human footprint of about 1% annually and even this rapid change understates the longer-term

extent of habitat loss in the northern Andean region where less than 10% of the original forest

cover remains (Henderson, Churchill, & Luteyn, 1991). In contrast, there has been a slight

reduction in the human footprint in landscapes prioritized for the highest Neotropical migrant

diversity.

When we changed the definition of our reactive conservation target to the number of

declining species there was a strong congruence with total diversity but little congruence with

recent anthropogenic change or areas prioritized for average severity of decline. This finding

stresses that even different conservation targets with a similar theme (i.e. declining species) can

lead to different outcomes for spatial conservation planning. We suspect that the lack of overlap

in regions with more severe average declines across the community and those with the largest

number of declining species is the combined result of two processes 1) the decrease in

Neotropical migrant diversity towards the southern portion of the non-breeding range and 2) the

influence of multiple stages of the annual cycle on population abundance and trends. We would

expect a high congruence between average severity of decline and declining species richness if

threats only occurred during the stationary non-breeding period but this is very unlikely to be the

case. Approximately 55% of the species in the northern portion of the non-breeding range still

had negative trends even if the majority of those were slight compared to species overwintering

in South America (see below). There is less evidence that conditions on the non-breeding

grounds have negatively impacted species overwintering in Mexico, Central America and the

Caribbean (see also NABCI 2012) and it is possible that species overwintering in these regions

are more evenly influenced by threats throughout the annual cycle.

Our study provides new evidence that recent land conversion in the South American

Andes has contributed to population declines of Neotropical migrants overwintering in the region

(Jones, Barg, Sillett, Veit, & Robertson, 2004; González-Prieto, Bayly, Colorado, & Hobson,

2017; Kramer et al. 2018, Wilson et al. 2018a). Indeed, 18 of 20 species in our analysis whose

winter range primarily includes northwest South America had declining trends with an average

decline for the community of approximately 51% between 1966 and 2015. Yet, because the

stationary portion of the non-breeding period represents only 50-60% of the annual cycle in most

species, other drivers may be involved. The group of declining Neotropical migrants in South

America occupy broad and often different regions on the breeding grounds and thus, a common

breeding ground threat for community wide declines in this group is unlikely. Yet, many of these

species use similar routes for at least part of the spring and/or fall migration and further study is

needed on the extent to which threats along shared migratory pathways are contributing to the

stronger declines for more southern overwintering Neotropical migrants. Steeper population

declines have previously been reported for long-distance compared to short-distance migrants

(Sanderson, Donald, Pain, Burfield, & van Bommel, 2006). This finding is consistent with

observations that high energy demands and unfamiliar landscapes make migration a particularly

risky stage of the annual cycle (Sillett & Holmes 2002, Klassen et al. 2014), especially where

migratory pathways intersect with human development (Berger, Young, & Berger, 2008;

Hewson, Thorup, Pearce-Higgins, Atkinson, 2016).

Conclusions

Considerable overlap between areas prioritized for the conservation of high species diversity and

declining species richness indicate a potential for conservation investment in landscapes likely to

support declining species without compromising conservation of diverse communities of

Neotropical migrant birds. Because contemporary declines often predict future declines (Keith et

al., 2015), area prioritizations based on the above targets are consistent with strategies aimed at

'keeping common species common' and stemming declines, including for species currently of

low conservation concern (Ceballos et al., 2017). However, as we have shown, an emphasis on

areas of high diversity or higher numbers of declining species will not incorporate areas where

the Neotropical migrant community is declining most strongly. Many of the species in these

communities are currently the focus of species-specific conservation efforts on the breeding

grounds (ESA 1973, SARA 2002) but our results point to the importance of protection and

restoration of habitat in the northern Andes for effective conservation of these species. Although

we found limited potential to meet all targets, we did identify ~2400km^2 of habitat in southern

Central America and northwest Colombia that met all three; about 30% of this area already is in

a National Park, another ~30% is managed with lower conservation emphasis. This type of

information also shows how the method we used in this study could inform planning at a local

scale. As an example, if the priority is to protect areas that meet proactive and reactive targets,

then the types of information provided in the approach we used could be combined with socio-

economic information (e.g. Naidoo et al. 2006; Wilson et al. 2006) to identify the costs and

benefits of allocating conservation status to sites within other landscapes that currently do not

have protected area status.

Authors' Contributions

372 SW, RS, ADR, JRB and PA designed the study; RS, SW and ACS analyzed data; SW wrote the

373 manuscript with feedback and contributions from all co-authors.

Data Accessibility

eBird STEM maps are in the process of being made publicly available (Fink et al. 2018,

https://doi.org/10.1101/251868; see also https://ebird.org/science/modeling). All other databases

used in this analysis are publicly available: Breeding Bird Survey trend data (Environment

Canada, 2017, https://www.canada.ca/en/environment-climate-change/services/bird-

surveys/landbird/north-american-breeding/overview.html; Sauer et al., 2017,

https://www.pwrc.usgs.gov/bbs/), Human Footprint Index (Venter et al., 2016,

https://doi.org/10.1038/ncomms12558), Land Cover (ESA Climate Change Initiative, 2017,

http://maps.elie.ucl.ac.be/CCI/viewer/download.php), Protected Areas (UNEP-WCMC, 2018,

https://www.protectedplanet.net/c/world-database-on-protected-areas).

Literature Cited

Bennett, J. R., & Arcese, P. (2013). Human influence and classical island biogeographic

predictors of rare species occurrence. Conservation Biology, 27, 417–421.

https://doi.org/10.1111/cobi.12015.

Berger, J., Young, J. K., & Berger, K. M. (2009). Protecting migration corridors: challenges and

optimism for Mongolian Saiga. PLOS Biology, 6, e165.

https://doi.org/10.1371/journal.pbio.0060165.

Buchanan, G. M., Donald, P. F., & Butchart, S. H. M. (2012). Identifying Priority Areas for

Conservation: A Global Assessment for Forest-Dependent Birds. PLoS One 12, e29080.

https://doi.org/10.1371/journal.pone.0029080

Carwardine, J., O'Connor, T., Legge, S., Mackey, B., Possingham, H. P., & Martin, T. G.

(2012). Prioritizing threat management for biodiversity conservation. Conservation Letters,

5, 196–204. https://doi.org/10.1111/j.1755-263X.2012.00228.x.

Ceballos, G., Ehrlich, P. R., & Dirzo, F. (2017). Biological annihilation via the ongoing sixth

mass extinction signaled by vertebrate population losses and declines. Proceedings of the

National Academy of Sciences, p. 201704949. https://doi.org/10.1073/pnas.1704949114.

Cofré, H., & Marquet, P. (1999). Conservation status, rarity, and geographic priorities for

conservation of Chilean mammals: an assessment. Biological Conservation, 88, 53–68.

https://doi.org/10.1016/S0006-3207(98)00090-1.

Environment Canada. (2017). North American Breeding Bird Survey - Canadian Trends

Website, Data-version 2015. Environment Canada, Gatineau, Quebec, K1A 0H3.

ESA Climate Change Initiative. (2017). Global land cover map 300m resolution for 2015.

Available at: http://maps.elie.ucl.ac.be/CCI/viewer/download.php.

[ESA] Endangered Species Act. (1973). Pub. L. No. 93–205, 87 Stat. 884

Fink, D., Hochachka, W. M., Zuckerberg, D., Winkler, D. W., Shaby, B., Munson, M. A.,…

Kelling, S. (2010). Spatiotemporal exploratory models for broad-scale survey data.

Ecological Applications, 20, 2131–2147. https://doi.org/10.1890/09-1340.1

Fink, D., Auer, T., Ruiz-Gutierrez, V., Hochachka, W.M., Johnston, A., La Sorte, F. A., &

Kelling, S. (2018). Modeling avian full annual cycle distribution and population trends with

citizen science data. bioRxiv 251868. https://doi.org/10.1101/251868.

González-Prieto, A. M., Bayly, N. J., Colorado, G. J. & Hobson, K. A. (2017). Topography of

the Andes Mountains shapes the wintering distribution of a migratory bird. Diversity and

Distributions 23, 118–129. https://doi.org/10.1111/ddi.12515.

Grenyer, R., Orme, C. D. L., Jackson, S. F., Thomas, G. H., Davies, R. G., Davies, T. J., …

Owens, I. P. F. (2006). Global distribution and conservation of rare and threatened

vertebrates. Nature, 444, 993–996. https://doi.org/10.1038/nature05237.

Hand, B. K., Cushman, S. A., Landguth, E. L., & Lucotch, J. (2014). Assessing multi-taxa

sensitivity to the human footprint, habitat fragmentation and loss by exploring alternative

scenarios of dispersal ability and population size: a simulation approach. Biodiversity and

Conservation, 23, 2761–2779. https://doi.org/10.1007/s10531-014-0747-x.

Harris, G., Thurgood, S., Hopcraft, J. G. C., Cromsight, J. P., & Berger, J. (2009). Global decline

in aggregated migrations of large terrestrial mammals. Endangered Species Research, 7, 55–

76. https://doi.org/10.3354/esr00173.

Henderson. A., Churchill, S. P., & Luteyn, J. L. (1991). Neotropical plant diversity. Nature 351,

21–22. 10.1038/351021e0.

Hewson, C. M., Thorup, K., Pearce-Higgins, J. W., & Atkinson, P. W. (2016). Population

decline is linked to migration route in the common cuckoo. Nature Communications, **7**,

12296. 10.1038/ncomms12296.

Hof, C., Araújo, M. B., Jetz, W., & Rahbek, C. (2011). Additive threats from pathogens, climate

and land-use change for global amphibian diversity. Nature, 480, 516–521.

https://doi.org/10.1038/nature10650.

Jenkins, C. N., Pimm, S. L., & Joppa, L. N. (2013). Global patterns of terrestrial vertebrate

diversity and conservation. Proceedings of the National Academy of Sciences, 110, E2602–

E2610. https://doi.org/10.1073/pnas.1302251110.

Jones, J., Barg, J. J., Sillett, T. S., Veit, L. M., & Robertson, R. J. (2004). Minimum estimates of

survival and population growth for Cerulean Warblers (*Dendroica cerulea*) breeding in

Ontario, Canada. The Auk: Ornithological Advances, 121(1), 15–22.

https://doi.org/10.1642/0004-8038(2004)121[0015:MEOSAP]2.0.CO;2.

Keith, D., Ackakaya, H. R., Butchart, S. H. M., Collen, B., Dulvy, N. K., Holmes, E. E.,…

Waples, R. S. (2015). Temporal correlations in population trends: Conservation implications

from time-series analysis of diverse animal taxa. Biological Conservation, 192, 247–257.

https://doi.org/10.1016/j.biocon.2015.09.021.

Klaassen, R. H. G., Hake, M., Strandberg, R., Koks, B. J., Trierweiler, C., Exo, K., …, Alerstam,

T. (2014). When and where does mortality occur in migratory birds? Direct evidence from

long-term satellite tracking of raptors. Journal of Animal Ecology, 83, 176–184.

https://doi.org/10.1111/1365-2656.12135.

Klein, C. J., Wilson, K. A., Watts, M., Stein, J., Carwardine, J., Mackey, B., Possingham, H. P.

(2009). Spatial conservation prioritization inclusive of wilderness quality: a case study of

Australia's biodiversity. Biological Conservation, 142, 1282–1290.

https://doi.org/10.1016/j.biocon.2009.01.035.

Kramer, G. R., Andersen, D. E, Buehler, D. A., Wood, P. B., Peterson, S. M., Lehman, J. A.,…

Streby, H. M. (2018). Population trends in *Vermivora* warblers are linked to strong

migratory connectivity. Proceedings of the National Academy of Sciences USA, 115,

E3192-E3200. https://doi.org/10.1073/pnas.1718985115.

La Sorte, F. A., Fink, D., Blancher, P. J., Rodewald, A. D., Ruiz-Gutierrez, V., Rosenberg, K. R.,

… Kelling, S. (2017). Global change and the distributional dynamics of migratory bird

populations wintering in Central America. Global Change Biology, 23, 5284–5296.

https://doi.org/10.1111/gcb.13794.

Martin, T. G., Chadès, I., Arcese, P., Marra, P. P., Possingham, H. P., & Norris R. (2007).

Optimal conservation of migratory species. PLoS ONE, 2, e751:

https://doi.org/10.1371/journal.pone.0000751.

Martin, T. G., Kehoe, L., Mantyka-Pringle, C. S., Chadès I., Wilson, S., Bloom, R.,… & Smith,

P. A. (2018). Prioritizing recovery funding to maximize conservation of endangered species.

Conservation Letters. In press.

Morrison, C. A., Robinson, R. A., Butler, S. J., Clark, J. A., & Gill, J. A. (2016). Demographic

drivers of decline and recovery in an Afro-Palearctic migratory bird population. Proceedings

of the Royal Society B, 283, 2016.1387. 10.1098/rspb.2016.1387.

Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., & Kent, J. (2000).

Biodiversity hotspots for conservation priorities. Nature, 403, 853–858.

https://doi.org/10.1038/35002501.

Naidoo, R., Balmford, A., Ferraro, P.J., Polasky, S., Ricketts, T.H., & Rouget, M. (2006).

Integrating economic costs into conservation planning. Trends in Ecology and Evolution, 21,

681−687. https://doi.org/10.1016/j.tree.2006.10.003.

North American Bird Conservation Initiative Canada (NABCI). 2012. The State of Canada's

Birds, 2012. Environment Canada, Ottawa, Canada. 36 pages.

North American Bird Conservation Initiative Canada (NABCI). (2016). The State of North

America's Birds, 2016. Environment and Climate Change Canada, Ottawa, Canada. 8 pages.

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D.,…Wagner, H.

(2017). vegan: Community Ecology Package. R package. version 2.4.3.

Orme, C. D. L., Davies, R. G., Burgess, M., Eigenbrod, F., Pickup, N., Olson, V. A.,…Owens, I.

P. F. (2005). Global hotspots of species richness are not congruent with endemism or threat.

Nature, 436, 1016–1019. https://doi.org/10.1038/nature03850.

Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N.,… Sexton,

J.O. (2014). The biodiversity of species and their rates of extinction, distribution and

protection. Science, 344, 1246752. 10.1126/science.1246752.

Possingham, H. P., & Wilson, K. A. (2005). Biodiversity: turning up the heat on hotspots.

Nature, 436, 919–920. https://doi.org/10.1038/436919a.

Prendergast, J. R., Quinn, R. M., Lawton, J. H., Eversham, B. C., & Gibbons, D. W. (1993).

Rare species, the coincidence of diversity hotspots and conservation strategies. Nature, 365,

335–337. https://doi.org/10.1038/365335a0.

Runge, C. A., Martin, T. G., Possingham, H. P., Willis, S. G., Fuller, R. A. (2014). Conserving

mobile species. Frontiers in Ecology and the Environment, 12, 395–402.

https://doi.org/10.1890/130237

Rushing, C. S., Ryder, T. B., Marra, P. P. (2016). Quantifying drivers of population dynamics for

a migratory bird throughout the annual cycle. Proceedings of the Royal Society B, 283,

2015.2846. 10.1098/rspb.2015.2846.

Sanderson, E. W., Jaiteh, M., Levy, M. A., Redford, K. H., Wannebo, A. V., Woolmer, G.

(2002). The human footprint and the last of the wild. Bioscience, 52, 891–904.

https://doi.org/10.1641/0006-3568(2002)052[0891:THFATL]2.0.CO;2.

Sanderson, F. J., Donald, P. F., Pain, D. J., Burfield, I. J., van Bommel, F. P. J. (2006). Long-

term population declines in Afro-Palearctic migrant birds. Biological Conservation, 131, 93–

105. https://doi.org/10.1016/j.biocon.2006.02.008.

[SARA] Species At Risk Act. (2002). Bill C-5, An act respecting the protection of wildlife

species at risk in Canada.

Sauer, J. R., Niven, D. K., Hines, J. E., Ziolkowski Jr., D. J., Pardieck, K. L., Fallon, J. E., &

Link, W. A. (2017). The North American Breeding Bird Survey, Results and Analysis 1966

- 2015. Version 01.30.2017 USGS Patuxent Wildlife Research Center, Laurel, MD.

Schipper, J., Chanson, J. S., Chiozza, F., Cox, N. A., Hoffmann, M., Katariya, V.,…Young, B. E.

(2008). The status of the world's land and marine mammals: diversity, threat and

knowledge. Science, 322, 225–230. DOI: 10.1126/science.1165115.

Sillett, T.S., & Holmes, R.T. (2002). Variation in survivorship of a migratory songbird

throughout its annual cycle. Journal of Animal Ecology, 71, 296–308.

https://doi.org/10.1046/j.1365-2656.2002.00599.x.

Somveille, M., Manica, A., Butchart, S.H.M., Rodrigues, A.S.L. (2013). Mapping global

distribution patterns for migratory birds. PLoS ONE, 8, e70907.

https://doi.org/10.1371/journal.pone.0070907.

Sullivan, B. L., Aycrigg, J. L., Barry, J. H., Bonney, R. E., Bruns, N., Cooper, C. B., … Kelling,

S. (2014). The eBird enterprise: An integrated approach to development and application of

citizen science. Biological Conservation, 169, 31–40.

https://doi.org/10.1016/j.biocon.2013.11.003.

Taylor, C. M., & Stutchbury, B. J. M. (2016). Effect of breeding versus winter habitat loss and

fragmentation on the population dynamics of a migratory songbird. Ecological Applications,

26, 424–437. https://doi.org/10.1890/14-1410.

UNEP-WCMC. (2018) World database on protected areas. UNEP-WCMC,Cambridge, United

Kingdom. Available from https://www.protectedplanet.net/c/world-database-on-protected-

areas. (accessed April 2018).

Venter, O., Sanderson, E. W., Magrach, A., Allan, J. R., Beher, J., Jones, K. R.,…Watson, J. M.

(2016). Sixteen years of change in the global terrestrial human footprint and implications for

biodiversity conservation. Nature Communications, **7**, 12558.

https://doi.org/10.1038/ncomms12558.

Watson, J. E. M., Fuller, R. A., Watson, A. W. T., Mackey, B. G., Wilson, K. A., Grantham, H.

S.,…Possingham, H. P. (2009). Wilderness and future conservation priorities in Australia.

Diversity and Distributions, 15, 1028–1036. https://doi.org/10.1111/j.1472-

4642.2009.00601.x.

Wilcove, D. S., & Wikelski, M. (2008). Going, Going, Gone: Is Animal Migration Disappearing.

PLoS Biology, 6, e188. https://doi.org/10.1371/journal.pbio.0060188.

Wilson, K. A., McBride, M. F., Bode, M., Possingham, H. P. (2006). Prioritizing global

conservation efforts. Nature, 440, 337–340. https://doi.org/10.1038/nature04366.

Wilson, S., Saracco, J. F., Krikun, R., Flockhart, D. T. T., Godwin, C. M., Foster, K. R. (2018a)

Drivers of decline across the annual cycle of a threatened migratory bird. Scientific Reports,

8, 7316. DOI.10.1038/s41598-018-25633-z.

Wilson, S., Smith, A. C., & Naujokaitis-Lewis, I. (2018b). Opposing responses to drought shape

spatial population dynamics of declining grassland birds. Diversity and Distributions. DOI:

10.1111/ddi.12811

Yackulic, C. B., Sanderson, E. W., & Uriarte, M. (2011). Anthropogenic and environmental

drivers of modern range loss in large mammals. Proceedings of the National Academy of

Sciences, 108, 4024–4029. https://doi.org/10.1073/pnas.1015097108.

568   Table 1. Comparison of landscape overlap for conservation targets based on diversity, average

569   severity of decline and number of declining species. Target landscapes are based on the

570   upper 10% of values for each case. Landscapes = number of overlapping landscapes in each

571   comparison, Percent of total = percent of all landscapes in the study region that overlap,

572   dissimilarity = Bray-Curtis dissimilarity index for the landscape pairs in each comparison.

Comparison	Landscapes	Percent of total	Dissimilarity
Diversity/severity of decline	35	0.04	0.996
Diversity/number declining	6622	6.89	0.355
Severity of decline/number declining	155	0.16	0.985
All targets	26	0.03	-

574   Table 2. Comparison of trends in the human footprint index, land cover diversity and proportion of forest cover for priority landscapes

575   selected for each target (means with 95% CI in brackets underneath). HF = average human footprint index, ΔHF = percent change

576   in HF between 1993 and 2009, LC diversity = Shannon index of land cover diversity, proportion forest = average proportion of

577   landscapes covered by forest land cover types.

Response	HF 1993	HF 2009	ΔHF	LC diversity	Proportion forest
Diversity	7.63	7.14	-6.38	0.636	0.763
	(7.53, 7.73)	(7.04, 7.25)		(0.627, 0.646)	(0.757, 0.768)
Severity of decline	5.72	6.35	11.04	0.701	0.663
	(5.60, 5.85)	(6.23, 6.48)		(0.688, 713)	(0.656, 0.670)
Number declining	10.62	9.94	-6.37	0.738	0.627
	(10.45, 10.78)	(9.78, 10.10)		(0.728, 0.748)	(0.620, 0.634)
Diversity/Severity of decline	7.20	8.41	16.79	0.269	0.890
	(4.87, 9.53)	(5.97, 10.85)		(0.127, 0.411)	(0.820, 0.960)
Diversity/number declining	7.94	7.37	-7.17	0.616	0.627
	(7.81, 8.07)	(7.25, 7.50)		(0.604, 0.628)	(0.618, 0.635)

Response	HF 1993	HF 2009	ΔHF	LC diversity	Proportion forest
Severity of decline/number declining	8.00	8.84	10.50	0.633	0.753
	(6.89, 9.12)	(7.69, 9.99)		(0.550, 0.716)	(0.709, 0.797)
All targets	8.15	9.48	16.31	0.339	0.856
	(5.12, 11.19)	(6.31, 12.64)		(0.156, 0.522)	(0.765, 0.947)
Region	9.09	9.01	-0.88	0.778	0.465
	(9.04, 9.14)	(8.96, 9.06)		(0.774, 0.781)	(0.463, 0.468)

Table 3. Comparison of protected area coverage for priority landscapes selected by each target.

Values show the percent of selected landscapes for each target that contained high (strict

nature reserve, wilderness area), medium (national park, national monument, habitat/species

management) and low (protected landscape/seascape, managed resource protected area)

categories. The number of selected landscapes for each target are shown after the target

name. See Appendix A for the proportion of target landscapes for all 7 individual protected

area classes.

Response	High	Medium	Low	Any
Diversity (9609)	1.63	6.68	13.34	19.44
Severity of decline (9726)	0.56	11.56	6.89	18.12
Number declining (10922)	1.64	6.92	12.42	18.86
Diversity/severity of decline (35)	0.00	28.57	28.57	51.43
Diversity/number declining (6622)	2.01	8.17	14.97	22.38
Severity of decline/number declining (155)	0.00	9.68	9.68	18.71
All targets (26)	0.00	26.92	26.92	50.00
Non-breeding region (96079)	2.52	5.53	9.78	15.93

587
588

Figure 1. An example eBird landscape approximately 8.3 × 8.3 km in southwest Costa Rica.

Using STEM distribution maps for 112 Neotropical migrants we predicted that this landscape

would have a Shannon diversity = 3.04, declining species richness = 26 and a median population

decline across species = 34.2% between 1966 and 2015. Forest cover in this unprotected

landscape was measured at 67% in 2016 and the human footprint index increased by 15.7%

between 1993 and 2009. We estimated these metrics for 96,079 landscapes across the non-

breeding range of Neotropical migrants in the western hemisphere.

Figure 2. Fine-scale patterns (top row) and priority landscapes (bottom row) for Neotropical

migrants based on high diversity (a,b), average severity of population decline (c,d) and declining

species richness (e,f). Priority landscapes in the bottom row are based on the top 10% of values

for each target. The stationary non-breeding region was defined by landscapes with at least 5

species present between Nov 14 and Mar 14 and is shown in light green in the bottom panel

figures.