Pathogen Detection and Microbiome Analysis of Infected Wheat Using a Portable DNA

Sequencer

Yiheng Hu and **Gamran S. Green**, Research School of Biology, The Australian National University,

Acton ACT2601, Australia; **Andrew W. Milgate**, Wagga Wagga Agricultural Institute, New South

Wales Department of Primary Industry, Wagga Wagga NSW 2650, Australia; **Eric A. Stone**, ANU-

CSIRO Centre for Genomics, Metabolomics and Bioinformatics, Acton ACT2601, Australia; **John**

P. Rathjen† and **Benjamin Schwessinger**†, Research School of Biology, The Australian National

University, Acton ACT2601, Australia.

†Corresponding authors: B. Schwessinger; E-mail: benjamin.schwessinger@anu.edu.au, and J. P.

Rathjen; Email: john.rathjen@anu.edu.au

Funding: This project was supported by The Hermon Slade Foundation, The Australian National

University and Australian Research Council.

Current address of G. S. Green: School of Biotechnology and Biomolecular Sciences, The

University of New South Wales, Kensington, NSW2052, Australia.

ABSTRACT

Fungal diseases of plants are responsible for major losses in agriculture, highlighting the need

for rapid and accurate identification of plant pathogens. Disease outcomes are often defined

not only by the main pathogen but are influenced by diverse microbial communities known as

the microbiome at sites of infection. Here we present the first use of whole genome sequencing

from a portable DNA sequencing device as a method for the detection of fungal pathogens

from wheat (*Triticum aestivum*). The data revealed that our method is robust and applicable to

the diagnosis of fungal diseases including wheat stripe rust (caused by *Puccinia striiformis* f. sp.

tritici), septoria tritici blotch (caused by *Zymoseptoria tritici*) and yellow leaf spot (caused by

Pyrenophora tritici repentis). We also identified a bacterial genus *Pseudomonas* co-present with

Puccinia and *Zymoseptoria* infections but not *Pyrenophora* infections. One limitation of the

method is the over-representation of redundant wheat genome sequences in the sample. This

could be addressed by amplicon-based sequencing approaches in future studies. Our work

outlines a new approach for the detection of a broad range of plant pathogens and associated

microbes using a portable sequencer, providing the basis for future development of a system

for on-site disease monitoring.

Plant diseases, especially those caused by fungal pathogens, jeopardise global crop biosecurity.

Increased global trade, human migration, environmental changes and the accelerated

emergence of virulence have been identified as causes for increasing prevalence of diseases in

plants and animals (Fisher et al. 2012). To prevent and manage diseases, rapid detection and

identification of their causal agents are crucial. This is particularly important for outbreak

management during the incursion of virulent new isolates that overcome classical methods of

control such as the use of resistant crop varieties (Bhattacharya 2017; Pretorius et al. 2000).

Traditional methods for crop disease diagnosis rely largely on the expertise of pathologists who

identify diseases initially by eye in the field. While molecular methods of pathogen detection

also exist, these are generally only capable of identifying a limited number of species (De Boer

and López 2012). Recently, other methods based on the use of optical sensors have been

developed for the detection, identification, and quantification of plant diseases (Mahlein 2015).

Conventional pathology, however, remains the primary means of pathogen identification and

harbors certain limitations: 1. It relies heavily on the physical appearance of disease symptoms;

2. It has difficulty detecting pathogens which do not infect aerial parts of the plant or prior to

the emergence of visible disease symptoms; 3. Co-infections may lead to conflicting visual

symptoms; 4. No method currently exists for the rapid identification of unknown pathogens

during outbreaks – a limitation epitomized by the recent, devastating incursion of South

American races of wheat blast fungus in Bangladesh (Callaway 2016).

DNA sequencing-based methods have been applied to address limitations to detection and

monitoring of environmental microbial communities (Nezhad 2014). Such assays rely largely on

amplification and sequencing of small conserved regions called amplicons from pathogen

genomes such as the internal transcribed spacer region (ITS), *Elongation Factor 1 alpha* (*EF1α*)

and *Beta-tubulin* genes for fungal species and 16S ribosomal RNA sequences for bacteria (Janda

and Abbott 2007; Raja et al. 2017). Closely-related species can be discriminated based on single

nucleotide polymorphisms (SNPs) within the amplicon sequences. This has been used for

example to separate risks and different epidemiological or regulatory outcomes posed by the

species in question (Chemidlin Prévost-Bouré et al. 2011; Thonar, Erb, and Jansa 2012).

Amplicon sequencing also allows characterisation of the microbial community - the microbiome

- within a sample. The microbiome can impact plant function for example on shifting flowering

time (Panke-Buisse et al. 2015) and promoting the colonization of fungal pathogen such as

Botrytis cinerea (Ritpitakphong et al. 2016). The microbiome is likely to contain organisms that

interact with disease-causing agents that could potentially be harnessed for biocontrol

strategies (Ellis 2017). However, restrictive read lengths generated by the Illumina sequencing

platform limits sequence-based identification of microbial species. Only 500 base pairs (bp) of

each DNA amplicon can be sequenced in each reaction. This means that only that genetic

variation present within the 500 bp is available for taxonomic assignment of the organism in

question (Benítez-Páez and Sanz 2017). Use of PCR also biases abundance of individual

amplicons, making estimation of population frequencies problematic (Kennedy et al. 2014).

Third-generation DNA sequencing platforms have developed rapidly during the last half-decade,

overcoming the read length issue inherent in second-generation sequencing methods and

facilitating the identification of microbial species (Benítez-Páez and Sanz 2017). Two competing

long-read sequencing technologies are currently in use. Single molecular real-time (SMRT)

sequencing (PacBio) enables detection of the incorporation of a fluorescent nucleotide into a

single molecule during DNA synthesis within a nanoscale observation chamber (Ardui et al.

2018). Nanopore sequencing (MinION; Oxford Nanopore Technologies (ONT)) enables direct

sequencing of native DNA by measuring the voltage change across an artificial membrane when

a single DNA molecule passes through a nanopore embedded in the membrane within a

flowcell (Jain et al. 2016). Advantages of nanopore sequencing include its relatively low cost,

pocket-size form, and real-time data processing. This means that it is a mobile technology

independent of large sequencing facilities, and as such has been useful for on-site sample

sequencing in extreme environments such as microbial paleo mats in the Antarctic (Johnson et

al. 2017), or providing rapid bacterial pathogen identification from clinical samples with a

turnaround time of six hours (Charalampous et al. 2018), both of which could not be achieved

with previous technologies. Most importantly, this technology provides dramatically longer

sequencing reads (~10 kb) than second-generation sequencing platforms such as Illumina.

Longer reads contain more genetic information, allowing more accurate identification of the

organism from which the sequence is derived (Benítez-Páez, Portune, and Sanz 2016; Benítez-

Páez and Sanz 2017). Although recent studies have shown important advances in identification

of bacterial pathogens using nanopore sequencing from clinical samples (Mitsuhashi et al. 2017;

Schmidt et al. 2017), little has been done in the detection of plant pathogens and microbiome

profiling of diseased plants using this approach.

Here, we tested nanopore sequencing for disease diagnosis in field-grown wheat and explored

its potential for microbiome profiling. We designed a whole-genome sequencing workflow for

the identification of major pathogen species from field samples, as well as the profiling of

microbes present in these samples to the species level. As a proof-of-concept, we applied our

sequencing workflow to wheat leaves infected with known fungal pathogens sampled from the

field, and a fungicide treated control. By mapping sequencing reads to databases, we

successfully detected all fungal diseases present in the infected samples including stripe rust

(caused by *Puccinia striiformis* f. sp. *tritici*), septoria tritici blotch (caused by *Zymoseptoria tritici*)

and yellow leaf spot (caused by *Pyrenophora tritici repentis*). We also conducted a preliminary

characterisation of the microbiomes associated with those diseases, and found bacterial genus

Pseudomonas co-present with stripe rust and septoria tritici blotch. Our results suggest this

technology has considerable potential for adaptation to a broad range of crop disease

diagnoses and environmental monitoring applications.

MATERIALS AND METHODS

Sample collection

Wheat cultivar Crusader (Advanta Seeds Pty. Ltd.) seeds were planted in the disease nursery

field on May 2016 at the Wagga Wagga Agricultural Institute (New South Wales (NSW),

35°02'24.3"S 147°19'09.5"E) with daily irrigation. Wheat straw infected the previous season

with septoria tritici blotch and yellow leaf spot separately then allowed to stand in the field to

promote development of pseudothecia were used to manually inoculate plants in the nursery

field at the tillering growth stage (4-5 weeks post emergence) while stripe rust infections

occurred naturally at jointing stage (7-9 weeks post emergence). Healthy or diseased plants

were diagnosed via symptomatology by a plant pathologist from the NSW Department of

Primary Industries before leaf samples were collected during September 2016. For sampling, an

entire tiller from each plant was removed and stored on ice. Samples from three separate

plants infected with the same disease were pooled and wrapped in aluminium foil before

shipping to the Rathjen laboratory, Research School of Biology, The Australian National

University. Two representative leaves from each symptomatology group containing visible

disease symptoms or fungicide treated control were cut into pieces roughly weighing 100 mg

and transferred 2 mL Eppendorf tubes containing one 5 mm metal bead. All tubes were labelled

and stored at -80^{o}C prior to DNA extraction.

DNA extraction

The DNA extraction protocol is described in (Hu 2016). Briefly, 100 mg leaf tissue was

homogenized and cells were lysed using cetyl trimethylammonium bromide (CTAB, Sigma-

Aldrich) buffer (added RNAse T1, Thermo Fisher, 1000 units per 1750 µL), followed by a

phenol:chloroform:isoamyl alcohol (25:24:1, Sigma-Aldrich) extraction to remove protein and

lipids. The DNA was precipitated with 700 µL isopropanol, washed with 1 mL 70 % ethanol

before dried for 5 minutes at room temperature and resuspended in TE buffer containing 1mM

Tris and 1mM EDTA at pH8. DNA extractions were performed four times independently, with

the operator blind to the identity of each sample. After extraction, DNA was purified with one

volume of Agencourt AMPure XP Beads (Beckman Coulter, Inc.) using the manufacturer's

protocol and stored at 4°C. Quality and average size of genomic DNA was visualized by gel

electrophoresis using 1% agarose gel for 1 hour at 100 volts. DNA was quantified by NanoDrop

and Qubit (Life Technologies) following the manufacture's protocol.

Library construction and sequencing

Sequencing libraries were constructed as previously described (Hu and Schwessinger 2018) with

the omission of the DNA shearing step. Briefly, a dA-tailing reaction was performed on 700 ng

genomic DNA from each sample using the NEBNext Ultra II End-repair/dA-tailing kit (New

England Biolabs). Repaired DNA was washed on AMPure XP Beads on a magnetic rack using 70 %

ethanol, and eluted with 31 µL nuclease-free water. Thirty µL eluate was used for barcode

adapter ligation via the NEB Blunt/TA Ligase Master Mix (New England Biolabs) and the Barcode

Adapter Mix (ONT), followed by another wash step on AMPure XP Beads. Twenty ng of the

adapter-ligated template in 2 µL of nuclease free water was added to each barcoding PCR

reaction with 50 µL LongAmp Taq 2X Master Mix (New England Biolabs), 46 µL water and 2 µL

PCR Barcodes (ONT). The PCR was performed as follows: Denaturation at 95°C for 3 min,

followed by 15 cycles of 95°C/15 s; 62°C/15 s; 65°C/150 s, and a final extension step at 65°C for

5 min. Barcoded DNAs were purified using 1 volume of AMPure XP Beads then pooled into 1 µg

fractions in a total of 45 µL to which 5 µL of DNA CS control (ONT) was added. A further dA-

tailing reaction was performed as above followed by purification on AMPure XP Beads, with the

DNA eluted in 31 µL nuclease free water. Adapter ligation was carried out using 30 µL end-

prepped DNA, 20 µL Adapter Mix (ONT) and 50 µL NEB Blunt/TA Ligase Master Mix. The

adapter-ligated DNA library was then purified on AMPure XP Beads using the Adapter Beads

Binding buffer (ONT) for washing, and eluted in 15 µL of Elution Buffer (ONT).

Sequencing reactions were performed on a MinION Flowcell (R9.4, ONT) connected to a MK1B

device (ONT) operated by the MinKNOW software (version 1.5.2 and version 1.6.11). Each

Flowcell was primed with 1 mL of priming buffer which consists of 480 µL Running Buffer Fuel

Mix (RBF, ONT) and 520 µL nuclease free water. 12 µL of amplicon library was added to a

loading mix including 35 µL RBF, 25.5 µL Library Loading Beads (ONT) and 2.5 µL water for a

final volume of 75 µL added to the Flowcell via the SpotON sample port. The

"NC_48Hr_sequencing_FLO-MIN106_SQK-LSK107" protocol was executed by the MinKNOW

after loading the library. The sequencing run was restarted after 12 h and stopped after 48 h.

Sequencing analysis

Raw fast5 files were processed by the Albacore 2.0.2 software (ONT) for basecalling, barcode

de-multiplexing and quality filtering (Phred quality score of >7). All reads that passed quality

filtering for each barcode were treated in parallel as follows: firstly, NanoLyse (De Coster et al.

2018) was used to remove lambda phage control DNA sequences; secondly, barcode and

adapter sequences were trimmed from the ends of reads using Porechop

(https://github.com/rrwick/Porechop). To identify middle adapter sequence by Porechop, a 95%

threshold was set, and reads with middle adapter sequences were split into two sequences;

thirdly, seqtk (https://github.com/lh3/seqtk) was used to convert the processed fastq file into

fasta format for further analysis. Two BLAST (Basic Local Alignment Search Tool, Altschul et al.,

1990) searches were performed on the fasta files for each barcode: first, against a wheat

reference genome (Clavijo et al. 2017, accessed 09/2017) and a custom database containing the

genomes of the four fungal species responsible for the diseases studied here: *Puccinia*

striiformis f. sp. *tritici*: Schwessinger et al. (2018), accessed 09/2017; *Z. tritici*:   Goodwin et al.

(2011), version 2, accessed 09/2017; *Pyrenophora tritici-repentis*: Manning et al. (2013),

unmasked assembly, accessed 09/2017. The genome of a further species suggested to be

absent from the studied wheat growing region, *Parastagonospora nordorum*, was included as a

negative control (Hane et al. 2007, accessed 09/2017). Fasta files representing raw sequence

reads were searched against the reference genomes using BLAST. The e-value cut-off for each

BLAST search was set to 0.01, and only the top hit was taken for further analysis. Reads that did

not hit the reference genomes were captured using the filterbyname.sh script from the bbmap

script package (https://sourceforge.net/projects/bbmap/). These reads were searched against

the National Center for Biotechnology Information (NCBI) nucleotide (nt) database

(ftp://ftp.ncbi.nlm.nih.gov/blast/db/) using BLAST with the same settings as previously. Those

reads providing hits were separated and retained again using filterbyname.sh script. Two

Python scripts (QC_and_BLAST.py and creating_final_dataframe.py) that combine all steps from

quality control to BLAST outputs and construct dataframes containing statistical and

information   from   BLAST   searches   for   each   read   were   deposited   at

https://github.com/Yiheng323/Pathogen_Detection_and_Microbiome_scripts. All   basecalled

fastq data for each sequencing run were deposited at NCBI Short Reads Archive (SRA) under the

accession number PRJNA493553. The sequencing summary files and final dataframes as .tab files

for each replicate were deposited at https://doi.org/10.6084/m9.figshare.7138262. The python

scripts used for generating Fig. 2 – 5, Supplementary Fig. S1 – S5 were also deposited at

https://github.com/Yiheng323/Pathogen_Detection_and_Microbiome_scripts.

RESULTS

Sequencing showed variations in yield and consistency in barcode classification

We designed a pipeline for identification of three known wheat pathogens from infected field

samples using MinION sequencing. We infected three experimental field plots as disease

nurseries with their corresponding pathogens (*Puccinia. striiformis* f. sp. *tritici* for stripe rust

disease, *Z. tritici* for Septorial tritici blotch and *Pyrenophora. tritici-repentis* for Yellow leaf spot)

commonly found in New South Wales, Australia, and conducted sampling during the 2016-17

growing season (September). The sequencing experiment included three steps (Fig. 1): 1.

Pathologists collected leaf samples from each disease nursery field, with visual confirmation of

disease symptoms from the three single infections plus one double infection (*Puccinia*

striiformis f. sp. *tritici* and *Z. tritici*), and a negative control which was the same wheat variety

treated with commercial fungicides at the 10- to 11-week growth stage (three months post

germination), before shipping to the laboratory on ice for processing. 2. We extracted DNA

without knowing the identity of each sample to eliminate potential bias from operators. 3. We

barcoded each sample to enable pooling of all five samples to reduce sequencing costs. The five

wheat leaf samples were barcoded as follows: leaves infected with one of the three fungal

pathogens (barcodes 1-3); leaves infected with two pathogens (barcode 4); and no visible

infection (barcode 5). We performed step 2 and 3 four times as biological replicates, using a

MinION 1D PCR barcoding workflow (EXP-PBC001 and SQK-LSK107). The raw data were

converted into sequence reads using the software Albacore, with a quality (Q) score for each

base estimating the likelihood of the identified base being correct using the Phred scale (Ewing

et al., 1998; Ewing and Green, 1998). Based on read mean Q scores, we grouped reads into pass

(Q score > 7) or fail (Q score <= 7). We binned reads based on the barcode sequences identified

by Albacore at both ends of each sequence. If no barcode sequence could be identified, we

named the corresponding read as unclassified. Overall, ~92-95% of the total sequence length of

all four sequencing runs passed quality filtering (Supplementary Table 1). The total sequence

length from the four runs varied from 682 to 5100 megabase pairs (Mbp). The mean read

length ranged from 1380 to 3054 bp. Proportions of each barcode in each run were relatively

even, although 23.3-25.7% of the sequences were unclassified from all four runs (Fig. 2).

Sequence-based identification correctly identifies the fungal species causing wheat diseases

We performed Basic Local Alignment Search Tool (BLAST, Altschul et al., 1990) searches for all

quality-filtered reads against a manually constructed reference database. This contained the

wheat genome and four fungal genomes including *Puccinia striiformis* f. sp. *tritici, Z. tritici,*

Pyrenophora tritici-repentis and *Parastagonospora nodorum*. These represent the host plant

genome, three of the fungal species present in the field samples, and one fungal species

(*Parastagonospora nordorum*) not expected to be present in the sampling area included as a

negative control. We took the hit with the lowest expectation value (e-value) from BLAST

analysis of each read as it represented the most significant match. The majority of reads are

wheat sequences, comprising total sequence length of 90.1% in flowcell 1, 80.1% in flowcell 2,

92.7% in flowcell 3 and 91.4% in flowcell 4 (Fig. 3). We found that healthy leaf samples (barcode

5) contained no or minor total length percentages of pathogen-related sequences in all four

replicates (< 0.01%, Fig. 4). This indicates that our analysis method is robust and does not lead

to high background signals in healthy wheat samples (barcode 5). Significantly higher

proportions of total sequencing length (0.5–5.7%) from infected samples (barcode 1-4) were

assigned as derived from pathogen genomes. In all cases, the pathogen genome identified by

the highest total sequence length corresponded to the pathogen causing the disease as

identified by symptoms (Fig. 4). This indicates that we are clearly able to identify the disease

causing pathogen via our sequencing workflow in the case of single infections. Infections with

two pathogens (barcode 4) were identified less clearly. In barcode 4 from all replicates, the

total sequence lengths assigned to *Puccinia striiformis* f. sp. *tritici* were between 0.1% and 0.4%,

which were close to or lower than the proportion of total length assigned to the negative

control species *Parastagonospora nodorum.* The percentages of total sequence length by

barcode that identified pathogen genomes were similar across all replicates, with around 0.6%

of barcode 1 sequences from *Puccinia striiformis* f. sp. *tritici*, 1.9-5.0% of barcode 2 sequences

from *Z. tritici*, 2.1-5.7% of barcode 3 sequences from *Pyrenophora tritici-repentis*, and 0.1-0.4%

of barcode 4 sequences contained *Puccinia striiformis* f. sp. *tritici* and 0.6-1.2% from *Z. tritici.*

Unexpectedly, we also identified *Parastagonospora nodorum* from different samples in all four

technical sequencing replicates. To ascertain whether *Parastagonospora nodorum* was truly

present, we extracted all reads that hit the *Parastagonospora nodorum* genome by BLAST

analysis against the reference database. We used these reads to perform an open BLAST search

against the entire NCBI non-redundant (nt) nucleotide database which contains all publically

available genome sequences. We found that 344 of 915 reads were assigned to genus

Parastagonospora, including 339 reads identified *Parastagonospora nodorum* SN15 genome

(Supplementary Fig. S1). A further 371 reads hit related species, including 226 from the same

Pleosporales order: *Leptosphaeria* spp. (84 reads), *Alternaria* spp. (81), *Pyrenophora* spp. (10),

Bipolaris spp. (9), *Shiraia* spp. (9) and other genus (Supplementary Fig. S1). The remaining 200

reads fell under the given e-value threshold (0.01) and failed to be classified. The difference

between both BLAST searches potentially stems from the relatively high error rate of the

MinION sequencer (Jain et al., 2018; Laver et al., 2015). Nevertheless, this result supports the

finding that *Parastagonospora nodorum* is present in the sample from the Wagga Wagga wheat

growing region despite the disease not purposefully inoculated in the trial.

Bacterial species associated with fungal disease development

To obtain an overall picture of the microbiome profile in wheat leaves infected by different

fungal pathogens, we attempted to identify other microbial species present on the leaves. To

do this, we searched the non-redundant NCBI nucleotide database with reads that were not

assigned to any of the five reference genomes present in the initial restricted database. We

found that the proportion of reads assigned to a genome in the restricted and the nt-NCBI

databases were high for flowcells 1, 3 and 4 (92.9, 94.3 and 93.8%, respectively), but only 84.6%

for flowcell 2, likely due to technical issues. Overall, fewer than 1.4% of passed reads from each

flowcell identified microbiota other than the major pathogens in the restricted database in and

on the sampled wheat leaves.

We pooled the reads from each barcode in all four biological replicates, and calculated the

percentage of total length that identified a specific genus in each sample using NCBI taxonomic

identifiers (Fig. 5A). Genus *Pseudomonas* took up the highest proportion of reads in all diseased

samples, ranging from 0.08% to more than 0.88%, but was nearly absent in healthy wheat

(barcode 5, 0.01%). Genus *Alternaria* is also found in all samples with a proportion between

0.05% to 0.2% in diseased samples and 0.005% in healthy wheat. Uniquely in barcode 1,

bacterial *Erwinia* comprised 0.09% of the total read length as represented by 219 reads and the

second highest proportion. In contrast, *Erwinia* only represented 0.003% and 0.001% in barcode

2 and 5, and none in barcode 3 and 4, while *Alternaria* ranked second highest in other diseased

samples (barcode 2 - 4). In barcodes 2 and 4, hits on *Z. tritici* were still present and may

represent the six *Z. tritici* strains genomes in the NCBI nt database not included in the *Z. tritici*

genome we used in the first BLAST analysis. At the species level (Fig. 5B), reads identifying

Pseudomonas syringae were much more frequent in samples infected with *Puccinia striiformis* f.

sp. *tritici* (barcode 1, 0.59%) and/or *Z. tritici* (barcode 2, 0.33%). This was not seen in the sample

infected with *Pyrenophora tritici-repentis* (barcode 3), where only 0.02% reads matched

Pseudomonas syringae. Populations of other *Pseudomonas* species such as *Pseudomonas poae*

followed a similar trend to the relative *Pseudomonas syringae* accross different wheat samples,

presenting much more in barcode 1, 2, 4 than in barcode 3 and 5. This finding corresponds to

other reports in which *Pseudomonas* spp. were associated with fungal infections of wheat (Al-

Sallami et al. 1997; Mehrabi et al. 2016). Considering the upper taxonomic classifications,

Proteobacteria and Ascomycota were the most abundant bacterial and fungal Phylums across

all samples (Supplementary Fig. S2) followed by Actinobacteria and Basidiomycota. At the Class

level (Supplementary Fig. S3), either Gammaproteobacteria or Dothiodeomycetes were the

most abundant. Overall, the microbiome structure of the upper taxonomy resembles the lower

taxonomy (Supplementary Fig. S2-S5), since most sequences were assigned to genus

Pseudomonas.

DISCUSSION

We have developed a WGS workflow which accurately identified three fungal pathogens

responsible for three major wheat diseases - stripe rust, septoria tritici blotch and yellow spot -

from field samples. We designed a two-step BLAST process to identify major agents in diseased

samples and characterise the associated microbiota. We identified a number of *Pseudomonas*

species in stripe rust and septoria tritici blotch infected wheat samples but not in yellow spot or

non-diseased samples.

WGS provides a wealth of information that could be used to accurately identify disease-causing

agents, and to begin to describe the composition of the microbial community (Brown et al.

2017; Thomas, Gilbert, and Meyer 2012). Without any amplification process which potentially

includes biases, the WGS data is obtained directly from the sample extracts, illustrated the

most close-to-reality features dependent only by sampling strategies. WGS requires less

processing time, omitting the PCR step, providing advantageous for on-site diagnostic purposes

where technical simplicity and fast processing time are prioritised. However, it has a significant

drawback in that most sequences are derived from the host (wheat) genome, representatively

90.1% in flowcell 1, 80.1% in flowcell 2, 92.7% in flowcell 3 and 91.4% in flowcell 4. This is due

to our DNA extraction method in which the whole leaf sample was grinded to retain all the

microbial population in and on the leaf. For the purpose of microbial species identification,

host sequences are unwanted and can overwhelm the community DNA (Sharpton 2014). To

enrich microbial sequences, one strategy is to apply intricate molecular methods such as

gradient centrifugation and sonication during the extraction process (Bulgarelli et al. 2012; Lim

et al. 2014). This can be challenging when applying to environmental samples, especially in field

conditions, as these techniques require sophisticated instruments. Another strategy to enrich

microbiome sequences involves sequencing microbial marker gene amplicons. With the

improvement of sequencing length, species level resolution using long read amplicon

sequencing can be achieved in few cases (Benítez-Páez and Sanz 2017; Martijn et al. 2017).

Amplicon sequencing seems to be a promising method for species identification, as it requires

less input DNA than WGS and marker gene databases contain information from many more

species than whole genome databases (Cruaud et al. 2017; Tessler et al. 2017; Ranjan et al.

2016). Amplicon sequencing is also possible in applied cases given remote PCR platforms are

rapidly developing, such as the miniPCR system (Amplyus, Cambridge, USA) and the Bento Lab

platform (Bento Bioworks Ltd, London, United Kingdom). Drawbacks of amplicon strategy

include biases related to PCR amplification, which lead to inaccuracy when calculating the

relative proportions of microbial species.

While many workflows are available for microbiome profiling using metagenomics data, there is

no standard method for analysing long read data. This makes validation of both methods and

results crucial (Quince et al. 2017). To validate the method, mock microbial communities or

samples containing known microbial species were commonly used for proof-of-concept. For

example, to verify the diagnosis of a specific pathogen strain based on SNPs called from MinION

reads, Votintseva et al. (2017) first identified pathogens using clinical methods then generated

sequencing data and mapped the data to the targeted species in real-time. The reactions were

stopped once enough SNPs were obtained to confidentially identify the targeted pathogen

strain. Similarly, our study used wheat samples manually inoculated with infections and visually

confirmed known pathogens prior to sequencing, while the same identifications from the first

BLAST analysis validated the capacity of our workflow for identification of those pathogen

species. To validate the result, different analysis approaches should be applied on the same

dataset. Brown et al. (2017) compared the performance of four bioinformatics pipelines on

identifying species within different synthetic bacterial communities using MinION. Compared to

the composition of mock community, even the pipeline that resulted in the most similar

classification still contains reads classified to species known not to be present. These false

positive results are due largely to sequencing errors, which illustrates the need for more

stringent filtering of the raw data.

Nanopore metagenomics will have great potential for quick diagnosis of suddenly emerging

crop disease and large-scale disease monitoring at centralized agricultural institutions. As

shown from other sequencing studies performed in a more extreme environment (Johnson et al.

2017; Pomerantz et al. 2018), all processes including DNA extraction and purification, library

preparation and sequencing can be completed within a few hours. A tailored database can be

created for each crop species containing the genomes of pathogens known to cause diseases to

minimise processing time. For rapidly evolving pathogens such as wheat rusts, a SNP-based

classification method as described by Votintseva et al. (2017) can be adapted to define rust

strains once enough SNPs are obtained. However, before running the sequencer, sampling

procedures and tissue processing strategies still need to be carefully curated. These vary in crop

species and tissue types, so economic efficiency and statistical significance should be

considered during the sampling process (Knight et al. 2012). Our study randomly sampled

multiple leaves for each sample and extracted four times as technical replicates. Consistency in

barcode classification and BLAST analysis through all four replicates indicated reliability of the

workflow. In centralized agricultural institutions, amplicon sequencing can potentially play a key

role and, to-date, there have been a few preliminary successes (Benítez-Páez, Portune, and

Sanz 2016; Benítez-Páez and Sanz 2017; Pomerantz et al. 2018). Given the multiplexing capacity

of the 96 native barcoding kit (ONT), hundreds of samples can be sequenced in parallel when

larger scale sequencing platform were employed (e.g. GridION, ONT). To achieve this goal, long

read databases, sequencing accuracy and access to large-scale sample processing power need

to be improved.

In conclusion, our workflow demonstrates the potential of this technology for plant pathogen

diagnosis with the potential for field applications, with the additional possibility of microbiome

characterization. A combination of on-site and centralized sequencing approaches would, in

future, revolutionize management of agricultural biosecurity and prevent crop loss to meet the

increasing food demand of human beings.

LITERATURE CITED

Al-Sallami, F., Karov, S., Vassileva, P., Popova, R., and Vassilev, V. 1997. Pseudomonas syringae

pv. atrofaciens associated with fungal black point of wheat (Triticum aestivum) grain. In

Pseudomonas Syringae Pathovars and Related Pathogens, Springer, p. 505–508.

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. 1990. Basic local alignment

search tool. J. Mol. Biol. 215:403–410.

Ardui, S., Ameur, A., Vermeesch, J. R., and Hestand, M. S. 2018. Single molecule real-time

(SMRT) sequencing comes of age: applications and utilities for medical diagnostics. Nucleic

Acids Res. 46:2159–2168.

Benítez-Páez, A., Portune, K. J., and Sanz, Y. 2016. Species-level resolution of 16S rRNA gene

amplicons sequenced through the MinION^{TM} portable nanopore sequencer. GigaScience. 5:4.

Benítez-Páez, A., and Sanz, Y. 2017. Multi-locus and long amplicon sequencing approach to

study microbial diversity at species level using the MinION^{TM} portable nanopore sequencer.

GigaScience. 6:1–12.

Bhattacharya, S. 2017. Deadly new wheat disease threatens Europe's crops. Nat. News. 542:145.

Brown, B. L., Watson, M., Minot, S. S., Rivera, M. C., and Franklin, R. B. 2017. MinION^{TM}

nanopore sequencing of environmental metagenomes: a synthetic approach. GigaScience. 6:1–

10.

Bulgarelli, D., Rott, M., Schlaeppi, K., Themaat, E. V. L. van, Ahmadinejad, N., Assenza, F., et al.

2012. Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial

microbiota. Nature. 488:91–95.

Callaway, E. 2016. Devastating wheat fungus appears in Asia for first time. Nat. News. 532:421.

Charalampous, T., Richardson, H., Kay, G. L., Baldan, R., Jeanes, C., Rae, D., et al. 2018. Rapid

Diagnosis of Lower Respiratory Infection using Nanopore-based Clinical Metagenomics.

bioRxiv. :387548.

Chemidlin Prévost-Bouré, N., Christen, R., Dequiedt, S., Mougel, C., Lelièvre, M., Jolivet, C., et al.

2011. Validation and application of a PCR primer set to quantify fungal communities in the soil

environment by real-time quantitative PCR. PloS One. 6:e24166.

Clavijo, B. J., Venturini, L., Schudoma, C., Accinelli, G. G., Kaithakottil, G., Wright, J., et al. 2017.

An improved assembly and annotation of the allohexaploid wheat genome identifies complete

families of agronomic genes and provides genomic evidence for chromosomal translocations.

Genome Res. 27:885–896.

Cruaud, P., Rasplus, J.-Y., Rodriguez, L. J., and Cruaud, A. 2017. High-throughput sequencing of

multiple amplicons for barcoding and integrative taxonomy. Sci. Rep. 7:41948.

De Boer, S. H., and López, M. M. 2012. New grower-friendly methods for plant pathogen

monitoring. Annu. Rev. Phytopathol. 50:197–218.

De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M., Van Broeckhoven, C., and Berger, B. 2018.

NanoPack: visualizing and processing long-read sequencing data. Bioinformatics. 34:2666–2669.

Ellis, J. G. 2017. Can Plant Microbiome Studies Lead to Effective Biocontrol of Plant Diseases?

Mol. Plant. Microbe Interact. 30:190–193.

Ewing, B., and Green, P. 1998. Base-Calling of Automated Sequencer Traces Using Phred. II.

Error Probabilities. Genome Res. 8:186–194.

Ewing, B., Hillier, L., Wendl, M. C., and Green, P. 1998. Base-Calling of Automated Sequencer

Traces UsingPhred. I. Accuracy Assessment. Genome Res. 8:175–185.

Fisher, M. C., Henk, D. A., Briggs, C. J., Brownstein, J. S., Madoff, L. C., McCraw, S. L., et al. 2012.

Emerging fungal threats to animal, plant and ecosystem health. Nature. 484:186–194.

Goodwin, S. B., M'Barek, S. B., Dhillon, B., Wittenberg, A. H. J., Crane, C. F., Hane, J. K., et al.

2011. Finished Genome of the Fungal Wheat Pathogen Mycosphaerella graminicola Reveals

Dispensome Structure, Chromosome Plasticity, and Stealth Pathogenesis. PLOS Genet.

7:e1002070.

Hane, J. K., Lowe, R. G. T., Solomon, P. S., Tan, K.-C., Schoch, C. L., Spatafora, J. W., et al. 2007.

Dothideomycete–Plant Interactions Illuminated by Genome Sequencing and EST Analysis of the

Wheat Pathogen Stagonospora nodorum. Plant Cell. 19:3347–3368.

Hu, Y. 2016. High quality DNA extraction from Fungi_small scale. protocols.io. Available at:

https://www.protocols.io/view/high-quality-dna-extraction-from-fungi-small-scale-exmbfk6

[Accessed September 26, 2018].

Hu, Y., and Schwessinger, B. 2018. Amplicon sequencing using MinION optimized from 1D

native barcoding genomic DNA. protocols.io. Available at:

https://www.protocols.io/view/amplicon-sequencing-using-minion-optimized-from-1d-

mhkc34w [Accessed September 27, 2018].

Jain, M., Koren, S., Miga, K. H., Quick, J., Rand, A. C., Sasani, T. A., et al. 2018. Nanopore

sequencing and assembly of a human genome with ultra-long reads. Nat. Biotechnol. 36:338–

345.

Jain, M., Olsen, H. E., Paten, B., and Akeson, M. 2016. The Oxford Nanopore MinION: delivery of

nanopore sequencing to the genomics community. Genome Biol. 17:239.

Janda, J. M., and Abbott, S. L. 2007. 16S rRNA Gene Sequencing for Bacterial Identification in

the Diagnostic Laboratory: Pluses, Perils, and Pitfalls. J. Clin. Microbiol. 45:2761–2764.

Johnson, S. S., Zaikova, E., Goerlitz, D. S., Bai, Y., and Tighe, S. W. 2017. Real-Time DNA

Sequencing in the Antarctic Dry Valleys Using the Oxford Nanopore Sequencer. J. Biomol. Tech.

JBT. 28:2–7.

Kennedy, K., Hall, M. W., Lynch, M. D. J., Moreno-Hagelsieb, G., and Neufeld, J. D. 2014.

Evaluating Bias of Illumina-Based Bacterial 16S rRNA Gene Profiles. Appl. Environ. Microbiol.

80:5717–5722.

Knight, R., Jansson, J., Field, D., Fierer, N., Desai, N., Fuhrman, J. A., et al. 2012. Unlocking the

potential of metagenomics through replicated experimental design. Nat. Biotechnol. 30:513–

520.

Laver, T., Harrison, J., O'Neill, P. A., Moore, K., Farbos, A., Paszkiewicz, K., et al. 2015. Assessing

the performance of the Oxford Nanopore Technologies MinION. Biomol. Detect. Quantif. 3:1–8.

Lim, Y. W., Haynes, M., Furlan, M., Robertson, C. E., Harris, J. K., and Rohwer, F. 2014. Purifying

the Impure: Sequencing Metagenomes and Metatranscriptomes from Complex Animal-

associated Samples. JoVE J. Vis. Exp. :e52117–e52117.

Mahlein, A.-K. 2015. Plant Disease Detection by Imaging Sensors – Parallels and Specific

Demands for Precision Agriculture and Plant Phenotyping. Plant Dis. 100:241–251.

Manning, V. A., Pandelova, I., Dhillon, B., Wilhelm, L. J., Goodwin, S. B., Berlin, A. M., et al. 2013.

Comparative Genomics of a Plant-Pathogenic Fungus, Pyrenophora tritici-repentis, Reveals

Transduplication and the Impact of Repeat Elements on Pathogenicity and Population

Divergence. G3 Genes Genomes Genet. 3:41–63.

Martijn, J., Lind, A. E., Spiers, I., Juzokaite, L., Bunikis, I., Pettersson, O. V., et al. 2017. Amplicon

sequencing of the 16S-ITS-23S rRNA operon with long-read technology for improved

phylogenetic classification of uncultured prokaryotes. bioRxiv. :234690.

Mehrabi, Z., McMillan, V. E., Clark, I. M., Canning, G., Hammond-Kosack, K. E., Preston, G., et al.

2016. *Pseudomonas* spp. diversity is negatively associated with suppression of the wheat take-

all pathogen. Sci. Rep. 6:29905.

Mitsuhashi, S., Kryukov, K., Nakagawa, S., Takeuchi, J. S., Shiraishi, Y., Asano, K., et al. 2017. A

portable system for rapid bacterial composition analysis using a nanopore-based sequencer and

laptop computer. Sci. Rep. 7:5657.

Nezhad, A. S. 2014. Future of portable devices for plant pathogen diagnosis. Lab. Chip.

14:2887–2904.

Panke-Buisse, K., Poole, A. C., Goodrich, J. K., Ley, R. E., and Kao-Kniffin, J. 2015. Selection on

soil microbiomes reveals reproducible impacts on plant function. ISME J. 9:980–989.

Pomerantz, A., Peñafiel, N., Arteaga, A., Bustamante, L., Pichardo, F., Coloma, L. A., et al. 2018.

Real-time DNA barcoding in a rainforest using nanopore sequencing: opportunities for rapid

biodiversity assessments and local capacity building. GigaScience. 7.

Pretorius, Z. A., Singh, R. P., Wagoire, W. W., and Payne, T. S. 2000. Detection of Virulence to

Wheat Stem Rust Resistance Gene Sr31 in Puccinia graminis. f. sp. tritici in Uganda. Plant Dis.

84:203–203.

Quince, C., Walker, A. W., Simpson, J. T., Loman, N. J., and Segata, N. 2017. Shotgun

metagenomics, from sampling to analysis. Nat. Biotechnol. 35:833–844.

Raja, H. A., Miller, A. N., Pearce, C. J., and Oberlies, N. H. 2017. Fungal Identification Using

Molecular Tools: A Primer for the Natural Products Research Community. J. Nat. Prod. 80:756–

770.

Ranjan, R., Rani, A., Metwally, A., McGee, H. S., and Perkins, D. L. 2016. Analysis of the

microbiome: Advantages of whole genome shotgun versus 16S amplicon sequencing. Biochem.

Biophys. Res. Commun. 469:967–977.

Ritpitakphong, U., Falquet, L., Vimoltust, A., Berger, A., Métraux, J.-P., and L'Haridon, F. 2016.

The microbiome of the leaf surface of Arabidopsis protects against a fungal pathogen. New

Phytol. 210:1033–1043.

Schmidt, K., Mwaigwisya, S., Crossman, L. C., Doumith, M., Munroe, D., Pires, C., et al. 2017.

Identification of bacterial pathogens and antimicrobial resistance directly from clinical urines by

nanopore-based metagenomic sequencing. J. Antimicrob. Chemother. 72:104–114.

Schwessinger, B., Sperschneider, J., Cuddy, W. S., Garnica, D. P., Miller, M. E., Taylor, J. M., et al.

2018. A Near-Complete Haplotype-Phased Genome of the Dikaryotic Wheat Stripe Rust Fungus

Puccinia striiformis f. sp. tritici Reveals High Interhaplotype Diversity. mBio. 9.

Sharpton, T. J. 2014. An introduction to the analysis of shotgun metagenomic data. Front. Plant

Sci. 5:209.

Tessler, M., Neumann, J. S., Afshinnekoo, E., Pineda, M., Hersch, R., Velho, L. F. M., et al. 2017.

Large-scale differences in microbial biodiversity discovery between 16S amplicon and shotgun

sequencing. Sci. Rep. 7:6589.

Thomas, T., Gilbert, J., and Meyer, F. 2012. Metagenomics - a guide from sampling to data

analysis. Microb. Inform. Exp. 2:3.

Thonar, C., Erb, A., and Jansa, J. 2012. Real-time PCR to quantify composition of arbuscular

mycorrhizal fungal communities--marker design, verification, calibration and field validation.

Mol. Ecol. Resour. 12:219–232.

Votintseva, A. A., Bradley, P., Pankhurst, L., Elias, C. del O., Loose, M., Nilgiriwala, K., et al. 2017.

Same-Day Diagnostic and Surveillance Data for Tuberculosis via Whole-Genome Sequencing of

Direct Respiratory Samples. J. Clin. Microbiol. 55:1285–1298.

529   **Figures and figure legends**

Fig. 1. The Nanopore sequencing workflow for fungal pathogen identification. 1) Sampling:

plant pathologists collected wheat leaf samples from disease nursery fields, with visual

confirmation of disease symptoms for four diseased sample (three single fungal infections and

one double fungal infection). A fungicide treated wheat sample from the same variety and

growth stage was collected as a negative control. 2) DNA sequencing: to minimise operator bias,

DNA was extracted from all five samples without knowledge of the causative agent(s) affecting

each sample. We labelled samples with DNA barcodes through a PCR reaction step before

sample pooling and 1D library preparations (SQK-LSK107); 3) Data analysis: We quality

controlled sequences based on Phred quality scores and trimmed adapter sequence using

Porechop (https://github.com/rrwick/Porechop). We performed BLAST (Altschul et al., 1990)

searches using processed reads and saved the information of the best hits. We summarized the

number of total sequence length from hits by each species using the python *matplotlib* module.

Images used in step 3 are from the official websites of each program.

Fig. 2. Uneven proportions of barcode-labelled DNA across replicates. Columns indicate four

sets of wheat leaf samples collected as biological replicates. The y-axis indicates relative

abundance of total sequence lengths labelled with one of five DNA barcodes within each

sample set. Each barcode represents one wheat sample. Reads without a detectable barcode

were labelled as 'unclassified'.

Fig. 3. Wheat is the predominant identified species in the whole genome sequencing analysis

using BLAST. The graph shows the relative sequence length per species identified using a

'winner-takes-all' approach when searching a targeted BLAST database Wheat (grey), fungal

pathogen genomes (blue), NCBI nucleotide (nt) database (red) and unmapped (light green) (see

results and methods for details). The numbers at the bottom of each bar show the total

sequence length [bp] of processed reads of each replicate. Replicate 1, 3, 4 showed similar

classifications while replicate 2 contains more unclassified sequences.

Fig. 4. The total sequence length per pathogen genome per barcode correlate with the

phenotypic disease identification. The graphs show the relative sequence length assigned to

each pathogen genome using a 'winner-takes-all' approach in the BLAST analysis. The

proportion of reads hitting the wheat host genome are listed in the top-right corner of each

graph. Unassigned reads are omitted from the visualization.

Fig. 5. Fungal pathogen infections of wheat increase the microbial population especially for stripe rust

infection and Septoria tritici blotch infection.   Percentage of sequences that did not hit the

restricted database but hit the non-redundant NCBI nucleotide database combined from all

replicates, and grouped by genus (A) or species (B). Different colours represented different

barcodes. Reads that are assigned to the restricted database was eliminated, and the top ten

highest classifications were displayed for each replicate in the y-axis. Genus *Pseudomonas*

represented the highest proportion across all barcodes containing Stripe rust (*Puccinia*

striiformis tritici) or Septoria tritici blotch (*Z. tritici*), with *Pseudomonas syringae* as the highest

at the species level.

Barcode01 Barcode02 Barcode03 Barcode04 Barcode05

Sampling

DNA sequencing

Quality control & Trimming

Porechop

Nucleotide BLAST

Nucleotide BLAST
nucleotide ▸ nucleotide

Visualization

matplotlib

Data analysis