Widespread gene duplication and adaptive evolution in the RNA interference pathways of the *Drosophila obscura* group

Danang Crysnanto[1,3]*

Darren J. Obbard[1,2]

Affiliation

1. Institute of Evolutionary Biology, University of Edinburgh, Charlotte Auerbach Road, Edinburgh, United Kingdom

2. Centre for Infection, Evolution and Immunity, University of Edinburgh, Edinburgh, United Kingdom

3. Present address: Animal Genomics, ETH Zurich, Zurich, Switzerland

Email

DC: danang.crysnanto@usys.ethz.ch

DJO: darren.obbard@ed.ac.uk

*Author for correspondence

ABSTRACT

Background: RNA interference (RNAi) related pathways provide defense against viruses and transposable elements, and have been implicated in the suppression of meiotic drive elements. Genes in these pathways often exhibit high levels of adaptive substitution, and over longer timescales show frequent gene duplication and loss—most likely as a consequence of their role in mediating conflict with these parasites. This is particularly striking for *Argonaute 2* (*Ago2*), which is ancestrally the key effector of antiviral RNAi in insects, but has repeatedly formed new testis-specific duplicates in the recent history of the *Drosophila obscura* group.

Results: Here we take advantage of publicly available genomic and transcriptomic data to identify six further RNAi-pathway genes that have duplicated in this clade of *Drosophila*, and examine their evolutionary history. As seen for *Ago2*, we observe high levels of adaptive amino-acid substitution and changes in sex-biased expression in many of the paralogs. However, our phylogenetic analysis suggests that co-duplications of the RNAi machinery were not synchronous, and our expression analysis fails to identify consistent male-specific expression.

Conclusions: These results confirm that RNAi genes, including genes of the antiviral and piRNA pathways, undergo frequent independent duplications and that their history has been particularly labile within the *Drosophila obscura* group. However, they also suggest that the selective pressures driving these changes have not been consistent, implying that more than one selective agent may be responsible.

Keywords: gene duplication, RNAi, RNA interference, adaptive evolution, neofunctionalization

Introduction

Gene duplication is an important process in molecular evolution, providing raw genetic material for evolutionary innovation. The subsequent evolutionary dynamics following gene duplication are often described in terms of two alternative models, 'neofunctionalization' and 'sub-functionalization' [1]. Under neofunctionalization, the functional redundancy following duplication provides relaxed selective constraint, and allows new mutations to accumulate through genetic drift. Most such mutations will reduce the functionality of the gene (resulting in pseudogenization), but some paralogs can be selected for novel or derived functions. Under sub-functionalization, the duplicates independently accumulate mutations that allow them to specialise in a subset of ancestral functions of a pleiotropic gene. Neo-functionalization leads to asymmetrical evolutionary rates among paralogs (with faster evolution in paralogs that gain derived function), whereas equal rates are expected for the latter [2]. It has been suggested that both processes have played an important role in the rapid evolution of RNA interference-related pathways, including the long- and short-term evolutionary history of the Argonautes, the effectors of RNAi [3–5].

The RNAi-related pathways comprise a range of small-RNA mechanisms best known for their roles in mediating the control of gene expression, antiviral responses, and defence against mobile genetic elements (respectively: the miRNA pathway; Dicer-1 and Argonaute-1 in insects [6]; the siRNA pathway; Dcr-2 and Argounate 2 in insects [7]; and the piRNA pathway; piwi-family Argonaute AGO3 and Piwi/Aub in insects [8, 9]). In addition, RNAi-related pathways have been implicated in a variety of biological processes, such as the control of dosage compensation [10–12] and the suppression of genetic drive [13–18], among others. Several genes involved in the defensive piRNA

and siRNA pathways, but not the miRNA pathway, display elevated rates of adaptive

protein evolution. This is best studied in *Drosophila* [19–21], but is also detectable in

other insects [22]. It has been hypothesised that this is a consequence of parasite-

mediated 'arms-race' coevolution [20, 23], either through conflict with parasite-

encoded immune suppressors—as widely seen in RNA viruses [24]— or in the case

of the piRNA pathway, through selection for 're-tuning' suppression mechanisms [25].

Adaptive evolution of RNAi pathways is also partly reflected in the gain, loss, and

functional divergence of Argonaute-family duplications [26]. For example, within the

Drosophilidae—an important model for RNAi-related pathways of animals—Piwi has

been duplicated in the lineages leading to *Phortica variegata* and *Scaptodrosophila*

deflexa [3], and *Ago2* has been duplicated in those leading to *S. deflexa*, *D. willistoni*,

D. melanogaster (where only one paralog remains – the canonical *Ago2*) and *D.*

pseudoobscura [4]. This is particularly striking in the *Drosophila obscura* species

group, which has experienced at least 6 independent duplications of *Ago2* over the

last 20 million years, with all but one of the resulting duplicates becoming testis-

specific, and most displaying evidence of recent and/or ongoing positive selection [4].

Very recently it has been noted that several accessory components of the siRNA and

piRNA pathways have also been duplicated in *D. pseudoobscura*, including

armitage, *asterix*, *cutoff*, *maelstrom*, *tejas* and *vreteno* [22]. In *D. melanogaster*, these

proteins are engaged in a number of roles in the piRNA pathway (**Table 1**). Here we

use publicly available data to reconfirm the history and expression of *Ago2* in the

obscura group, and to test whether duplications in the other genes also show male-

specific expression, whether they are contemporaneous with those of *Ago2*, and

whether they too show strong signatures of adaptive protein evolution. We find no

clear pattern of these duplications being coincident with *Ago2* duplications, but both

asterix and *cutoff* duplications display increased sexual dimorphism relative to their ancestral copies, through decreased female expression. In addition, several of the gene duplicates show evidence of adaptive protein evolution in *D. pseudoobscura*, including both copies of *cutoff*, the ancestral copy of *asterix*, and the new duplicates of *tejas*, *maelstrom* and *vreteno*.

RESULTS

Obscura group *Argonaute 2* are duplicated and show male-biased expression

The *Drosophila obscura* group has experienced multiple duplications of *Ago2* and it has previously been shown that these are associated with positive selection and testis-specific expression [4]. Here we reanalyzed the expression patterns and evolutionary history of these genes using publicly available RNAseq and genomic data, additionally including newly available genomic sequences from *D. algonquin*, *D. athabasca*, *D. bifasciata* and *D. miranda*. In contrast to the previous qPCR analysis that failed to identify strong expression of the ancestral copy in *D. pseudoobscura* (*Ago2d* [4]), we found that all the *Ago2* homologs in *D. pseudoobscura* were detectable at a high level in RNAseq data, and that that all show significant male-bias (**Figure 1**). The *Ago2d* expression detected here is unlikely to be an artefact of cross-mapping between paralogs as we observed the reads that mapped uniquely across the gene. The male bias was largest for *Ago2e*, where expression in males is approximately 1000-fold higher than females (pMCMC<0.001; **Figure 1**), and smallest in *Ago2c* and the ancestral copy *Ago2d*, consistent with the *ca.* two-fold enrichment of the single copy of *Ago2* in male *D. melanogaster*. We also confirmed that *D. miranda*, a close relative of *D. pseudoobscura* that has not previously been analyzed, displayed a qualitatively similar pattern among those paralogs represented (**Figure 1**). In *D. obscura* and we

125 found the ancestral copy (*Ago2a*) again showed slightly, but marginally significantly,

126 higher expression in males (pMCMC=-0.014), but that other *Ago2* proteins showed a

127 strong male biased expression, with the largest effect for *Ago2f*, where male

128 expression was 2000-fold higher (**Figure 1**; pMCMC<0.001).

Six piRNA pathway genes are duplicated, and *asterix* and *cutoff* duplicates show

increased male-bias in their expression

Palmer *et al.* [22] recently identified six accessory piRNA pathway genes that have

also experienced duplication in the obscura group (**Additional file 1 Table S1**). We

could locate the duplicates for all genes in *D. pseudoobscura*, except for *armitage*

where we instead identified a duplicate in the affinis subgroup but not in the obscura

or subobscura subgroups. Where new chromosomal locations could be determined

by synteny in *D. pseudoobscura*, we found that *cutoff*, *maelstrom* and *vreteno* were

duplicated from an autosome to the X chromosome, *asterix* duplicated from the X

chromosome to an autosome, and *tejas* duplicated between autosomal locations. Two

duplicates (*asterix* and *tejas*) lack introns, suggesting they are retro-transcribed copies

created through an mRNA intermediate.

Using public RNAseq data from *D. pseudoobscura*, *D. miranda*, and *D. obscura*, we

found that all of the gene duplicates were expressed (**Figure 2**). *Armitage*, which was

not duplicated within the newly examined lineages for which RNAseq data were

available, did not show strong sex-biased expression. Similarly, *tejas*, *maelstrom*, and

vreteno were not strongly differentially expressed between the sexes, and nor were

their duplicates in *D. pseudoobscura* and *D. miranda*. In contrast, both *asterix* and

cutoff duplicates displayed substantially reduced expression in females and slightly

increased expression in males (**Figure 2**). For example, as previously reported from

qPCR analysis [27] the paralog of *asterix* in *D. pseudoobscura* displays ca. 1000-fold higher expression in males than females. In both *D. pseudoobscura* and *D. miranda*, those genes with overall strongly increased male-biased expression (*Argonaute 2*, *asterix*, *cutoff,* and their paralogs) had the highest expression in testis, and had reduced their expression in ovaries (**Additional file 2 Figure S1**).

Adaptive amino-acid substitutions are generally more common in the duplicates

Using population genetic data from *D. pseudoobscura*, and *D. miranda* as an outgroup, we used the McDonald-Kreitman framework and a maximum-likelihood extension to estimate the rate of adaptive substitution in protein sequences, and to test whether this was different between the ancestral and duplicated copies [28, 29]. Treating genes individually, we found evidence for positive selection acting on at least one paralog for each of the genes except *asterix* and *Ago2c* ($p < 0.05$; **Additional file 3 Table S2**). Among the ancestral copies, only *cutoff* displayed evidence of positive selection. We then tested whether the paralogs generally showed a different pattern of selection to the ancestral copies by dividing the genes into two classes (6 ancestral copies and 8 paralogs) and comparing the likelihood of models that allowed the classes to differ in the adaptive rate α (**Table 2**) [29]. The best supported model allowed α to differ between ancestral and duplicate copies (Akaike weight: 0.81), and the second best supported model was that in which α=0 for the ancestral copies (Akaike weight: 0.19), providing overall evidence that the paralogs have experienced more adaptive protein evolution. In the best-supported model, the α was estimated to be 0.68 for the duplicate group, which more than three times larger than the ancestral group (0.20). In case segregating weakly-deleterious variants had led to a downward bias in estimates of α, we repeated this analysis excluding all alleles with a minor allele

frequency <0.125 [30], although this reduced power to the extent that few genes

remained individually significant (**Additional file 6 Table S3**). We also repeated the

analysis with a larger dataset PRJNA326536 [31] (**Additional file 6 Table S3**), and

obtained qualitatively similar results (R^2=0.946 for α estimate between the analyses;

the second dataset, while larger, is less suitable for analysis as only the third

chromosome is a direct sample from a wild population).

Gene duplications were unlikely to be contemporaneous

Given the multiple duplications of *Ago2* and the piRNA pathway components in the

obscura group, we hypothesized that some duplications may have occurred near-

simultaneously, duplicating whole components of a pathway together. We therefore

used relaxed-clock phylogenetic methods to estimate the relative timings of each

duplication. In agreement with the previous analysis of Ago2 [4], we found that the

duplications giving rise to *Ago2e* and *Ago2f* predated the split between the obscura

and pseudoobscura subgroups, with a subsequent loss of *Ago2f* from the

pseudoobscura subgroup (**Figure 3**). In contrast, we found that duplications in five of

the other six genes unambiguously occurred after the obscura/pseudoobscura split,

with the timing of duplication in *maelstrom* being uncertain. Briefly, *armitage* displayed

a single duplication shared by members of the affinis subgroup, *asterix* and *tejas* a

single duplication each in the lineage leading to *D. pseudoobscura* (which were

subsequently lost in the affinis subgroup), *cutoff* a single duplication recently in the

pseudoobscura subgroup, and *vreteno* a single duplication at the base of the obscura

group (**Figure 4**). For *maelstrom*, the maximum clade credibility tree suggests

duplication occurred very slightly prior to this split, followed by subsequent loss of one

paralog the obscura subgroup (**Figure 4**). However, this was poorly supported, and

similar pattern to the others genes may be more parsimonious. We used the posterior

distributions of split times, relative to the divergence time of the obscura and pseudoobscura subgroups, to infer whether or not duplications occurred at approximately the same time (**Figure 5**). Although the small amount of information available from single genes made relative timings highly uncertain, it is clear that few of the *Ago2* duplications could have been concurrent with the piRNA-pathway duplications (**Additional file 5 Figure S2**). However, the recent and rapid duplications within the piRNA could have been concurrent, with *vreteno*, *tejas, maelstrom* and *asterix* not differing significantly, all having duplicated very close to the split between *D. obscura* and *D. pseudoobscura* (posterior overlap >0.1 in each case; **Additional file 5 Figure S2**).

DISCUSSION

Although four of the six piRNA pathway duplicates did not display altered tissue specificity compared to the ancestral copy, *asterix* and *cutoff* both became significantly more male biased, as did each of the *Ago2* duplicates [4]; **Figure 1**, **Figure 2**). In each case, this was due to higher (or exclusive) expression in the testis. The duplicated genes also showed higher rates of adaptive amino acid substitution, together and individually, whereas only two (*asterix* and *armitage*) displayed evidence of positive selection when single-copy in *D. melanogaster* (**Additional file 6 Table S3**).

This new tissue specificity and the rapid evolution of duplicated copies broadly suggest that gene duplication in these pathways may be associated with functional diversification through neofunctionalization, for example by testis-specific selective pressure. Three main selective pressures seem likely candidates to have driven this. First, given the role of *Ago2* in antiviral defense in *Drosophila* [7], and the role of the piRNA pathway in antiviral defense in mosquitoes [32], it is possible that these

duplications have specialized to a virus that is active in the male germline, such as *D.*

obscura and *D. affinis* Sigmaviruses [33]. Second, given the role of all of these genes

in the suppression of transposable elements (TEs), their evolution may have been

shaped by the invasion of TEs that are more active in testis, as seen for Penelope [34]

and copia [35]. Such a 'duplication arms-race' in response to TE invasion is thought

to occur in mammals, where repeated duplications of KRAB-ZNF family are selected

following the invasion of novel TEs, and subsequently provide defence [36, 37].

Alternatively, duplicates may quantitatively enhance the pre-existing response to TEs,

as suggested for another rapidly-evolving piRNA-pathway component, Rhino [38].

The third, and arguably most compelling hypothesis, is that selection is mediated by

conflict with meiotic drive elements, such as sex ratio distorting X-chromosomes [39,

40]. Most directly, meiotic drive elements are common in *Drosophila*, and RNAi-related

pathways have been widely implicated in their action and suppression [15, 16, 18]. In

addition, sex-chromosome drive is widespread in the *Drosophila obscura* group. X

chromosome drive was first described in *D. obscura*, and has also been reported in

pseudoobscura, *persimilis*, *affinis*, *azteca*, *subobscura* and *athabasca* [39] and is

mediated through a testis-specific function (Y-bearing sperm have reduced function).

Finally, a testis-specific class of hairpin (endo) siRNAs is required for male fertility in

D. melanogaster [41], testes-restricted clustered miRNAs show rapid evolutionary

turnover and are represented in large numbers in *D. pseudoobscura* [42], and

suppression of sex-specific duplicates of S-Lap1 via a small-RNA mechanism has very

recently been implicated in the meiotic drive mechanism of *D. pseudoobscura* [17]. In

this context, it is also interesting to note that *Ago2* is involved in directing

heterochromatin formation in *Drosophila* dosage compensation [10–12], and that in *D.*

melanogaster the sex-ratio distorting *Spiroplasma* achieves male-killing through the

disruption of dosage compensation, although this acts at the embryonic stage [43].

Nevertheless, in the absence of mechanistic studies, this remains highly speculative.

Testis is generally more permissive to gene expression and testis-specific expression

may be a transient state (i.e. the "Out of Testis Hypothesis" [44]). In addition, the

application of MK-like analyses to paralogs is inherently flawed [1], as the MK

framework implicitly assumes that the selective regime has been consistent across all

(group and outgroup) sequences analyzed. If gene duplicates experience an early but

transient period of relaxed constraint, a high proportion of the amino-acid fixations may

have occurred as a result of genetic drift that is no longer detectable from current

patterns of polymorphism.

258 Methods

259 Sequence collation and paralog identification

The full-length sequences for 7 RNA inteference genes; *armitage* (*armi*), *asterix* (*arx*),

cutoff (*cuff*), *tejas (tej)*, *vreteno* (*vret*), *maelstrom* (*mael*), and *Argonaute 2* (*Ago2*) from

12 obscura group species were identified using tBLASTn (BLAST+ 2.6.0) [45] with a

local BLAST database (see below for the details of the construction of local genomic

database). Known gene sequences from *D. pseudoobscura* and *D. melanogaster*

were used as a query with a stringent e-value threshold (1e-40). Genes were inferred

to have been duplicated when BLAST indicated that there were multiple full-length hits

located in different genomic regions. The sequences were manually inspected, introns

removed and the coding frame identified using Bioedit v 7.2 [46]. Genes in *D.*

pseudoobscura were classified as ancestral or duplicate copies based on the syntenic

orthology with *D. melanogaster* using Flybase Genome Browser [47]. High quality

genomes are not available for other members of the obscura groups, and in those

cases ancestral/derived status was assigned based on homology with *D.*

pseudoobscura. To provide a comprehensive overview of the evolution of the RNAi

paralogs, we included 24 *Drosophila* species outside of *obscura* group with assembled

genomes already available in public databases. The Flybase and NCBI tblastn online

portal were used to identify the target genes with queries from *D. melanogaster* or

closely related species.

Five obscura group species had assembled genomes at the time of this study: *D.*

pseudoobscura (assembly Dpse_3.0 [48]), *D. miranda* (assembly DroMir_2.2 [49]

[50]), *D. persimilis* (assembly dper_caf1 [51]), *D. affinis* (Drosophila affinis Genome

Release 1.0 [52]) and *D. lowei* (Drosophila lowei Genome Release 1.0 [52, 53]) and

in these cases the genome was directly used for local BLAST database. For four

species (*D. obscura*, *D. subobscura*, *D. subsilvestris*, *D. tristis*) we used de novo

assembled transcriptomes based on paired RNA-seq reads data from wild-collected

males [54] (Accession: PRJNA312496). Assembly was performed using Trinity [55]

with '--trimmomatric' and otherwise default parameters, and the assembled

transcriptome was searched locally using BLAST. For three other species: *D.*

athabasca, *D. Algonquin* [56] (Accession: PRJNA274695) and *D. bifasciata*

(Accession: PRJDB4817), only unassembled genomic reads were available. For these

species we applied a targeted assembly approach as follows: (i) reads that had local

similarity with all known duplicated RNAi proteins were identified using Diamond [57]

with relaxed e-value of 1; (ii) hits from Diamond were then retained and used for

assembly using Spades v3.10.1 [58]; and (iii) scaffolds produced by Spades were then

used as references in local BLAST database.

Phylogenetic analysis and the relative timing of duplications

Bayesian relaxed clock trees were used to infer the evolutionary relationship among paralogs. First, the sequences were aligned as translated nucleotide in Clustal W [59] with default parameters. Regions with ambiguous alignment were identified and removed manually by eye. A total of 7 gene trees were then inferred using Beast v1.7.0 [60]. Inference used a relaxed clock model with an uncorrelated lognormal distribution among branches, and an HKY substitution model with empirical base frequencies and rate variation among sites was modelled as a gamma distribution with four categories. The site model allowed for third codon position to have different substitution model from the other positions.

The trees were scaled by setting the time to most recent common ancestor of the *D. obscura* group to have lognormal distribution with a data-scale mean of 1, and a very small standard deviation of 0.01. This had the advantage of scaling all duplications to the same relative timescale, while allowing different genes and different paralogs to vary in their rate. To record the posterior ages of duplication, we specified the ancestral and duplicated genes as a distinct taxon set. The Monte Carlo Markov Chain analysis was run for at least 100 million states and posterior sample was recorded every 10000 states. Log files were then inspected in Tracer v1.6 [61] for parameter stationarity, and adequate sampling as indicated by an effective sample size over 200. Finally, 25% of initial trees were discarded as burn-in, and maximum clade credibility trees were summarized using Tree Annotator. Parameter MCMC files were processed using a custom R script [62] to infer the posterior distribution the age of duplication for each gene and to quantify the degree overlapping between these age distributions.

Differential expression analysis of the duplicated RNAi genes

For this analysis, we used obscura group transcriptome datasets available in EBI ENA (European Nucleotide Archive, http://www.ebi.ac.uk/ena) and DDBJ (DNA DataBank of Japan, http://www.ddbj.nig.ac.jp/) that included the sex and tissue annotation. The datasets comprised 163, 42 and 34 RNA-seqs datasets of *D. pseudoobscura, D. miranda* and *D. obscura* respectively; Bioproject: DRA004463, PRJEB1227 [63], PRJNA226598, PRJNA219224 [64], PRJNA326536 [31], PRJNA74723, PRJNA321079, PRJNA291085 [65], PRJNA268967 [66]. Since our main interest was the comparison expression between sex, but not its absolute expression value, we a performed a simple read-counting analysis. In outline, each RNA-seq dataset was mapped to the full-length CDS using Bowtie2 v2.3.2 [67] with mode '--very sensitive' and otherwise default parameters. The reads mapped to reference were counted using combination of SAMtools view flag -F 4 and SAMtools idxstats v1.4 [68]. The count data were then normalized by gene length and read depth, where it was then scaled relative to the expression of RpL32. To determine the statistical significance of difference gene expression, generalised linear mixed models were fitted using R package MCMCglmm [69] with sex as fixed effect and tissue as a random effect, and log-transformed normalised expression as the response variable. The natural logarithm transformation ($\log_e$) was used to reduce the skewness of the distributions. To allow for zero value for non-expressed genes, the genes with read count 0 was replaced with 1.

$$\mathbf{Y} \sim \boldsymbol{\mu} + \mathbf{sex} + \mathbf{tissue\ (random)} + \boldsymbol{\varepsilon}$$

Where Y is $\log_e$ transformed normalized expression data (response variable), μ is mean of $\log_e$ transformed expression and ε is residual error. The random effects

343   (tissue) and the residual were assumed to be distributed multivariate normal with mean

344   0 and uncorrelated covariance matrix MVN (0, $I\sigma^2$). Sex was modelled as a factor with

345   2 variables (male-female) and tissue contained 13 variables of different tissue.

Population Genetic Analysis of the RNAi Duplicated Genes

We used the McDonald-Krietman test [28] to compare the rate of adaptive evolution

between ancestral and duplicate genes using polymorphism data from publicly-

available sequencing datasets: Pseudobase (12 strains of *pseudoobscura*, Accession

list: SRP007802 [53]) and 12 strains *D. miranda* (Bioproject: PRJNA277849 [70]).

Genomic reads for each strain were mapped to the genomic reference using Bowtie2

with '--very-sensitive' mode and otherwise default parameters and reads mapped to

the genes of interest were extracted using SAMtools view (flag -F 4). Duplicate reads

were marked using MarkDuplicates (Picard Tools [71]). To reduce the excessive

variants surrounding indel, we then applied GATK IndelRealigner [71], which discards

the original mapping and performs local-realignment around indel. The output was

then sorted and indexed and the BAM file was used for 'mpileup' variant calling

(SAMtools v1.4 [72]). The output VCF files were then filtered to only include SNP

(GATK SelectVariants [71]), and variants that were covered by less than five reads

were masked with 'N' (undetermined bases, --snpmask GATK v3.5 [71]). The variant

files   were   then   converted   to   FASTA   format   using   GATK

FastaAlternateReferenceMaker, which replaced genomic reference with variants

defined in VCF files [73] and output the heterozygous calls with IUPAC ambiguous

code. Finally, FastPHASE [74] was used to generate pseudo-haplotypes, although

haplotype information was not utilized by the analysis.

MK tests were performed for each gene on *D. pseudoobscura-D miranda.* DNAsp v5.0

[75] was used to estimate the statistics for the MK test and Fisher's exact test was

used to calculate the statistical significance for single-gene analyses. Genes were

then grouped into ancestral and duplicate genes, and a cross-gene analysis was

performed using a maximum likelihood extension of the MK test [29]. Five different

models were fitted that differed in the constraint of α (proportion of non-synonymous

subtitutions estimated to be adaptive), and the relative support between models was

compared using Akaike Weights.

List of Abbreviations

Ago2: Agronaute 2 **armi**: armitage **arx**: asterix **BAM**: Binary Alignment Map **BLAST**:

Basic Alignment Search Tool **cuff**: cutoff **GATK**: Genome Analysis Toolkit **HKY**:

Hasegawa, Kishino and Yano model **mael**: maelstrom **MCMC**: Monte Carlo Markov

Chain **MK:** McDonald-Kreitman test **RNAi**: RNA interference **SNP**: Single Nucleotide

Polymorphism **TE:** Transposable Elements **tej**: tejas **VCF**: Variant Call Format **vret**:

vreteno

Declarations

Ethics approval and consent to participate

Not applicable

Consent to publication

Not applicable

Availability of data and material

Fasta alignment (both for phylogenetic and MK analysis) and raw expression data are

available via Figshare (DOI: 10.6084/m9.figshare.7145720)**.**

Competing interests

The authors declare that they have no competing interests.

Funding

DC was financially supported by a Master's Training Scholarship from the Indonesian Endowment Fund for Education (LPDP) and the University of Edinburgh School of Biological Science Bursary for MSc in Quantitative Genetics and Genome Analysis.

Authors' contributions

DJO and DC conceived the study and designed the analysis, DC analyzed the data and wrote the first draft of the manuscript. Both authors read and approved the final manuscript.

Acknowledgements

We thank Billy Palmer for initial discussion on the variant calling, and Billy Palmer and Samuel Lewis for comments on an earlier version of this manuscript. We thank the many people who made their published data publicly available, and Shu Kondo for permission to use unpublished data from *D. bifasciata*.

References

1. Hahn MW. Distinguishing Among Evolutionary Models for the Maintenance of Gene Duplicates. J Hered. 2009;100:605–17. doi:10.1093/jhered/esp047.

2. Innan H, Kondrashov F. The evolution of gene duplications: classifying and distinguishing between models. Nat Rev Genet. 2010;11:97–108. doi:10.1038/nrg2689.

3. Lewis SH, Salmela H, Obbard DJ. Duplication and diversification of dipteran argonaute genes, and the evolutionary divergence of Piwi and Aubergine. Genome Biol Evol. 2016;8:507–18.

4. Lewis SH, Webster CL, Salmela H, Obbard DJ. Repeated duplication of Argonaute2 is associated with strong selection and testis specialization in Drosophila. Genetics. 2016;204:757–69.

5. Singh RK, Gase K, Baldwin IT, Pandey SP. Molecular evolution and diversification of the Argonaute family of proteins in plants. BMC Plant Biol. 2015;15:23. doi:10.1186/s12870-014-0364-6.

6. Vidigal JA, Ventura A. The biological functions of miRNAs: lessons from in vivo studies. Trends Cell Biol. 2015;25:137–47. doi:10.1016/j.tcb.2014.11.004.

7. Bronkhorst AW, van Rij RP. The long and short of antiviral defense: small RNA-based immunity in insects. Curr Opin Virol. 2014;7:19–28. doi:10.1016/J.COVIRO.2014.03.010.

8. Czech B, Hannon GJ. One Loop to Rule Them All: The Ping-Pong Cycle and piRNA-Guided Silencing. Trends Biochem Sci. 2016;41:324–37. doi:10.1016/j.tibs.2015.12.008.

9. Lewis SH, Quarles KA, Yang Y, Tanguy M, Frézal L, Smith SA, et al. Pan-arthropod analysis reveals somatic piRNAs as an ancestral defence against transposable elements. Nat Ecol Evol. 2018;2:174–81. doi:10.1038/s41559-017-0403-4.

10. Menon DU, Meller VH. A role for siRNA in X-chromosome dosage compensation in Drosophila melanogaster. Genetics. 2012;191:1023–8. doi:10.1534/genetics.112.140236.

11. Tang W, Seth M, Tu S, Shen E-Z, Li Q, Shirayama M, et al. A Sex Chromosome piRNA Promotes Robust Dosage Compensation and Sex Determination in C. elegans. Dev Cell. 2018;44:762–770.e3. doi:10.1016/j.devcel.2018.01.025.

12. Deshpande N, Meller VH. Chromatin That Guides Dosage Compensation Is Modulated by the siRNA Pathway in Drosophila melanogaster. Genetics. 2018;209:1085–97. doi:10.1534/genetics.118.301173.

13. Tao Y, Masly JP, Araripe L, Ke Y, Hartl DL. A sex-ratio Meiotic Drive System in Drosophila simulans. I: An Autosomal Suppressor. PLoS Biol. 2007;5:e292. doi:10.1371/journal.pbio.0050292.

14. Tao Y, Araripe L, Kingan SB, Ke Y, Xiao H, Hartl DL. A sex-ratio Meiotic Drive System in Drosophila simulans. II: An X-linked Distorter. PLoS Biol. 2007;5:e293. doi:10.1371/journal.pbio.0050293.

15. Gell SL, Reenan RA. Mutations to the piRNA pathway component aubergine enhance meiotic drive of segregation distorter in Drosophila melanogaster. Genetics. 2013;193:771–84. doi:10.1534/genetics.112.147561.

16. Aravin AA, Klenov MS, Vagin V V, Bantignies F, Cavalli G, Gvozdev VA. Dissection of a natural RNA silencing process in the Drosophila melanogaster germ line. Mol Cell Biol. 2004;24:6742–50. doi:10.1128/MCB.24.15.6742-6750.2004.

17. Ellison C, Leonard C, Landeen E, Gibilisco L, Phadnis N, Bachtrog D. Rampant cryptic
sex chromosome drive in Drosophila. doi:10.1101/324368.

18. Lin C-J, Hu F, Dubruille R, Smibert P, Loppin B, Correspondence ECL. The
hpRNA/RNAi Pathway Is Essential to Resolve Intragenomic Conflict in the Drosophila Male
Germline. 2018. doi:10.1016/j.devcel.2018.07.004.

19. Obbard DJ, Jiggins FM, Halligan DL, Little TJ. Natural selection drives extremely rapid
evolution in antiviral RNAi genes. Curr Biol. 2006;16:580–5.

20. Obbard DJ, Gordon KHJ, Buck AH, Jiggins FM. The evolution of RNAi as a defence
against viruses and transposable elements. Philos Trans R Soc Lond B Biol Sci.
2009;364:99–115. doi:10.1098/rstb.2008.0168.

21. Kolaczkowski B, Hupalo DN, Kern AD. Recurrent Adaptation in RNA Interference Genes
Across the Drosophila Phylogeny. Mol Biol Evol. 2011;28:1033–42.
doi:10.1093/molbev/msq284.

22. Palmer WH, Hadfield JD, Obbard DJ. RNA-Interference Pathways Display High Rates of
Adaptive Protein Evolution in Multiple Invertebrates. Genetics. 2018;208:1585–99.
doi:10.1534/genetics.117.300567.

23. Marques JT, Carthew RW. A call to arms: coevolution of animal viruses and host innate
immune responses. Trends Genet. 2007;23:359–64. doi:10.1016/J.TIG.2007.04.004.

24. van Mierlo JT, Overheul GJ, Obadia B, van Cleef KWR, Webster CL, Saleh M-C, et al.
Novel Drosophila Viruses Encode Host-Specific Suppressors of RNAi. PLoS Pathog.
2014;10:e1004256. doi:10.1371/journal.ppat.1004256.

25. Blumenstiel JP, Erwin AA, Hemmer LW. What drives positive selection in the Drosophila
piRNA machinery? The genomic autoimmunity hypothesis. Yale J Biol Med. 2016;89:499–
512. http://www.ncbi.nlm.nih.gov/pubmed/28018141. Accessed 24 Jul 2017.

26. Dowling D, Pauli T, Donath A, Meusemann K, Podsiadlowski L, Petersen M, et al.
Phylogenetic Origin and Diversification of RNAi Pathway Genes in Insects. Genome Biol
Evol. 2017;1:evw281. doi:10.1093/gbe/evw281.

27. Meisel RP, Hilldorfer BB, Koch JL, Lockton S, Schaeffer SW. Adaptive Evolution of
Genes Duplicated from the Drosophila pseudoobscura neo-X Chromosome. Mol Biol Evol.
2010;27:1963–78. doi:10.1093/molbev/msq085.

28. McDonald JH, Kreitman M. Adaptive protein evolution at the Adh locus in Drosophila.
Nature. 1991;351:652–654. doi:10.1038/350055a0.

29. Welch JJ. Estimating the genomewide rate of adaptive protein evolution in drosophila.
Genetics. 2006;173:821–37. doi:10.1534/genetics.106.056911.

30. Charlesworth J, Eyre-Walker A. The McDonald-Kreitman Test and Slightly Deleterious
Mutations. Mol Biol Evol. 2008;25:1007–15. doi:10.1093/molbev/msn005.

31. Fuller ZL, Haynes GD, Richards S, Schaeffer SW. Genomics of Natural Populations:
How Differentially Expressed Genes Shape the Evolution of Chromosomal Inversions in
Drosophila pseudoobscura. Genetics. 2016;204:287–301. doi:10.1534/genetics.116.191429.

32. Campbell CL, Black WC, Hess AM, Foy BD. Comparative genomics of small RNA
regulatory pathway components in vector mosquitoes. BMC Genomics. 2008;9:425.
doi:10.1186/1471-2164-9-425.

33. Longdon B, Wilfert L, Obbard DJ, Jiggins FM. Rhabdoviruses in two species of
drosophila: Vertical transmission and a recent sweep. Genetics. 2011;188:141–50.
doi:10.1534/genetics.111.127696.

34. Rozhkov N V., Aravin AA, Zelentsova ES, Schostak NG, Sachidanandam R, McCombie
WR, et al. Small RNA-based silencing strategies for transposons in the process of invading
Drosophila species. RNA. 2010;16:1634–45. doi:10.1261/rna.2217810.

35. Pasyukova, S. Nuzhdin, W. Li E, Nuzhdin S, Li W, Flavell AJ. Germ line transposition of
the copia retrotransposon in Drosophila melanogaster is restricted to males by tissue-
specific control of copia RNA levels. Mol Gen Genet MGG. 1997;255:115–24.
doi:10.1007/s004380050479.

36. Thomas JH, Schneider S. Coevolution of retroelements and tandem zinc finger genes.
Genome Res. 2011;21:1800–12. doi:10.1101/gr.121749.111.

37. Jacobs FMJ, Greenberg D, Nguyen N, Haeussler M, Ewing AD, Katzman S, et al. An
evolutionary arms race between KRAB zinc-finger genes ZNF91/93 and SVA/L1
retrotransposons. Nature. 2014;516:242–5. doi:10.1038/nature13760.

38. Levine MT, Wende Vander HM, Hsieh E, Baker EP, Malik HS. Recurrent gene
duplication diversifies genome defense repertoire in Drosophila. Mol Biol. 2016;33:1–13.

39. Jaenike J. Sex Chromosome Meiotic Drive. Annu Rev Ecol Syst. 2001;32:25–49.
doi:10.1146/annurev.ecolsys.32.081501.113958.

40. Lindholm AK, Dyer KA, Firman RC, Fishman L, Forstmeier W, Holman L, et al. The
Ecology and Evolutionary Dynamics of Meiotic Drive. Trends Ecol Evol. 2016;31:315–26.
doi:10.1016/J.TREE.2016.02.001.

41. Wen J, Duan H, Bejarano F, Okamura K, Fabian L, Brill JA, et al. Adaptive regulation of
testis gene expression and control of male fertility by the Drosophila hairpin RNA pathway.
[Corrected]. Mol Cell. 2015;57:165–78. doi:10.1016/j.molcel.2014.11.025.

42. Mohammed J, Flynt AS, Panzarino AM, Mondal MMH, DeCruz M, Siepel A, et al. Deep
experimental profiling of microRNA diversity, deployment, and evolution across the
Drosophila genus. Genome Res. 2018;28:52–65. doi:10.1101/gr.226068.117.

43. Harumoto T, Anbutsu H, Lemaitre B, Fukatsu T. Male-killing symbiont damages host's
dosage-compensated sex chromosome to induce embryonic apoptosis. Nat Commun.
2016;7:12781. doi:10.1038/ncomms12781.

44. Kaessmann H. Origins, evolution, and phenotypic impact of new genes. Genome Res.
2010;20:1313–26. doi:10.1101/gr.101386.109.

45. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool.
J Mol Biol. 1990;215:403–10. doi:10.1016/S0022-2836(05)80360-2.

46. Hall Thomas. BioEdit: a user-firendly biological sequence alignment editor and analysis
program for Windows 95/95/NT. Oxford Univ. 1999;41:95–8. doi:citeulike-article-id:691774.

47. St. Pierre SE, Ponting L, Stefancsik R, McQuilton P. FlyBase 102—advanced
approaches to interrogating FlyBase. Nucleic Acids Res. 2014;42:D780–8.
doi:10.1093/nar/gkt1092.

48. English AC, Richards S, Han Y, Wang M, Vee V, Qu J, et al. Mind the Gap: Upgrading
Genomes with Pacific Biosciences RS Long-Read Sequencing Technology. PLoS One.
2012;7:e47768. doi:10.1371/journal.pone.0047768.

49. Alekseyenko AA, Ellison CE, Gorchakov AA, Zhou Q, Kaiser VB, Toda N, et al.
Conservation and de novo acquisition of dosage compensation on newly evolved sex
chromosomes in Drosophila. Genes Dev. 2013;27:853–8. doi:10.1101/gad.215426.113.

50. Zhou Q, Bachtrog D. Sex-specific adaptation drives early sex chromosome evolution in
Drosophila. Science (80-). 2012;337:341–5. doi:10.1126/science.1225385.

51. Drosophila 12 Genomes Consortium AG, Clark AG, Eisen MB, Smith DR, Bergman CM,
Oliver B, et al. Evolution of genes and genomes on the Drosophila phylogeny. Nature.
2007;450:203–18. doi:10.1038/nature06341.

52. Palmieri N, Kosiol C, Schlötterer C. The life cycle of Drosophila orphan genes. Elife.
2014;3:e01311. doi:10.7554/eLife.01311.

53. McGaugh SE, Heil CSS, Manzano-Winkler B, Loewe L, Goldstein S, Himmel TL, et al.
Recombination Modulates How Selection Affects Linked Sites in Drosophila. PLoS Biol.
2012;10:e1001422. doi:10.1371/journal.pbio.1001422.

54. Webster CL, Waldron FM, Robertson S, Crowson D, Ferrari G, Quintana JF, et al. The
Discovery, Distribution, and Evolution of Viruses Associated with Drosophila melanogaster.
PLOS Biol. 2015;13:e1002210. doi:10.1371/journal.pbio.1002210.

55. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length
transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol.
2011;29:644–52.

56. Wong Miller KM, Bracewell RR, Eisen MB, Bachtrog D. Patterns of Genome-Wide
Diversity and Population Structure in the Drosophila athabasca Species Complex. Mol Biol
Evol. 2017. doi:10.1093/molbev/msx134.

57. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. Nat
Methods. 2014;12:59–60. doi:10.1038/nmeth.3176.

58. Nurk S, Bankevich A, Antipov D, Gurevich A, Korobeynikov A, Lapidus A, et al.
Assembling genomes and mini-metagenomes from highly chimeric reads. In: Lecture Notes
in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture
Notes in Bioinformatics). Springer, Berlin, Heidelberg; 2013. p. 158–70. doi:10.1007/978-3-
642-37195-0_13.

59. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: Improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific gap
penalties and weight matrix choice. Nucleic Acids Res. 1994;22:4673–80.
doi:10.1093/nar/22.22.4673.

60. Drummond AJ, Rambaut A. BEAST: Bayesian evolutionary analysis by sampling trees.
BMC Evol Biol. 2007;7:214. doi:10.1186/1471-2148-7-214.

61. Rambaut A. Tracer v1.6. http://tree.bio.ed.ac.uk/software/tracer/. 2013.

62. R Core Team. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. 2016. https://www.r-project.org/.

63. Chen Z-X, Sturgill D, Qu J, Jiang H, Park S, Boley N, et al. Comparative validation of the
D. melanogaster modENCODE transcriptome annotation. Genome Res. 2014;24:1209–23.
doi:10.1101/gr.159384.113.

64. VanKuren NW, Vibranovski MD. A novel dataset for identifying sex-biased genes in
Drosophila. J genomics. 2014;2:64–7. doi:10.7150/jgen.7955.

65. Nyberg KG, Machado CA. Comparative Expression Dynamics of Intergenic Long
Noncoding RNAs in the Genus *Drosophila*. Genome Biol Evol. 2016;8:1839–58.
doi:10.1093/gbe/evw116.

66. Gomes S, Civetta A. Hybrid male sterility and genome-wide misexpression of male
reproductive proteases. Sci Rep. 2015;5:11976. doi:10.1038/srep11976.

67. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment
of short DNA sequences to the human genome. Genome Biol. 2009;10:R25. doi:10.1186/gb-

2009-10-3-r25.

68. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.
doi:10.1093/bioinformatics/btp352.

69. Hadfield JD. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The
MCMCglmm R Package. J Stat Softw. 2010;33:1–22. doi:10.18637/jss.v033.i02.

70. Smukowski Heil CS, Ellison C, Dubin M, Noor MAF. Recombining without Hotspots: A
Comprehensive Evolutionary Portrait of Recombination in Two Closely Related Species of
Drosophila. Genome Biol Evol. 2015;7:2829–42. doi:10.1093/gbe/evv182.

71. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, et
al. From fastQ data to high-confidence variant calls: The genome analysis toolkit best
practices pipeline. Curr Protoc Bioinforma. 2013; SUPL.43:11.10.1-11.10.33.
doi:10.1002/0471250953.bi1110s43.

72. Li H. A statistical framework for SNP calling, mutation discovery, association mapping
and population genetical parameter estimation from sequencing data. Bioinformatics.
2011;27:2987–93. doi:10.1093/bioinformatics/btr509.

73. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant
call format and VCFtools. Bioinformatics. 2011;27:2156–8.
doi:10.1093/bioinformatics/btr330.

74. Scheet P, Stephens M. A Fast and Flexible Statistical Model for Large-Scale Population
Genotype Data: Applications to Inferring Missing Genotypes and Haplotypic Phase. Am J
Hum Genet Am J Hum Genet. 2006;7878:629–44. www.ajhg.org. Accessed 17 Jul 2017.

75. Librado P, Rozas J. DnaSP v5: a software for comprehensive analysis of DNA
polymorphism data. Bioinformatics. 2009;25:1451–2. doi:10.1093/bioinformatics/btp187.

76. Leader DP, Krause SA, Pandit A, Davies SA, Dow JAT. FlyAtlas 2: a new version of the
Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq and sex-specific data.
Nucleic Acids Res. 2018;46:D809–15. doi:10.1093/nar/gkx976.

77. Mohn F, Sienski G, Handler D, Brennecke J. The Rhino-Deadlock-Cutoff Complex
Licenses Noncanonical Transcription of Dual-Strand piRNA Clusters in Drosophila. Cell.
2014;157:1364–79. doi:10.1016/j.cell.2014.04.031.

78. Zamparini AL, Davis MY, Malone CD, Vieira E, Zavadil J, Sachidanandam R, et al.
Vreteno, a gonad-specific protein, is essential for germline development and primary piRNA
biogenesis in Drosophila. Development. 2011;138:4039–50. doi:10.1242/dev.069187.

79. Saito K, Ishizu H, Komai M, Kotani H, Kawamura Y, Nishida KM, et al. Roles for the Yb
body components Armitage and Yb in primary piRNA biogenesis in Drosophila. Genes Dev.
2010;24:2493–8. doi:10.1101/gad.1989510.

80. Vourekas A, Zheng K, Fu Q, Maragkakis M, Alexiou P, Ma J, et al. The RNA helicase
MOV10L1 binds piRNA precursors to initiate piRNA processing. Genes Dev. 2015;29:617–
29. doi:10.1101/gad.254631.114.

81. Patil VS, Kai T. Repression of Retroelements in Drosophila Germline via piRNA Pathway
by the Tudor Domain Protein Tejas. Curr Biol. 2010;20:724–30.
doi:10.1016/j.cub.2010.02.046.

82. Ohtani H, Iwasaki YW, Shibuya A, Siomi H, Siomi MC, Saito K. DmGTSF1 is necessary
for Piwi-piRISC-mediated transcriptional transposon silencing in the Drosophila ovary.
Genes Dev. 2013;27:1656–61. doi:10.1101/gad.221515.113.

83. Dönertas D, Sienski G, Brennecke J. Drosophila Gtsf1 is an essential component of the
Piwi-mediated transcriptional silencing complex. Genes Dev. 2013;27:1693–705.
doi:10.1101/gad.221150.113.

84. Sato K, Siomi MC. Functional and structural insights into the piRNA factor Maelstrom.
FEBS Lett. 2015;589:1688–93. doi:10.1016/j.febslet.2015.03.023.

85. Li WH. Unbiased estimation of the rates of synonymous and nonsynonymous
substitution. Journal of Molecular Evolution. 1993;36:96–9. doi:10.1007/BF02407308.

Figures

Figure 1 Expression Profile of Argonaute 2

Plots show the difference in expression between female (red) and male (blue) flies, based on public RNAseq data, normalized to rpL32 and plotted on a natural log scale. The significance of differences between the sexes was assessed using a linear model fitted with MCMCglmm and is denoted by asterisks: * $0.001 < pMCMC < 0.05$; ** $pMCMC <= 0.001$). Sample size (n) represents the number of RNAseq datasets used (combined across tissues). *Ago2d* is the ancestral copy in Dpse and Dmir, *Ago2a* is the ancestral copy in *D. obscura* and *Ago2a* is recently duplicated in Dpse become *Ago2a1* (*Ago2a*) and *Ago2a3*.

Figure 2 The expression profile of RNAi-accessory protein genes

Plots show the expression pattern between sex (male: blue, female:red) for genes other than *Ago2*; 'anc' ancestral copy, 'dup' duplicate copy as inferred by synteny. The y-axis is the natural log of normalized expression. The significance between sexes is denoted by * (0.001 < pMCMC < 0.05) and ** (pMCMC < 0.001). Sample sizes are the same as **Figure 1**.

Figure 3 Bayesian Relaxed Clock Gene Tree for Argonaute2

The duplication events are marked by yellow diamonds, species other than the obscura group are collapsed (brown triangle), and paralog clades are colored. Bayesian posterior supports are only shown for the nodes with support less than 1. Genes not previously included in the analysis of [4] are marked in bold. Time is expressed relative to the split between the obscura and subobscura subgroups (orange boxes), which was constrained to be 1 using a strongly informative prior.

Figure 4 Bayesian Relaxed Clock Trees for 6 RNAi accessory protein genes
Ancestral genes are marked by bold blue, duplicates in bold red. Yellow diamonds indicate duplication events. Species other than *obscura* group are collapsed (green triangle; melanogaster group and brown triangle: other *Drosophila* species). Posterior Bayesian Supports are only shown in the nodes with support less than 1. Duplicated genes which could not be assigned as ancestral or duplicate is marked by _1 or _2. Scale axis is in the time relative to the *obscura* speciation, which was set to 1.

Figure 5 Density plot for posterior distributions of the duplication age

The MCMC posterior of the age of duplication node after 25% burn-in. (**A**) *Argounate 2* and (**B**) RNAi accessory proteins. The broken red line denotes speciation event in *obscura* group which was normalised to be 1.

Tables

Table 1 The details of RNAi accessory genes duplicated in the obscura group as reported by Palmer *et al.* [22]
*The tissue gene expression in *Dmel* is based on the FlyAtlas2 [76]. We report tissues with enrichment > 0.4.

Gene	Involvement in piRNA pathway	Function of the protein product	Tissue expression in *Dmel*	Reference
cutoff (*cuff*)	piRNA transcription	Forms a complex with Rhino-Deadlock-Cutoff (Rhi-Del-Cuff) to protect uncapped non-canonical (dual-strand cluster) piRNA transcript from degradation, splicing and transcription termination	testis	[77]
vreteno (*vret*)	piRNA biogenesis	A Tudor-domain protein which essential for an early primary piRNA processing	larval brain, adult female salivary gland, ovary, testis	[78]
armitage (*armi*)	piRNA biogenesis	A RNA helicase which unwinds the piRNA intermediates before loading into Piwi	ubiquitously expressed	[79] [80]
tejas (*tej*)	Secondary piRNA production (Ping-pong cycle)	A Tudor-domain protein which physically interact with *Vas*, *Spn-E* and *Aub* for a proper ping-pong cycle in the *nuage*	testis, accessory glands, adult female salivary gland, ovary	[81]
asterix (*arx*)	TGS (Transcriptional Gene Silencing)	A zinc-finger protein which directly interacts with Piwi to scan and identify the transposon transcriptions as target for histone modifications	ubiquitously expressed	[82] [83]
maelstrom (*mael*)	TGS	Act downstream of Piwi to establish histone modification and prevent the spreading of the silencing marker to the surrounding genes	brain, testis, adult female salivary gland, ovary	[84]

Table 2 Joint estimates of adaptive evolution across genes

Maximum-likelihood extension of MK test model fitted with different constraints on α [29]. LnL is the log likelihood of the model, AIC is the Akaike Information Criterion with corresponding relative probability as Akaike Weight (w_i). The most supported model is in bold.

Model	Model description	LnL	AICc	Akaike weight(w_i)	Maximum likelihood α estimate	
					ancestral	duplicate
M0	α-anc=0, α-dup=0	-300.61	633.2207	4.00×10^{-12}	0	0
M1	α-anc > 0, α-dup>0 α-anc = α-dup	-284.105	602.2092	2.6×10^{-05}	0.539	0.539
M2	α-anc>0, α-dup=0	-301.857	637.7133	5.10×10^{-13}	0.047	0
M3	α-anc=0, α-dup>0	-275.249	584.4975	0.186	0	0.618
M4	**α-anc > 0, α-dup>0 α-anc ≠ α-dup**	**-272.774**	**581.5471**	**0.813**	**0.2**	**0.676**

Additional files

Additional file 1 Table S1

File format: xlsx
Title: **The detailed RNAi genes and its duplicate in** *D. pseudoobscura*
Description: The genomic position is based on the D. pseudoobscura assembly 3.0

Gene	Flybase ID	Locus Tag	Chromosomal location	Muller Element	Start Position	Gene length	Duplication mechanism
Armi	FBgn0246685	GA25304	4_group1	D	981803	3507	
Arx-ancestral	FBgn0077765	GA17756	XR_group8	D	6958861	495	
Arx-duplicate	FBgn0247462	GA26086	4_group3	B	10387204	501	Retrotransposition
Cuff-ancestral	FBgn0246456	GA25073	3	C	17023607	1113	
Cuff-duplicate	FBgn0244163	GA22760	XL_group1a	A	3011154	1119	Direct DNA duplication
Tejas-ancestral	FBgn0081173	GA21185	3	C	17440474	1782	
Tejas-duplicate	FBgn0248235	GA26863	2	E	1079477	1413	Retrotransposition
Vret-ancestral	FBgn0078422	GA18420	2	E	6266048	2085	
Vret-duplicate	FBgn0244928	GA23527	XL_group1e	A	10126307	2110	
Mael-ancestral	FBgn0248264	GA26892	2	E	54104	1314	
Mael-duplicate	FBgn0249827	GA28467	XR_group8	D	396866	1113	Direct DNA duplication
Ago2a	FBgn0249477	GA28114	Unknown_group_265		16703	2008	Direct DNA duplication
Ago2b	FBgn0248821	GA27454	2	E	23856171	2940	Direct DNA duplication
Ago2c	FBgn0248778	GA27411	2	E	21862037	2915	Direct DNA duplication
Ago2d(ancestral)	FBgn0245029	GA23629	XR_group6	D	200896	2839	
Ago2e	FBgn0247385	GA23629	4_group3	B	6716295	2285	Direct DNA duplication

Additional file 2 Figure S1

File format: pdf
Title: **The expression profile of RNAi across tissue**
Description: The normalized expression plotted across tissues. The error bars denote the standard error for the given tissue. Blue bar indicates male and female is indicated by red bar. The plot is shown for *D. pseudoobscura* (n=163), *D. miranda* (n=42) and *D. obscura* (n=34), respectively.

D. pseudoobscura

D. miranda

D. obscura

Additional file 3 Table S2

File format: xlsx
Title: **MK test of duplicated RNAi genes in** *D. pseudoobscura-D. miranda*
Description: Ds is synonymous divergence, Dn is non-synonomous divergence, Pn is the non-synonomous polymorphisms, Ps is the synonymous polymorphisms, α represents proportion of substitutions that are adaptive, a is the absolute number of adaptive substitution. Ln is the number of non-synonymous sites, Ls is the number of synonymous sites. Ka is the number of non-synonymous mutations per non-synonymous sites and Ks the number of synonymous mutations per synonymous sites from single randomly chosen strain for each species (Li, 1993 [85] calculated using R package seqinr). Parameter ω_a is identical to Ka/Ks ratio except that the numerator only takes adaptive divergence (α * Dn)/Ln)/(Ds/Ls).

Gene	Ds	Dn	Ps	Pn	alpha(α)	Fisher p-value	A	Ln	Ls	Ka	Ks	Ka/Ks	ω_a
Armi	34	72	22	30	0.356	0.220	26	2639.83	810.17	0.0380	0.0538	0.707	0.252
Arx-ancestral	2	9	7	4	**0.905**	**0.017**	8	383.67	102.33	0.0246	0.0232	**1.059**	**0.958**
Arx-duplicate	3	5	10	10	0.400	0.680	2	386.67	105.33	0.0204	0.0400	0.509	0.204
Cuff-ancestral	5	29	10	10	**0.828**	**0.010**	24	863.67	255.33	0.0352	0.0275	**1.281**	**1.060**
Cuff-duplicate	6	39	14	17	**0.813**	**0.003**	32	856.83	250.17	0.0297	0.0417	**0.714**	**0.580**
Tejas-ancestral	11	24	10	23	-0.054	1.000	0	1082.17	312.83	0.0312	0.0293	1.067	0.000
Tejas-duplicate	16	68	9	13	**0.660**	**0.046**	45	1264.83	379.17	0.0222	0.0352	**0.631**	**0.416**
Mael-ancestral	8	27	9	22	0.276	0.586	7	1001.67	303.33	0.0312	0.0293	1.067	0.294
Mael-duplicate	4	28	11	13	**0.831**	**0.007**	23	865.67	247.33	0.0323	0.0162	**2.000**	**1.662**
Vret-ancestral	17	34	37	70	0.054	1.000	2	1610.33	471.67	0.0211	0.0360	0.586	0.032
Vret-duplicate	25	65	23	26	**0.565**	**0.027**	37	1582.83	466.17	0.0517	0.0617	**0.838**	**0.473**
Ago2b	41	52	3	12	**-2.154**	**0.097**	0	882.00	264.00	0.0590	0.1553	**0.380**	**0.000**
Ago2c	2	34	2	22	0.353	1.000	12	1571.83	483.17	0.0216	0.0041	5.226	1.845
Ago2d	27	44	5	16	-0.964	0.300	0	1228.83	367.17	0.0358	0.0735	0.487	0.000
Ago2e	3	18	54	16	**0.951**	**0.000**	17	1410.17	464.83	0.0128	0.0065	**1.978**	**1.881**

Additional file 5 Figure S2

File format: pdf
Title: **The heat-map p-value of the difference between pairwise posterior distributions**
Description: Large p-value (>0.05) indicates the overlapping distribution and the duplication time might be shared (blue box). Red boxes denote comparison with p-value < 0.05 which indicate non-overlapping posterior distribution and an asynchronous duplication events . Pink colored box indicates the marginally significant (0.01 < p-value <0.05).

Additional file 6 Table S3

File format: multiple-tab xlsx
Title: **Additional MK test analysis**
Description: Table the results of additional MK test analysis where minor allele frequency is removed, repetition with larger dataset and results in *D. melanogaster*

Table 1 MK test result in *D. pseudoobscura* with MAF < 12.5% removed

Gene	Ds	Dn	Ps	Pn	α	Fisher p-value
Armi	38	74	7	11	0.193	0.79
Arx-ancestral	**2**	**9**	**3**	**0**	**1**	**0.0274**
Arx-duplicate	3	5	1	1	0.4	1
Cuff-ancestral	5	29	2	8	0.31	1
Cuff-duplicate	**6**	**41**	**5**	**4**	**0.883**	**0.0099**
Tejas-ancestral	11	24	5	11	-0.008	1
Tejas-duplicate	17	69	1	2	0.507	0.496
Mael-ancestral	8	28	4	4	0.714	0.1846
Mael-duplicate	**4**	**31**	**2**	**1**	**0.935**	**0.0592**
Vret-ancestral	18	35	16	27	0.132	0.831
Vret-duplicate	**25**	**68**	**8**	**7**	**0.665**	**0.069**
Ago2b	41	53	2	6	-1.321	0.4621
Ago2c	2	34	0	5	0	0
Ago2d	27	44	2	8	-1.455	0.318
Ago2e	**4**	**19**	**13**	**2**	**0.968**	**0.000037**

Table 2 MK test result for larger pseudoobscura dataset Fuller *et. al* [31]

Gene	Ds	Dn	Ps	Pn	NI	α	Fisher p-value
Armi	**33**	**71**	**36**	**38**	**0.491**	**0.509**	**0.02259**
Arx-ancestral	**2**	**7**	**14**	**4**	**0.082**	**0.918**	**0.011**
Arx-duplicate	3	5	14	11	0.471	0.529	0.438
Cuff-ancestral	**4**	**29**	**18**	**16**	**0.123**	**0.877**	**0.000569**
Cuff-duplicate	**6**	**39**	**15**	**18**	**0.185**	**0.815**	**0.002**
Tejas-ancestral	12	23	11	28	1.328	-0.328	0.62
Tejas-duplicate	**14**	**67**	**16**	**23**	**0.3**	**0.7**	**0.0069**
Mael-ancestral	8	27	7	22	0.724	0.276	0.568
Mael-duplicate	**4**	**28**	**11**	**15**	**0.195**	**0.805**	**0.015**
Vret-ancestral	17	33	44	85	0.995	0.005	1
Vret-duplicate	**25**	**65**	**25**	**27**	**0.415**	**0.585**	**0.018**

Table 3 MK test from DGRP Freeze 1 Dataset (*D. melanogaster*** - ***D. simulans***)**

Gene	Ds	Dn	Ps	Pn	NI	α	Fisher p-value
Arx	**9**	**3**	**3**	**3**	**0**	**0.333**	**0.04**
Ago2	53	100	1	4	2.12	-1.12	0.66
Tejas	35	87	4	20	2.011	-1.011	0.314
Mael	22	36	4	5	0.764	0.236	0.72
Armi	**54**	**64**	**17**	**6**	**0.298**	**0.702**	**0.02**
Vret	45	30	13	17	1.962	-0.962	0.134
Cuff	33	62	5	5	0.532	0.468	0.49