Stage-specific transcriptomes and DNA methylomes indicate an early and

transient loss of transposon control in Arabidopsis shoot stem cells

Ruben Gutzat[1], Klaus Rembart[1], Thomas Nussbaumer[2], Rahul Pisupati[1], Falko Hofmann[1],

Gabriele Bradamante[1], Nina Daubel[1#], Angelika Gaidora[1§], Nicole Lettner[1], Mattia Donà[1],

Magnus Nordborg[1], Michael Nodine[1], Ortrun Mittelsten Scheid [1]

[1] *Gregor Mendel Institute (GMI), Austrian Academy of Sciences, Vienna BioCenter (VBC),*

Dr. Bohr-Gasse 3, 1030 Vienna, Austria.

[2] *Institute of Environmental Medicine, UNIKA-T, Helmholtz Center Munich, Augsburg,*

Germany.

[#] *present address: Uppsala University, Uppsala, Sweden*

[§] *present address: University of Natural Resources and Life Sciences, Vienna, Austria*

In contrast to animals, postembryonic development in plants is modular, and aerial organs originate from stem cells in the center of the shoot apical meristem (SAM) throughout life. Descendants of SAM stem cells in the subepidermal layer (L2) will give also rise to male and female gametes [1] and therefore can be considered primordial germ cells. In these cells, transmission of somatic mutations including virus and TE insertions must be avoided. Despite their essential role for plant development and intergenerational continuity, a comprehensive molecular analysis of SAM plant stem cells has been missing, due to their low number, deep embedding among non-stem cells and difficult isolation. Here we present a comprehensive analysis of stage-specific gene expression and DNA methylation dynamics in Arabidopsis SAM stem cells. This revealed that stem cell expression signatures are mostly defined by development, but we also identified a core set of differentially expressed stemness genes. Surprisingly, vegetative SAM stem cells showed increased expression of transposable elements (TEs) relative to surrounding cells, despite high expression of genes connected to epigenetic silencing. We also find increasing methylation at CHG and a drop in CHH methylation at TEs before stem cells enter the reproductive lineage, indicating an onset of epigenetic reprogramming at an early stage. Transiently elevated TE expression is reminiscent of that in animal primordial germ cells (PGCs) [2] and demonstrates commonality of transposon biology. Our results connect SAM stem cells with germline development and transposon evolution and will allow future experiments to determine the degree of epigenetic heritability between generations.

In contrast to short and straight germlines in animals, plants have multiple germlines in the form of meristems, and, if necessary, these can even be generated *de novo* from differentiated cells. Meristems are tissue in regions of plant growth and contain a few stem cells that give rise to various derivatives, including gametes. In Arabidopsis, the SAM stem cell niche is marked

by expression of the *CLAVATA3 (CLV3)* gene. Other transcription factors, signaling molecules

(including *CLV3*), and receptors (for review see e.g. [3,4]) are necessary for stem cell

maintenance, however, our knowledge of the characteristics of "stemness" and the molecular

signatures of plant stem cells remains very limited. In order to overcome technical difficulties

associated with stem cell isolation, the number of *CLV3*-expressing cells was increased using

an *ap1-1;cal1-1* double-mutant and enabled the first stem cell transcriptome [5,6]. However these

plants show stem cell-related phenotypic aberrations, and the analysis was limited to floral

meristems. A recent study used the INTACT method to gain insights into gene expression and

histone H3 modification dynamics of the entire SAM, but nonetheless included different cell

types and gene expression domains [7]. With the same method, nuclei from SAM stem cells and

differentiated leaves were isolated and compared for chromatin accessibility [8]. We wanted to

obtain information about gene expression and DNA methylation of pure SAM stem cell

fractions and generated Arabidopsis plants expressing a transcriptional fusion of the *CLV3*

promoter [9] and mCherry-labelled histone H2B. Microscopic analysis demonstrated correct and

specific expression of the p*CLV3:mCherry-H2B* marker in nuclei of approximately 20-40 stem

cells in 14-day-old seedlings (Fig. 1a). We applied fluorescence-activated nuclear sorting

(FANS) [10] to nuclei isolated from tissue manually enriched for shoot apical meristems and

collected mCherry-positive and -negative nuclei, with non-transgenic plants as controls (Fig.

1b, Supplementary Fig. 1a, Supplementary Table 1). Microscopic analysis confirmed that all

sorted nuclei from the positive channel appeared intact and displayed red fluorescence,

validating the purity of the fraction (Supplementary Fig. 1b). The transcript level of endogenous

CLV3 was more than 1000-fold higher in mCherry-positive versus mCherry-negative nuclei or

whole seedlings (Fig. 1c). General RNA expression of sorted nuclei was highly correlated with

total RNA expression (Pearson correlation coefficient = 0.94; Supplementary Fig. 1c),

indicating that nuclear RNA represents well the transcriptome of whole cells. Therefore, we

will refer to samples from mCherry/*CLV3*-positive nuclei as stem cells and to mCherry/*CLV3*-

negative samples from the closest neighboring tissue as non-stem cells.

We generated and sequenced RNA expression libraries from stem and non-stem cells isolated

from heart through torpedo stage embryos (E), 7 day- (D7), 14 day- (D14), and 35 day- (D35)

old plants (Supplementary Fig. 2a,b, Supplementary Table 2), aiming (i) to identify expression

signatures in stem cells preceding major developmental switches, (ii) to find genes that are

involved in epigenetic resetting and germline formation, and (iii) to detect "stemness core

genes" whose expression would characterize stem cells independent of development.

Normalized read counts demonstrated high enrichment of *CLV3* and *mCherry* transcripts in

stem cells at all developmental stages (Fig. 2a). High expression of the meristem marker genes

STM and *KNAT1* relative to nuclei of 14-day-old whole seedlings (S14) confirmed the

meristematic features of the non-stem cells (Fig. 2a).

Transcriptome-wide clustering analysis showed that the expression signature of stem cells is

dominated by the respective developmental stage rather than by cell type (Fig. 2b). Pairwise

comparison between stem cells with the respective non-stem cells revealed differentially

expressed genes (DEGs, q <0.05) at all four timepoints, respectively (Fig. 2c), the majority

upregulated in stem cells (with the exception of the embryo samples). Interestingly, GO term

analysis revealed that GOs describing reproductive processes, floral organ development, and

inflorescence development were already enriched in E, D7, and D14 (Supplementary Table 3,

Supplementary Fig. 3), while their absence in D35 stem cells was likely due to the low number

of DEGs.

Overlap analysis between samples (Fig. 2d, Supplementary Fig. 4) revealed many stage-

specific DEGs but also identified a set of 10 core genes (including *CLV3*) that were more highly

expressed in stem cells of all four stages (Fig. 2a, Supplementary Fig. 4 and 5, Supplementary

Table 4), and 23 genes with elevated expression in three out of the four stages (Supplementary

Table 4). Twelve out of these 33 genes encode transcription factors (p-value for enrichment:

1.24e-08). Seven have already been connected with a meristem- or stem cell-related function

(Supplementary Table 4) leaving the remaining 26 as candidates for a potential role in stem cell

maintenance.

We could not detect significant overlap with transcript analysis in the SAM during flower

induction [7], probably due to differences in experimental set up and tissue type. The meristem

transcriptome of the *ap1-1;cal1-1* double-mutant [6] had limited but significant overlap for

upregulated genes (Supplementary Table 5). Comparison with transcriptome data for different

types of root meristem cells [11] resulted in an overlap especially with upregulated genes with

WOX5-expressing cells of the quiescent center (Supplementary Table 5). Also noticeable was

an overlap between upregulated stem cell DEGs with genes related to DNA methylation or

siRNAs highly expressed in meristematic tissue [12] (Fig. 3a). Among these are transcripts of two

Argonaute proteins (AGO5 and AGO9), two histone methyltransferases (SUVH4 and SUVR2),

the nucleosome remodeler DDM1, and three putative RNA-dependent RNA polymerases

(RDR3, 4, and 5). This indicated that specific family members of prominent epigenetic

components were upregulated in stem cells. Since AGO9, SUVH4, and DDM1 (among others)

are necessary for TE repression [13-15], we asked whether TEs were downregulated in stem cells

relative to the surrounding cells. Indeed, several Arabidopsis TE families [16] were 2-fold less

expressed in stem versus non-stem cells through the four stages (Fig. 4 and Supplementary

Table 6). Surprisingly, with the same significance threshold, we found other TE families that

were more highly expressed in stem than in non-stem cells (Fig. 4, Supplementary Table 6).

Strikingly, at D7, the number of highly expressed TE groups coincided with the lowest number

of downregulated other groups, indicating a transient loss of control over TE expression in stem

cells at this early stage of vegetative growth, followed by resilencing towards generational

transition.

TEs overexpressed in D7 were mostly COPIA LTR-retroelements and Mutator-like DNA transposons but also included Helitrons, gypsy-like LTR elements, and SINEs (Supplementary Table 6). As LTR retroelements are more prevalent within pericentromeric regions, whereas SINEs and Helitrons are distributed on chromosome arms [17], TE expression in stem cells occurred independently of chromosomal localization. We could not find a bias for TEs that were recently mobile in natural populations [18], nor for transposons with new insertions in DNA methylation-deficient mutants [19,20].

To determine whether TE expression was influenced by changing DNA methylation, we performed whole-genome bisulfite sequencing of genomic DNA from D7, D14, and D35 stem and non-stem nuclei, with material from 7 d- and 14 d-old seedlings as reference. Modification of cytosines in plants (reviewed in [21]) at CG sites (mCG) is mainly achieved by MET1 and occurs in repetitive sequences as well as along the gene body of protein-coding genes. Cytosine methylation at CHG sites (mCHG) (H = A, C, or T) is installed by CMT2 and CMT3. Cytosine methylation at CHH sites (mCHH) is established by DRM1 and DRM2 as well as CMT2. mCHG and mCHH are mostly restricted to repetitive sequences and important for TE silencing. mCHG is recognized by the histone methyltransferase SUVH4 which methylates histone H3 on lysine 9, a binding site for CMT3, and thereby reinforces DNA methylation in heterochromatic domains [22].

Analysis of DNA methylation distribution revealed pronounced differences around the centromeres for mCHG and mCHH, with the highest mCHG and lowest mCHH portion in stem cells of D35 (Fig. 5a and Supplementary Fig. 6). Congruent with the distribution along the chromosomes, metaplot analyses revealed that these methylation differences were found at TEs, while protein-coding genes were not affected (Fig. 5b). mCHG levels increased with developmental age, and TEs in stem cells had consistently higher mCHG levels than the

respective non-stem cells, reaching a maximum at D35. Conversely, mCHH decreased with

developmental age, most pronounced in stem cells (Fig. 5b).

While TE groups varying in genomic location, cytosine content, structure and localization of

repeats, and siRNA targeting sites [17] showed similarly increasing mCHG and decreasing

mCHH in stem cells over developmental time (Supplementary Fig. 7), there was a correlation

with their length: plotting methylation levels of TEs against their size range (Fig. 5c) revealed

that mCHG in older meristems increased more in long TEs (>2.5 kb), parallel to decreasing

mCHH. This suggests a contribution of DDM1, as this chromatin remodeler mediates

methylation preferentially at long TEs [23].

In order to understand which DNA methylation components are involved in methylation

dynamics in stem cells, we identified differentially methylated regions (DMRs) for each

timepoint and compared them with DMRs of mutants lacking different epigenetic

components [24]. Increased mCHG in stem cells was especially pronounced at D14 in hypo-

DMRs of *ddm1*, *suvh4*, *cmt3*, and *suvh456;* DMRs with reduced mCHH overlapped with those

of *cmt2*, *suvh456*, *ddm1,* and *met1* (Fig. 6). This suggested a concerted action between the

chromatin remodeler DDM1 and the reinforcing heterochromatin formation of CMT3 and

histone methyltransferases to establish strong CHG methylation in stem cells entering the

reproductive phase. Furthermore, the reduction of CHH methylation in *cmt2* and *suvh456*

DMRs indicated a functional interference of CMT3 activity in stem cells with the related

CMT2.

The elevated TE expression at D7 correlated with a minimum of *AGO5* and *AGO9* transcript

expression (Fig. 3b). While they belong to different clusters of the AGO clade [25], both have

been previously identified to be expressed in meristematic tissue of embryos [26,27] or in gametes

or gametophytes [28,29]. AGO5 has not been connected with RNA-directed DNA methylation

(RdDM) of TEs [30], and neither *ago5* nor *ago9* showed many DMRs in DNA of whole

seedlings [24]. However, AGO9 can restore methylation in an *ago4* mutant if accordingly

expressed [26], suggesting that it can substitute RdDM-related functions. Although their exact

molecular functions and the subpopulation of bound small RNAs in the stem cells remain to be

determined, their expression anticorrelated with active TEs could hint to a specific protection

of germline precursor cells from virus and/or TE invasions. The transient loss of TE control in

early vegetative stages might even provide the sequence-specific information via small RNAs

that are then available for stem cell-enriched or -specific silencing components at later stages.

It has often been noted, but not explained, that mutants lacking major components of the RNA-

directed DNA methylation (RdDM) pathway have no or only mild developmental defects.

Reinforced silencing in stem cells during development, involving additional specific factors

like RDR3, 4, or 5, may be responsible for this resilience. Such a barrier might break down only

upon special conditions, as indicated by stress-induced transposition prior to flower formation

that occurs only in RdDM-compromised mutants [19]. Interestingly, a recent study showed that

male premeiotic meiocytes also exhibit high mCG and mCHG methylation and low mCHH [31].

This raises the intriguing possibility that cells of the central zone of the SAM enter a germline

DNA methylation state long before they can be cytologically distinguished. Alternatively, our

data could also suggest the presence of several cell types within the central domain of the SAM.

The possibility to extend the isolation of stem cells at different stages, from mutants and under

different external conditions, will enable future experiments to shed more light on epigenetic

maintenance and dynamics in germline precursor cells.

Data access

DNA bisulfite and RNA-seq data have been deposited in the ArrayExpress database at EMBL-EBI (www.ebi.ac.uk/arrayexpress) under accession number E-MTAB-5478 and E-MTAB-5479.

Methods

Plant material

All experiments were performed with *Arabidopsis thaliana* ecotype Col-0, wild type or transgenic for p*CLV3:H2B-mCherry*. The pCLV3:H2B-mCherry construct was generated as follows: the coding sequence of the H2B gene was PCR-amplified with primer H2B-forward and H2B-reverse (Supplementary Table 7) from cDNA prepared from 14 d-old seedlings. The vector pCLV3:erCFP [9] was cut with *BamH*I and *Sac*I, and the H2B amplicon was inserted (In-Fusion, Clontech) into the open vector. The resulting plasmid was opened with *Sac*I and In-Fusion-filled with a PCR-amplified mCherry-coding fragment using the primers mCherry-fusion-F1 and mCherry-fusion-R1 (Supplementary Table 7). Correct sequence of the resulting vector pCLV3:H2B-mCherry was confirmed by Sanger-sequencing. The construct was used to generate transgenic plants by the floral dip method [32]. Primary transformants were selected with glufosinate (Merck) and their progeny screened for lines with a segregation ratio of 3 resistant to 1 sensitive plant. Homozygous offspring were propagated for seed amplification.

Growth conditions

All plants were grown either *in vitro* on GM medium with or without selection or on soil under a 16 h light/8 h dark regime at 21°C. Material was always harvested at the same time of the light period.

Microscopic analysis and immunostaining

For wide-field microscopy, plant material was immersed in PBS buffer and imaged with a Zeiss

Axio Imager epifluorescence microscope. Isolated nuclei were imaged with an LSM780 Axio

Observer, and Images were deconvolved using Hyugens Core (Scientific Volume Imaging)

with a theoretical PSF. Immunostaining was performed according to [33], with an additional

clearing step using ScaleA [34] and DAPI as counterstain. Anti-mCherry nanobodies were

purchased from Chromotek (#rba594-100). Immunostains of meristems were imaged using the

Airyscan mode on an LSM880 Axio Observer.

Fluorescence-activated nuclei sorting (FANS)

For 7D/14C/35D samples, 200-800 apexes (depending on size) of soil-grown plants with the

corresponding age were collected. For embryo samples, ovules from siliques of a few

representative plants were analyzed to contain early heart till early torpedo stage embryos, and

developmentally identical siliques were used to dissect 3000-4000 ovules. Collected material

was immediately transferred into nuclei isolation buffer on ice (NIB: 500 mM sucrose, 100 mM

KCl, 10 mM Tris-HCl pH 9.5, 10 mM EDTA, 4 mM spermidine, 1 mM spermine and 0.1% v/v

2-mercaptoethanol, prepared just before use [35]). The material was then transferred into a tube

containing 1.8 ml of nuclear extraction buffer (NEB of the Sysmex CyStain® PI Absolute P kit

(#05-5022) plus 1% v/v 2-mercaptoethanol) and disrupted with the TissueRuptor (Qiagen) at

the lowest speed for 1 min. The suspension was filtered (30 µm filter nylon mesh, Sysmex #

04-0042-2316) and centrifuged for 10 min at 4000 rcf at 4°C. The nuclear pellet was

resuspended in Precise P staining buffer (Sysmex #05-5022; plus 1% v/v 2-mercaptoethanol

and DAPI to a final concentration of 5 µg/ul), incubated for 15 min and again filtered (30 µm)

into tubes (Sarstedt #55.484.001). Sorting was performed on a BD FACSAria™ III cell sorter

(70 µm nozzle). Forward/Side scatter and DAPI and mCherry gating were adjusted with wild

type nuclei (DAPI-positive, mCherry-negative) as reference. The mCherry gate was adjusted

so that a maximum of 1/10 of mCherry events occurred in wild type compared to the

p*CLV3:mCherry-H2B* line. For DNA extraction, nuclei were directly sorted into Genomic

Lysis Buffer (Quick-DNA Microprep Kit, Zymo Research, #D3020,), and DNA was purified

according to the suppliers' protocol for whole blood and serum samples. DNA was quantified

using pico-green on a NanoDrop fluorospectrometer (Thermo Scientific). For RNA isolation,

NIB, NEB, and staining buffer were complemented with RiboLock RNase inhibitor (Thermo

Scientific #EO0381, final concentration 1 U/µl) and nuclei were directly sorted into TRIzol LS

(Ambion, #10296028). RNA was prepared according to the manufacturers' recommendation,

except that nuclease-free glycogen (Thermo Scientific) was added during an overnight

precipitation at -20°C. Amount and quality of RNA was determined on an RNA 6000 pico-chip

(Bioanalyzer/Agilent Technologies). For DNA and RNA extraction, DNA-LoBind tubes

(Eppendorf, #022431021) were used.

qPCR analysis

For qPCR and enrichment analysis, RNA was extracted with TRIzol LS (Ambion) either from

sorted nuclei or from shock-frozen and ground tissue material. RNA was treated with DNAse

(Thermo Scientific, #79254) and reverse-transcribed with iScript (Biorad, #172-5038). qPCR

assays were performed with Universal ProbeLibrary (UPL) assays (Roche, # 06402682001)

with primers and probes described in Supplementary Table 7.

Library preparation and sequencing

For RNA library preparation, total RNA of biological duplicates was extracted either from

nuclei directly sorted into TRIzol LS or from shock-frozen ground material and used to generate

cDNA libraries with the SMART-Seq v4 Ultra Low Input RNA Kit (Clontech). For the

comparison with the nuclear RNA transcriptome, RNA was extracted from DAPI-stained

FANSed nuclei isolated from 14 d-old p*CLV3:mCherry-H2B* seedlings with the same protocol

as for cDNA production. cDNA populations were paired-end sequenced on a HiSeq 2500

Illumina sequencing platform. For bisulfite library preparation, at least 200 pg of DNA was

used. Libraries were prepared with the Pico Methyl-Seq Library Prep Kit (Zymo Research

#D5456) according to the manufacturer's protocol.

262   Analysis of the RNA-sequencing data

For the analysis of nuclear to total RNA expression correlation, Tophat [36] was used for mapping

to the TAIR10 reference genome after removal of low-quality bases with Trimmomatic [37]

(parameters: LEADING:8 TRAILING:10 SLIDINGWINDOW:4:15 MINLEN:50;).

Cuffdiff [38] was used for normalization.

For all other analysis, RNA-seq reads were first adapter- and quality-trimmed with Trim

Galore! (Krueger F. Trim galore, v0.4.1, with default parameters). The reads were then aligned

to the TAIR10 reference genome (including the mCherry sequence) with STAR [39] (v2.5.2a)

(Supplementary Table 8). Alignment parameters for STAR were set by the quantifier RSEM [40]

(v1.2.30), which are based on previous ENCODE standards. The annotation used for

quantification was Araport11. RSEM was run with default settings. To correct for possible

positional biases in the data, we activated RSEM's positional bias correction option (--estimate-

rspd). The resulting gene expression tables were imported into R (v3.4) via the tximport

package [41] (v1.4.0). Consecutive differential gene expression analysis was performed with

DESeq2 [42] (v1.16). Samples of the same stages were analyzed pairwise via DESeq2s Wald test

(FDR < 0.05). To detect genes that are differentially regulated in stem cells across all

timepoints, we made use of DESeq2's model-based likelihood ratio test (LRT, FDR < 0.05).

The LRT allowed us to investigate how well the expression of a gene is recapitulated by

different models. DESeq2 compares two models, one full model and a reduced model. Our full

model factored in the cell type, the stage, and the interaction of both, while our reduced model

did not factor in the interaction, leaving us with a set of differentially expressed genes whose

variation can be explained by a combination of cell type and time. The RNA-seq pipeline is

available under https://gitlab.com/nodine-lab/rsem-rna-seq-pipeline/. GO enrichments were

calculated using the AmiGO2 tool and the PANTHER classification system

(http://amigo.geneontology.org/rte) [43]. Visualization and clustering of the data was achieved

using the R packages "gplots" and "gclus".

DEG TE-Families

All RNAseq samples were quality-trimmed using cutadap (v1.14) (Marcel Martin; Cutadapt

removes adapter sequences from high-throughput sequencing reads; EMBnet.journal; Vol17,

No1) and trimmomatic [37] (v0.36). STAR [39] (v2.5.2a) (Col-0 Arabidopsis reference genome, the

Araport11 gene and TE annotations) was used as reference to map the reads, allowing multiple

hits (--outFilterMultimapNmax 100 and --winAnchorMultimapNmax 100). TEtranscripts from

the TEToolkit [44] (v1.5.1) was used in multi-mode to find DEG TE-families.

Analysis of the bisulfite-sequencing data

Illumina HiSeq 2500 sequencing data was obtained from three stages (D7, D14, and D35) each

in three different settings (+: FANS-sorted stem cell tissue, -: non-stem cell but meristematic

tissue, s; whole seedling). Samples D14 and D35 were sequenced with 125 bp paired end reads,

D7 with 50 bp paired end reads (Supplementary Table 8). The data were quality-checked

(fastqc) and trimmed with TrimGalore (Krueger F. Trim galore, v0.4.1, default settings with

stringency = 1) and trimmomatic [37] (v0.36, sliding window: 4:20, leading: 20). Bismark [45]

(v0.18.1 with Bowtie2 v2.2.9) was used to map the reads to the *Arabidopsis thaliana* Col-0

reference genome (including mitochondria and chloroplast genomes) in the non-directional

mode with a mapping stringency of L,0,-0.6. A mapping-position-based removal of duplicates

(Bismark) was applied, and the C-to-T conversion rate was calculated using the reads mapped

to the chloroplast genome (ranging from 98.9 to 99.5%). Methylation was called (Bismark),

ignoring the first bases according to the M-Bias plots. Samples with same stages and settings

were pooled to a single sample, resulting in genome coverages for the nuclear genome from

16,4x to 53,9x.

DMR analysis

Differentially methylation positions (DMP) were identified by Fisher's exact test. Their positions were clustered together based on a minimum distance of 50 bp between DMPs to call a differential methylated region (DMR). DMR calling was done using methylpy (https://github.com/yupenghe/methylpy.git) version 1.1.9. We used custom R and python scripts for further analysis of these DMRs.

Author contributions

R.G. and O.M.S. designed the study and wrote the manuscript, R.G., N.D., A.G., G.B., N.L., and M.D. performed the experiments, K.R., T.N., R.P., and F.H. analyzed the data, Ma.N. and Mi.N. discussed the results and commented on the manuscript.

Acknowledgments

We are grateful for excellent support by the GMI/IMP/IMBA Biooptics facilities, the Next Generation Sequencing and Plant Sciences units of the Vienna BioCenter Core Facilities (VBCF). We thank Claude Becker, Frederic Berger, and James Matthew Watson (all GMI, Vienna) for helpful comments on the manuscript, and Thomas Laux for the CLV3 promoter. We gratefully acknowledge financial support from the Austrian Science Fund (FWF I489, I1477 to O.M.S and I3687 to R.G.) and the Plant Fellows Program (to R.G.).

Competing interests

The authors declare no competing financial interest.

 Figure legends

 Figure 1: Establishment of FANS for stem cells of the shoot apical meristem (SAM).

(**a**) Expression of H2B-mCherry under control of the *CLV3* promoter in 14 d-old seedlings.

Whole-mount immunostaining using α-mCherry antibodies and laser scanning microscopy

(scale bar 10 μm). (**b**) Example of a FANS experiment: mCherry-positive (+) and mCherry-

negative (-) gates of DAPI-gated nuclei. Numbers indicate total number and percent of DAPI

events. (**c**) Enrichment of *CLV3* transcript in mCherry-positive nuclei determined by qRT-PCR.

Figure 2: Differential RNA expression in SAM stem cells during development.

(**a**) Expression of *CLV3*, *mCherry*, and the meristem marker genes *STM* and *KNAT1*. (**b**)

Hierarchical clustering of expression data. (**c**) Number of DEGs between stem and non-stem

cells at each timepoint. The banded portion of the bars indicates the number of transcription

factor genes (also in parenthesis). (**d**) Overlap of genes with higher expression in stem cells

(excluding *mCherry)*. s = stem cells; n = non-stem cells.

Figure 3: Expression analysis of genes related to epigenetic regulation.

(**a**) Expression heatmap (in alphabetical order of gene acronyms). (**b**) Expression of

significantly upregulated DNA methylation-related genes in stem cells, marked with # in (a).

Asterisks indicate timepoints of significantly different expression between stem and non-stem

cells. s = stem cells; n = non-stem cells.

Figure 4: Expression analysis of transposable elements.

(**a**) Heatmap of expression differences for all 318 Arabidopsis TE groups in stem cells relative

to non-stem cells at different timepoints. (**b**) Number of TE groups with at least 2x expression

difference at the different timepoints.

Figure 5: DNA methylation analysis of stem cells at different developmental stages.

(**a**) CG, CHG, and CHH methylation at chromosome 3 in stem and non-stem cells. (**b**)

Metaplots of DNA methylation at CG, CHG, and CHH for genes and transposons. (**c**) Locally

weighted scatterplot smoothing fit of CG, CHG, and CHH methylation levels in stem cells and

non-stem cells plotted on TE size. D7, D14 and D35 = sorted nuclei 7, 14, and 35 d.a.g., S7 and

S14 = above-ground seedlings 7 and 14 d. a. g., s = stem cells; n = non-stem cells.

Figure 6: DMR analysis of stem cells.

DNA methylation differences between stem and non-stem cell nuclei within DMRs of different

epigenetic mutants. For each category, the scale denotes the number of standard deviations of

differential methylation in relation to the rest of the genome.

Supplementary Figure 1: Isolation of stem cell nuclei and RNA comparison.

(**a**) Gating strategy used for FANS of stem cells. Representative FANS plots are shown. Events

are gated for DAPI (top row) and next either for mCherry+ or mCherry- (bottom row). For

numbers see also Supplementary Table 1. (**b**) Examples of mCherry-positive nuclei after FANS

(scale bar 5 μm). (**c**) Correlation of log10-normalized FPKM values of nuclear and total RNA

extracted from 14 d-old seedlings.

Supplementary Figure 2: Growth stages used for genome-wide expression and DNA

methylation analysis in stem and non-stem cells.

(**a**) Developmental stages of representative plants (scale bars 1 cm). (**b**) Wide-field microscopic

images with RFP filters. (**c**) LSM pictures of representative plants. For better visualization

DAPI was used as counterstain in E, D7 and D14. IM = Inflorescence meristem. FM = Floral

meristem. Scale bars in (b): 60 μm for the embryo; 1 mm for the other three stages. Scale bars

in (c): 20 μm.

Supplementary Figure 3: Clustered heatmap displaying GO-term enrichment.

Color codes represent the negative ln of the Bonferroni corrected p-value for enrichment of

each GO-term. A p-value of 0.05 corresponds approximately to 3. See also Supplementary

Table 3 for exact values. s = stem cells; n = non-stem cells.

Supplementary Figure 4: Overlap of DEGs at different timepoints.

(**a**) Venn diagrams for genes up- and (**b**) downregulated in stem cells, respectively. (**c**) p-values

(hypergeometric tests) for likelihood of overlap of upregulated genes in different pairs of

timepoints.

Supplementary Figure 5: Expression of core stemness genes.

Bar plots of expression of genes that are significantly upregulated in SAM stem cells throughout

development. s = stem cells; n = non-stem cells.

Supplementary Figure 6: DNA methylation analysis of stem cells on all five Arabidopsis

chromosomes in stem and non-stem cells at different developmental stages.

s = stem cells; n = non-stem cells.

Supplementary Figure 7: DNA methylation analysis of different TE classes in stem and

non-stem cells at different developmental stages.

s = stem cells; n = non-stem cells.

Supplementary Table 1: Examples of FANS data

Supplementary Table 2: RNA expression data

Supplementary Table 3: GO-term annotations

Supplementary Table 4: Overlapping DEGs

Supplementary Table 5: Comparison with other data sets

Supplementary Table 6: TE expression data

Supplementary Table 7: Primer sequences

Supplementary Table 8: Mapping statistics

References

406
407   1.   d'Amato, F. Role of somatic mutations in the evolution of higher plants *Caryologia* **50**, 1-15
408   (1997).
409   2.   Haig, D. Transposable elements: Self-seekers of the germline, team-players of the soma.
410   *Bioessays* **38**, 1158-1166 (2016).
411   3.   Greb, T. & Lohmann, J.U. Plant stem cells. *Current Biology* **26**, R816-R821 (2016).
412   4.   Soyars, C.L., James, S.R. & Nimchuk, Z.L. Ready, aim, shoot: stem cell regulation of the
413   shoot apical meristem. *Current Opinion in Plant Biology* **29**, 163-168 (2016).
414   5.   Yadav, R.K., Tavakkoli, M., Xie, M., Girke, T. & Reddy, G.V. A high-resolution gene
415   expression map of the Arabidopsis shoot meristem stem cell niche. *Development* **141**, 2735-
416   2744 (2014).
417   6.   Yadav, R.K., Girke, T., Pasala, S., Xie, M. & Reddy, G.V. Gene expression map of the
418   Arabidopsis shoot apical meristem stem cell niche. *Proc Natl Acad Sci U S A* **106**, 4941-6
419   (2009).
420   7.   You, Y. *et al.* Temporal dynamics of gene expression and histone marks at the Arabidopsis
421   shoot meristem during flowering. *Nat Commun* **8**, 15120 (2017).
422   8.   Sijacic, P., Bajic, M., McKinney, E.C., Meagher, R.B. & Deal, R.B. Changes in chromatin
423   accessibility between Arabidopsis stem cells and mesophyll cells illuminate cell type-specific
424   transcription factor networks. *Plant Journal* **94**, 215-231 (2018).
425   9.   Tucker, M.R. *et al.* Vascular signalling mediated by ZWILLE potentiates WUSCHEL
426   function during shoot meristem stem cell development in the Arabidopsis embryo.
427   *Development* **135**, 2839-43 (2008).
428   10.   Zhang, C., Barthelson, R.A., Lambert, G.M. & Galbraith, D.W. Global characterization of
429   cell-specific gene expression through fluorescence-activated sorting of nuclei. *Plant Physiol*
430   **147**, 30-40 (2008).
431   11.   de Luis Balaguer, M.A. *et al.* Predicting gene regulatory networks by combining spatial and
432   temporal gene expression data in Arabidopsis root stem cells. *Proc Natl Acad Sci U S A* **114**,
433   E7632-E7640 (2017).
434   12.   Baubec, T., Finke, A., Mittelsten Scheid, O. & Pecinka, A. Meristem-specific expression of
435   epigenetic regulators safeguards transposon silencing in Arabidopsis. *EMBO Rep* **15**, 446-52
436   (2014).
437   13.   Duran-Figueroa, N. & Vielle-Calzada, J.P. ARGONAUTE9-dependent silencing of
438   transposable elements in pericentromeric regions of Arabidopsis. *Plant Signal Behav* **5**.
439   14.   Hirochika, H., Okamoto, H. & Kakutani, T. Silencing of retrotransposons in arabidopsis and
440   reactivation by the ddm1 mutation. *Plant Cell* **12**, 357-368 (2000).
441   15.   Lippman, Z., May, B., Yordan, C., Singer, T. & Martienssen, R. Distinct mechanisms
442   determine transposon inheritance and methylation via small interfering RNA and histone
443   modification. *Plos Biology* **1**, 420-428 (2003).
444   16.   Arabidopsis_Genome_Initiative. Analysis of the genome sequence of the flowering plant
445   Arabidopsis thaliana. *Nature* **408**, 796-815 (2000).
446   17.   Underwood, C.J., Henderson, I.R. & Martienssen, R.A. Genetic and epigenetic variation of
447   transposable elements in Arabidopsis. *Curr Opin Plant Biol* **36**, 135-141 (2017).
448   18.   Quadrana, L. & Colot, V. Plant transgenerational epigenetics. in *Annual Review of Genetics,*
449   *Vol 50*, Vol. 50 (ed. Bonini, N.M.) 467-491 (2016).
450   19.   Ito, H. *et al.* An siRNA pathway prevents transgenerational retrotransposition in plants
451   subjected to stress. *Nature* **472**, 115-U151 (2011).
452   20.   Tsukahara, S. *et al.* Bursts of retrotransposition reproduced in Arabidopsis. *Nature* **461**, 423-6
453   (2009).
454   21.   Zhang, H., Lang, Z. & Zhu, J.K. Dynamics and function of DNA methylation in plants.
455   *Nature Reviews Molecular Cell Biology* **19**, 489-506 (2018).
456   22.   Du, J.M. *et al.* Mechanism of DNA methylation-directed histone methylation by
457   KRYPTONITE. *Molecular Cell* **55**, 495-504 (2014).
458   23.   Zemach, A. *et al.* The Arabidopsis nucleosome remodeler DDM1 allows DNA
459   methyltransferases to access H1-containing heterochromatin. *Cell* **153**, 193-205 (2013).

460   24.   Stroud, H., Greenberg, M.V.C., Feng, S.H., Bernatavichute, Y.V. & Jacobsen, S.E.
461   Comprehensive analysis of silencing mutants reveals complex regulation of the Arabidopsis
462   methylome. *Cell* **152**, 352-364 (2013).
463   25.   Vaucheret, H. Plant ARGONAUTES. *Trends in Plant Science* **13**, 350-358 (2008).
464   26.   Havecker, E.R. *et al.* The Arabidopsis RNA-directed DNA Methylation Argonautes
465   functionally diverge based on their expression and interaction with target loci. *Plant Cell* **22**,
466   321-334 (2010).
467   27.   Scutt, C.P. *et al.* The identification of candidate genes for a reverse genetic analysis of
468   development and function in the Arabidopsis gynoecium. *Plant Physiology* **132**, 653-665
469   (2003).
470   28.   Olmedo-Monfil, V. *et al.* Control of female gamete formation by a small RNA pathway in
471   Arabidopsis. *Nature* **464**, 628-32 (2010).
472   29.   Tucker, M.R. *et al.* Somatic small RNA pathways promote the mitotic events of
473   megagametogenesis during female reproductive development in Arabidopsis. *Development*
474   **139**, 1399-1404 (2012).
475   30.   Havecker, E.R., Wallbridge, L.M., Fedito, P., Hardcastle, T.J. & Baulcombe, D.C. Metastable
476   differentially methylated regions within Arabidopsis inbred populations are associated with
477   modified expression of non-coding transcripts. *Plos One* **7**(2012).
478   31.   Walker, J. *et al.* Sexual-lineage-specific DNA methylation regulates meiosis in Arabidopsis.
479   *Nat Genet* **50**, 130-137 (2018).
480   32.   Clough, S.J. & Bent, A.F. Floral dip: a simplified method for Agrobacterium-mediated
481   transformation of Arabidopsis thaliana. *Plant Journal* **16**, 735-743 (1998).
482   33.   Pasternak, T. *et al.* Protocol: an improved and universal procedure for whole-mount
483   immunolocalization in plants. *Plant Methods* **11**, 50 (2015).
484   34.   Hama, H. *et al.* Scale: a chemical approach for fluorescence imaging and reconstruction of
485   transparent mouse brain. *Nat Neurosci* **14**, 1481-8 (2011).
486   35.   Pavlova, P., Tessadori, F., de Jong, H.J. & Fransz, P. Immunocytological analysis of
487   chromatin in isolated nuclei. *Methods Mol Biol* **655**, 413-32 (2010).
488   36.   Trapnell, C., Pachter, L. & Salzberg, S.L. TopHat: discovering splice junctions with RNA-
489   Seq. *Bioinformatics* **25**, 1105-1111 (2009).
490   37.   Bolger, A.M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
491   data. *Bioinformatics* **30**, 2114-2120 (2014).
492   38.   Trapnell, C. *et al.* Differential gene and transcript expression analysis of RNA-seq
493   experiments with TopHat and Cufflinks. *Nat Protoc* **7**, 562-78 (2012).
494   39.   Dobin, A. & Gingeras, T.R. Mapping RNA-seq reads with STAR. *Current Protocols in*
495   *Bioinformatics* **51**(2015).
496   40.   Li, B. & Dewey, C.N. RSEM: accurate transcript quantification from RNA-Seq data with or
497   without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).
498   41.   Soneson, C., Love, M.I. & Robinson, M.D. Differential analyses for RNA-seq: transcript-level
499   estimates improve gene-level inferences. *F1000Res* **4**, 1521 (2015).
500   42.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
501   RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
502   43.   Mi, H., Muruganujan, A., Casagrande, J.T. & Thomas, P.D. Large-scale gene function
503   analysis with the PANTHER classification system. *Nat Protoc* **8**, 1551-66 (2013).
504   44.   Jin, Y., Tam, O.H., Paniagua, E. & Hammell, M. TEtranscripts: a package for including
505   transposable elements in differential expression analysis of RNA-seq datasets. *Bioinformatics*
506   **31**, 3593-3599 (2015).
507   45.   Krueger, F. & Andrews, S.R. Bismark: a flexible aligner and methylation caller for Bisulfite-
508   Seq applications. *Bioinformatics* **27**, 1571-1572 (2011).

Figure 1 | Establishment of FANS for stem cells of the shoot apical meristem (SAM).
(**a**) Expression of H2B-mCherry under control of the *CLV3* promoter in 14 d-old seedlings.
Whole-mount immunostaining using α-mCherry antibodies and laser scanning microscopy
(scale bar 10 μm). (**b**) Example of a FANS experiment: mCherry-positive (+) and mCher-
ry-negative (-) gates of DAPI-gated nuclei. Numbers indicate total number and percent of
DAPI events. (**c**) Enrichment of *CLV3* transcript in mCherry-positive nuclei determined by
qRT-PCR.

Figure 2 | Differential RNA expression in SAM stem cells during development. (**a**) Expression of *CLV3*, *mCherry* and the meristem marker genes *STM* and *KNAT1*. (**b**) Hierarchical clustering of expression data. (**c**) Number of DEGs between stem and non-stem cells at each timepoint. The banded portion of the bars indicates the number of transcription factor genes (also in parenthesis). (**d**) Overlap of genes with higher expression in stem cells (excluding *mCherry*). s = stem cells; n = non-stem cells.

Figure 3 | Expression analysis of genes related to epigenetic regulation. (**a**) Expression heatmap (in alphabetical order of gene acronym). (**b**) Expression of significantly upregulated DNA methylation-related genes in stem cells, marked with # in (a). Asterisks indicate time-points of significantly different expression between stem and non-stem cells. s = stem cells; n = non stem cells.

Figure 4 | Expression analysis of transposable elements. (**a**) Heatmap of expression differences for all 318 Arabidopsis TE groups in stem cells relative to non-stem cells at different timepoints. (**b**) Number of TE groups with at least 2x expression difference at the different timepoints.

Figure 5 | DNA methylation analysis of stem cells at different developmental stages. (**a**) CG, CHG and CHH methylation of chromosome 3 in stem and non-stem cells. (**b**) Metaplots of DNA methylation at CG, CHG, and CHH for genes and transposons. (**c**) Locally weighted scatterplot smoothing fit of CG, CHG and CHH methylation levels in stem cells and non-stem cells plotted on TE size. D7, D14 and D35 = sorted nuclei 7, 14, and 35 d.a.g., S7 and S14 = above-ground seedlings 7 and 14 d. a. g., s = stem cells; n = non-stem cells.

Figure 6| DMR analysis of stem cells.
DNA methylation differences between stem and non-stem cell nuclei within DMRs of different epigenetic mutants. For each category, the scale denotes the number of standard deviations of differential methylation in relation to the rest of the genome.

Supplementary Figure 1 | Isolation of stem cell nuclei and RNA comparison. (**a**) Gating strategy used for FANS of stem cells. Representative FANS plots are shown. Events are gated for DAPI (top row) and next either for mCherry+ or mCherry- (bottom row). For numbers see also Supplementary Table S1. (**b**) Examples of mCherry-positive nuclei after FANS (scale bar 5 μm). (**c**) Correlation of log10-normalized FPKM values of nuclear and total RNA extracted from 14 d-old seedlings.

heart-torpedo stage embryos (E) | 7 d. a. g. (D7) | 14 d. a. g. (D14) | 35 d. a. g. (D35) | 14 d. a. g. Col-0 seedling

Supplementary Figure 2 | Growth stages used for genome-wide expression and DNA methylation analysis in stem and non-stem cells. (**a**) Developmental stages of representative plants (scale bars 1 cm). (**b**) Wide-field microscopic images with RFP filters. (**c**) LSM pictures of representative plants. For better visualization DAPI was used as counterstain in E, D7 and D14. IM = Inflorescence meristem. FM = Floral meristem. Scale bars in (b): 60 μm for the embryo; 1 mm for the other three stages. Scale bars in (c): 20 μm.

Supplementary Figure 3 | Clustered heatmap displaying GO-term enrichment. Color codes represent the negative ln of the Bonferroni corrected p-value for enrichment of each GO-term. A p-value of 0.05 corresponds approximately to 3. See also Supplementary Table S3 for exact values. s = stem cells; n = non stem cells.

a

UP

Transcription factor
Epigenetic regulator
Other gene

b

DOWN

c

Supplementary Figure 4 | Overlap of DEGs at different timepoints. (**a**) Venn diagrams for genes up- and (**b**) downregulated in stem cells, respectively. (**c**) p-values (hypergeometric tests) for likelyhood of overlap of upregulated genes in different pairs of timepoints.

Supplementary Figure 5 | Expression of core stemness genes. Bar plots of expression of genes that are significantly upregulated in SAM stem cells throughout development. s = stem cells; n = non stem cells.

Supplementary Figure 6| DNA methylation analysis of stem cells on all five Arabidopsis chromosomes in stem and non-stem cells at different developmental stages. s = stem cells; n = non-stem cells.

Supplementary figure 7| DNA methylation analysis of different TE classes in stem and non-stem cells at different developmental stages. s = stem cells; n = non-stem cells.