1

Characterizing the pathogenic, genomic, and chemical traits of *Aspergillus fischeri*, a close

relative of the major human fungal pathogen *Aspergillus fumigatus*

Matthew E. Mead[a], Sonja L. Knowles[b], Huzefa A. Raja[b], Sarah R. Beattie[c#], Caitlin H.

Kowalski[c], Jacob L. Steenwyk[a], Lilian P. Silva[d], Jessica Chiaratto[d], Laure N.A. Ries[d], Gustavo

H. Goldman[d], Robert A. Cramer[c], Nicholas H. Oberlies[b], and Antonis Rokas[a,*]

[a]Department of Biological Sciences, Vanderbilt University, Nashville, Tennessee, USA

[b]Department of Chemistry and Biochemistry, University of North Carolina at Greensboro,

Greensboro, North Carolina, USA

[c]Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth,

Hanover, New Hampshire, USA

[d]Faculdade de Ciencias Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Brazil

Running Title: Comparing *Aspergillus fischeri* with *A. fumigatus*

[*]Address correspondence to Antonis Rokas, Antonis.rokas@vanderbilt.edu

[#] Department of Pediatrics, University of Iowa Carver College of Medicine, Iowa City, Iowa,

USA

Keywords: fungal pathogenesis, comparative genomics, secondary metabolism, specialized

metabolism, evolution of virulence, *laeA*

Abstract

   Aspergillus fischeri is closely related to *Aspergillus fumigatus*, the major cause of invasive mold infections. Even though *A. fischeri* is commonly found in diverse environments, including hospitals, it rarely causes invasive disease; why that is so is unclear. Comparison of *A. fischeri* and *A. fumigatus* for diverse pathogenic, genomic, and secondary metabolic traits revealed multiple differences for pathogenesis-related phenotypes, including that *A. fischeri* is less virulent than *A. fumigatus* in multiple animal models of disease, grows slower in low oxygen environments, and is more sensitive to oxidative stress. In contrast, the two species exhibit high genomic similarity; ~90% of the *A. fumigatus* proteome is conserved in *A. fischeri*, including 48/49 genes known to be involved in *A. fumigatus* virulence. However, only 10/33 *A. fumigatus* biosynthetic gene clusters (BGCs) likely involved in secondary metabolite production are conserved in *A. fischeri* and only 13/48 *A. fischeri* BGCs are conserved in *A. fumigatus*. Detailed chemical characterization of *A. fischeri* cultures grown on multiple substrates identified multiple secondary metabolites, including two new compounds and one never before isolated as a natural product. Interestingly, an *A. fischeri* deletion mutant of *laeA*, a master regulator of secondary metabolism, produced fewer secondary metabolites and in lower quantities, suggesting that regulation of secondary metabolism is at least partially conserved. These results suggest that the non-pathogenic *A. fischeri* possesses many of the genes important for *A. fumigatus* pathogenicity but is divergent with respect to its ability to thrive under host-relevant conditions and its secondary metabolism.

Importance

Aspergillus fumigatus is the primary cause of aspergillosis, a devastating ensemble of

diseases associated with severe morbidity and mortality worldwide. *A. fischeri* is a close relative

of *A. fumigatus*, but is not generally observed to cause human disease. To gain insights into the

underlying causes of this remarkable difference in pathogenicity, we compared two

representative strains (one from each species) for a range of host-relevant biological and

chemical characteristics. We found that disease progression in multiple *A. fischeri* mouse models

was slower and caused less mortality than *A. fumigatus*. The two species also exhibited different

growth profiles when placed in a range of stress-inducing conditions encountered during

infection, such as low levels of oxygen and the presence of reactive oxygen species-inducing

agents. Interestingly, we also found that the vast majority of *A. fumigatus* genes known to be

involved in virulence are conserved in *A. fischeri*, whereas the two species differ significantly in

their secondary metabolic pathways. These similarities and differences that we identified are the

first step toward understanding the evolutionary origin of a major fungal pathogen.

Introduction

Aspergillosis is a major cause of human morbidity and mortality, resulting in over 200,000 life-threatening infections each year worldwide, and is primarily caused by the fungal pathogen *Aspergillus fumigatus* (1). Multiple virulence traits related to Invasive Aspergillosis (IA) are known for *A. fumigatus*, including thermotolerance, the ability to grow under low oxygen conditions, the ability to acquire micronutrients such as iron and zinc in limiting environments, and the ability to produce a diverse set of secondary metabolites (2).

A. fumigatus thermotolerance, a key trait for its survival inside mammalian hosts, is likely to have arisen through adaptation to the warm temperatures present in decaying compost piles, one of the organism's ecological niches (3-5). The primary route of *A. fumigatus* colonization and infection is through the lung, where oxygen levels have been observed to be as low as 2/3 of atmospheric pressure, and a successful response to this hypoxic environment is required for pathogenesis (6, 7). *A. fumigatus* produces a diverse set of small, bioactive molecules, known as secondary metabolites, which are biosynthesized in pathways that exist outside of primary metabolism. Some of these secondary metabolites and their regulators have been shown to be required for severe disease in mouse models (8-10). Furthermore, a master regulator of secondary metabolism, *laeA*, is also required for full virulence in IA mouse model studies (11, 12).

Other species closely related to *A. fumigatus* are also capable of causing disease, but they are rarely observed in the clinic (2, 13-15). For example, *A. fischeri* is the closest evolutionary relative to *A. fumigatus* for which a genome has been sequenced (16, 17), but is only rarely

reported to cause human disease (2). Recent evolutionary genomic analyses suggest that *A.*

fischeri and *A. fumigatus* last shared a common ancestor approximately 4 million years ago (95%

credible interval: 2 – 7 million years ago) (17). Why *A. fischeri*-mediated disease is less common

than *A. fumigatus*-mediated disease remains an open question. Non-mutually exclusive

possibilities include differences in ecological abundance, lack of species level diagnosis in the

clinic of disease-causing strains, and innate differences in pathogenicity and virulence between

the two species.

Previous studies have suggested that the difference in the frequencies with which the two

species cause disease is unlikely to be solely due to ecological factors, as both can be isolated

from a variety of locales, including soils, fruits, and hospitals (18-20). For example,

approximately 2% of the fungi isolated from the respiratory intensive care unit at Beijing

Hospital were *A. fischeri* compared to approximately 23% of fungal species identified as *A.*

fumigatus (20). While *A. fischeri* is easily isolated from a variety of environments, only a few

cases of human infections have been reported (21-24). Furthermore, numerous recent

epidemiological studies from multiple countries that used state-of-the-art molecular typing

methods were able to identify several rarely isolated pathogenic species closely related to *A.*

fumigatus, such as *A. lentulus* and *A. udagawae*, as the source of 10-15% of human infections but

did not identify *A. fischeri* in any patient sample (13, 14, 25-27).

If ecological factors and lack of precision in species identification cannot explain why *A.*

fischeri is non-pathogenic and *A. fumigatus* is pathogenic, other factors must be responsible. An

early genomic comparison between *A. fumigatus*, *A. fischeri*, as well as the more distantly related

Aspergillus clavatus identified 818 genes that were *A. fumigatus*-specific (16). These genes were

enriched for functions associated with carbohydrate transport and catabolism, secondary

metabolite biosynthesis, and detoxification (16), raising the possibility that the observed

differences in pathogenicity observed between *A. fischeri* and *A. fumigatus* have a molecular

basis.

To gain further insight into why *A. fischeri*-mediated disease is less abundant than *A.*

fumigatus-mediated disease, we took a multi-pronged approach to investigate phenotypic,

genomic, and chemical differences between *A. fischeri* strain NRRL 181 and *A. fumigatus* strain

CEA10. We observed that while *A. fischeri* is able to cause fatal disease in multiple animal

models, its disease progression and response to multiple host-relevant stresses is markedly

different than that of *A. fumigatus*. We also found that while the two organisms' genomes are in

general very similar, the sets of secondary metabolite pathways in each of them exhibit a

surprisingly low level of overlap. Examination of the secondary metabolite profile of *A. fischeri*

identified both previously isolated as well as novel compounds. Finally, construction of a mutant

A. fischeri strain that lacked the *laeA* gene, a master regulator of secondary metabolism, and

examination of its chemical profile suggested that LaeA-mediated regulation of secondary

metabolism in *A. fischeri* closely resembles that of *A. fumigatus*. These results begin to reveal the

molecular differences between *A. fischeri* and *A. fumigatus* related to fungal pathogenesis and

suggest that a functional evolutionary genomic comparison between pathogenic and non-

pathogenic species closely related to *A. fumigatus* harbors great promise for generating insights

into the evolution of fungal disease.

Results

<u>*A. fischeri* is significantly less virulent than *A. fumigatus* in multiple animal models of Invasive</u>

<u>Pulmonary Aspergillosis (IPA)</u>

In contrast to *A. fumigatus*-mediated disease, only a handful of cases of invasive fungal

infections have been reported to be caused by *A. fischeri* (21-24). Given this contrast, we utilized

two immunologically distinct murine IPA models to assess differences in pathogenicity and

virulence between the two species. In a leukopenic murine model, *A. fischeri* NRRL 181 is

significantly less virulent than *A. fumigatus* CEA10, in a dose dependent manner (Fig. 1). Using

an inoculum of $1x10^5$ conidia, *A. fischeri* is completely attenuated in virulence, with 100%

murine survival by day 15 post-fungal challenge. In contrast, inoculation with *A. fumigatus*

results in 100% murine mortality by day 15 (Fig. 1A). Using a higher dose ($2x10^6$) of conidia,

both strains cause 90% mortality by day 14; however, the disease progression is markedly

different. 80% of mice inoculated with *A. fumigatus* succumb to infection by day 4, whereas for

mice inoculated with *A. fischeri*, the first mortality event occurs on day 5, and then one or two

mice succumb each day until day 14 (Fig. 1B). Thus, despite the similar overall mortality at

higher fungal challenge doses, *A. fischeri* is significantly less virulent than A. fumigatus in a

leukopenic murine IPA model.

As the patient population at risk for IA continues to change (28), we also tested a non-

leukopenic triamcinolone (steroid)-induced immune suppression model and observed a

significant reduction in virulence of *A. fischeri* compared to *A. fumigatus* (p<0.0001 by Log-

Rank and Gehan-Breslow-Wilcoxon tests). All mice inoculated with *A. fumigatus* succumbed to

infection by day 3; however, similar to the leukopenic model, mice inoculated with *A. fischeri*

had slower disease progression as monitored by Kaplan-Meier analyses (Fig. 1C).

We observed similar pathogenicity and virulence results when using the *Galleria*

mellonella insect larvae model of aspergillosis (Fig. 1DE). Both low (1×10^6 conidia) and high

(1×10^9 conidia) inoculum experiments showed significant differences between the disease

progression of *A. fischeri* (slower) and *A. fumigatus* (faster) in this insect model of fungal

pathogenicity.

To better understand what is happening *in vivo* during disease progression with *A.*

fischeri and *A. fumigatus,* histological analyses on lungs from the triamcinolone model 3 days

post inoculation were utilized. Histological sections were stained with Gomori methenamine

silver (GMS) to visualize fungal burden and with hematoxylin and eosin (H&E) stain to visualize

host related pathology (Fig. 1F). Overall, mice inoculated with *A. fischeri* had similar numbers of

fungal lesions as those inoculated with *A. fumigatu*s but the lesions caused by the two species

were phenotypically distinct (Fig. 1F). In larger terminal bronchioles infected with *A. fumigatus*,

there was greater fungal growth per lesion, and the growth was observed throughout the

bronchiole itself, extending well into the lumen. These lesions are accompanied by substantial

granulocytic inflammation and obstructs the airways surrounding the hyphae (Fig. 1F). In the

lesions containing *A. fischeri*, the fungal growth is contained to the epithelial lining of the

bronchioles. This pattern of growth is accompanied by inflammation at the airway epithelia,

leaving the airway lumen largely unobstructed (Fig. 1F). The lack of airway obstruction during

A. fischeri infection may contribute to the reduced virulence compared to *A. fumigatus*.

Although the distribution of the fungal lesions varies, there is still significant fungal

growth in mice infected with *A. fischeri*, suggesting that *A. fischeri* is capable of growing within

the immune compromised host. Indeed, we tested the growth rate of *A. fischeri* and *A. fumigatus*

in lung homogenate as a proxy for growth capability within the nutrient environment of the host

and observed no difference between the two strains (Fig. S1). These experiments show that in

multiple models of fungal disease, *A. fischeri* is less virulent than *A. fumigatus*, although *A.*

fischeri is still capable of causing disease using a higher dose and importantly, is able to grow

within the immune compromised murine lung.

When compared to *A. fumigatus*, *A. fischeri* differs in its response to several host-relevant

stresses

Our *in vivo* experiments suggested that the lower virulence of *A. fischeri* is not solely a

result of its inability to grow within the host. Therefore, we hypothesized that an additional

contributing factor was the inability of *A. fischeri* to mitigate host stress. Nutrient fluctuation is a

stress encountered *in vivo* during *A. fumigatus* infection (29). To assess differences in metabolic

plasticity between the two species, we measured the two organisms' growth on media

supplemented with glucose, fatty acids (Tween-80), or casamino acids. Because low oxygen

tension is a significant stress encountered during infection (6), and recently, fitness in low

oxygen has been correlated to virulence of *A. fumigatus* (30), we measured growth of both

species at 37°C in both normoxic (ambient air) and hypoxia-inducing (0.2% O_2, 5% CO_2)

conditions. In normoxia with glucose, fatty acids (Tween-80), or casamino acids supplied as the

carbon source, radial growth of *A. fischeri* was lower than that of *A. fumigatus* (Fig. 2). However,

on rich media both organisms grew equally well (Fig. 2). We also observed a slower growth rate

of *A. fischeri* compared to *A. fumigatus* in the first 16 hours of culture in liquid media supplied

with glucose at 37°C. At 30°C, *A. fischeri* grew the same as, or better than, *A. fumigatus* except

on Tween-80 where *A. fumigatus* had a slight advantage (Fig. S2). Also, *A. fischeri* grew

substantially worse than *A. fumigatus* when grown at 44°C (Fig. S3). To determine relative

fitness in hypoxic liquid environments, we measured the ratio of biomass in liquid culture in

ambient air (normoxia) versus hypoxic (0.2% O_2, 5%CO_2) conditions. *A. fischeri* showed

significantly lower fitness in hypoxic conditions, with about an 8.5-fold lower biomass than *A.*

fumigatus (Fig. 3A). These data suggest that *A. fischeri* is less fit than *A. fumigatus* at 37°C and

in low oxygen conditions, both of which have been shown to impact fungal virulence.

205   Metabolic flexibility, or the ability for an organism to utilize multiple carbon sources

simultaneously, has been suggested to provide a fitness advantage to *Candida albicans* during *in*

vivo growth (31). Metabolic flexibility can be characterized using the glucose analog, 2-

deoxyglucose (2-DG), in combination with an alternative carbon source available *in vivo*, such as

lactate. 2-DG triggers carbon catabolite repression, which shuts down alternative carbon

utilization pathways. However, in *C. albicans* this shut down is delayed and growth occurs on

lactate with 2-DG (31, 32). We tested the metabolic flexibility of both *A. fumigatus* and *A.*

fischeri and observed that while both species can grow in the presence of 2-DG on lactate, the

growth inhibition of *A. fischeri* is higher (~60%) than that of *A. fumigatus* (~35%; Fig. 3B). Even

under low oxygen conditions (5% and 2%), *A. fumigatus* maintains this metabolic flexibility

except under extremely low oxygen conditions (0.2%), whereas *A. fischeri* shows even greater

inhibition at all oxygen tensions of 5% or below. These data suggest that while both species

exhibit some level of metabolic flexibility, *A. fumigatus* appears more metabolically flexible

under a wider range of conditions than *A. fischeri*.

Next, we measured the susceptibility of *A. fischeri* to oxidative stress, cell wall stress, and

antifungal drugs. Interestingly, we observed that *A. fischeri* is more resistant to the intracellular

oxidative stress agent menadione than *A. fumigatus* but more susceptible to the external oxidative

stress agent hydrogen peroxide (Fig. 3CD). As the *in vivo* levels of inflammation caused by the

two species appeared different, we indirectly tested for differences in cell wall pathogen-

associated molecular patterns using the cell wall perturbing agents Congo Red and Calcofluor

White. *A. fumigatus* was significantly more resistant to both agents than *A. fischeri* (Fig. 3C),

suggesting differences in the response to cell wall stress or in the composition and organization

of the cell wall between the two species. These differences are likely important for host immune

cell recognition and interaction, which in turn influences pathology and disease outcome.

Lastly, *A. fischeri* showed enhanced resistance relative to *A. fumigatus* for three of the

four antifungal drugs tested (Table 1), consistent with previous experiments (33). Overall, our

phenotypic data show that the response of *A. fischeri* to host-related stresses and antifungals is

substantially different from that of *A. fumigatus*. Furthermore, our results suggest that increased

growth capability of *A. fumigatus* in low oxygen and in high temperatures are two important

attributes that likely contribute to its pathogenic potential compared to *A. fischeri*.

The proteomes of *A. fumigatus* and *A. fischeri* are highly similar, but their secondary metabolic

pathways show substantial divergence

The large differences in virulence and virulence-related traits we observed between *A.*

fumigatus and *A. fischeri* led us to investigate the genotypic differences that could be

responsible. To describe the genomic similarities and differences between *A. fumigatus* and *A.*

fischeri, we determined how many orthologous proteins and how many species-specific proteins

were present in each genome using a Reciprocal Best BLAST Hit approach (34). We identified

8,737 proteins as being shared between the two species (Fig. S4), representing 88% and 84% of

the *A. fumigatus* and *A. fischeri* proteomes, respectively, and 1,684 *A. fischeri*-specific proteins

(16% of its proteome) and 1,189 *A. fumigatus*-specific proteins (12% of its proteome). To narrow

our search for genes that are absent in *A. fischeri* but are important for *A. fumigatus* disease, we

compiled a list of 49 *A. fumigatus* genes considered to be involved in virulence (Table S1) based

on two previously published articles (35, 36) and extensive literature searches of our own. We

observed that all but one of these virulence-associated genes were also present in *A. fischeri,* a

surprising finding considering the substantial differences observed between the two species in

our animal models of infection. The virulence-associated gene not present in *A. fischeri* is *pesL*

(Afu6g12050), a non-ribosomal peptide synthase that is essential for the synthesis of the

secondary metabolite fumigaclavine C and required for virulence in the *Galleria* model of *A.*

fumigatus infection (37).

Since the only previously described *A. fumigatus* virulence-associated gene not present in

the *A. fischeri* genome (i.e. *pesL*) is involved in secondary metabolism, we investigated the

differences between the repertoire of secondary metabolic pathways present in *A. fumigatus* and

A. fischeri. Using the program antiSMASH (38), we identified 598 secondary metabolic cluster

genes distributed amongst 33 clusters in *A. fumigatus* (Table S2) and 786 secondary metabolite

cluster genes spread out over 48 clusters in *A. fischeri* (Table S3). Of these 598 *A. fumigatus*

genes, 407 (68%) had an orthologous gene that was part of an *A. fischeri* secondary metabolic

gene cluster. This level of conservation of secondary metabolic cluster genes (68%) is much

lower than the amount of conservation observed for the rest of the proteome (88%), illustrating

the rapid rate at which fungal metabolic pathways evolve (39, 40).

We next directly compared the secondary metabolic gene clusters of the two organisms.

An *A. fumigatus* gene cluster was considered conserved in *A. fischeri* if $\geq 90\%$ of its genes were

also present in an *A. fischeri* gene cluster and vice versa. We found that only 10 / 33 *A. fumigatus*

gene clusters are conserved in *A. fischeri* and only 13 / 48 *A. fischeri* gene clusters are conserved

in *A. fumigatus* (Fig. 4), a finding consistent with the low conservation of individual secondary

metabolic genes between the two species. While only 10 *A. fumigatus* gene clusters were

conserved in *A. fischeri*, many other clusters contained one or more orthologs of genes in *A.*

fischeri secondary metabolic gene clusters.

Only one gene cluster (Cluster 18) was completely *A. fumigatus*-specific. Interestingly,

our previous examination of the genomes of 66 *A. fumigatus* strains showed that this cluster was

a "jumping cluster", as it was found to be present in only 5 strains and to reside in three distinct

genomic locations (40). Conversely, there are 10 *A. fischeri*-specific gene clusters that do not

have orthologs in secondary metabolic gene clusters in *A. fumigatus*. One of these gene clusters

is responsible for making helvolic acid [a gene cluster known to be absent from the *A. fumigatus*

strain CEA10 but present in strain Af293 (40)], but the other 9 have not been biochemically

connected to any metabolite.

All the genes required for the production of the mycotoxin gliotoxin are located in a gene

cluster in *A. fischeri* (Fig. S5), and are in fact similar to their *A. fumigatus* orthologs (41), even

though *A. fischeri* is not known to produce this mycotoxin (42). Both the gliotoxin and

acetylaszonalenin gene clusters are adjacent to one another in the *A. fischeri* genome (Fig. S5). In

A. fumigatus, the gliotoxin gene cluster is immediately next to what appears to be a truncated

version of the acetylaszonalenin cluster that lacks portions of the nonribosomal peptide synthase

and acetyltransferase genes as well as the entire indole prenyltransferase gene required for

acetylaszonalenin production. The close proximity of these two gene clusters is noteworthy, as it

is similar to previously reported "super clusters" in *A. fumigatus* and *A. fumigatus*-related strains

(43). These super clusters have been hypothesized to be "evolutionary laboratories" that may

give rise to new compounds and pathways (40).

<u>Isolation and characterization of three new compounds from *A. fischeri*</u>

The relatively low level of conservation of secondary metabolic gene clusters we

observed between *A. fumigatus* and *A. fischeri* led us to characterize the secondary metabolites

produced by *A. fischeri* (Fig. S6) (44-48). The one strain-many compounds (OSMAC) approach

was used to alter the secondary metabolites being biosynthesized in order to produce a diverse

set of molecules (49-52). Depending on the media on which it was grown, *A. fischeri* produced

as few as 4 (Yeast Extract Soy Peptone Dextrose Agar - YESD) or as many as 10 compounds

(Oatmeal Agar – OMA) (Fig. S7). These results showed that culture media influences the

biosynthesis of secondary metabolites in *A. fischeri*, a phenomenon observed in many other fungi

(50, 53).

To characterize the peaks of interest we observed when *A. fischeri* was grown on OMA,

we increased the size of our fungal cultures; doing so yielded seven previously isolated

compounds (sartorypyrone A (**1**), aszonalenin (**4**), acetylaszonalenin (**5**), fumitremorgin A (**6**),

fumitremorgin B (**7**), verruculogen (**8**), and the C-11 epimer of verruculogen TR2 (**9**)) and three

newly biosynthesized secondary metabolites (sartorypyrone E (**2**), 14-epi-aszonapyrone A (**3**),

and 13-*O*-fumitremorgin B (**10**). Two of the secondary metabolites were new compounds (**2** and

3) and one was a new natural product (**10**)) (Fig. 5B). The structures for all 10 compounds were

determined using a set of spectroscopic (1 and 2D NMR) and spectrometric techniques (HRMS).

Our data for sartorypyrone A (**1**) (54), aszonalenin (**4**) (55, 56), acetylaszonalenin (**5**) (54, 57),

fumitremorgin A (**6**) (58, 59), fumitremorgin B (**7**) (60-62), verruculogen (**8**) (63, 64), and the C-

11 epimer of verruculogen TR2 (**9**) (64) correlated well with literature values. The structures of

14-epi-aszonapyrone A (**3**), and 13-*O*-prenyl fumitremorgin B (**10**) were fully characterized in

this study (see Figshare document: https://doi.org/10.6084/m9.figshare.7149167); the structure

elucidation of sartorypyrone E (**2**) is ongoing and will be reported in detail in a forthcoming

manuscript.

Since four secondary metabolites (**5**-**8**) from *A. fischeri* had also been reported from *A.*

fumigatus, we hypothesized that the mechanisms *A. fischeri* employs to regulate its secondary

metabolism would also be similar to those used by *A. fumigatus*. To test this hypothesis, we

constructed a deletion mutant of *laeA* in *A. fischeri* (Fig. S8). LaeA is a master regulator of

secondary metabolism in *A. fumigatus* and a variety of other fungi (65-67). Both the wild type

and Δ*laeA* strains of *A. fischeri* were subjected to LC-MS analysis. The chromatographic profile

of Δ*laeA* showed mass data that corresponded to sartorypyrone A (**1**), sartorypyrone E (**2**), 14-

epi-aszonapyrone A (**3**), aszonalenin (**4**), acetylaszonalenin (**5**), fumitremorgin A (**6**),

verruculogen (**8**), and the C-11 epimer of verruculogen TR2 (**9**). However, the relative

abundance of compounds present was very low compared to the wild type (Fig. 5C).

Fumitremorgin B (**7**) and 13-*O*-prenyl-fumitremorgin B (**10**) were not produced by the Δ*laeA*

mutant at all.

Discussion

A. fumigatus is a major human fungal pathogen, yet its close relative *A. fischeri* is rarely

an agent of human disease. A number of traits that contribute to the virulence of *A. fumigatus*

have been characterized, but their distribution and potential role in *A. fischeri*-mediated disease

was largely unknown. In this study, we thoroughly characterized *A. fischeri* (strain NRRL 181)

and compared it to *A. fumigatus* (strain CEA10) for multiple disease-relevant biological and

chemical differences. Our data shows that *A. fischeri* can grow in a mammalian host but is much

less fit and causes a disease progression quite different than that observed during *A. fumigatus*

infections (Figs. 1 and 2). Further investigations revealed that secondary metabolic genes are

much less conserved than genes in the rest of the genome (Fig. S4), and a chemical analysis of *A.*

fischeri resulted in the identification of both previously identified and new compounds (Fig. 5).

While the biosynthetic pathways producing secondary metabolites in *A. fischeri* and *A. fumigatus*

appear to be quite different, our data suggest that a master regulator of secondary metabolism in

A. fumigatus (*laeA*) possesses a similar role in *A. fischeri* (Fig. 5C).

In order to cause disease, a microbe must be able to respond to the set of diverse and

stressful environments presented by its host. Based on our data, *A. fischeri* is unable to respond

to many of these stresses as well as *A. fumigatus* (Figs. 2-3). We hypothesize that this inability to

thrive under stress contributes to the varying disease progressions observed during our animal

model experiments (Fig. 1). Some or all of the genetic determinants responsible for this

discrepancy in stress response and virulence could reside in the 9 *A. fumigatus*-specific genes we

identified (Fig. S4); alternatively, some of the ~1,700 *A. fischeri*-specific genes we identified

may inadvertently facilitate control of *A. fischeri* in a mammalian host. Importantly, our analyses

herein were conducted on single strains of the two species and further studies are needed to

determine how representative these observed trait and genomic differences are across strains.

Even though more than 10% of the genes in each species lack an ortholog in the other

species, only ~2% (1/49) of previously identified genetic determinants of virulence in *A.*

fumigatus are not conserved in *A. fischeri* (Table S1). This result, and our observation that many

of the pathways of secondary metabolism are quite different between *A. fischeri* and *A.*

fumigatus, support a multifactorial model of *A. fumigatus* virulence (1, 68, 69) and suggest a

need to investigate virulence on multiple levels of biological complexity. In order to cause

disease in a host, *A. fumigatus* (and other species closely related to it) must adhere and germinate

in the lung (69), survive inherently stressful conditions presented by host environments (ex.

severe lack of metals and oxygen) (6, 70, 71), and modulate or endure actions of the host

immune system (72). Given the diversity of these activities, it is unlikely that single genes or

pathways will be responsible for the totality of *A. fumigatus*-derived disease, even though not all

genes in the genome have been characterized for their role in pathogenicity. We hypothesize that

multiple pathways (including those involved in secondary metabolism) have changed during the

evolution of *A. fischeri* and *A. fumigatus*, resulting in their differing ability to cause disease.

A. fumigatus and *A. fischeri* are members of *Aspergillus* section *Fumigati*, a clade that

includes multiple closely related species, some of which are pathogens (e.g., *A. fumigatus*, *A.*

lentulus, and *A. udagawae*) and some of which are considered non-pathogens (e.g., *A. fischeri*, *A.*

aureolus, and *A. turcosus*) (2, 42, 73, 74). The ability to cause disease in humans appears to have

either arisen or been lost (or both) multiple times independently during the evolution of this

lineage, as pathogenic species are spread throughout the phylogeny (17, 75). A broader,

phylogenetically-informed comparison of pathogenic and non-pathogenic species in section

Fumigati would provide far greater resolution in identifying (or dismissing) factors and pathways

that may contribute or prevent the ability of these organisms to cause disease. Also, leveraging

the diversity of section *Fumigati* would give researchers a better understanding of the nature and

evolution of human fungal pathogenesis as the appreciation for the health burden caused by fungi

increases (76).

An important caveat to our experiments is that we only analyzed a single, representative

strain from each species. Several recent studies have identified a wide variety of differences

between *A. fumigatus* strains, which have in turn been shown to contribute to physiological

differences, including but not limited to secondary metabolism and virulence (30, 40, 72, 73).

While the genome of only one isolate of *A. fischeri* has so far been sequenced (16) and the

organism has only been reported to cause human disease a few times (21-24), it would be of

great interest to compare patient-derived and environment-derived isolates at the genomic,

phenotypic, and chemical levels. Although it appears that clinical and environmental isolates do

not stem from separate lineages in *A. fumigatus* (77), whether this is also the case for largely

non-pathogenic species, such as *A. fischeri*, or for rarely isolated pathogenic species, such as *A.*

lentulus or *A. udagawae*, remains largely unknown.

Materials and Methods

<u>Strains and growth media</u>

A. fischeri strain NRRL 181 was acquired from the ARS Culture Collection (NRRL). *A.*

fumigatus strain CEA10 (CBS 144.89) was obtained from the CBS. All strains were grown on

glucose minimal media (GMM) from conidial glycerol stocks stored at -80°C. All strains were

grown in the presence of white light at 37°C. Conidia were collected in 0.01% Tween-80 and

enumerated with a hemocytometer.

<u>Murine virulence studies</u>

For the chemotherapeutic (leukopenic) murine model, outbred CD-1 female mice

(Charles River Laboratories, Raleigh, NC, USA), 6-8 weeks old, were immunosuppressed with

intraperitoneal (i.p.) injections of 150 mg/kg cyclophosphamide (Baxter Healthcare Corporation,

Deerfield, IL, USA) 48 hours before and 72 hours after fungal inoculation, along with

subcutaneous (s.c.) injections of 40 mg/kg Kenalog-10 (triamcinolone acetonide, Bristol-Myer

Squibb, Princeton, NJ, USA) 24 hours before and 6 days after fungal inoculation. For the murine

triamcinolone model outbred CD-1 female mice, 6-8 weeks old, were treated with 40 mg/kg

Kenalog-10 by s.c. injection 24 hours prior to fungal inoculation.

423    For both models, conidial suspensions of $2x10^6$ conidia were prepared in 40 μL sterile

424 PBS and administered to mice intranasally while under isoflourine anesthesia. Mock mice were

425 given 40 μL PBS. Mice were monitored three times a day for signs of disease for 14 or 18 days

426 post-inoculation. Survival was plotted on Kaplan-Meir curves and statistical significance

427 between curves was determined using Mantel-Cox Log-Rank and Gehan Breslow-Wilcoxon

428 tests. Mice were housed in autoclaved cages at 4 mice per cage with HEPA filtered air and

429 autoclaved food and water available at libitum.

431 *Galleria mellonella* virulence studies

432    *G. mellonella* larvae were obtained by breeding adult moths (78). *G. mellonella* larvae of

433 a similar size were selected (approximately 275–330 mg) and kept without food in glass

434 container (Petri dishes), at 37°C, in darkness for 24 h prior to use. *A. fumigatus* and *A.*

435 *fischeri* conidia were obtained by growing on YAG media culture for 2 days. The conidia were

436 harvested in PBS and filtered through a Miracloth (Calbiochem). The concentration of conidia

437 was estimated by using hemocytometer, and resuspended at a concentration of 2.0 x 10^8

438 conidia/ml. The viability of the conidia was determined by incubating on YAG media culture, at

439 37°C, 48 hours. Inoculum (5 µl) of conidia from both strains were used to investigate the

440 virulence of *A. fumigatus* and *A. fischeri* against *G. mellonella*. Ten *G. mellonella* in the final

441 (sixth) instar larval stage of development were used per condition in all assays. The control

442 group was the larvae inoculated with 5 µl of PBS to observe the killing due to physical trauma.

443 The inoculum was performed by using Hamilton syringe (7000.5KH) and 5 µl into the haemocel

444 of each larva via the last left proleg. After, the larvae were incubated in glass container (Petri

dishes) at 37ºC in the dark. The larval killing was scored daily. Larvae were considered dead by

presenting the absence of movement in response to touch.

Histopathology

Outbred CD-1 mice, 6-8 weeks old, were immunosuppressed and intranasally inoculated

with $2x10^6$ conidia as described above for the chemotherapeutic and corticosteroid murine

models. Mice were sacrificed 72 hours post inoculation. Lungs were perfused with 10% buffered

formalin phosphate before removal, then stored in 10% buffered formalin phosphate until

embedding. Paraffin embedded sections were stained with haematoxylin and eosin (H&E) and

Gömöri methenamine silver (GMS). Slides were analyzed microscopically with a Zeiss Axioplan

2 imaging microscope (Carl Zeiss Microimaging, Inc. Thornwood, NY, USA) fitted with a

Qimiging RETIGA-SRV Fast 1394 RGB camera. Analysis was performed in Phylum Live 4

imaging software.

Ethics Statement

We carried out our mouse studies in strict accordance with the recommendations in the

Guide for the Care and Use of Laboratory Animals of the National Research Council (Council,

1996). The mouse experimental protocol was approved by the Institutional Animal Care and Use

Committee (IACUC) at Dartmouth College (Federal-Wide Assurance Number: A3259-01).

Growth Assays

Radial growth was quantified by point inoculation of $1x10^3$ conidia in 2 μL on indicated

media; plates were incubated at 37°C in normoxia (~21% O_2, 5% CO_2) or hypoxia (0.2% O_2, 5%

CO_2). Colony diameter was measured every 24 hours for 4 days and reported as the average of

three biological replicates per strain.

For 2-DG experiments, $1x10^3$ conidia in 2 μL were spotted on 1% lactate minimal media

with or without 0.1% 2-deoxyglucose (2-DG; Sigma, D8375). Plates were incubated for 3 days

at 37°C in normoxia or hypoxia with 5% CO_2. Percent inhibition was calculated by dividing

radial growth on 2-DG plates by the average radial growth of biological triplicates on plates

without 2-DG.

Fungal biomass was quantified by measuring the dry weight of fungal tissue from $5x10^7$

conidia grown in 100 mL liquid GMM shaking at 200 rpm for 48 hours in normoxia (~21% O_2)

and hypoxia (0.2% O_2, 5% CO_2). Liquid biomass is reported as the average of three biological

replicates per strain. Hypoxic conditions were maintained using an $INVIVO_2$ 400 Hypoxia

Workstation (Ruskinn Technology Limited, Bridgend, UK) with a gas regulator and 94.8% N_2.

Liquid growth curves were performed with conidia adjusted to $2x10^4$ conidia in 20 μL

0.01% Tween-80 in 96-well dishes, then 180 μL of media (GMM or lung homogenate) was

added to each well. Plates were incubated at 37°C for 7 hours, then Abs_{405} measurements were

taken every 10 minutes for the first 16 hours of growth with continued incubation at 37°C. Lung

homogenate media was prepared as follows: lungs were harvested from healthy CD-1 female

mice (20-24 g) and homogenized through a 100 μM cell strainer in 2 mL PBS/lung. Homogenate

was diluted 1:4 in sterile PBS, spun down to remove cells, then filter sterilized through 22 μM

PVDF filters.

Cell wall and oxidative stresses

493   Congo Red (0.5 mg/mL), Menadione (20 µM), or calcofluor white (CFW, 25 µg/mL)

494 were added to GMM plates. $1x10^3$ conidia (Calcofluor white and Menadione) or $1x10^5$ conidia

495 (Congo Red) were point inoculated and plates were incubated for 96 hours at 37°C with 5% CO_2.

Orthology Determination and Analyses

Genomes for *A. fumigatus* CEA10 and *A. fischeri* NRRL 181 were downloaded from

NCBI (Accession numbers of GCA_000150145.1 and GCF_000149645.1, respectively). To

identify putative orthologous genes between *A. fischeri* and *A. fumigatus,* a reciprocal best

BLAST hit (RBBH) approach was used. We blasted the proteome of *A. fischeri* to *A. fumigatus*

and vice versa using an e-value cutoff of 10^{-3} and then filtered for RBBHs according to bitscore

(79). A pair of genes from each species was considered orthologous if their best blast hit was to

each other. Species-specific and orthologous protein sets were visualized using version 3.0.0 of

eulerAPE (80).

Secondary Metabolism Cluster Prediction and Analyses

Version 4.2.0 of antiSMASH (38) was used with its default settings to identify secondary

metabolite clusters. Orthologous cluster genes were identified using our RBBH results and

visualized using version 0.69 of Circos (81). Chromosomes were identified for *A. fischeri*

NRRL1 and *A. fumigatus* CEA10 using NUCMER (82)and chromosomal sequences from *A.*

fumigatus strain AF293 from NCBI (Accession number GCA_000002655.1). Syntenic clusters

were visualized using easyfig version 2.2.2 (83).

Secondary Metabolite Extraction and Identification

Secondary metabolites were extracted from *A. fischeri* using techniques well established

in the Natural Products literature (84, 85). This was done by adding a 1:1 mixture of

$CHCl_3:CH_3OH$ and left to shake overnight. The resulting slurry was partitioned twice, first with

a 4:1:5 $CHCl_3:CH_3OH:H_2O$ solution, with the organic layer drawn off and evaporated to dryness

in vaccuo, and secondly reconstituting 1:1:2 $CH_3CN:CH_3OH$:hexanes, where the organic layer

was drawn off and evaporated to dryness. The extract then underwent chromatographic

separation (flash chromatography and HPLC) using varied gradient systems. The full structural

characterization of the new secondary metabolites is provided in the Figshare document

(https://doi.org/10.6084/m9.figshare.7149167).

Construction of the *A. fischeri ΔlaeA* mutant

The gene replacement cassettes were constructed by *''in vivo''* recombination in *S.*

cerevisiae as previously described by (86, 87). Approximately 2.0 kb from the 5'-UTR and 3'-

UTR flanking regions of the targeted ORF regions were selected for primer design. The primers

pRS NF010750 5´fw (5´-

GTAACGCCAGGGTTTTCCCAGTCACGACGCAGTCTAACGCTGGGCCCTTCC-3´) and

pRS NF010750 3´rv (5´-

GCGGTTAACAATTTCTCTCTGGAAACAGCTACGGCGTTTGACGGCACAC-3´) contained

a short homologous sequence to the Multicloning site (MCS) of the plasmid pRS426. Both the

5'- and 3'-UTR fragments were PCR-amplified from *A. fischeri* genomic DNA (gDNA). The

prtA gene, conferring resistance to pyrithiamine, which was placed within the cassette as a

dominant marker, was amplified from the pPRT1 plasmid by using the primers prtA NF010750

5´rv (5´-GTAATCAATTGCCCGTCTGTCAGATCCAGGTCGAGGAGGTCCAATCGG-3´)

and prtA NF010750 3´fw (5´-

CGGCTCATCGTCACCCCATGATAGCCGAGATCAATCTTGCATCC-3´). The deletion

cassette was generated by transforming each fragment along with the plasmid pRS426 cut with

*Bam*HI/*Eco*RI into the *S. cerevisiae* strain SC94721, using the lithium acetate method (88). The

DNA from the transformants was extracted by the method described by Goldman et al. (89). The

cassette was PCR-amplified from these plasmids utilizing TaKaRa Ex Taq™ DNA Polymerase

(Clontech Takara Bio) and used for *A. fisheri* transformation according to the protocol described

by Malavazi and Goldman (87). Southern blot and PCR analyses were used to demonstrate that

the cassette had integrated homologously at the targeted *A. fischeri* locus. Genomic DNA from

A. fischeri was extracted by grinding frozen mycelia in liquid nitrogen and then gDNA was

extracted as previously described (87). Standard techniques for manipulation of DNA were

carried out as described (90). For Southern blot analysis, restricted chromosomal DNA fragments

were separated on 1% agarose gel and blotted onto Hybond N$^+$ nylon membranes (GE

Healthcare). Probes were labeled using [α-^{32}P]dCTP using the Random Primers DNA Labeling

System (Life Technologies). Labeled membranes were exposed to X-ray films, which were

scanned for image processing. Southern blot and PCR schemes are shown in Fig. S8.

Acknowledgements

Computational infrastructure was provided by The Advanced Computing Center for

Research and Education (ACCRE) at Vanderbilt University. MEM, JS, and AR were supported

by a Vanderbilt University Discovery Grant. Research in AR's lab is also supported by the

National Science Foundation (DEB-1442113), the Guggenheim Foundation, and the Burroughs

Wellcome Fund. RAC holds an Investigator in the Pathogenesis of Infectious Diseases Award

supported by the Burroughs Wellcome Fund and is also supported by a National Institute of

Allergy and Infectious Diseases award 1R01AI130128. SRB was supported, in part, by the

National Institute of General Medical Sciences of the National Institutes of Health under Award

Number T32GM008704. SLK was supported by the National Center for Complementary and

Integrative Health, a component of the National Institutes of Health, under award number T32

AT008938. GHG was supported by grants from Fundação de Amparo à Pesquisa do Estado de

São Paulo (FAPESP) and Conselho Nacional de Desenvolvimento Científico e Tecnológico

(CNPq), both from Brazil.

References

1.   **Latgé JP**. 1999. *Aspergillus fumigatus* and aspergillosis. Clin Microbiol Rev **12**:310–350.

2.   **Lamoth F**. 2016. *Aspergillus fumigatus*-Related Species in Clinical Practice. Frontiers in
Microbiology **7**:683.

3.   **Tekaia F**, **Latgé J-P**. 2005. *Aspergillus fumigatus*: saprophyte or pathogen? Current
Opinion in Microbiology **8**:385–392.

4.   **Bhabhra R**, **Miley MD**, **Mylonakis E**, **Boettner D**, **Fortwendel J**, **Panepinto JC**,
Postow M, **Rhodes JC**, **Askew DS**. 2004. Disruption of the *Aspergillus fumigatus* gene
encoding nucleolar protein CgrA impairs thermotolerant growth and reduces virulence.
Infect Immun **72**:4731–4740.

5.   **Wagener J**, **Echtenacher B**, **Rohde M**, **Kotz A**, **Krappmann S**, **Heesemann J**, **Ebel F**.
2008. The Putative α-1,2-Mannosyltransferase AfMnt1 of the Opportunistic Fungal
Pathogen *Aspergillus fumigatus* Is Required for Cell Wall Stability and Full Virulence.
Eukaryotic Cell **7**:1661–1673.

6.   **Grahl N**, **Puttikamonkul S**, **Macdonald JM**, **Gamcsik MP**, **Ngo LY**, **Hohl TM**,
Cramer RA. 2011. *In vivo* hypoxia and a fungal alcohol dehydrogenase influence the
pathogenesis of invasive pulmonary aspergillosis. PLoS Pathog **7**:e1002145.

7.   **Willger SD**, **Puttikamonkul S**, **Kim K-H**, **Burritt JB**, **Grahl N**, **Metzler LJ**, **Barbuch
R**, **Bard M**, **Lawrence CB**, **Cramer RA**. 2008. A sterol-regulatory element binding
protein is required for cell polarity, hypoxia adaptation, azole drug resistance, and
virulence in *Aspergillus fumigatus*. PLoS Pathog **4**:e1000200.

8.   **Rokas A**, **Wisecaver JH**, **Lind AL**. 2018. The birth, evolution and death of metabolic
gene clusters in fungi. Nature Reviews Microbiology **163**:150.

9.   **Keller NP**, **Turner G**, **Bennett JW**. 2005. Fungal secondary metabolism — from
biochemistry to genomics. Nature Reviews Microbiology **3**:937–947.

10.   **Bignell E**, **Cairns TC**, **Throckmorton K**, **Nierman WC**, **Keller NP**. 2016. Secondary
metabolite arsenal of an opportunistic pathogenic fungus. Phil Trans R Soc B
371:20160023–9.

11.   **Sugui JA**, **Pardo J**, **Chang YC**, **Müllbacher A**, **Zarember KA**, **Galvez EM**, **Brinster
L**, **Zerfas P**, **Gallin JI**, **Simon MM**, **Kwon-Chung KJ**. 2007. Role of *laeA* in the
Regulation of *alb1*, *gliP*, Conidial Morphology, and Virulence in *Aspergillus fumigatus*.
Eukaryotic Cell **6**:1552–1561.

12.   **Bok JW**, **Balajee SA**, **Marr KA**, **Andes D**, **Nielsen KF**, **Frisvad JC**, **Keller NP**. 2005.
LaeA, a regulator of morphogenetic fungal virulence factors. Eukaryotic Cell **4**:1574–
1582.

13. **Balajee SA**, **Kano R**, **Baddley JW**, **Moser SA**, **Marr KA**, **Alexander BD**, **Andes D**,
Kontoyiannis DP, **Perrone G**, **Peterson S**, **Brandt ME**, **Pappas PG**, **Chiller T**. 2009.
Molecular identification of *Aspergillus* species collected for the Transplant-Associated
Infection Surveillance Network. J Clin Microbiol **47**:3138–3141.

14. **Alastruey-Izquierdo A**, **Mellado E**, **Peláez T**, **Pemán J**, **Zapico S**, **Alvarez M**,
Rodríguez-Tudela JL, **Cuenca-Estrella M**, **FILPOP Study Group**. 2013. Population-
based survey of filamentous fungi and antifungal resistance in Spain (FILPOP Study).
Antimicrob Agents Chemother **57**:3380–3387.

15. **van der Linden JWM**, **Arendrup MC**, **Warris A**, **Lagrou K**, **Pelloux H**, **Hauser PM**,
Chryssanthou E, **Mellado E**, **Kidd SE**, **Tortorano AM**, **Dannaoui E**, **Gaustad P**,
Baddley JW, **Uekötter A**, **Lass-Flörl C**, **Klimko N**, **Moore CB**, **Denning DW**,
Pasqualotto AC, **Kibbler C**, **Arikan-Akdagli S**, **Andes D**, **Meletiadis J**, **Naumiuk L**,
Nucci M, **Melchers WJG**, **Verweij PE**. 2015. Prospective multicenter international
surveillance of azole resistance in *Aspergillus fumigatus*. Emerging Infect Dis **21**:1041–
1044.

16. **Fedorova ND**, **Khaldi N**, **Joardar VS**, **Maiti R**, **Amedeo P**, **Anderson MJ**, **Crabtree J**,
Silva JC, **Badger JH**, **Albarraq A**, **Angiuoli S**, **Bussey H**, **Bowyer P**, **Cotty PJ**, **Dyer**
PS, **Egan A**, **Galens K**, **Fraser-Liggett CM**, **Haas BJ**, **Inman JM**, **Kent R**, **Lemieux S**,
Malavazi I, **Orvis J**, **Roemer T**, **Ronning CM**, **Sundaram JP**, **Sutton G**, **Turner G**,
Venter JC, **White OR**, **Whitty BR**, **Youngman P**, **Wolfe KH**, **Goldman GH**, **Wortman**
JR, **Jiang B**, **Denning DW**, **Nierman WC**. 2008. Genomic islands in the pathogenic
filamentous fungus *Aspergillus fumigatus*. PLoS Genet **4**:e1000046.

17. **Steenwyk JL**, **Shen X-X**, **Lind AL**, **Goldman GG**, **Rokas A**. 2018. A robust
phylogenomic timetree for biotechnologically and medically important fungi from
Aspergillaceae (Eurotiomycetes, Ascomycota). bioRxiv 370429.

18. **Hong S-B**, **Kim D-H**, **Park I-C**, **Samson RA**, **Shin H-D**. 2010. Isolation and
Identification of *Aspergillus* Section *Fumigati* Strains from Arable Soil in Korea.
Mycobiology **38**:1.

19. **Frąc M**, **Jezierska-Tys S**, **YAGUCHI T**. 2015. Occurrence, Detection, and Molecular
and Metabolic Characterization of Heat- Resistant Fungi in Soils and Plants and Their
Risk to Human Health. Advances in Agronomy. Elsevier Ltd.

20. **Tong X**, **Xu H**, **Zou L**, **Cai M**, **Xu X**, **Zhao Z**, **Xiao F**, **Li Y**. 2017. High diversity of
airborne fungi in the hospital environment as revealed by meta-sequencing-based
microbiome analysis. Sci Rep **7**:39606.

21. **Summerbell RC**, **de Repentigny L**, **Chartrand C**, **St Germain G**. 1992. Graft-related
endocarditis caused by *Neosartorya fischeri var. spinosa*. J Clin Microbiol **30**:1580–1582.

22. **Lonial S**, **Williams L**, **Carrum G**, **Ostrowski M**, **McCarthy P**. 1997. *Neosartorya*
fischeri: an invasive fungal pathogen in an allogeneic bone marrow transplant patient.
Bone Marrow Transplant **19**:753–755.

23.   **Gerber J**, **Chomicki J**, **Brandsberg JW**, **Jones R**, **Hammerman KJ**. 1973. Pulmonary
645   aspergillosis caused by *Aspergillus fischeri var. spinosus*: report of a case and value of
646   serologic studies. Am J Clin Pathol **60**:861–866.

647   24.   **Coriglione G**, **Stella G**, **Gafa L**, **Spata G**, **Oliveri S**, **Padhye AA**, **Ajello L**. 1990.
648   *Neosartorya fischeri var fischeri* (Wehmer) Malloch and Cain 1972 (anamorph:
649   *Aspergillus fischerianus* Samson and Gams 1985) as a cause of mycotic keratitis. Eur J
650   Epidemiol **6**:382–385.

651   25.   **Escribano P**, **Peláez T**, **Muñoz P**, **Bouza E**, **Guinea J**. 2013. Is azole resistance in
652   *Aspergillus fumigatus* a problem in Spain? Antimicrob Agents Chemother **57**:2815–2820.

653   26.   **Negri CE**, **Gonçalves SS**, **Xafranski H**, **Bergamasco MD**, **Aquino VR**, **Castro PTO**,
654   **Colombo AL**. 2014. Cryptic and rare *Aspergillus* species in Brazil: prevalence in clinical
655   samples and in vitro susceptibility to triazoles. J Clin Microbiol, 7 ed. **52**:3633–3640.

656   27.   **Sabino R**, **Verissimo C**, **Parada H**, **Brandao J**, **Viegas C**, **Carolino E**, **Clemons KV**,
657   **Stevens DA**. 2014. Molecular screening of 246 Portuguese *Aspergillus* isolates among
658   different clinical and environmental sources. Med Mycol **52**:519–529.

659   28.   **Abers MS**, **Ghebremichael MS**, **Timmons AK**, **Warren HS**, **Poznansky MC**, **Vyas
660   JM**. 2016. A Critical Reappraisal of Prolonged Neutropenia as a Risk Factor for Invasive
661   Pulmonary Aspergillosis. Open Forum Infect Dis **3**:ofw036.

662   29.   **Beattie SR**, **Mark KMK**, **Thammahong A**, **Ries LNA**, **Dhingra S**, **Caffrey-Carr AK**,
663   **Cheng C**, **Black CC**, **Bowyer P**, **Bromley MJ**, **Obar JJ**, **Goldman GH**, **Cramer RA**.
664   2017. Filamentous fungal carbon catabolite repression supports metabolic plasticity and
665   stress responses essential for disease progression. PLoS Pathog **13**:e1006340–29.

666   30.   **Kowalski CH**, **Beattie SR**, **Fuller KK**, **McGurk EA**, **Tang Y-W**, **Hohl TM**, **Obar JJ**,
667   **Cramer RA**. 2016. Heterogeneity among Isolates Reveals that Fitness in Low Oxygen
668   Correlates with *Aspergillus fumigatus*. Virulence. mBio **7**.

669   31.   **Childers DS**, **Raziunaite I**, **Mol Avelar G**, **Mackie J**, **Budge S**, **Stead D**, **Gow NAR**,
670   **Lenardon MD**, **Ballou ER**, **MacCallum DM**, **Brown AJP**. 2016. The Rewiring of
671   Ubiquitination Targets in a Pathogenic Yeast Promotes Metabolic Flexibility, Host
672   Colonization and Virulence. PLoS Pathog **12**:e1005566.

673   32.   **Sandai D**, **Yin Z**, **Selway L**, **Stead D**, **Walker J**, **Leach MD**, **Bohovych I**, **Ene IV**,
674   **Kastora S**, **Budge S**, **Munro CA**, **Odds FC**, **Gow NAR**, **Brown AJP**. 2012. The
675   evolutionary rewiring of ubiquitination targets has reprogrammed the regulation of carbon
676   assimilation in the pathogenic yeast *Candida albicans*. mBio **3**.

677   33.   **Houbraken J**, **Weig M**, **Groß U**, **Meijer M**, **Bader O**. 2016. *Aspergillus
678   oerlinghausenensis*, a new mould species closely related to *A. fumigatus*. FEMS Microbiol
679   Lett **363**:fnv236.

34. **Salichos L**, **Rokas A**. 2011. Evaluating ortholog prediction algorithms in a yeast model
clade. PLoS ONE **6**:e18755.

35. **Abad A**, **Fernández-Molina JV**, **Bikandi J**, **Ramírez A**, **Margareto J**, **Sendino J**,
Hernando FL, **Pontón J**, **Garaizar J**, **Rementeria A**. 2010. What makes *Aspergillus*
fumigatus a successful pathogen? Genes and molecules involved in invasive aspergillosis.
Rev Iberoam Micol **27**:155–182.

36. **Kjærbølling I**, **Vesth TC**, **Frisvad JC**, **Nybo JL**, **Theobald S**, **Kuo A**, **Bowyer P**,
Matsuda Y, **Mondo S**, **Lyhne EK**, **Kogle ME**, **Clum A**, **Lipzen A**, **Salamov A**, **Ngan**
CY, **Daum C**, **Chiniquy J**, **Barry K**, **LaButti K**, **Haridas S**, **Simmons BA**, **Magnuson**
JK, **Mortensen UH**, **Larsen TO**, **Grigoriev IV**, **Baker SE**, **Andersen MR**. 2018.
Linking secondary metabolites to gene clusters through genome sequencing of six diverse
Aspergillus species. Proc Natl Acad Sci USA.

37. **O'Hanlon KA**, **Gallagher L**, **Schrettl M**, **Jöchl C**, **Kavanagh K**, **Larsen TO**, **Doyle S**.
2012. Nonribosomal Peptide Synthetase Genes *pesL* and *pes1* Are Essential for
Fumigaclavine C Production in *Aspergillus fumigatus*. Appl Environ Microbiol **78**:3166–
3176.

38. **Blin K**, **Wolf T**, **Chevrette MG**, **Lu X**, **Schwalen CJ**, **Kautsar SA**, **Suarez Duran HG**,
de los Santos ELC, **Kim HU**, **Nave M**, **Dickschat JS**, **Mitchell DA**, **Shelest E**, **Breitling**
R, **Takano E**, **Lee SY**, **Weber T**, **Medema MH**. 2017. antiSMASH 4.0—improvements
in chemistry prediction and gene cluster boundary identification. Nucleic Acids Res
45:W36–W41.

39. **Lind AL**, **Wisecaver JH**, **Smith TD**, **Feng X**, **Calvo AM**, **Rokas A**. 2015. Examining
the evolution of the regulatory circuit controlling secondary metabolism and development
in the fungal genus *Aspergillus*. PLoS Genet **11**:e1005096.

40. **Lind AL**, **Wisecaver JH**, **Lameiras C**, **Wiemann P**, **Palmer JM**, **Keller NP**, **Rodrigues**
F, **Goldman GH**, **Rokas A**. 2017. Drivers of genetic diversity in secondary metabolic
gene clusters within a fungal species. PLoS Biol **15**:e2003583.

41. **Gardiner DM**, **Howlett BJ**. 2005. Bioinformatic and expression analysis of the putative
gliotoxin biosynthetic gene cluster of *Aspergillus fumigatus*. FEMS Microbiol Lett
248:241–248.

42. **Frisvad JC**, **Larsen TO**. 2016. Extrolites of *Aspergillus fumigatus* and Other Pathogenic
Species in *Aspergillus* Section *Fumigati*. Frontiers in Microbiology **6**:1485.

43. **Wiemann P**, **Guo C-J**, **Palmer JM**, **Sekonyela R**, **Wang CCC**, **Keller NP**. 2013.
Prototype of an intertwined secondary-metabolite supercluster. Proc Natl Acad Sci USA
110:17065–17070.

44. **El-Elimat T**, **Raja HA**, **Day CS**, **Chen W-L**, **Swanson SM**, **Oberlies NH**. 2014.
Greensporones: Resorcylic Acid Lactones from an Aquatic *Halenospora* sp. J Nat Prod
77:2088–2098.

45.   **El-Elimat T**, **Raja HA**, **Figueroa M**, **Falkinham JO**, **Oberlies NH**. 2014.
719   Isochromenones, isobenzofuranone, and tetrahydronaphthalenes produced by *Paraphoma*
720   *radicina*, a fungus isolated from a freshwater habitat. Phytochemistry **104**:114–120.

721   46.   **El-Elimat T**, **Raja HA**, **Day CS**, **McFeeters H**, **McFeeters RL**, **Oberlies NH**. 2017. α-
722   Pyrone derivatives, tetra/hexahydroxanthones, and cyclodepsipeptides from two
723   freshwater fungi. Bioorg Med Chem **25**:795–804.

724   47.   **Raja HA**, **Paguigan ND**, **Fournier J**, **Oberlies NH**. 2017. Additions to *Lindgomyces*
725   (Lindgomycetaceae, Pleosporales, Dothideomycetes), including two new species
726   occurring on submerged wood from North Carolina, USA, with notes on secondary
727   metabolite profiles. Mycol Progress **16**:535–552.

728   48.   **Rivera-Chávez J**, **Raja HA**, **Graf TN**, **Gallagher JM**, **Metri P**, **Xue D**, **Pearce CJ**,
729   **Oberlies NH**. 2017. Prealamethicin F50 and related peptaibols from *Trichoderma*
730   *arundinaceum*: Validation of their authenticity via *in situ* chemical analysis. RSC Adv
731   **7**:45733–45751.

732   49.   **Sanchez JF**, **Somoza AD**, **Keller NP**, **Wang CCC**. 2012. Advances in *Aspergillus*
733   secondary metabolite research in the post-genomic era. Nat Prod Rep **29**:351–371.

734   50.   **Bode HB**, **Bethe B**, **Höfs R**, **Zeeck A**. 2002. Big effects from small changes: possible
735   ways to explore nature's chemical diversity. Chembiochem **3**:619–627.

736   51.   **Vandermolen KM**, **Raja HA**, **El-Elimat T**, **Oberlies NH**. 2013. Evaluation of culture
737   media for the production of secondary metabolites in a natural products screening
738   program. AMB Express **3**:71.

739   52.   **Hemphill CFP**, **Sureechatchaiyan P**, **Kassack MU**, **Orfali RS**, **Lin W**, **Daletos G**,
740   **Proksch P**. 2017. OSMAC approach leads to new fusarielin metabolites from *Fusarium*
741   *tricinctum*. J Antibiot **70**:726–732.

742   53.   **Frisvad JC**, **Andersen B**, **Thrane U**. 2008. The use of secondary metabolite profiling in
743   chemotaxonomy of filamentous fungi. Mycol Res **112**:231–240.

744   54.   **Eamvijarn A**, **Gomes NM**, **Dethoup T**, **Buaruang J**, **Manoch L**, **Silva A**, **Pedro M**,
745   **Marini I**, **Roussis V**, **Kijjoa A**. 2013. Bioactive meroditerpenes and indole alkaloids from
746   the soil fungus *Neosartorya fischeri* (KUFC 6344), and the marine-derived fungi
747   *Neosartorya laciniosa* (KUFC 7896) and *Neosartorya tsunodae* (KUFC 9213).
748   Tetrahedron **69**:8583–8591.

749   55.   **Kimura Y**, **Hamasaki T**, **Nakajima H**, **Isogai A**. 1982. Structure of aszonalenin, a new
750   metabolite of *Aspergillus zonatus*. Tetrahedron Letters **23**:225–228.

751   56.   **Ruchti J**, **Carreira EM**. 2014. Ir-Catalyzed Reverse Prenylation of 3-Substituted Indoles:
752   Total Synthesis of (+)-Aszonalenin and (−)-Brevicompanine B. J Am Chem Soc
753   **136**:16756–16759.

57. **Ellestad GA**, **Mirando P**, **Kunstmann MP**. 1973. Structure of the metabolite LL-
S490.beta. from an unidentified *Aspergillus* species. J Org Chem **38**:4204–4205.

58. **Yamazaki M**, **Fujimoto H**, **Kawasaki T**. 1980. Chemistry of tremorogenic metabolites.
I. Fumitremorgin A from *Aspergillus fumigatus*. Chem Pharm Bull **28**:245–254.

59. **Feng Y**, **Holte D**, **Zoller J**, **Umemiya S**, **Simke LR**, **Baran PS**. 2015. Total Synthesis of
Verruculogen and Fumitremorgin A Enabled by Ligand-Controlled C–H Borylation. J Am
Chem Soc **137**:10160–10163.

60. **Pohland AE**, **Schuller PL**, **Steyn PS**, **Van Egmond HP**. 1982. Physicochemical data for
some selected mycotoxins. Pure and Applied Chemistry **54**:2219–2284.

61. **Afiyatullov SS**, **Kalinovskii AI**, **Pivkin MV**, **Dmitrenok PS**, **Kuznetsova TA**. 2005.
Alkaloids from the Marine Isolate of the Fungus *Aspergillus fumigatus*. Chemistry of
Natural Compounds **41**:236–238.

62. **Mundt K**, **Wollinsky B**, **Ruan H-L**, **Zhu T**, **Li S-M**. 2012. Identification of the
verruculogen prenyltransferase FtmPT3 by a combination of chemical, bioinformatic and
biochemical approaches. Chembiochem **13**:2583–2592.

63. **Fayos J**, **Lokensgard D**, **Clardy J**, **Cole RJ**, **Kirksey JW**. 1974. Letter: Structure of
verruculogen, a tremor producing peroxide from *Penicillium verruculosum*. J Am Chem
Soc **96**:6785–6787.

64. **Fill TP**, **Asenha HBR**, **Marques AS**, **Ferreira AG**, **Rodrigues-Fo E**. 2013. Time course
production of indole alkaloids by an endophytic strain of *Penicillium brasilianum*
cultivated in rice. Nat Prod Res **27**:967–974.

65. **Bok JW**, **Keller NP**. 2004. LaeA, a regulator of secondary metabolism in *Aspergillus* spp.
Eukaryotic Cell **3**:527–535.

66. **Hoff B**, **Kamerewerd J**, **Sigl C**, **Mitterbauer R**, **Zadra I**, **Kürnsteiner H**, **Kück U**.
2010. Two components of a velvet-like complex control hyphal morphogenesis,
conidiophore development, and penicillin biosynthesis in *Penicillium chrysogenum*.
Eukaryotic Cell **9**:1236–1250.

67. **Wiemann P**, **Brown DW**, **Kleigrewe K**, **Bok JW**, **Keller NP**, **Humpf H-U**, **Tudzynski
B**. 2010. FfVel1 and FfLae1, components of a velvet-like complex in *Fusarium fujikuroi*,
affect differentiation, secondary metabolism and virulence. Mol Microbiol **77**:972–994.

68. **Dagenais TRT**, **Keller NP**. 2009. Pathogenesis of *Aspergillus fumigatus* in Invasive
Aspergillosis. Clin Microbiol Rev **22**:447–465.

69. **Ben-Ami R**, **Lewis RE**, **Kontoyiannis DP**. 2010. Enemy of the (immunosuppressed)
state: an update on the pathogenesis of *Aspergillus fumigatus* infection. Br J Haematol
150:406–417.

70.   **Hissen AHT**, **Wan ANC**, **Warwas ML**, **Pinto LJ**, **Moore MM**. 2005. The *Aspergillus*
fumigatus siderophore biosynthetic gene *sidA*, encoding L-ornithine N5-oxygenase, is
required for virulence. Infect Immun **73**:5493–5503.

71.   **Moreno MA**, **Ibrahim-Granet O**, **Vicentefranqueira R**, **Amich J**, **Ave P**, **Leal F**,
Latgé J-P, **Calera JA**. 2007. The regulation of zinc homeostasis by the ZafA
transcriptional activator is essential for *Aspergillus fumigatus* virulence. Mol Microbiol
64:1182–1197.

72.   **Cramer RA**, **Rivera A**, **Hohl TM**. 2011. Immune responses against *Aspergillus*
fumigatus: what have we learned? Curr Opin Infect Dis **24**:315–322.

73.   **Mellado E**, **Alcazar-Fuoli L**, **Cuenca-Estrella M**, **Rodriguez-Tudela JL**. 2011. Role of
Aspergillus lentulus 14-α sterol demethylase (Cyp51A) in azole drug susceptibility.
Antimicrob Agents Chemother **55**:5459–5468.

74.   **Sugui JA**, **Vinh DC**, **Nardone G**, **Shea YR**, **Chang YC**, **Zelazny AM**, **Marr KA**,
Holland SM, **Kwon-Chung KJ**. 2010. *Neosartorya udagawae* (*Aspergillus udagawae*),
an emerging agent of aspergillosis: how different is it from *Aspergillus fumigatus*? J Clin
Microbiol **48**:220–228.

75.   **Hubka V**, **Barrs V**, **Dudová Z**, **Sklenář F**, **Kubátová A**, **Matsuzawa T**, **Yaguchi T**,
Horie Y, **Nováková A**, **Frisvad JC**, **Talbot JJ**, **Kolařík M**. 2018. Unravelling species
boundaries in the *Aspergillus viridinutans* complex (section *Fumigati*): opportunistic
human and animal pathogens capable of interspecific hybridization. Persoonia.

76.   2017. Stop neglecting fungi. Nat Microbiol **2**:17120.

77.   **Abdolrasouli A**, **Rhodes J**, **Beale MA**, **Hagen F**, **Rogers TR**, **Chowdhary A**, **Meis JF**,
Armstrong-James D, **Fisher MC**. 2015. Genomic Context of Azole Resistance
Mutations in *Aspergillus fumigatus* Determined Using Whole-Genome Sequencing. mBio,
2nd ed. **6**:e00536.

78.   **Fuchs BB**, **O'Brien E**, **Khoury JBE**, **Mylonakis E**. 2010. Methods for using *Galleria*
mellonella as a model host to study fungal pathogenesis. Virulence **1**:475–482.

79.   **Madden T**. 2013. The BLAST Sequence Analysis Tool. *In* The NCBI Handbook
[Internet]. 2nd edition. National Center for Biotechnology Information (US).

80.   **Micallef L**, **Rodgers P**. 2014. eulerAPE: Drawing Area-Proportional 3-Venn Diagrams
Using Ellipses. PLoS ONE **9**:e101717–18.

81.   **Krzywinski M**, **Schein J**, **Birol I**, **Connors J**, **Gascoyne R**, **Horsman D**, **Jones SJ**,
Marra MA. 2009. Circos: an information aesthetic for comparative genomics. Genome
Res **19**:1639–1645.

82.   **Kurtz S**, **Phillippy A**, **Delcher AL**, **Smoot M**, **Shumway M**, **Antonescu C**, **Salzberg**
SL. 2004. Versatile and open software for comparing large genomes. Genome Biol **5**:R12.

83.   **Sullivan MJ**, **Petty NK**, **Beatson SA**. 2011. Easyfig: a genome comparison visualizer.
Bioinformatics **27**:1009–1010.

84.   **El-Elimat T**, **Figueroa M**, **Ehrmann BM**, **Cech NB**, **Pearce CJ**, **Oberlies NH**. 2013.
High-resolution MS, MS/MS, and UV database of fungal secondary metabolites as a
dereplication protocol for bioactive natural products. J Nat Prod **76**:1709–1716.

85.   **Paguigan ND**, **El-Elimat T**, **Kao D**, **Raja HA**, **Pearce CJ**, **Oberlies NH**. 2017.
Enhanced dereplication of fungal cultures via use of mass defect filtering. J Antibiot
70:553–561.

86.   **Colot HV**, **Park G**, **Turner GE**, **Ringelberg C**, **Crew CM**, **Litvinkova L**, **Weiss RL**,
Borkovich KA, **Dunlap JC**. 2006. A high-throughput gene knockout procedure for
Neurospora reveals functions for multiple transcription factors. Proc Natl Acad Sci USA
103:10352–10357.

87.   **Malavazi I**, **Goldman GH**. 2012. Gene disruption in *Aspergillus fumigatus* using a PCR-
based strategy and in vivo recombination in yeast. Methods Mol Biol **845**:99–118.

88.   **Schiestl RH**, **Gietz RD**. 1989. High efficiency transformation of intact yeast cells using
single stranded nucleic acids as a carrier. Curr Genet **16**:339–346.

89.   **Goldman GH**, **Reis Marques dos E**, **Duarte Ribeiro DC**, **de Souza Bernardes LA**,
Quiapin AC, **Vitorelli PM**, **Savoldi M**, **Semighini CP**, **de Oliveira RC**, **Nunes LR**,
Travassos LR, **Puccia R**, **Batista WL**, **Ferreira LE**, **Moreira JC**, **Bogossian AP**,
Tekaia F, **Nobrega MP**, **Nobrega FG**, **Goldman MHS**. 2003. Expressed sequence tag
analysis of the human pathogen *Paracoccidioides brasiliensis* yeast phase: identification
of putative homologues of *Candida albicans* virulence and pathogenicity genes.
Eukaryotic Cell **2**:34–48.

90.   **Sambrook J**, **Russell DW**. 2001. Molecular Cloning, 3rd ed. Cold Spring Harbor
Laboratory Press, London.

91.   **Yin W-B**, **Grundmann A**, **Cheng J**, **Li S-M**. 2009. Acetylaszonalenin biosynthesis in
Neosartorya fischeri. Identification of the biosynthetic gene cluster by genomic mining
and functional proof of the genes by biochemical investigation. J Biol Chem **284**:100–109.

Tables

Table 1. *A. fischeri* **shows enhanced resistance relative to** *A. fumigatus* **for several**

antifungal drugs.

Strain	Posaconazole [µg/ml]	Voriconazole [µg/ml]	Itraconazole [µg/ml]	Caspofungin [µg/ml]
A. fumigatus	0.7	0.8	5	0.09
A. fischeri	2.4	> 4	> 24	0.06

 Figure Legends

Figure 1: *A. fischeri* is significantly less virulent than *A. fumigatus* in multiple murine

models of invasive pulmonary aspergillosis. AB) Cumulative survival of mice inoculated with

$1x10^5$ (A) or $2x10^6$ (B) conidia in a leukopenic model of IPA. A) n=10/group B) n=12/group,

4/PBS. *p=0.0098 by Log-Rank test, p=0.0002 by Gehan-Breslow-Wilcoxon test. C) Cumulative

survival of mice inoculated with 2e6 conidia in a triamcinolone model of IPA. n=12/group,

4/PBS. *p=<0.0001 by Log-Rank and Gehan-Breslow-Wilcoxon tests. D) and E) Cumulative

survival of *G. mellonella* larvae inoculated with $1x10^6$ (D) or $1x10^9$ (E) conidia. 10 larvae were

used per condition in all assays. Survival curves for *A. fischeri* and *A. fumigatus* were

significantly different (p<0.003) in both Log-Rank and Gehan-Breslow-Wilcoxon tests for both

inoculums. F) Histological sections from 3 days post inoculation in a triamcinolone model of

IPA stained with H&E and GMS. Images were acquired at 100x.

Figure 2: *A. fischeri* is unable to thrive under suboptimal metabolic conditions at 37°C.

$1x10^3$ conidia were point inoculated on each plate then plates were incubated at 37°C in

normoxia (N; ~21% oxygen, 5%CO_2) or hypoxia (H; 0.2% O_2, 5%CO_2); colony diameter was

measured every 24 hours. Mean and SEM of triplicates. CAA – Casamino acids; GMM –

glucose minimal media.

Figure 3: *A. fischeri* is more susceptible to multiple host-relevant stresses than *A. fumigatus*.

A) Fitness ratio of *A. fumigatus* or *A. fischeri* during hypoxic vs normoxic growth (measured as

the dry weight of cultures). Data represent mean and SEM of biological triplicates; ***p=0.0006

by Student's t-test. B) Growth inhibition of strains grown on 1% lactate minimal media with

0.1% 2-deoxyglucose (2-DG) under a range of low oxygen conditions. C) *A. fumigatus* and *A.*

fischeri were grown in the presence of the cell wall perturbing agent Congo Red (0.5mg/mL), the

oxidative stressor Menadione (20 µM), or the chitin perturbing agent calcofluor white (CFW,

25µg/mL). Plates were grown for 96 hours at 37°C and 5% CO_2. For all plates except Congo Red

and its GMM control, 1e3 spores were plated. For Congo red and the control GMM plate 1e5

spores were plated. Student's t-test was performed where *: $p<0.05$, **: $p<0.01$. D) Strains were

grown for 48 h at 37°C in liquid complete medium supplemented with increasing concentrations

of hydrogen peroxide.

Figure 4: Secondary Metabolite Clusters of *A. fumigatus* and *A. fischeri* show substantial

evolutionary divergence. Predicted secondary metabolite gene clusters are shown in the inner

track, are alternatively colored dark and light gray, and their size is proportional to the number of

genes in them. Black ticks on the exterior of the cluster track indicate a gene that possesses an

ortholog in the other species but is not in a secondary metabolite gene cluster in the second

species. White dots indicate species-specific clusters. Solid bars on the exterior correspond to the

chromosome on which the clusters below them reside. Genes are connected to their orthologs in

the other species with dark lines if >90% of the cluster genes in *A. fumigatus* are conserved in the

same cluster in *A. fischeri*. Lighter lines connect all other orthologs that are present in both

species' sets of secondary metabolite clusters. Image was made using Circos version 0.69-4 (81).

Figure 5: Secondary metabolite production in *A. fischeri*. A) Compounds isolated from *A.*

fischeri: (**1**) sartorypyrone A, (**2**) sartorypyrone E, (**3**) 14-epimer aszonapyrone A, (**4**)

aszonalenin, (**5**) acetylaszonalenin, (**6**) fumitremorgin A, (**7**) fumitremorgin B, (**8**) verruculogen,

(9) C-11 epimer verruculogen TR2, and **(10)** 13-*O*-prenyl-fumitremorgin B. The color coding

indicates which putative class the molecule belongs to; e.g., terpenes, PKS, or NRPS. B) Top,

Aspergillus fischeri was initially grown on rice for two weeks, and then extracted using methods

outlined in Fig. S6. The rice culture yielded compounds **1**, **4**, and **5**. Middle, *A. fischeri* was

grown on multigrain Cheerios for two weeks, which yielded compounds **1** and **4-9**. Bottom, *A.*

fischeri on Quaker oatmeal for two weeks. All compounds that were previously isolated in rice

and multigrain cheerios cultures in addition to three new compounds (**2**, **3**, and **10**) were found in

the oatmeal culture. All pictures depict fungi growing in 250 mL Erlenmeyer flasks; left panel

indicates top view, while the right panel shows bottom view. All chromatographic profiles have

been normalized to the highest μAU value. C) *Aspergillus fischeri* WT and Δ*laeA* were grown on

solid breakfast oatmeal for two weeks and extracted using organic solvents as indicated

previously. The crude de-sugared and de-fatted extracts were run using UPLC-MS at a

concentration of 2 mg/mL with 5 μL being injected for analysis. The chromatographic profiles

were normalized to the highest μAU value. Mass spec analysis indicated the presence of

secondary metabolites **1**–**10** within the wild type, and only **1-6**, **8**, and **9** were seen in the *ΔlaeA*

mutant. All pictures show *A. fischeri* grown on oatmeal agar in Petri plates.

Supplementary Material

Figure S1: *A. fumigatus* **grows slower than** *A. fischeri* **in glucose minimal media (GMM),**

but at the same speed as *A. fischeri* **in lung homogenate media.** *A. fumigatus* CEA10 or *A.*

fischeri NRRL181 were cultured in flat-bottom 96 well plates at $2x10^4$ conidia per well. Conidia

were added in a 20 µL of 0.01% Tween-80 and media was carefully pipetted over the inoculum

into each well. Lung homogenate was generated according to (29). Plates were incubated for 7

hours at 37°C before measurements at 405 nm were taken every 10 min. Mean and SEM of

eight technical replicates; data is representative of three biological replicates.

Figure S2: *A. fischeri* **and** *A. fumigatus* **exhibit similar growth patterns at 30°C.** $1x10^3$

conidia were point inoculated on each plate then plates were incubated at 30°C in normoxia

(~21% oxygen, 5%CO_2); colony diameter was measured every 24 hours. Mean and SEM of

triplicates. Tween-80 – 1% Tween-80 provided as sole carbon source; CAA – Casamino acids;

GMM – glucose minimal media.

Figure S3: In contrast to *A. fumigatus***,** *A. fischeri* **fails to thrive at 44°C.** Error bars indicate

standard deviations between biological duplicates (**P-value < 0.005 in a paired, equal variance

student t-test).

Figure S4: The genomes of *A. fumigatus* **and** *A. fischeri* **are largely similar, but their**

secondary metabolic pathways are quite divergent. Left, Venn diagram showing the sets of *A.*

fischeri-specific proteins, shared orthologous proteins, and *A. fumigatus*-specific proteins

encoded in each genome. Numbers below each species name indicate the total number of

proteins encoded in that genome. Right, Venn diagram showing the sets of *A. fischeri*-specific

secondary metabolite cluster proteins, shared secondary metabolite cluster genes, and *A.*

fumigatus-specific secondary metabolite cluster genes. Numbers below each species name

indicate the total number of secondary metabolite cluster proteins encoded in that genome. In

each diagram, circles are proportional to the number of proteins they contain.

Figure S5: The acetylaszonalenin and gliotoxin clusters in *A. fumigatus* and *A. fischeri* are

located immediately next to one another. The portions of Clusters 37 and 25 from *A. fischeri*

and *A. fumigatus*, respectively, that are known to contain the previously characterized

acetylaszonalenin (91) and gliotoxin (41) clusters is shown. Genes colored in shades of green are

involved in the acetylaszonalenin biosynthetic pathway. Dark green, *anaPS* (nonribosomal

peptide synthase). Light green, *anaAT* (acetyltransferase). Green, *anaPT* (prenyltransferase).

Orange, gliotoxin biosynthetic genes. Gray arrow, syntenic gene in both species not involved in

gliotoxin synthesis. Sequences that are similar to one another (based on blastn scores) are

marked by gray parallelograms. Image was made using EasyFig version 2.2.2 (83).

Figure S6: A custom chemical analysis protocol was developed for studying the metabolites

produced by *A. fischeri*. Approximately 60 mL of 1:1 $CH_3OH:CH_3Cl$ was added to cultures of

Aspergillus fischeri grown on solid-state fermentation for two weeks. The cultures were then

chopped thoroughly with a large scalpel and shaken for 16 hours using an orbital shaker. The

liquid culture was then vacuum filtered and concentrated using 90 mL CH_3Cl and 150 mL water

and transferred into a separatory funnel. The organic (bottom) layer was drawn off and

evaporated to dryness. The dried, de-sugared extract was reconstituted in 100 mL of 1:1

$CH_3OH:CH_3CN$ and 100 mL of hexane. The biphasic solution was shaken vigorously and

transferred to a separatory funnel. The $CH_3OH:CH_3CN$ layer was evaporated to dryness under

vacuum, producing a de-fatted extract. The extract was then subdivided into several peaks or

fractions using flash chromatography. The subfractions were further separated using HPLC until

pure compounds were isolated. The pure compounds were subjected to UPLC-MS analysis to

establish the molecular formula and fragmentation patterns. Finally, pure compounds were

identified using both NMR analysis as well as information from UPLC-MS data.

Figure S7: *A. fischeri* **produces different numbers of metabolites, depending on the media it**

is grown on. Base peak chromatograms as measured by LC-MS, illustrating how the chemistry

profiles varied based on growth conditions. PDA + ab was used as the chemical control to

observe the differences in the secondary metabolites, due to it being the media that *A. fischeri* is

stored. There were overall no chemical differences observed between the different variations of

PDA media. Each peak (which indicates different chemical entities) was observed in the three

PDA variations, albeit at fluctuating intensities. SDA, PYG, and YESD produced the majority of

the peaks observed in PDA, but it also lacked some observed peaks, indicating that these growth

conditions were not chemically favored. CYA produced the majority of the peaks, as well as an

additional peak that was observed at a much lower intensity in PDA. However, this peak was

similarly observed in OMA. OMA produced similar peaks to those observed in PDA, but with

higher intensity. Due to this, OMA was selected to further study. The gray boxes indicate

differences in the observed peaks compared to PDA. See Figshare document

(https://doi.org/10.6084/m9.figshare.7149167) for more information.

Figure S8: Southern blot confirms construction of the Δ*laeA* mutant. A 1kb probe recognizes

a single DNA band (~4.4kb) in the wild type strain and a single DNA band (~2.7kb) in the Δ*laeA*

mutant.

All supplemental tables can be found on Figshare (https://doi.org/10.6084/m9.figshare.7149167)

Table S1: Virulence-associated genes in *A. fumigatus* and *A. fischeri*.

Table S2: Bioinformatically predicted secondary metabolite clusters in *A. fumigatus* strain

CEA10.

1000

1001   **Table S3: Bioinformatically predicted secondary metabolite clusters in *A. fischeri* strain**

1002   **NRRL 181.**

1003

1004   **Table S4: Different Types of Growth Media used for *Aspergillus fischeri*.**

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5