Dynamic network partnerships and social contagion drive cooperation

Short title: Cooperation begets cooperation

Roslyn Dakin[a,1] roslyn.dakin@gmail.com and T. Brandt Ryder[a,1,]* rydert@si.edu

[a] Migratory Bird Center
Smithsonian Conservation Biology Institute
National Zoological Park
MRC 5503 Box 37012
Washington, DC 20013, USA

[1] RD and TBR contributed equally to this work

* To whom correspondence should be addressed:
T. Brant Ryder
Migratory Bird Center
Smithsonian Conservation Biology Institute
National Zoological Park
MRC 5503 Box 37012
Washington, DC 20013, USA
Phone: 202-633-4181
Email: rydert@si.edu

ETHICS
All methods were approved by the Smithsonian ACUC (protocols #12-23, 14-25, and 17-11) and the Ecuadorean Ministry of the Environment (MAE-DNB-CM-2015-0008).

DATA ACCESSIBILITY
All materials necessary to reproduce the results and figures, including data, R scripts and model results, are available for reviewers at the following link and will be made public upon acceptance: https://figshare.com/s/e73a3cb48e4714898d55

AUTHOR CONTRIBUTIONS
TBR and RD designed the study and developed the methods. TBR collected the data. RD and TBR analyzed the data and wrote and revised the manuscript.

COMPETING INTERESTS
We have no competing interests.

KEYWORDS (max 6): cooperation, social contagion, dynamic social networks, behavioral plasticity, indirect genetic effects, wire-tailed manakin

44 **ABSTRACT (200 word limit)**

Both reciprocity and positive assortment (like with like) are predicted to promote the evolution

46 of cooperation, yet how partners influence each other's behavior within dynamic networks is not

well understood. One way to test this question is to partition phenotypic variation into

48 differences among individuals in the expression of cooperative behavior (the "direct effect"), and

plasticity within individuals in response to the social environment (the "indirect effect"). A

50 positive correlation between these two sources of variation, such that more cooperative

individuals elicit others to cooperate, is predicted to facilitate social contagion and selection on

52 cooperative behavior. Testing this hypothesis is challenging, however, because it requires

repeated measures of behavior across a dynamic social landscape. Here, we use an automated

54 data-logging system to quantify the behavior of 179 wire-tailed manakins, birds that form

cooperative male-male coalitions, and we use multiple-membership models to test the hypothesis

56 that dynamic network partnerships shape within-individual variation in cooperative behavior.

Our results show strong positive correlations between a bird's own sociality and his estimated

58 effect on his partners, consistent with the hypothesis that cooperation begets cooperation. These

findings support the hypothesis that social contagion can facilitate selection for cooperative

60 behavior within social networks.

INTRODUCTION

62   Cooperation is an emergent property of social interactions within a network, yet our

understanding of how cooperative behavior emerges and is maintained is still major problem in

64   evolutionary biology. Two processes that favor the evolution of cooperation are reciprocity,

wherein cooperative behavior is socially contagious, and positive assortment, wherein non-

66   random structured interactions create clusters of cooperators in the network [1–4]. These

processes are non-exclusive, because any source of reciprocity that causes individuals to act

68   more like their cooperative neighbors will also contribute to the process of positive assortment.

Hence, a major question is how these social processes within complex networks influence

70   variation in cooperative behavior and the emergence of sociality.

Answering this question requires partitioning phenotypic variation in cooperation into

72   two major sources (Fig. 1a): (1) differences among individuals in the expression of a behavior

due to their intrinsic biology (the "direct effect"), as well as (2) plasticity within individuals

74   induced by the social environment (the "indirect effect"). This approach, largely derived from

quantitative genetics (i.e., interacting phenotypes and indirect genetic effects [5–8]), accounts for

76   the multilevel nature of behavior and can be used to quantify social influence while accounting

for other sources of variation [9,10]. A key feature of this framework is its focus on repeatable

78   individual differences [9]. This is important because repeatable phenotypic variation is the raw

material upon which selection acts [11].

80   Variance partitioning approaches can also be used to quantify among-individual variation

in social influence (i.e., the extent to which individuals differentially alter the expression of their

82   partner's phenotype). This is important given growing evidence that the social environment can

alter phenotypic expression and subsequent selection dynamics (reviewed in [9,10]). Moreover,

84   estimates of social influence as a trait are key to understanding how behaviors are transmitted or

      inhibited within social networks [12]. When an individual's average phenotypic expression

86   positively covaries with his social influence (e.g., more cooperative individuals elicit their

      partners to cooperate, Fig. 1b), behaviors can be transmitted via social contagion among partners

88   within a network [12]. This social contagion is hypothesized to promote the spread of

      cooperative behavior and facilitate selection for cooperation [7,13–15]. Conversely, negative

90   covariance between an individual's average expression of a trait and his social influence (Fig.

      1c) creates social inhibition which prevents the spread of, and constrains selection for, a specific

92   behavioral trait within the network [16–18]. Finally, there may be no relationship between

      average expression and social influence among interacting phenotypes (Fig. 1d).

94   These theoretical principles highlight how social influence can promote and/or constrain

      the emergence of population-level processes such as cooperation [10,19,20], but empirical tests

96   of these hypotheses have been limited. Recent experiments have demonstrated that cooperative

      and social behaviors can be transmitted via social contagion and dynamic network partnerships

98   in flies, fish, and humans [12,14,15,21–23]. To date, however, these laboratory studies have

      considered simple dyadic or small group interactions that do not reflect realistic social

100   complexity [but see 24], and may overestimate social influence [8,25]. Thus, a complete

      understanding of social influence within in a dynamic social network will require collecting

102   repeated measures of behavior while accounting for who interacts with whom, and the frequency

      with which these interactions occur.

104   Here, we address this gap by testing how the social environment influences cooperative

      behavior in a lek-breeding bird, the wire-tailed manakin (*Pipra filicauda*). The social system of

106   this species is well-studied and consists of two male status classes defined as territorial and non-

territorial (or 'floater') males [26]. Both status classes engage in repeated social interactions in

108   the context of male-male display coalitions which function both in the maintenance of

dominance hierarchies and in attracting females [27; see also Movie 1]. At any given time, a

110   male will have multiple coalition partners and these heterogeneous partnerships create complex

reticulate social networks [28]. Moreover, although manakin coalitions are relatively stable on an

112   annual basis [27], partnerships are dynamic on shorter daily or weekly timescales (see ESM 1)

and males show substantial variation in cooperation [28,29]. Previous work on wire-tailed

114   manakins has established that cooperative behavior within the social network has clear fitness

benefits: territorial males with more coalition partners have greater reproductive success [30],

116   and floaters with more coalition partners have an increased probability of territorial inheritance

[28]. Attaining more coalition partners is thus essential for floater fitness because territoriality is

118   a prerequisite for access to mating and reproductive success [30].

       To evaluate the influence of social network interactions (i.e., coalition partnerships) on

120   social behavior, we leveraged our knowledge of wire-tailed manakin natural history to develop

an autonomous proximity data-logging system (Fig. 2). The proximity approach allowed us to

122   obtain repeated measures of behavioral phenotype for a population of 179 males and to

characterize the identity and frequency of cooperative coalition partnerships within the social

124   network (Fig. 3). We chose the phenotypes illustrated in Fig. 3 because they capture a range of

social behaviors that are expected to promote reproductive success and thus be under selection.

126   "Effort" is a measure of daily attendance on territories where the coalitions are formed and is a

behavior known predict male mating success in other lekking taxa [31]. "Degree" and "strength"

128   measure the number of cooperative coalition partners and frequency of those partnerships within

the network, respectively. These measures of cooperation are known to affect social ascension

130 and reproductive success in manakins and other organisms [28,30,32]. Social "importance"

measures a male's position with the broader social network by characterizing the exclusivity of

132 his coalition partnerships. Exclusivity is contingent on the proportion of direct and indirect social

relationships (i.e., how frequently your partner displays with you vs. with his other coalition

134 partners); recent work suggests that these indirect network connections often capture among-

individual differences in social behavior [33]. Moreover, in manakins, the exclusivity and

136 subsequent stability of coalition partnerships can increase elements of display coordination and

signal intensity, both of which are preferred by females [27,34].

138   Here, we use multiple membership models to evaluate the extent to which an individual's

intrinsic biology (direct effects) and his repeated social network interactions (indirect effects) can

140 explain variation in cooperative behavior. Our analysis had three aims. First, we estimated the

direct effect as the proportion of total phenotypic variance attributed to differences among

142 individuals (i.e., the repeatable component of X's own behavior in Fig. 1a). Second, we

estimated the indirect effect as strength of social influence on phenotypic variation within

144 individuals (i.e., the repeatable component of Y and Z's influence on other individuals in Fig 1a).

Our analysis also captures the reciprocal nature of social influence in each coalition partnership.

146 A significant indirect effect in this analytical framework has two main implications: on the one

hand, it tests how much of the total phenotypic variance can be explained by social influence, but

148 also, it measures how individuals vary in their social influence within the network by accounting

for the frequency of interactions between coalition partners. Third, we examined the transmission

150 of behaviors among coalition partners by testing for social contagion (Fig. 1b-d). Given that

social contagion is defined as the spread of a particular behavior among partners within a social

152 network [12], we estimated the correlation between an individual's average expression of a trait

and his social influence on that same trait in his coalition partners. A positive covariance

154 between these repeatable components of variation is consistent with the hypothesis that the

expression of a given behavior is social contagious. We compared all of these results to null

156 models of the network data in which the direct effects and indirect effects had been removed by

permuting individual ID labels.

MATERIAL AND METHODS

160 **Study System and Field Methods**

We studied a color-banded population of male wire-tailed manakins (*P. filicauda*) at the Tiputini

162 Biodiversity Station, Orellana Province, Ecuador (0° 38' S, 76° 08' W, approx. 200 m elevation)

across three breeding seasons (November to March 2015-18; hereafter years '15-16', '16-17',

164 and '17-18'). The males at this site have been observed and color-banded annually since 2002.

Male status (i.e., territorial or floater) was determined by direct observation in the field following

166 methods outlined in Ryder et al. [28]. All methods were approved by the Smithsonian ACUC

(protocols #12-23, 14-25, and 17-11) and the Ecuadorean Ministry of the Environment (MAE-

168 DNB-CM-2015-0008).

To collect repeated measures of the behavior and social environment of male manakins,

170 we developed the automated proximity system shown in Fig. 2 [35]. Each individual was

outfitted with a coded nano-tag (NTQB-2, Lotek Wireless; $n_{15-16} = 100$ males that were tagged

172 and color-banded, $n_{16-17} = 114$, $n_{17-18} = 82$; and $n_{total} = 179$). Annual sample sizes are non-

exclusive because many males were tagged in multiple years (see also 'Sample Sizes' below).

174 The sample size was not pre-determined, but instead the aim was to capture all individuals and

monitor as many social network interactions as possible. Proximity data-loggers (hereafter DL;

176 SRX-DL800, Lotek Wireless) were placed within the lek territories where the manakins engage

in their cooperative displays (Fig. 2a; see ESM 1). DLs recorded continuously for 10 hours from

178 0600 to 1600, such that each day constitutes a repeated measure of behavior. DLs were deployed

at a given lek for an average of ~6 days (± 1 SD) per sampling interval and moved among leks

180 on a rotating schedule. In total, we recorded 29,760 hours of proximity data over 249 days (15-

16: 49 territories, mean = 16 recording days per territory; 16-17: 54 territories, mean = 21 days;

182 17-18: 48 territories, mean = 22 days, SD = ±4 days in all years).

184 **Data Processing**

To define distinct social interactions from the proximity data, we used a rule set with spatial and

186 temporal thresholds illustrated in Fig. 2a-d. This rule set was based on 16 years of previous study

demonstrating that spatial co-occurrence within a manakin territory is a both a necessary

188 prerequisite for, and an accurate predictor of, cooperative display interactions among males

[28,35]. We also used a ground-truthing experiment to confirm that the spatial threshold would

190 identify birds in close proximity (Fig. 2b-c; see ESM 1).

After removing any detections that were recorded at more than one site simultaneously

192 (see ESM 1 for details), we used the proximity data to calculate repeated measures of four

behavioral phenotypes thought to be important for manakin reproductive success, illustrated in

194 Fig. 3a. (1) Effort is the number of unique DL pings by a focal individual. (2) Strength is the sum

of a male's edge weights within the network. (3) Degree is the number of unique edges or direct

196 links a male has within the network. Strength and degree are thus measures of the frequency of

cooperative behavior and the number of cooperative coalition partners, respectively. (4)

198 Importance is a measure of the exclusivity of a male's coalition partners, on a scale from 0 to 1.

To calculate importance, we first found the proportion of each partner's interactions that were
200   with a focal male. Then, we took a weighted average using the focal male's interaction
frequencies. Thus, a male whose partners often interact with other individuals would obtain a
202   low score for importance, whereas a male whose partners interact with him exclusively would
obtain a score of 1. Importance is thus akin to an indirect network metric that accounts for both
204   direct and indirect social interactions (reviewed in [33]). Additional details of the data processing
methods and descriptive statistics for the behavioral phenotypes and their correlations are
206   provided in ESM 1 (Tables S1, S2).

208   **Statistical Analysis**

Analyses were performed using R 3.5.1 [36]. All materials necessary to reproduce our results
210   and figures, including data, R scripts, and model results are provided at: [37].

212   *Assortment*

As a general description of network assortment, we calculated the assortativity coefficient, ρ, for
214   each of the four phenotypes. This analysis was based on a single average value for each
individual (Fig. 3b) and a single social network compiled over all three years of study. We first
216   derived Newman's assortativity coefficients using the igraph package 1.1.2 [38], and verified
that the results were identical to weighted network assortment coefficients for continuous
218   phenotypes using the assortnet package 0.12 [39,40]. To test whether phenotypic assortment was
statistically significant, we determined the 95% confidence interval using the jackknife
220   resampling method in assortnet.

222   *Direct and Indirect Effects*

A major challenge when analyzing dynamic networks is that individuals can have multiple

224   partners and variable interaction frequencies (Fig. 2). We therefore developed an analytical

approach based on the multiple-membership structure modeling framework (i.e., multi-

226   membership models; hereafter, MMMs) for partitioning the phenotypic variance between the

direct and indirect effects. MMMs are an extension of multi-level models that incorporate a

228   heterogeneous weighted random effect [41], thus making it possible to capture variation in

dynamic network partnerships. The key advantage of MMMs is that the analysis can be weighted

230   according to the frequency with which different groupings or social relationships occurred.

We fit MMMs using the brms package 2.5.0 [42] which employs the Stan language in R,

232   using the model syntax shown in Box 1. The response variable was an individual's behavioral

phenotype on a particular day (t_1); we also included the bird's identity as a random effect, as well

234   as his top four most frequent partners on the previous day (t_0) in the multiple-membership

structure, weighted by their interaction frequencies (Box 1). Thus, our analysis asks, how is a

236   bird's behavior influenced by his own identity and who he interacted with on the previous day?

Although a few birds had more than four partners in a single day, we chose to limit our analysis

238   to the top four because additional partners comprised less than 2% of a bird's daily total in our

data. When an individual had fewer than 4 partners on a given day, we distributed one, two, or

240   three of his realized partner IDs and their interaction frequencies over the remaining empty

slot(s), while maintaining each partner's total weight as a constant. As long as the total weight of

242   each partner remains unchanged, the exact redistribution procedure does not influence the

results, because MMM analysis normalizes weights within the focal individuals [41].

244   Prior to modelling, three of the four behavioral phenotypes were log-transformed (effort,

strength, and degree) so that Gaussian model assumptions were met, and all phenotypes were

246   standardized (mean 0, SD 1). We used the default uninformative priors and stored 2,000 samples

from each of four independently-seeded chains for each phenotype, verifying that the

248   convergence statistics in the model output were all equal to 1. We then obtained estimates of the

variance components for each phenotype by pooling the posterior values from all four chains.

250   Posterior distributions for the direct and indirect effects were calculated as Var_{focal}/Var_{total} and

Var_{social}/Var_{total}, respectively [9], wherein Var_{focal} is the variance due to the identity of the focal

252   individuals; Var_{social} is the variance due to the MMM structure (i.e., the social environment); and

Var_{total} is the sum of all variance components (including focal, social, date, lek, and the

254   residual/unexplained variance).

       To account for additional sources of variation (Table S1), we also included fixed effects

256   of focal status (territorial vs. floater), study year, and an individual's number of years of tenure in

the present study, as well as random effects of date and lek in all models. By including year,

258   date, and lek, our estimates of the direct and indirect effects account for spatial and temporal

variation in the shared non-social environment that may influence focal and partner behavior

260   (e.g., [43–45]). Years of tenure was included instead of age, because age was not known for most

(69%) of the social interactions. Hence, our analysis computes the proportion of phenotypic

262   variance due to direct and indirect effects after accounting for status and any spatial/temporal

dynamics. Although feedback loops are implied [8], we do not consider the influence of second-

264   or higher-order connections among focal and partner phenotypes [46,47].

Box 1. Syntax for weighted MMM analysis of direct and indirect effects in brms

scale(Focal.phenotype) ~ focal.status + study year + tenure, random + (1|Focal.ID) + (1|mm(Partner.ID1,

Partner.ID2, Partner.ID3, Partner.ID4), weights=cbind(frequency.1, frequency.2, frequency.3, frequency.4)) +

(1|date) + (1|lek)

---

268 *Null Models*

Social network data are by definition non-independent, so they violate the assumptions of

270 traditional statistical inference [48,49]. We therefore used a null model framework to compare

our posterior estimates with the same analyses performed on randomly permuted data [48,49].

272 The null permutation preserved all structure in the data but permuted the ID labels, which were

swapped among individuals in the same daily recording session. This is equivalent to a node-

274 label network permutation. Given that territorial and floater birds differ [35] (Table S1), the

permutation was also constrained to swap ID labels among status-matched individuals recorded

276 on the same day. Hence, the null data preserved spatiotemporal sources of variance/covariance,

but disrupted sources of variation that could be attributed specifically to identity. This provides a

278 strong test of the identity-based effects that are the focus of our analysis.

To analyze the null (permuted) data, we fit MMMs using the same syntax presented in

280 Box 1. After verifying initial model diagnostics on a small number of model runs for each

phenotype, we generated 1,000 null datasets, and fit a single model chain to each dataset storing

282 2,000 iterations. Because the direct and indirect effects are bounded by 0, we used one-sided

tests to compare the observed posterior estimates with the median value obtained from these null

284 permutations.

Finally, as an additional check on our results, we also examined null models that

286 separately permuted either the Focal.ID labels or the Partner.ID labels alone. As expected,

permuting Focal.ID reduced the direct effect to near zero, but gave the same indirect effect

288   observed in the actual data, whereas permuting Partner.ID reduced the indirect effect to near zero

but gave the same direct effect observed in the actual results.

Contagion

292   Social contagion generates a positive covariance between a bird's own phenotypic expression

and his social influence on others' expression of the same trait (Fig. 1b-d). To determine this

294   covariance, we obtained posterior estimates of phenotypic expression and social influence for

each male in the population from the fitted MMMs. These individual estimates are akin to the

296   random intercepts or BLUPs in a frequentist framework. We used the posterior medians to test

the correlation between an individual's own phenotypic expression and his social influence on

298   each of the four behavioral phenotypes. That is, do individuals that express consistently high

levels of a phenotype also elicit that same phenotype in others? To verify that these associations

300   were significant even after accounting for error, we used Deming regression analyses with the

posterior standard deviations as the error in each variable. For inference, we also compared the

302   observed correlation statistic (r) with that derived from the null models above, which preserved

spatiotemporal sources of covariance. Hence, this null provided a more conservative test of the

304   covariance. Finally, because directly estimating covariance from the MMMs is not currently

possible ([42,50]; Bürkner and Hadfield, pers. comm.), we also performed a sensitivity analysis

306   to validate this approach (ESM 1). The sensitivity analysis demonstrated that in models with a

large sample size like ours, the bias in the direct and indirect effects caused by the covariance is

308   very small, and much smaller than the uncertainty inherent in these estimates (Fig. S2). Thus,

any bias is not expected to affect our conclusions.

Sample Sizes

312   A small number of individuals (n = 9) had tags that either stopped transmitting to the receivers

shortly after they were released, or they left the study population, and hence they are not

314   included in this study. Twenty-two of the successfully tagged males were also implanted with

hormone and sham pellets in part of years 2 and/or 3 for a separate study of the endocrine

316   mechanisms of cooperative behavior. We consequently limited our statistical analysis here to

focal individuals at times when they had no implants; nevertheless, the implant birds still acted

318   within the social networks and their interactions are retained when calculating the behavioral

phenotype of other individuals. Our sample size for analysis was 2,935 daily measures of 144

320   focal individuals. Note that this is smaller than the number of tagged birds, because a focal bird

had to have a known social environment on the previous day to be included in the MMM

322   analysis. Of these 144 males, 69 were tracked in one year, 49 in two years, and 26 in all three

study years; 8 floater individuals also attained territorial status during the study.

RESULTS

326   Three of the four phenotypes, strength, degree, and importance, were positively assorted in the

social network (Fig. 3b). All four phenotypes also had statistically significant individual

328   differences in expression, such that the direct effects could explain ~12-30% of the total variance

(Fig. 4a, Table S3, all p < 0.006). Because these percentages are moderate, this result also

330   demonstrates substantial within-individual plasticity in all four behavioral phenotypes.

We next asked if the social environment could explain some of this within-individual

332   variation in behavior. We found that between 12-24% of the total variance could be attributed to

indirect effects caused by the dynamic network interactions on the previous day (Fig. 4b, Table

334   S3, all p < 0.005). The phenotypes with the larger direct effects (effort and strength) also tended

to have smaller indirect effects, and vice versa (inset, Fig. 4b). These significant indirect effects

336   also imply that social influence is itself a trait that varies among individual male manakins.

Moreover, our null models demonstrate that indirect effects of this magnitude do not arise by

338   chance in permuted data with the exact same phenotypes and network topologies.

To further visualize these indirect effects, we also plotted within-individual plasticity in

340   phenotypic expression in relation to the social environment, as defined by average partner social

influence from the previous day (Fig. 4c). These plots show how individuals adjust their

342   behavior based on their dynamic social interactions on the previous day.

Finally, we asked if indirect effects as our measure of social influence could drive the

344   social contagion of behavior. The key parameter to test this hypothesis is the correlation between

an individual's own phenotypic expression and his social influence on others (Fig. 1). We found

346   strong positive correlations between expression and social influence for three of the four

phenotypes: strength, degree, and importance (Fig. 5), all of which exhibited positive assortment

348   (Fig. 3b). Moreover, these estimates of covariance were greater than expected in our permuted

data (Fig. S3). These results suggest that individual manakins with consistently high strength,

350   degree, and importance tend to elicit increases in those respective phenotypes in their partners. In

contrast, although greater effort was also elicited by particular individuals (Fig. 4), its lack of a

352   significant positive correlation suggests that effort is not socially contagious (Fig. 5).

354   **DISCUSSION**

Social contagion facilitates the evolution of positive assortment and cooperation [4,12], yet

356   understanding how dynamic network interactions shape variation in cooperative behavior has

remained challenging outside of the laboratory. Here, we monitored a large population of

358   cooperative wire-tailed manakins, and we applied a multiple-membership analysis to partition

direct and indirect sources of variation in four behavioral phenotypes. We found that cooperative

360   behavior was socially contagious: those individuals who had on average more cooperative

coalition partnerships (degree) and higher frequency of interactions (strength) tended to elicit

362   greater levels of those behaviors in their partners. Moreover, individuals who exhibited more

exclusive and stable partnerships (high importance) elicited their partners to have more exclusive

364   coalition partnerships as well. These results are notable because social connectivity as measured

by the number of coalition partners has a direct link to reproductive success in this population

366   and other species in the same family [28–30]. In contrast, we find no evidence that a male's

effort in attending leks was socially contagious (although it was influenced by partner identity).

368   Viewed cumulatively, our work demonstrates how multiple sources of phenotypic variation can

be parsed in a network to quantify the role of social influence on behavioral plasticity in the wild.

370   Furthermore, we demonstrate that a given behavior can be horizontally transmitted through the

network via social contagion, and that this process can contribute to positive network assortment

372   reinforcing cooperative behavior.

Over the last five years, social network analysis has rapidly advanced our understanding

374   of the causes and consequences of social behavior [reviewed in 51]. For example, recent work

has characterized individual differences in network position [52,53], patterns of phenotypic

376   assortment [54,55], and the fitness consequences of repeated social interactions [29,30,56]. Only

recently, however, have we begun to integrate network theory into our thinking about selection

378   dynamics and how structured interactions modulate phenotypic variance [19,20]. Here, we

continue to advance that theme with the first empirical evidence that dynamic network

380   interactions and social influence can have profound effects on variation in cooperative behavior.

We expect that these social processes also contribute to the emergence and maintenance of

382   cooperation in other animal systems [10], yet empirical validation is needed.

If the observed correlation between the expression of cooperation (strength and degree)

384   and social influence in wire-tailed manakins has some genetic basis (i.e., heritability), then our

results imply that indirect effects can enhance the response to selection on cooperative behavior

386   [5,7,13]. This is because selection acting on the genetic architecture for cooperative behavior

would also facilitate a social environment that reinforces the benefits of cooperation [4]. The

388   genetic basis of social influence could be tested in future studies that build on our approach by

using genotyping to resolve the population pedigree. This will require analytical methods that

390   can account for genetic structure in the social environment [16,18,24].

Biologists have long recognized that no phenotype is expressed in a vacuum. It is

392   important to note that in natural settings, the non-social environment is also variable, and thus

interacting individuals may share features of the environment that can be mutually influential.

394   Although this possibility cannot be excluded here or in any observational study, our analyses

account for sources of shared annual, daily, and spatial variation. Hence, the results are expected

396   to be independent of factors that vary in space and time, such as resource availability, female

activity, and climatic variation [43–45].

398   Another important point is that in natural settings, individuals have the capacity to choose

their coalition partners and these decisions can directly impact the costs and benefits of engaging

400   in cooperative acts [21,22,57]. For example, young wire-tailed manakins are known to increase

coalition stability with age as they ascend the queue to territorial status [27]. The long-term

402   nature of male social partnerships and evidence of behavioral plasticity in response to the

dynamic social landscape [i.e., social competence, *sensu* 58] both likely facilitate the choice of

404   similar partners (homophily) in this study system. Furthermore, homophily of socially

contagious traits may further amplify or accelerate their transmission within well connected

406   social networks [20]. Determining the relative contribution of these two processes (contagion

and homophily) in a natural social network system remains a major challenge for future work

408   and may require social transplant experiments to resolve.

Finally, our results raise several key questions for further work. First, what are the

410   sensory mechanisms of social contagion for cooperative behavior (e.g., calls, odors, and/or

gestural displays)? Our analysis here considered each behavioral phenotype separately, but in

412   reality, these behaviors are not all independent (Table S2). Thus, the causal pathways for social

transmission may be complex. Second, what explains the persistence of individual differences in

414   social influence [20]? Third, what are the underlying physiological and mechanistic processes,

and how do these mechanisms facilitate or constrain social contagion as a form of cooperative

416   reciprocity? Work examining the hormone signaling pathways that control cooperative behavior

will further elucidate the links between genotype, phenotype, and partner interactions [59].

418   Moreover, because behavior, hormones, and gene expression can mutually affect one another,

characterizing how these processes interact to modulate the social dynamics is a pivotal next step

420   [60]. Given that repeated heterogeneous interactions underlie virtually all animal social systems,

our results provide a framework that can be broadly applied to test how social influence changes

422   in response to physiological manipulations. Ultimately, understanding how selection shapes

interactive phenotypes like cooperation will require integrative approaches that consider both

424   mechanism and social context.

FIGURE LEGENDS

Fig. 1. A conceptual framework for social influence. (a) A phenotype "P" varies within and
428 among individuals. In this example, yellow indicates high expression of the trait whereas blue
indicates low expression of the trait. The upper histogram shows the hypothetical phenotypic
430 distribution for a population, whereas the lower histogram represents the narrower range of
phenotypic values expressed by a given individual, X, within that population. Individual
432 phenotypes are the result of both intrinsic biology (the "direct effect") and plasticity induced by
the social environment (e.g., the "indirect effect" as X interacts with conspecifics Y and Z). This
434 indirect effect represents how interactions with Y and Z at an earlier time, t_0, can differentially
influence X's phenotype at time t_1. Note that X would also influence Y and Z's phenotype at
436 time t_1 in the analysis, but for simplicity, we show only one direction of social influence here. (b-
d) Because Y and Z also express phenotype P, there may be covariance between and individual's
438 average phenotypic expression (t_{avg}) and average social influence. For example, (b) Y and Z's
average phenotypic expression may be positively correlated with their social influence on X's
440 phenotype. This positive covariance is predicted to generate social contagion. Alternatively,
phenotypic expression may be (d) unrelated or (c) negatively related to social influence, the latter
442 of which would generate social inhibition.

444 **Fig. 2. Social interactions were tracked using an automated proximity data-logging system.**
(a) Each bird was tagged with a small transmitter that broadcasts a unique digital code at 20 s
446 intervals. Tag pings were received by stationary data loggers (DLs) set with a fixed detection
radius to encompass a male's territory where cooperative displays take place. (b) The tag signal
448 strength, RSSI, decays with distance from the DL. Hence, we define an interacting dyad using a
conservative spatiotemporal threshold wherein individuals had to be detected within 45 s of each
450 other and with a ΔRSSI < 10. (c) A ground-truthing analysis demonstrates that tag pairs that
meet this threshold are typically ≤ 5 m apart (vertical line = median), corresponding to a range
452 that would permit both visual and acoustic contact. For comparison, events with ΔRSSI ≥ 10
have a greater median inter-tag distances of 15 m. The data in (b) and (c) are based on the same
454 ground-truthing experiment. (d) An example of detection data for DLs at two neighboring
territories. The resulting time series illustrates two interactions, dyad 1 and dyad 2, that are
456 delimited by horizontal brackets. Note that birds occasionally ping the same DL around the same
time without meeting the ΔRSSI spatial threshold. (e) Overall, this automated system recorded
458 approx. 37,000 social interactions among 179 birds. (f) An example of daily interactions within a
subset of the population social network illustrating the dynamic nature of the study system. Each
460 node in the network represents an individual male sized according to degree. Data are shown for
14 of the males recorded from January 1-6 in 2016, with solid edges representing interactions
462 that occurred on that day. Dotted edges are interactions that occurred but not on the given
sampling day.

Fig. 3. We analyzed four behavioral phenotypes that are important for manakin
466 **reproductive success.** (a) These behaviors occurred on territories within the leks and were
quantified on a daily basis. The cartoon illustrates an increase in the phenotype from left to right.
468 Note that in the illustration for importance, the focal male (gray node) maintains the same
strength and degree, but increases his importance on the right, because each of his partners
becomes more exclusive. Meanwhile, his partners' importance actually decreases on the right,

demonstrating that changes in this metric are not necessarily matched among partners (i.e., many scenarios are possible). (b) Observed distributions of average phenotypic expression for floater and territory-holding males. Dashed vertical lines show medians for each status class. For effort,

1,000 pings are approximately 5.5 hours of attendance. Strength, degree, and importance are all positively assorted (ρ) within the social network. See Tables S1-S2 and Fig. S1 for additional

data.

Fig. 4. Manakins differ both in their social behavior and their influence on others. (a-b) The proportion of total phenotypic variance in four behavioral phenotypes attributed to (a) individual

differences (the direct effect) and (b) social influence (the indirect effect), as defined based on partners from the previous day. For each phenotype, the observed result is significantly greater

than the null expectation. The vertical lines show the posterior median. Phenotypes with larger direct effects tend to have weaker social influence (inset, b). The axes on the inset span 0 to 0.4.

Note that the tails of all distributions are extended beyond the range of the data for visual continuity. (c) Within individuals, phenotypic expression depends on the previous day's social

environment. On the y-axes, individual phenotypic expression values are centered to show change relative to an individual's mean expression; values for effort, strength, and degree were

also log-transformed prior to centering. The x-axis represents the average influence of a bird's partners on the previous day. A separate linear fit is shown for each focal bird observed on at

least 10 days (n = 95). See Tables S3 and S4 for detailed results.

Fig. 5. Positive covariance indicates that cooperative social behaviors are contagious. Each graph shows an individual's estimated social influence (y-axis) in relation to his own phenotypic

expression (x-axis) for that same behavior (n = 142 individuals analyzed as both focal and partner). Strength, degree, and importance all have strong positive correlations that are

significantly greater than 0 and greater than the null expectation from permuted data, indicating that indirect effects can facilitate the spread of these behaviors through the social network. In

contrast, the correlation for effort is not statistically significant. Data points and error bars are derived from the posterior medians and standard deviations, respectively. The regression fit lines

account for posterior estimates of the error in both variables. Pearson's correlation coefficients are also given on the lower right of each panel.

ACKNOWLEDGEMENTS

504   We thank Ignacio Moore, Ben J. Vernasco, and Camilo Alfonso for their tireless effort in project

organization and collecting field data. David and Consuelo Romo, Kelly Swing, Diego Mosquera

506   and Gabriela Vinueza and staff provided amazing logistical support at Tiputini Biodiversity

Station. We thank Brian S. Evans for his brilliance in data architecture and the development of R

508   functions. Brent Horton, Mike Webster, Joel McGlothlin, Niels Dingemanse, and one

anonymous reviewer provided valuable comments on an earlier version of this manuscript. This

510   work was supported by the National Science Foundation (NSF) IOS 1353085 and the

Smithsonian Migratory Bird Center.

SUPPORTING INFORMATION

ESM 1. Additional details of the methods and data tables.

Movie 1. Male wire-tailed manakins perform cooperative displays in Ecuador. These video clips

518   were taken during the 2015-16 breeding season and show examples of the coded tags used to

track social interactions.

REFERENCES

1. Trivers RL. 1971 The evolution of reciprocal altruism. *The Quarterly Review of Biology* **46**, 35–57. (doi:10.1086/406755)

2. Ohtsuki H, Hauert C, Lieberman E, Nowak MA. 2006 A simple rule for the evolution of cooperation on graphs and social networks. *Nature* **441**, 502–505. (doi:10.1038/nature04605)

3. Rankin DJ, Taborsky M. 2009 Assortment and the evolution of generalized reciprocity. *Evolution* **63**, 1913–1922. (doi:10.1111/j.1558-5646.2009.00656.x)

4. van Doorn GS, Taborsky M. 2012 The evolution of generalized reciprocity on social interaction networks. *Evolution* **66**, 651–664. (doi:10.1111/j.1558-5646.2011.01479.x)

5. Moore AJ, Brodie ED, Wolf JB. 1997 Interacting phenotypes and the evolutionary process. I. Direct and indirect genetic effects of social interactions. *Evolution* **51**, 1352–1362. (doi:10.2307/2411187)

6. Wolf JB, Brodie ED, Moore AJ. 1999 Interacting phenotypes and the evolutionary process. II. Selection resulting from social interactions. *The American Naturalist* **153**, 254–266. (doi:10.1086/303168)

7. McGlothlin JW, Moore AJ, Wolf JB, Brodie ED. 2010 Interacting phenotypes and the evolutionary process. III. Social evolution. *Evolution* **64**, 2558–2574. (doi:10.1111/j.1558-5646.2010.01012.x)

8. Bijma P. 2014 The quantitative genetics of indirect genetic effects: a selective review of modelling issues. *Heredity* **112**, 61–69. (doi:10.1038/hdy.2013.15)

9. Dingemanse NJ, Araya-Ajoy YG. 2015 Interacting personalities: behavioural ecology meets quantitative genetics. *Trends in Ecology & Evolution* **30**, 88–97. (doi:10.1016/j.tree.2014.12.002)

10. Bailey NW, Marie-Orleach L, Moore AJ, Simmons L. 2018 Indirect genetic effects in behavioral ecology: does behavior play a special role in evolution? *Behav Ecol* **29**, 1–11. (doi:10.1093/beheco/arx127)

11. Endler JA. 1986 *Natural Selection in the Wild*. Princeton University Press.

12. Christakis NA, Fowler JH. 2013 Social contagion theory: examining dynamic social networks and human behavior. *Statistics in Medicine* **32**, 556–577. (doi:10.1002/sim.5408)

13. Wolf JB, Brodie ED, Cheverud JM, Moore AJ, Wade MJ. 1998 Evolutionary consequences of indirect genetic effects. *Trends in Ecology & Evolution* **13**, 64–69. (doi:10.1016/s0169-5347(97)01233-0)

14. Bleakley BH, Brodie ED. 2009 Indirect genetic effects influence antipredator behavior in guppies: estimates of the coefficient of interation psi and the inheritance of reciprocity. *Evolution* **63**, 1796–1806. (doi:10.1111/j.1558-5646.2009.00672.x)

15. Edenbrow M, Bleakley BH, Darden SK, Tyler CR, Ramnarine IW, Croft DP. 2017 The evolution of cooperation: interacting phenotypes among social partners. *The American Naturalist* **189**, 630–643. (doi:10.1086/691386)

16. Wilson AJ, Gelin U, Perron MC, Reale D. 2009 Indirect genetic effects and the evolution of aggression in a vertebrate system. *Proceedings of the Royal Society B* **276**, 533–541. (doi:10.1098/rspb.2008.1193)

17. Wilson AJ, Morrissey MB, Adams MJ, Walling CA, Guinness FE, Pemberton JM, Clutton-Brock TH, Kruuk LEB. 2011 Indirect genetics effects and evolutionary constraint: an analysis of social dominance in red deer, *Cervus elaphus*. *Journal of Evolutionary Biology* **24**, 772–783. (doi:10.1111/j.1420-9101.2010.02212.x)

18. Santostefano F, Wilson AJ, Niemelä PT, Dingemanse NJ. 2017 Indirect genetic effects: a key component of the genetic architecture of behaviour. *Scientific Reports* **7**, 10235. (doi:10.1038/s41598-017-08258-6)

19. Fisher DN, McAdam AG. 2017 Social traits, social networks and evolutionary biology. *Journal of Evolutionary Biology* **30**, 2088–2103. (doi:10.1111/jeb.13195)

20. Montiglio P, McGlothlin JW, Farine DR. 2017 Social structure modulates the evolutionary consequences of social plasticity: A social network perspective on interacting phenotypes. *Ecology and Evolution* **8**, 1451–1464. (doi:10.1002/ece3.3753)

21. Fehl K, Post DJ van der, Semmann D. 2011 Co-evolution of behaviour and social network structure promotes human cooperation. *Ecology Letters* **14**, 546–551. (doi:10.1111/j.1461-0248.2011.01615.x)

22. Rand DG, Arbesman S, Christakis NA. 2011 Dynamic social networks promote cooperation in experiments with humans. *PNAS* **108**, 19193–19198. (doi:10.1073/pnas.1108243108)

23. Anderson BB, Scott A, Dukas R. 2017 Indirect genetic effects on the sociability of several group members. *Animal Behaviour* **123**, 101–106. (doi:10.1016/j.anbehav.2016.10.028)

24. Fisher DN, Wilson AJ, Boutin S, Dantzer B, Lane JE, Coltman DW, Gorrell JC, McAdam AG. 2018 Indirect effects of territorial neighbors on the timing of spring breeding may counteract changes in selection in North American red squirrels. *bioRxiv* (doi:10.1101/329276)

25. Hadfield JD, Wilson AJ. 2007 Multilevel selection 3: Modeling the effects of interacting individuals as a function of group size. *Genetics* **177**, 667–668. (doi:10.1534/genetics.107.075622)

26. Schwartz P, Snow DW. 1978 Display and related behavior of the wire-tailed manakin. *Living Bird* **17**, 51–78.

590   27. Ryder TB, Blake JG, Parker PG, Loiselle BA. 2011 The composition, stability, and kinship of reproductive coalitions in a lekking bird. *Behav Ecol* **22**, 282–290. (doi:10.1093/beheco/arq213)

592   28. Ryder TB, McDonald DB, Blake JG, Parker PG, Loiselle BA. 2008 Social networks in the lek-mating wire-tailed manakin (*Pipra filicauda*). *Proceedings of the Royal Society B* **275**, 594   1367–1374. (doi:10.1098/rspb.2008.0205)

   29. McDonald DB. 2007 Predicting fate from early connectivity in a social network. *PNAS* **104**, 596   10910–10914. (doi:10.1073/pnas.0701159104)

   30. Ryder TB, Parker PG, Blake JG, Loiselle BA. 2009 It takes two to tango: reproductive skew 598   and social correlates of male mating success in a lek-breeding bird. *Proceedings of the Royal Society B* **276**, 2377–2384. (doi:10.1098/rspb.2009.0208)

600   31. Hoglund J, Alatalo RV. 1995 *Leks*. Princeton University Press.

   32. Formica VA, Wood CW, Larsen WB, Butterfield RE, Augat ME, Hougen HY, Brodie ED. 602   2011 Fitness consequences of social network position in a wild population of forked fungus beetles (*Bolitotherus cornutus*). *Journal of Evolutionary Biology* **25**, 130–137. 604   (doi:10.1111/j.1420-9101.2011.02411.x)

   33. Firth JA, Sheldon BC, Brent LJN. 2017 Indirectly connected: simple social differences can 606   explain the causes and apparent consequences of complex social network positions. *Proc. R. Soc. B* **284**, 20171939. (doi:10.1098/rspb.2017.1939)

608   34. Trainer JM, McDonald DB. 1995 Singing performance, frequency matching and courtship success of long-tailed manakins (*Chiroxiphia linearis*). *Behav Ecol Sociobiol* **37**, 249–254. 610   (doi:10.1007/BF00177404)

   35. Ryder TB, Horton BM, Tillaart M van den, Morales JDD, Moore IT. 2012 Proximity data-612   loggers increase the quantity and quality of social network data. *Biology Letters* **8**, 917–920. (doi:10.1098/rsbl.2012.0536)

614   36. R Core Team. 2018 *R 3.5.1: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. See https://www.R-project.org.

616   37. Dakin R, Ryder TB. 2018 https://figshare.com/s/e73a3cb48e4714898d55 Supplementary Materials for: Dynamic network partnerships and social contagion drive cooperation.

618   38. Csardi G, coauthors. 2017 *igraph 1.1.2: network analysis and visualization.* See https://cran.r-project.org/web/packages/igraph/index.html.

620   39. Farine DR. 2014 Measuring phenotypic assortment in animal social networks: weighted associations are more robust than binary edges. *Animal Behaviour* **89**, 141–153. 622   (doi:10.1016/j.anbehav.2014.01.001)

40. Farine DR. 2016 *assortnet 0.12: calculate the assortativity coefficient of weighted and binary networks*. See https://cran.r-project.org/web/packages/assortnet/index.html.

41. Buerkner P-C. 2017 brms: an R package for Bayesian multilevel models Using Stan. *Journal of Statistical Software* **80**. (doi:10.18637/jss.v080.i01)

42. Buerkner P-C. 2018 *brms 2.5.0: Bayesian regression models using Stan*. See https://cran.r-project.org/web/packages/brms/index.html.

43. Ryder TB, Sillett TS. 2016 Climate, demography and lek stability in an Amazonian bird. *Proc. R. Soc. B* **283**, 20152314. (doi:10.1098/rspb.2015.2314)

44. Ryder TB, Blake JG, Loiselle BA, Vega Rivera JH. 2006 A test of the environmental hotspot hypothesis for lek placement in three species of manakins (Pipridae) in Ecuador. *The Auk* **123**, 247–258. (doi:10.1642/0004-8038(2006)123[0247:ATOTEH]2.0.CO;2)

45. Loiselle BA, Blake JG, Durães R, Ryder TB, Tori W, Vega Rivera JH. 2007 Environmental and spatial segregation of leks among six co-occurring species of manakins (Pipridae) in eastern Ecuador. *The Auk* **124**, 420–431. (doi:10.1642/0004-8038(2007)124[420:EASSOL]2.0.CO;2)

46. McGlothlin JW, Brodie ED. 2009 How to measure indirect genetic effects: the congruence of trait-based and variance-partitioning approaches. *Evolution* **63**, 1785–1795. (doi:10.1111/j.1558-5646.2009.00676.x)

47. Brent LJN. 2015 Friends of friends: are indirect connections in social networks important to animal behaviour? *Animal Behaviour* **103**, 211–222. (doi:10.1016/j.anbehav.2015.01.020)

48. Farine DR, Whitehead H. 2015 Constructing, conducting and interpreting animal social network analysis. *Journal of Animal Ecology* **84**, 1144–1163. (doi:10.1111/1365-2656.12418)

49. Farine DR, Freckleton R, Rands S. 2017 A guide to null models for animal social network analysis. *Methods in Ecology and Evolution* **8**, 1309–1320. (doi:10.1111/2041-210X.12772)

50. Hadfield J. 2018 *MCMCglmm 2.26: MCMC generalised linear mixed models*. See https://cran.r-project.org/web/packages/MCMCglmm/index.html.

51. Kurvers RHJM, Krause J, Croft DP, Wilson ADM, Wolf M. 2014 The evolutionary and ecological consequences of animal social networks: emerging issues. *Trends in Ecology & Evolution* **29**, 326–335. (doi:10.1016/j.tree.2014.04.002)

52. Jacoby DMP, Fear LN, Sims DW, Croft DP. 2014 Shark personalities? Repeatability of social network traits in a widely distributed predatory fish. *Behav Ecol Sociobiol* **68**, 1995–2003. (doi:10.1007/s00265-014-1805-9)

53. Aplin LM *et al.* 2015 Consistent individual differences in the social phenotypes of wild great tits, *Parus major*. *Animal Behaviour* **108**, 117–127. (doi:10.1016/j.anbehav.2015.07.016)

658   54. Croft DP, Krause J, Darden SK, Ramnarine IW, Faria JJ, James R. 2009 Behavioural trait
       assortment in a social network: patterns and implications. *Behav Ecol Sociobiol* **63**, 1495–
660   1503. (doi:10.1007/s00265-009-0802-x)

       55. McDonald GC, Farine DR, Foster KR, Biernaskie JM. 2017 Assortment and the analysis of
662   natural selection on social traits. *Evolution* **71**, 2693–2702. (doi:10.1111/evo.13365)

       56. Oh KP, Badyaev AV. 2010 Structure of social networks in a passerine bird: consequences
664   for sexual selection and the evolution of mating strategies. *The American Naturalist* **176**,
       E80–E89. (doi:10.1086/655216)

666   57. Bergmuller R, Schurch R, Hamilton IM. 2010 Evolutionary causes and consequences of
       consistent individual variation in cooperative behaviour. *Philosophical Transactions of the*
668   *Royal Society B* **365**, 2751–2764. (doi:10.1098/rstb.2010.0124)

       58. Taborsky B, Oliveira RF. 2012 Social competence: an evolutionary approach. *Trends in*
670   *Ecology & Evolution* **27**, 679–688.

       59. Soares MC, Bshary R, Fusani L, Goymann W, Hau M, Hirschenhauser K, Oliveira RF. 2010
672   Hormonal mechanisms of cooperative behaviour. *Philosophical Transactions of the Royal*
       Society B **365**, 2737–2750. (doi:10.1098/rstb.2010.0151)

674   60. Oliveira RF. 2009 Social behavior in context: hormonal modulation of behavioral plasticity
       and social competence. *Integr Comp Biol* **49**, 423–440. (doi:10.1093/icb/icp055)

Fig. 1

Fig. 1. A conceptual framework for social influence. (a) A phenotype "P" varies within and among individuals. In this example, yellow indicates high expression of the trait whereas blue indicates low expression of the trait. The upper histogram shows the hypothetical phenotypic distribution for a population, whereas the lower histogram represents the narrower range of phenotypic values expressed by a given individual, X, within that population. Individual phenotypes are the result of both intrinsic biology (the "direct effect") and plasticity induced by the social environment (e.g., the "indirect effect" as X interacts with conspecifics Y and Z). This indirect effect represents how interactions with Y and Z at an earlier time, t0, can differentially influence X's phenotype at time t1. Note that X would also influence Y and Z's phenotype at time t1 in the analysis, but for simplicity, we show only one direction of social influence here. (b-d) Because Y and Z also express phenotype P, there may be covariance between and individual's average phenotypic expression (tavg) and average social influence. For example, (b) Y and Z's average phenotypic expression may be positively correlated with their social influence on X's phenotype. This positive covariance is predicted to generate social contagion. Alternatively, phenotypic expression may be (d) unrelated or (c) negatively related to social influence, the latter of which would generate social inhibition.

Fig. 2

Fig. 2. Social interactions were tracked using an automated proximity data-logging system. (a) Each bird was tagged with a small transmitter that broadcasts a unique digital code at 20 s intervals. Tag pings were received by stationary data loggers (DLs) set with a fixed detection radius to encompass a male's territory where cooperative displays take place. (b) The tag signal strength, RSSI, decays with distance from the DL. Hence, we define an interacting dyad using a conservative spatiotemporal threshold wherein individuals had to be detected within 45 s of each other and with a ∆RSSI < 10. (c) A ground-truthing analysis demonstrates that tag pairs that meet this threshold are typically ≤ 5 m apart (vertical line = median), corresponding to a range that would permit both visual and acoustic contact. For comparison, events with ∆RSSI ≥ 10 have a greater median inter-tag distances of 15 m. The data in (b) and (c) are based on the same ground-truthing experiment. (d) An example of detection data for DLs at two neighboring territories. The resulting time series illustrates two interactions, dyad 1 and dyad 2, that are delimited by horizontal brackets. Note that birds occasionally ping the same DL around the same time without meeting the ∆RSSI spatial threshold. (e) Overall, this automated system recorded approx. 37,000 social interactions among 179 birds. (f) An example of daily interactions within a subset of the population social network illustrating the dynamic nature of the study system. Each node in the network represents an individual male sized according to degree. Data are shown for 14 of the males recorded from January 1-6 in 2016, with solid edges representing interactions that occurred on that day. Dotted edges are interactions that occurred but not on the given sampling day.

Fig. 3

Fig. 3. We analyzed four behavioral phenotypes that are important for manakin reproductive success. (a) These behaviors occurred on territories within the leks and were quantified on a daily basis. The cartoon illustrates an increase in the phenotype from left to right. Note that in the illustration for importance, the focal male (gray node) maintains the same strength and degree, but increases his importance on the right, because each of his partners becomes more exclusive. Meanwhile, his partners' importance actually decreases on the right, demonstrating that changes in this metric are not necessarily matched among partners (i.e., many scenarios are possible). (b) Observed distributions of average phenotypic expression for floater and territory-holding males. Dashed vertical lines show medians for each status class. For effort, 1,000 pings are approximately 5.5 hours of attendance. Strength, degree, and importance are all positively assorted ( ) within the social network. See Tables S1-S2 and Fig. S1 for additional data.

Fig. 4

Fig. 4. Manakins differ both in their social behavior and their influence on others. (a-b) The proportion of total phenotypic variance in four behavioral phenotypes attributed to (a) individual differences (the direct effect) and (b) social influence (the indirect effect), as defined based on partners from the previous day. For each phenotype, the observed result is significantly greater than the null expectation. The vertical lines show the posterior median. Phenotypes with larger direct effects tend to have weaker social influence (inset, b). The axes on the inset span 0 to 0.4. Note that the tails of all distributions are extended beyond the range of the data for visual continuity. (c) Within individuals, phenotypic expression depends on the previous day's social environment. On the y-axes, individual phenotypic expression values are centered to show change relative to an individual's mean expression; values for effort, strength, and degree were also log-transformed prior to centering. The x-axis represents the average influence of a bird's partners on the previous day. A separate linear fit is shown for each focal bird observed on at least 10 days (n = 95). See Tables S3 and S4 for detailed results.

Fig. 5

Fig. 5. Positive covariance indicates that cooperative social behaviors are contagious. Each graph shows an individual's estimated social influence (y-axis) in relation to his own phenotypic expression (x-axis) for that same behavior (n = 142 individuals analyzed as both focal and partner). Strength, degree, and importance all have strong positive correlations that are significantly greater than 0 and greater than the null expectation from permuted data, indicating that indirect effects can facilitate the spread of these behaviors through the social network. In contrast, the correlation for effort is not statistically significant. Data points and error bars are derived from the posterior medians and standard deviations, respectively. The regression fit lines account for posterior estimates of the error in both variables. Pearson's correlation coefficients are also given on the lower right of each panel.