Dynamic network partnerships and social contagion drive cooperation

Short title: Cooperation begets cooperation

Roslyn Dakin[a,1] roslyn.dakin@gmail.com and T. Brandt Ryder[a,1,]* rydert@si.edu

[a] Migratory Bird Center

8   Smithsonian Conservation Biology Institute
National Zoological Park

10 MRC 5503 Box 37012
Washington, DC 20013, USA

[1] RD and TBR contributed equally to this work

* To whom correspondence should be addressed:

16 T. Brant Ryder
Migratory Bird Center

18 Smithsonian Conservation Biology Institute
National Zoological Park

20 MRC 5503 Box 37012
Washington, DC 20013, USA

22 Phone: 202-633-4181
Email: rydert@si.edu

ETHICS

26 All methods were approved by the Smithsonian ACUC (protocol #17-11) and the Ecuadorean
Ministry of the Environment (MAE-DNB-CM-2015-0008).

DATA ACCESSIBILITY

30 All materials necessary to reproduce the results and figures, including data, R scripts and
posterior model results, are available for reviewers at the following link and will be made public

32 upon acceptance: https://figshare.com/s/e73a3cb48e4714898d55

34 AUTHOR CONTRIBUTIONS
TBR and RD designed the study and developed the methods. TBR collected the data. RD and

36 TBR analyzed the data and wrote and revised the manuscript.

38 COMPETING INTERESTS
We have no competing interests.

KEYWORDS (max 6): cooperation, social contagion, dynamic social networks, behavioral

42 plasticity, indirect genetic effects, wire-tailed manakin

44 **ABSTRACT (200 word limit)**

Both reciprocity and positive assortment (like with like) are predicted to promote the evolution

46   of cooperation, yet how partners influence each other's behavior within dynamic networks is not

well understood. One way to test this question is to partition phenotypic variation into

48   differences among individuals in the expression of cooperative behavior (the "direct effect"), and

plasticity within individuals in response to the social environment (the "indirect effect"). A

50   positive correlation between these two sources of variation, such that more cooperative

individuals elicit others to cooperate, is predicted to facilitate social contagion and selection on

52   cooperative behavior. Testing this hypothesis is challenging, however, because it requires

repeated measures of behavior across a dynamic social landscape. Here, we use an automated

54   data-logging system to quantify the behavior of 179 wire-tailed manakins, birds that form

cooperative male-male coalitions, and we use multiple-membership models to test the hypothesis

56   that dynamic network partnerships shape within-individual variation in cooperative behavior.

Our results show strong positive correlations between a bird's own sociality and his estimated

58   effect on his partners, consistent with the hypothesis that cooperation begets cooperation. These

findings support the hypothesis that social contagion can facilitate selection for cooperative

60   behavior within social networks.

INTRODUCTION

Cooperation is an emergent property of social interactions within a network, yet our understanding of how cooperative behavior emerges and is maintained is still major problem in

evolutionary biology. Two processes that favor the evolution of cooperation are reciprocity, wherein cooperative behavior is socially contagious, and positive assortment, wherein non-

random structured interactions create clusters of cooperators in the network (Trivers, 1971; Ohtsuki et al., 2006; Rankin and Taborsky, 2009; van Doorn and Taborsky, 2012). These

processes are non-exclusive, because any source of reciprocity that causes individuals to act more like their cooperative neighbors will also contribute to the process of positive assortment.

Hence, a major question is how these social processes within complex networks influence variation in cooperative behavior and the emergence of sociality.

Answering this question requires partitioning phenotypic variation in cooperation into two major sources (Fig. 1a): (1) differences among individuals in the expression of a behavior

due to their intrinsic biology (the "direct effect"), as well as (2) plasticity within individuals induced by the social environment (the "indirect effect"). This approach, largely derived from

quantitative genetics (i.e., interacting phenotypes and indirect genetic effects; Moore et al., 1997; Wolf et al., 1999; McGlothlin et al., 2010; Bijma, 2014), accounts for the multilevel nature of

behavior and can be used to quantify social influence while accounting for other sources of variation (Dingemanse and Araya-Ajoy, 2015; Bailey et al., 2018). A key feature of this

framework is its focus on repeatable individual differences (Dingemanse and Araya-Ajoy, 2015). This is important because repeatable phenotypic variation is the raw material upon which

selection acts (Endler, 1986).

Variance partitioning approaches can also be used to quantify among-individual variation

84 in social influence (i.e., the extent to which individuals differentially alter the expression of their

partner's phenotype). This is important given growing evidence that the social environment can

86 alter phenotypic expression and subsequent selection dynamics (reviewed in: Dingemanse and

Araya-Ajoy, 2015; Bailey et al., 2018). Moreover, estimates of social influence as a trait are

88 important for determining social contagion, which we define as positive transmission of behavior

among interacting individuals within a network (Christakis and Fowler, 2013). Specifically, a

90 positive covariance between an individual's phenotypic expression and his or her social

influence would occur, for example, when more cooperative individuals elicit others to cooperate

92 (Fig. 1b). This social contagion would promote the spread of cooperative behavior and facilitate

selection for cooperation (Wolf et al., 1998; Bleakley and Brodie, 2009; McGlothlin et al., 2010;

94 Edenbrow et al., 2017). Conversely, negative covariance occurs when more cooperative

individuals inhibit others from cooperating (Fig. 1c). This social inhibition would prevent the

96 spread of, and constrain selection for, a specific behavioral trait like cooperation (or aggression;

see for example: Wilson et al., 2009; Wilson et al., 2011; Santostefano et al., 2017). Finally,

98 there may be no relationship between expression and influence among interacting phenotypes

(Fig. 1d).

100 These theoretical principles highlight how social influence can promote the emergence of

population-level processes such as cooperation (Fisher and McAdam, 2017; Montiglio et al.,

102 2017; Bailey et al., 2018), but empirical tests of these hypotheses have been limited. Recent

experiments have demonstrated that cooperative and social behaviors can be transmitted via

104 social contagion and dynamic network partnerships in flies, fish, and humans (Bleakley and

Brodie, 2009; Fehl et al., 2011; Rand et al., 2011; Christakis and Fowler, 2013; Edenbrow et al.,

106   2017; Anderson et al., 2017). To date, however, these laboratory studies have considered simple

dyadic or small group interactions that do not reflect realistic social complexity (but see Fisher et

108   al., 2018), and may overestimate social influence (Hadfield and Wilson, 2007; Bijma, 2014).

Thus, a complete understanding of social influence within in a dynamic social network will

110   require collecting repeated measures of behavior while accounting for who interacts with whom,

and the frequency with which these interactions occur.

112   Here, we address this gap by testing how the social environment influences cooperative

behavior in a lek-breeding bird, the wire-tailed manakin (*Pipra filicauda*). The social system of

114   this species is well-studied and consists of two male status classes defined as territorial and non-

territorial (or 'floater') males (Schwartz and Snow, 1978). Wire-tailed manakins form male-male

116   coalitions to perform cooperative displays that attract females and form the basis for dynamic

social networks (Movie 1). Territoriality is a strong prerequisite for access to mating and thus

118   reproductive success in this system is highly skewed towards a subset of territorial males (Ryder

et al., 2009). Previous work on wire-tailed manakins has established that cooperative behavior

120   has clear fitness benefits: territorial males with high social connectivity have greater reproductive

success (Ryder et al., 2009), and floaters with high connectivity have an increased probability of

122   territorial inheritance (Ryder et al., 2008).

        To evaluate the influence of the social environment on cooperative behavior, we

124   leveraged our knowledge of the social system to develop an autonomous proximity data-logging

system to obtain repeated measures of behavioral phenotype for a population of 179 males (Figs.

126   2, 3). We chose the phenotypes illustrated in Fig. 3 because they capture a range of social

behaviors that are expected to promote reproductive success and thus be under selection: "effort"

128   is known predict male mating success in other taxa (Hoglund and Alatalo, 1995); "strength" and

"degree" are particularly good predictors of social ascension and reproductive success in

130   manakins and other organisms (Ryder et al., 2008; Ryder et al., 2009; Formica et al., 2011); and

social "importance" may predict individual quality and signal output (Trainer and McDonald,

132   1995; Ryder et al., 2011).

We used variance partitioning to evaluate the impact of direct and indirect effects on

134   variation manakin cooperative behavior. The manakin system is ideal for this question because,

although manakin coalitions are relatively stable on an annual basis (Ryder et al., 2011),

136   partnerships are dynamic on shorter daily or weekly timescales and males show substantial

variation in cooperation (e.g., Fig. 2f; see also the supplementary materials). Our analysis had

138   three aims. First, we estimated the proportion of total phenotypic variance attributed to

differences among individuals (the direct effect). Second, we estimated the strength of social

140   influence on phenotypic variation within individuals (the indirect effect). A significant indirect

effect has two main implications: on the one hand, it tests how much of the total phenotypic

142   variance can be explained by social influence, but also, it implies that individuals differ in the

effect they have on others (Fig. 1a). We compared these results to null models of the network

144   data where the direct effects and indirect effects had been removed by permuting individual ID

labels. Third, we tested for social contagion (Fig. 1b-d) by estimating the correlation between an

146   individual's own phenotypic expression (the repeatable component of a specific individual's

behavior) and his social influence (the repeatable component of his influence on others). A

148   positive covariance is consistent with the hypothesis that the expression of behavior is social

contagious.

152   **MATERIAL AND METHODS**

Study System and Field Methods

154   We studied a color-banded population of male wire-tailed manakins (*P. filicauda*) at the Tiputini

Biodiversity Station, Orellana Province, Ecuador (0° 38' S, 76° 08' W, approx. 200 m elevation)

156   across three breeding seasons (November to March 2015-18; hereafter years '15-16', '16-17',

and '17-18'). The males at this site have been observed and color-banded annually since 2002.

158   Male status (i.e., territorial or floater) was determined by direct observation in the field following

methods outlined in Ryder et al. (2008).

160   To collect repeated measures of the behavior and social environment of male manakins,

we developed the automated proximity system shown in Fig. 2 (Ryder et al., 2012). Each

162   individual was outfitted with a coded nano-tag (NTQB-2, Lotek Wireless; $n_{15-16} = 100$ males that

were tagged and color-banded, $n_{16-17} = 114$, $n_{17-18} = 82$, and $n_{total} = 179$). The sample size was not

164   pre-determined, but instead the aim was to capture all individuals and monitor as many social

network interactions as possible. Proximity data-loggers (hereafter DL; SRX-DL800, Lotek

166   Wireless) were placed within the lek territories where the manakins engage in their cooperative

displays (Fig. 2a). DLs recorded continuously for 10 hours from 0600 to 1600, such that each

168   day constitutes a repeated measure of behavior. DLs were deployed at a given lek for an average

of ~6 days (± 1 SD) per sampling interval and moved among leks on a rotating schedule. In total,

170   we recorded 29,760 hours of proximity data over 249 days (15-16: 49 territories, mean = 16

recording days per territory; 16-17: 54 territories, mean = 21 days; 17-18: 48 territories, mean =

172   22 days, SD = ±4 days in all years).

Data Processing

176   To define distinct social interactions from the proximity data, we used a rule set with spatial and

temporal thresholds illustrated in Fig. 2a-d. This rule set was based on 16 years of previous study

178   demonstrating that spatial co-occurrence within a manakin territory is a both a necessary

prerequisite for, and an accurate predictor of, cooperative display interactions among males

180   (Ryder et al., 2008; Ryder et al., 2012). We also used a ground-truthing experiment to confirm

that the spatial threshold would identify birds in close proximity (Fig. 2b-c).

182   After removing any detections that were recorded at more than one site simultaneously

(see supplement for details), we used the proximity data to calculate repeated measures of four

184   behavioral phenotypes thought to be important for manakin reproductive success, illustrated in

Fig. 3a. (1) *Effort,* a measure of attendance on the lek, is the number of unique pings by a focal

186   individual. (2) *Strength* is the frequency of social interactions summed across all partners. (3)

Degree is the number of unique partners with whom a male interacted. Strength and degree are

188   thus measures of the frequency of cooperative behavior and the number of cooperative partners,

respectively. (4) *Importance* is a measure of the exclusivity of a male's partners, on a scale from

190   0 to 1. To calculate importance, we first found the proportion of each partner's interactions that

were with a focal male. Then, we took a weighted average using the focal male's interaction

192   frequencies. Thus, a male whose partners often interact with other individuals would obtain a

low score for importance, whereas a male whose partners interact with him exclusively would

194   obtain a score of 1. Additional details of the data processing methods and descriptive statistics

for the behavioral phenotypes and their correlations are provided in the supplement (Tables S1,

196   S2).

198 **Statistical Analysis**

Analyses were performed using R 3.5.1 (R Core Team, 2018). All materials necessary to

200 reproduce our results and figures, including data, R scripts, and posterior model results are

provided at: https://figshare.com/s/e73a3cb48e4714898d55

Assortment

204 As a general description of network assortment, we calculated the assortativity coefficient, ρ, for

each of the four phenotypes. This analysis was based on a single average value for each

206 individual (Fig. 3) and a single social network compiled over all three years of study. We first

derived Newman's assortativity coefficients using the igraph package 1.1.2 (Csardi and coathors,

208 2017), and verified that the results were identical to weighted network assortment coefficients for

continuous phenotypes using the assortnet package 0.12 (Farine, 2014; Farine, 2016). To test

210 whether phenotypic assortment was statistically significant, we determined 95% confidence

intervals using the jackknife resampling method in assortnet.

Direct and Indirect Effects

214 A major challenge when analyzing dynamic networks is that individuals can have multiple

partners and variable interaction frequencies (Fig. 2). We therefore developed an analytical

216 approach based on the multiple-membership structure modeling framework (i.e., multi-

membership models; hereafter, MMMs) for partitioning the phenotypic variance between the

218 direct and indirect effects. MMMs are an extension of multi-level models that incorporate a

heterogeneous weighted random effect (Buerkner, 2017), thus making it possible to capture

220 variation in dynamic network partnerships. The key advantage of MMMs is that the analysis can

be weighted according to the frequency with which different groupings or social relationships

222   occurred.

We fit MMMs using the brms package 2.4.0 (Buerkner, 2018) which employs the Stan

224   language in R, using the model syntax shown in Box 1. The response variable was an

individual's behavioral phenotype on a particular day; we also included bird's identity as a

226   simple random effect, as well as his top four most frequent partners on the previous day in the

multiple-membership structure, weighted by their interaction frequencies (Box 1). Thus, our

228   analysis asks, how is a bird's behavior influenced by his own identity and who he interacted with

on the previous day? Although a few birds had more than four partners in a single day, we chose

230   to limit our analysis to the top four because additional partners comprised less than 2% of a

bird's daily total in our data. When an individual had fewer than 4 partners on a given day, we

232   distributed one, two, or three of his realized partner IDs and their interaction frequencies over the

remaining empty slot(s), while maintaining each partner's total weight as a constant. As long as

234   the total weight of each partner remains unchanged, the exact redistribution procedure does not

influence the results, because MMM analysis normalizes weights within the focal individuals

236   (Buerkner, 2017).

Prior to modelling, three of the four behavioral phenotypes were log-transformed (effort,

238   strength, and degree) so that Gaussian model assumptions were met, and all phenotypes were

standardized (mean 0, SD 1). We used the default uninformative priors and stored 2,000 samples

240   from each of four independently-seeded chains for each phenotype, verifying that the

convergence statistics in the model output were all equal to 1. We then obtained estimates of the

242   variance components for each phenotype by pooling the posterior values from all four chains.

Posterior distributions for the direct and indirect effects were calculated as Var_{focal}/Var_{total} and

244 Var_{social}/Var_{total}, respectively (Dingemanse and Araya-Ajoy, 2015), wherein Var_{focal} is the

variance due to the identity of the focal individuals; Var_{social} is the variance due to the MMM

246 structure (i.e., the social environment); and Var_{total} is the sum of all variance components

(including focal, social, date, lek, and the residual/unexplained variance).

248   To account for additional sources of variation (Table S1), we also included fixed effects

of focal status (territorial vs. floater) and study year as well as random effects of date and lek in

250 all models. By including year, date, and lek, our estimates of the direct and indirect effects

account for spatial and temporal variation in the shared non-social environment that may

252 influence focal and partner behavior (e.g., Ryder et al., 2006; Loiselle et al., 2007; Ryder and

Sillett, 2016). Hence, our analysis computes the proportion of phenotypic variance due to direct

254 and indirect effects after accounting for status and any spatial/temporal dynamics. We do not

consider the influence of second- or higher-order connections or positive feedback loops among

256 focal and partner phenotypes, although these may also occur (McGlothlin and Brodie, 2009;

Brent, 2015).

Box 1. Syntax for weighted MMM analysis of direct and indirect effects in brms

scale(Focal.phenotype) ~ focal.status + year, random + (1|Focal.ID) + (1|mm(Partner.ID1, Partner.ID2,

Partner.ID3, Partner.ID4), weights=cbind(frequency.1, frequency.2, frequency.3, frequency.4)) + (1|date) +

(1|lek)

Null Models

260 Social network data are by definition non-independent, so they violate the assumptions of

traditional statistical inference (Farine and Whitehead, 2015; Farine et al., 2017). We therefore

262 used a null model framework to compare our posterior estimates of the direct and indirect effects

with the same analyses performed on randomly permuted data (Farine and Whitehead, 2015;

264   Farine et al., 2017). The null permutation preserved all structure in the data but permuted the ID

labels, which were swapped among individuals within days. This is equivalent to a node-label

266   network permutation. Given that territorial and floater birds differ (Ryder et al., 2012) (Table

S1), the permutation was also constrained to swap ID labels among status-matched individuals

268   recorded on the same day. Hence, the null model preserved the observed phenotypic distributions

and the spatiotemporal structure in the data, but broke any associations with identity, to provide a

270   strong test of the direct and indirect effects.

To analyze the null (permuted) data, we fit MMMs using the same syntax presented in

272   Box 1. After verifying initial model diagnostics on a small number of model runs for each

phenotype, we generated 1,000 null datasets, and fit a single model chain storing 2,000 iterations

274   to each one. Because the direct and indirect effects are bounded by 0, we used one-sided tests to

compare the observed posterior estimates with the median value obtained from these null

276   permutations.

Finally, as an additional check on our results, we also examined null models that

278   separately permuted either the Focal.ID labels or the Partner.ID labels alone. As expected,

permuting Focal.ID reduced the direct effect to near zero but gave the same indirect effect

280   observed in the actual data, whereas permuting Partner.ID reduced the indirect effect to near zero

but gave the same direct effect observed in the actual results.

Contagion

284   Social contagion results in a positive covariance between a bird's own phenotypic expression

and his social influence on others (Fig. 1b-d). To determine this covariance, we obtained

286   posterior estimates of phenotypic expression and social influence for each male in the population

from the fitted MMMs. These individual estimates are akin to the random intercepts or BLUPs in

288   a frequentist framework. We used the posterior medians to test the correlation between an

individual's own phenotypic expression and his social influence on each of the four behavioral

290   phenotypes. That is, do individuals that express consistently high levels of a phenotype also elicit

that same phenotype in others? To verify that these associations were significant even after

292   accounting for error, we used Deming regression analyses with the posterior standard deviations

as the error in x and y.

Sample Sizes

296   A small number of individuals (n = 9) had tags that either stopped transmitting to the receivers

shortly after they were released, or they left the study population, and hence they are not

298   included in this study. Twenty-two of the successfully tagged males were also implanted with

hormone and sham pellets in part of years 2 and/or 3 for a separate study of the endocrine

300   mechanisms of cooperative behavior. We consequently limited our statistical analysis here to

focal individuals at times when they had no implants; nevertheless, the implant birds still acted

302   within the social networks and their interactions are retained when calculating the behavioral

phenotype of other individuals. Our sample size for analysis was 2,935 daily measures of 144

304   focal individuals. Note that this is smaller than the number of tagged birds, because a focal bird

had to have a known social environment on the previous day to be included in the MMM

306   analysis.

310   **RESULTS**

Three of the four phenotypes, strength, degree, and importance, were positively assorted in the

312   social network (Fig. 3b). All four phenotypes also had statistically significant individual

differences in expression, such that the direct effects could explain ~12-30% of the total variance

314   (Fig. 4a, Table S3, all $p < 0.004$). Because these estimates are moderate, this result demonstrates

substantial within-individual plasticity in all four behavioral phenotypes.

316   We next asked if the social environment could explain some of this within-individual

variation in behavior. We found that between 12-24% of the total variance could be attributed to

318   indirect effects caused by the dynamic network interactions on the previous day (Fig. 4b, Table

S3, all $p < 0.006$). The phenotypes with the larger direct effects (effort and strength) also tended

320   to have smaller indirect effects, and vice versa (inset, Fig. 4b). These significant indirect effects

also imply that social influence is itself a trait that varies among individual male manakins.

322   Moreover, our null models demonstrate that indirect effects of this magnitude do not arise by

chance in permuted data with the exact same phenotypes and network topologies.

324   To further visualize these indirect effects, we also plotted within-individual plasticity in

phenotypic expression in relation to the social environment, as defined by average partner social

326   influence from the previous day (Fig. 4c). These plots show how individuals adjust their

behavior based on their dynamic social interactions on the previous day.

328   Finally, we asked if indirect effects as our measure of social influence could drive the

social contagion of behavior. The key parameter to test this hypothesis is the correlation between

330   an individual's own phenotypic expression and his social influence on others (Fig. 1). We found

strong positive correlations between expression and social influence for three of the four

332   phenotypes: strength, degree, and importance (Fig. 5), all of which exhibited positive assortment

(Fig. 3b). This suggests that individual manakins with consistently high strength, degree, and

334   importance tend to elicit increases in those respective phenotypes in their partners. In contrast,

although greater effort was also elicited by particular individuals (Fig. 4), its lack of a significant

336   positive correlation demonstrates that effort is not socially contagious (Fig. 5).

338   **DISCUSSION**

Social contagion facilitates the evolution of positive assortment and cooperation (van Doorn and

340   Taborsky, 2012; Christakis and Fowler, 2013), yet understanding how dynamic network

interactions shape variation in cooperative behavior has remained challenging outside of the

342   laboratory. Here, we monitored a large population of cooperative wire-tailed manakins, and we

applied a variance-partitioning approach to characterize sources of variation in four behavioral

344   phenotypes. We found that individuals who are consistently more connected (high strength and

degree) tend to elicit increased connectivity in others, and individuals who repeatedly attract

346   more exclusive partnerships (high importance) tend to elicit greater importance in others. This is

notable because social connectivity has a direct link to reproductive success in this population

348   and related species in the same family (McDonald, 2007; Ryder et al., 2008; Ryder et al., 2009).

In contrast, we find no evidence that a male's effort in attending leks is socially contagious

350   (although it is influenced by partner identity). Viewed cumulatively, our work demonstrates how

multiple sources of phenotypic variation can be parsed in a network to quantify the social

352   influence on behavioral plasticity in the wild. Furthermore, we demonstrate that social contagion

can contribute to positive network assortment in a cooperative system.

354   Over the last five years, social network analysis has rapidly advanced our understanding

of the causes and consequences of social behavior (reviewed in Kurvers et al., 2014). For

356   example, recent work has characterized individual differences in network position (Jacoby et al.,

       2014; Aplin et al., 2015), patterns of phenotypic assortment (Croft et al., 2009; McDonald et al.,

358   2017), and the fitness consequences of repeated social interactions (McDonald, 2007; Ryder et

       al., 2009; Oh and Badyaev, 2010). Only recently, however, have we begun to integrate network

360   theory into our thinking about selection dynamics and how structured interactions modulate

       phenotypic variance (Fisher and McAdam, 2017; Montiglio et al., 2017). Here, we continue to

362   advance that theme with the first empirical evidence that dynamic network interactions and

       social influence can have profound effects on variation in cooperative behavior. We expect that

364   these social processes will also contribute to the emergence and maintenance of cooperation in

       other animal systems (Bailey et al., 2018).

366   If the observed correlation between the expression of degree and downstream social

       influence in wire-tailed manakins has some genetic basis (i.e., heritability), then our results

368   imply that indirect effects can enhance the response to selection on cooperative behavior (Moore

       et al., 1997; Wolf et al., 1998; McGlothlin et al., 2010). This is because selection acting on the

370   genetic architecture for cooperative behavior would also facilitate a social environment that

       reinforces the benefits of cooperation (van Doorn and Taborsky, 2012). The genetic basis of

372   social influence could be tested in future studies that build on our approach by using genotyping

       to resolve the population pedigree. This will require analytical methods that can simultaneously

374   account for genetic structure among identities in the social environment (Wilson et al., 2009;

       Santostefano et al., 2017; Fisher et al., 2018).

376   Biologists have long recognized that no phenotype is expressed in a vacuum. It is

       important to note that in natural settings, the non-social environment is also variable, and thus

378   interacting individuals may share features of the environment that can be mutually influential.

Although this possibility cannot be excluded here or in any observational study, our analyses

380   account for sources of shared annual, daily, and spatial variation. Hence, the results are expected

to be independent of factors that vary in space and time, such as resource availability, female

382   activity, and climatic variation (Ryder et al., 2006; Loiselle et al., 2007; Ryder and Sillett, 2016).

Another important point is that in natural settings, individuals have the capacity to choose

384   their partners and these decisions can directly impact the costs and benefits of engaging in

cooperative acts (Bergmuller et al., 2010; Fehl et al., 2011; Rand et al., 2011). For example,

386   young wire-tailed manakins are known to increase coalition stability with age as they ascend the

queue to territorial status (Ryder et al., 2011). The long-term nature of male social partnerships

388   and evidence of behavioral plasticity in response to the dynamic social landscape (i.e., social

competence, *sensu* Taborsky and Oliveira, 2012) both likely facilitate the choice of similar

390   partners (homophily) in this study system. Furthermore, homophily of socially contagious traits

may further amplify or accelerate its transmission (Montiglio et al., 2017). Determining the

392   relative contribution of these two processes (contagion and homophily) in a natural system

remains a major challenge for future work and may require social transplant experiments to

394   resolve.

Finally, our results raise several key questions for further work. First, what are the

396   sensory mechanisms of social contagion for cooperative behavior (e.g., calls, odors, and/or

gestural displays)? Second, what explains the persistence of individual differences in social

398   influence (Montiglio et al., 2017)? Third, what are the underlying physiological and mechanistic

processes, and how do these mechanisms facilitate or constrain social contagion as a form of

400   cooperative reciprocity? Work examining the hormone signaling pathways that control

cooperative behavior will further elucidate the links between genotype, phenotype, and partner

402 interactions (Soares et al., 2010). Moreover, because behavior, hormones, and gene expression

can mutually affect one another, characterizing how these processes interact to modulate the

404 social dynamics is a pivotal next step (Oliveira, 2009). Given that repeated heterogeneous

interactions underlie virtually all animal social systems, our results provide a framework that can

406 be broadly applied to test how social influence changes in response to physiological

manipulations. Ultimately, understanding how selection shapes interactive phenotypes like

408 cooperation will require integrative approaches that consider both mechanism and social context.

FIGURE LEGENDS

Fig. 1. A conceptual framework for social influence. (a) A phenotype "P" varies within and
412 among individuals. The upper histogram shows the hypothetical phenotypic distribution for a
population, whereas the lower histograms represents the narrower range of phenotypic values
414 expressed by a given individual (e.g., X) within that population. Individual phenotypes are the
result of both intrinsic biology (the "direct effect") and plasticity induced by the social
416 environment (e.g., the "indirect effect" as X interacts with conspecifics Y and Z). This indirect
effect represents how Y and Z can differentially influence the phenotype of X through social
418 influence. (b-d) Because Y and Z also express phenotype P, there may be covariance between
social influence and phenotypic expression. For example, (b) phenotypic expression may be
420 positively correlated with social influence on X's phenotype. This positive covariance is
predicted to generate social contagion. Alternatively, phenotypic expression may be (c) unrelated
422 or (d) negatively related to social influence, the latter of which would generate social inhibition.

424 **Fig. 2. Social interactions were tracked using an automated proximity data-logging system.**
(a) Each bird was tagged with a small transmitter that broadcasts a unique digital code at 20 s
426 intervals. Tag pings were received by stationary data loggers (DLs) set with a fixed detection
radius to encompass a male's territory where cooperative displays take place. (b) The tag signal
428 strength, RSSI, decays with distance from the DL. Hence, we define an interacting dyad using a
conservative spatiotemporal threshold wherein individuals had to be detected within 45 s of each
430 other and with a ΔRSSI < 10. (c) A ground-truthing analysis demonstrates that tag pairs that
meet this threshold are typically $\leq$ 5 m apart (vertical line = median), corresponding to a range
432 that would permit both visual and acoustic contact. For comparison, events with ΔRSSI $\geq$ 10
have a greater median inter-tag distances of 15 m. The data in (b) and (c) are based on the same
434 ground-truthing experiment. (d) An example of detection data for DLs at two neighboring
territories. The resulting time series illustrates two interactions, dyad 1 and dyad 2, that are
436 delimited by horizontal brackets. Note that birds occasionally ping the same DL around the same
time without meeting the ΔRSSI spatial threshold. (e) Overall, this automated system recorded
438 approx. 37,000 social interactions among 179 birds. (f) An example of daily interactions within a
subset of the population social network illustrating the dynamic nature of the study system. Each
440 node in the network represents an individual male sized according to degree. Data are shown for
14 of the males recorded from January 1-6 in 2016, with solid edges representing interactions
442 that occurred on that day. Dotted edges are interactions that occurred but not on the given
sampling day.

**Fig. 3. We analyzed four behavioral phenotypes that are important for manakin
446 reproductive success.** (a) Four behavioral phenotypes occurred on territories within the leks and
were quantified on a daily basis. For effort, 1,000 pings are approximately 5.5 hours of
448 attendance. Note that in the illustration for importance, the focal male (gray node) maintains the
same strength and degree but increases his importance on the right because each of his partners is
450 exclusive. (b) Observed distributions of average phenotypic expression for floater and territory-
holding males. Dashed vertical lines show medians for each status class. Strength, degree, and
452 importance are all positively assorted within the social network. See Tables S1-S2 and Figure S1
for additional data.

Fig. 4. Manakins differ both in their social behavior and their influence on others. (a-b) The proportion of total phenotypic variance in four behavioral phenotypes attributed to (a) individual differences (the direct effect) and (b) social influence (the indirect effect). For each phenotype, the observed result is significantly greater than the null expectation. The vertical lines show the posterior median. Phenotypes with larger direct effects tend to have weaker social influence (inset, b). The axes on the inset span 0 to 0.4. Note that the tails of all distributions are extended beyond the range of the data for visual continuity. (c) Within individuals, phenotypic expression depends on the social environment. On the y-axes, individual phenotypic expression are centered to show change relative to an individual's mean expression; values for effort, strength, and degree were also log-transformed prior to centering. The x-axis represents the average influence of a bird's partners on the previous day. A separate linear fit is shown for each focal bird observed on at least 10 days (n = 95). See Table S3 for detailed results.

Fig. 5. Positive covariance indicates that cooperative social behaviors are contagious. Each graph shows an individual's estimated social influence (y-axis) in relation to his own phenotypic expression (x-axis) for that same behavior (n = 142 individuals analyzed as both focal and partner). Strength, degree, and importance all have strong positive correlations, indicating that indirect effects can facilitate the spread of these behaviors through the social network. In contrast, the correlation for effort is not statistically significant. Data points and error bars are derived from the posterior medians and standard deviations, respectively. The regression fit lines account for posterior estimates of the error in both variables. Pearson's correlation coefficients are also given on the lower right of each panel.

478 **ACKNOWLEDGEMENTS**

We thank Ignacio Moore, Ben J. Vernasco, and Camilo Alfonso for their tireless effort in project

480 organization and collecting field data. David and Consuelo Romo, Kelly Swing, Diego Mosquera

and Gabriela Vinueza and staff provided amazing logistical support at Tiputini Biodiversity

482 Station. We thank Brian S. Evans for his brilliance in data architecture and the development of R

functions. Brent Horton, Mike Webster, and Joel McGlothlin provided valuable comments on an

484 earlier version of this manuscript. This work was supported by the National Science Foundation

(NSF) IOS 1353085 and the Smithsonian Migratory Bird Center.

SUPPORTING INFORMATION

Supplementary Materials. Additional details of the methods and data tables.

Movie 1. Male wire-tailed manakins perform cooperative displays in Ecuador. These video clips

492 were taken during the 2015-16 breeding season and show examples of the coded tags used to

track social interactions.

494   **REFERENCES**

Anderson, B. B., Scott, A. and Dukas, R. (2017). Indirect genetic effects on the sociability of
496   several group members. *Animal Behaviour* **123**, 101–106.

Aplin, L. M., Firth, J. A., Farine, D. R., Voelkl, B., Crates, R. A., Culina, A., Garroway, C.
498   **J., Hinde, C. A., Kidd, L. R., Psorakis, I., et al.** (2015). Consistent individual
differences in the social phenotypes of wild great tits, *Parus major*. *Animal Behaviour*
500   **108**, 117–127.

Bailey, N. W., Marie-Orleach, L., Moore, A. J. and Simmons, L. (2018). Indirect genetic
502   effects in behavioral ecology: does behavior play a special role in evolution? *Behav Ecol*
29, 1–11.

Bergmuller, R., Schurch, R. and Hamilton, I. M. (2010). Evolutionary causes and
504   consequences of consistent individual variation in cooperative behaviour. *Philosophical*
506   *Transactions of the Royal Society B* **365**, 2751–2764.

Bijma, P. (2014). The quantitative genetics of indirect genetic effects: a selective review of
508   modelling issues. *Heredity* **112**, 61–69.

Bleakley, B. H. and Brodie, E. D. (2009). Indirect genetic effects influence antipredator
510   behavior in guppies: estimates of the coefficient of interation psi and the inheritance of
reciprocity. *Evolution* **63**, 1796–1806.

Brent, L. J. N. (2015). Friends of friends: are indirect connections in social networks important
512   to animal behaviour? *Animal Behaviour* **103**, 211–222.

Buerkner, P.-C. (2017). brms: an R package for Bayesian multilevel models Using Stan.
514   *Journal of Statistical Software* **80**,.

Buerkner, P.-C. (2018). *brms 2.4.0: Bayesian regression models using Stan*.
516

Christakis, N. A. and Fowler, J. H. (2013). Social contagion theory: examining dynamic social
518   networks and human behavior. *Statistics in Medicine* **32**, 556–577.

Croft, D. P., Krause, J., Darden, S. K., Ramnarine, I. W., Faria, J. J. and James, R. (2009).
520   Behavioural trait assortment in a social network: patterns and implications. *Behav Ecol*
Sociobiol **63**, 1495–1503.

Csardi, G. and coathors (2017). *igraph 1.1.2: network analysis and visualization*.
522

Dingemanse, N. J. and Araya-Ajoy, Y. G. (2015). Interacting personalities: behavioural
524   ecology meets quantitative genetics. *Trends in Ecology & Evolution* **30**, 88–97.

Edenbrow, M., Bleakley, B. H., Darden, S. K., Tyler, C. R., Ramnarine, I. W. and Croft, D.
526   **P.** (2017). The evolution of cooperation: interacting phenotypes among social partners.
The American Naturalist **189**, 630–643.

528 **Endler, J. A.** (1986). *Natural Selection in the Wild*. Princeton University Press.

Farine, D. R. (2014). Measuring phenotypic assortment in animal social networks: weighted
530   associations are more robust than binary edges. *Animal Behaviour* **89**, 141–153.

Farine, D. R. (2016). *assortnet 0.12: calculate the assortativity coefficient of weighted and
532   binary networks*.

Farine, D. R. and Whitehead, H. (2015). Constructing, conducting and interpreting animal
534   social network analysis. *Journal of Animal Ecology* **84**, 1144–1163.

Farine, D. R., Freckleton, R. and Rands, S. (2017). A guide to null models for animal social
536   network analysis. *Methods in Ecology and Evolution* **8**, 1309–1320.

Fehl, K., Post, D. J. van der and Semmann, D. (2011). Co-evolution of behaviour and social
538   network structure promotes human cooperation. *Ecology Letters* **14**, 546–551.

Fisher, D. N. and McAdam, A. G. (2017). Social traits, social networks and evolutionary
540   biology. *Journal of Evolutionary Biology* **30**, 2088–2103.

Fisher, D. N., Wilson, A. J., Boutin, S., Dantzer, B., Lane, J. E., Coltman, D. W., Gorrell, J.
542   **C. and McAdam, A. G.** (2018). Indirect effects of territorial neighbors on the timing of
        spring breeding may counteract changes in selection in North American red squirrels.
544   *bioRxiv*.

Formica, V. A., Wood, C. W., Larsen, W. B., Butterfield, R. E., Augat, M. E., Hougen, H.
546   **Y. and Brodie, E. D.** (2011). Fitness consequences of social network position in a wild
        population of forked fungus beetles (*Bolitotherus cornutus*). *Journal of Evolutionary*
548   *Biology* **25**, 130–137.

Hadfield, J. D. and Wilson, A. J. (2007). Multilevel selection 3: Modeling the effects of
550   interacting individuals as a function of group size. *Genetics* **177**, 667–668.

Hoglund, J. and Alatalo, R. V. (1995). *Leks*. Princeton University Press.

552 **Jacoby, D. M. P., Fear, L. N., Sims, D. W. and Croft, D. P.** (2014). Shark personalities?
        Repeatability of social network traits in a widely distributed predatory fish. *Behav Ecol*
554   *Sociobiol* **68**, 1995–2003.

Kurvers, R. H. J. M., Krause, J., Croft, D. P., Wilson, A. D. M. and Wolf, M. (2014). The
556   evolutionary and ecological consequences of animal social networks: emerging issues.
        Trends in Ecology & Evolution **29**, 326–335.

558 **Loiselle, B. A., Blake, J. G., Durães, R., Ryder, T. B., Tori, W. and Vega Rivera, J. H.**
        (2007). Environmental and spatial segregation of leks among six co-occurring species of
560   manakins (Pipridae) in eastern Ecuador. *The Auk* **124**, 420–431.

McDonald, D. B. (2007). Predicting fate from early connectivity in a social network. *PNAS* **104**, 10910–10914.

McDonald, G. C., Farine, D. R., Foster, K. R. and Biernaskie, J. M. (2017). Assortment and the analysis of natural selection on social traits. *Evolution* **71**, 2693–2702.

McGlothlin, J. W. and Brodie, E. D. (2009). How to measure indirect genetic effects: the congruence of trait-based and variance-partitioning approaches. *Evolution* **63**, 1785–1795.

McGlothlin, J. W., Moore, A. J., Wolf, J. B. and Brodie, E. D. (2010). Interacting phenotypes and the evolutionary process. III. Social evolution. *Evolution* **64**, 2558–2574.

Montiglio, P., McGlothlin, J. W. and Farine, D. R. (2017). Social structure modulates the evolutionary consequences of social plasticity: A social network perspective on interacting phenotypes. *Ecology and Evolution* **8**, 1451–1464.

Moore, A. J., Brodie, E. D. and Wolf, J. B. (1997). Interacting phenotypes and the evolutionary process. I. Direct and indirect genetic effects of social interactions. *Evolution* **51**, 1352–1362.

Oh, K. P. and Badyaev, A. V. (2010). Structure of social networks in a passerine bird: consequences for sexual selection and the evolution of mating strategies. *The American Naturalist* **176**, E80–E89.

Ohtsuki, H., Hauert, C., Lieberman, E. and Nowak, M. A. (2006). A simple rule for the evolution of cooperation on graphs and social networks. *Nature* **441**, 502–505.

Oliveira, R. F. (2009). Social behavior in context: Hormonal modulation of behavioral plasticity and social competence. *Integr Comp Biol* **49**, 423–440.

R Core Team (2018). *R 3.5.1: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.

Rand, D. G., Arbesman, S. and Christakis, N. A. (2011). Dynamic social networks promote cooperation in experiments with humans. *PNAS* **108**, 19193–19198.

Rankin, D. J. and Taborsky, M. (2009). Assortment and the evolution of generalized reciprocity. *Evolution* **63**, 1913–1922.

Ryder, T. B. and Sillett, T. S. (2016). Climate, demography and lek stability in an Amazonian bird. *Proc. R. Soc. B* **283**, 20152314.

Ryder, T. B., Blake, J. G., Loiselle, B. A. and Vega Rivera, J. H. (2006). A test of the environmental hotspot hypothesis for lek placement in three species of manakins (Pipridae) in Ecuador. *The Auk* **123**, 247–258.

594   **Ryder, T. B., McDonald, D. B., Blake, J. G., Parker, P. G. and Loiselle, B. A.** (2008). Social
networks in the lek-mating wire-tailed manakin (*Pipra filicauda*). *Proceedings of the*
596   *Royal Society B* **275**, 1367–1374.

Ryder, T. B., Parker, P. G., Blake, J. G. and Loiselle, B. A. (2009). It takes two to tango:
598   reproductive skew and social correlates of male mating success in a lek-breeding bird.
Proceedings of the Royal Society B **276**, 2377–2384.

600   **Ryder, T. B., Blake, J. G., Parker, P. G. and Loiselle, B. A.** (2011). The composition,
stability, and kinship of reproductive coalitions in a lekking bird. *Behav Ecol* **22**, 282–
602   290.

Ryder, T. B., Horton, B. M., Tillaart, M. van den, Morales, J. D. D. and Moore, I. T. (2012).
604   Proximity data-loggers increase the quantity and quality of social network data. *Biology
Letters* **8**, 917–920.

606   **Santostefano, F., Wilson, A. J., Niemelä, P. T. and Dingemanse, N. J.** (2017). Indirect genetic
effects: a key component of the genetic architecture of behaviour. *Scientific Reports* **7**,
608   10235.

Schwartz, P. and Snow, D. W. (1978). Display and related behavior of the wire-tailed manakin.
610   *Living Bird* **17**, 51–78.

**Soares, M. C., Bshary, R., Fusani, L., Goymann, W., Hau, M., Hirschenhauser, K. and
612   Oliveira, R. F.** (2010). Hormonal mechanisms of cooperative behaviour. *Philosophical
Transactions of the Royal Society B* **365**, 2737–2750.

614   **Taborsky, B. and Oliveira, R. F.** (2012). Social competence: an evolutionary approach. *Trends
in Ecology & Evolution* **27**, 679–688.

616   **Trainer, J. M. and McDonald, D. B.** (1995). Singing performance, frequency matching and
courtship success of long-tailed manakins (*Chiroxiphia linearis*). *Behav Ecol Sociobiol*
618   **37**, 249–254.

Trivers, R. L. (1971). The evolution of reciprocal altruism. *The Quarterly Review of Biology* **46**,
620   35–57.

van Doorn, G. S. and Taborsky, M. (2012). The evolution of generalized reciprocity on social
622   interaction networks. *Evolution* **66**, 651–664.

Wilson, A. J., Gelin, U., Perron, M. C. and Reale, D. (2009). Indirect genetic effects and the
624   evolution of aggression in a vertebrate system. *Proceedings of the Royal Society B* **276**,
533–541.

626   **Wilson, A. J., Morrissey, M. B., Adams, M. J., Walling, C. A., Guinness, F. E., Pemberton,
J. M., Clutton-Brock, T. H. and Kruuk, L. E. B.** (2011). Indirect genetics effects and
628   evolutionary constraint: an analysis of social dominance in red deer, *Cervus elaphus*.
Journal of Evolutionary Biology **24**, 772–783.

630 **Wolf, J. B., Brodie, E. D., Cheverud, J. M., Moore, A. J. and Wade, M. J.** (1998).
Evolutionary consequences of indirect genetic effects. *Trends in Ecology & Evolution* **13**,
632 64–69.

Wolf, J. B., Brodie, E. D. and Moore, A. J. (1999). Interacting phenotypes and the evolutionary
634 process. II. Selection resulting from social interactions. *The American Naturalist* **153**,
254–266.

Fig. 1

Fig. 1. A conceptual framework for social influence. (a) A phenotype "P" varies within and among individuals. The upper histogram shows the hypothetical phenotypic distribution for a population, whereas the lower histograms represents the narrower range of phenotypic values expressed by a given individual (e.g., X) within that population. Individual phenotypes are the result of both intrinsic biology (the "direct effect") and plasticity induced by the social environment (e.g., the "indirect effect" as X interacts with conspecifics Y and Z). This indirect effect represents how Y and Z can differentially influence the phenotype of X through social influence. (b-d) Because Y and Z also express phenotype P, there may be covariance between social influence and phenotypic expression. For example, (b) phenotypic expression may be positively correlated with social influence on X's phenotype. This positive covariance is predicted to generate social contagion. Alternatively, phenotypic expression may be (c) unrelated or (d) negatively related to social influence, the latter of which would generate social inhibition.

Fig. 2

Fig. 2. Social interactions were tracked using an automated proximity data-logging system. (a) Each bird was tagged with a small transmitter that broadcasts a unique digital code at 20 s intervals. Tag pings were received by stationary data loggers (DLs) set with a fixed detection radius to encompass a male's territory where cooperative displays take place. (b) The tag signal strength, RSSI, decays with distance from the DL. Hence, we define an interacting dyad using a conservative spatiotemporal threshold wherein individuals had to be detected within 45 s of each other and with a ΔRSSI < 10. (c) A ground-truthing analysis demonstrates that tag pairs that meet this threshold are typically ≤ 5 m apart (vertical line = median), corresponding to a range that would permit both visual and acoustic contact. For comparison, events with ΔRSSI ≥ 10 have a greater median inter-tag distances of 15 m. The data in (b) and (c) are based on the same ground-truthing experiment. (d) An example of detection data for DLs at two neighboring territories. The resulting time series illustrates two interactions, dyad 1 and dyad 2, that are delimited by horizontal brackets. Note that birds occasionally ping the same DL around the same time without meeting the ΔRSSI spatial threshold. (e) Overall, this automated system recorded approx. 37,000 social interactions among 179 birds. (f) An example of daily interactions within a subset of the population social network illustrating the dynamic nature of the study system. Each node in the network represents an individual male sized according to degree. Data are shown for 14 of the males recorded from January 1-6 in 2016, with solid edges representing interactions that occurred on that day. Dotted edges are interactions that occurred but not on the given sampling day.

Fig. 3

Fig. 3. We analyzed four behavioral phenotypes that are important for manakin reproductive success. (a) Four behavioral phenotypes occurred on territories within the leks and were quantified on a daily basis. For effort, 1,000 pings are approximately 5.5 hours of attendance. Note that in the illustration for importance, the focal male (gray node) maintains the same strength and degree but increases his importance on the right because each of his partners is exclusive. (b) Observed distributions of average phenotypic expression for floater and territory-holding males. Dashed vertical lines show medians for each status class. Strength, degree, and importance are all positively assorted within the social network. See Tables S1-S2 and Figure S1 for additional data.

Fig. 4

Fig. 4. Manakins differ both in their social behavior and their influence on others. (a-b) The proportion of total phenotypic variance in four behavioral phenotypes attributed to (a) individual differences (the direct effect) and (b) social influence (the indirect effect). For each phenotype, the observed result is significantly greater than the null expectation. The vertical lines show the posterior median. Phenotypes with larger direct effects tend to have weaker social influence (inset, b). The axes on the inset span 0 to 0.4. Note that the tails of all distributions are extended beyond the range of the data for visual continuity. (c) Within individuals, phenotypic expression depends on the social environment. On the y-axes, individual phenotypic expression are centered to show change relative to an individual's mean expression; values for effort, strength, and degree were also log-transformed prior to centering. The x-axis represents the average influence of a bird's partners on the previous day. A separate linear fit is shown for each focal bird observed on at least 10 days (n = 95). See Table S3 for detailed results.

Fig. 5

Fig. 5. Positive covariance indicates that cooperative social behaviors are contagious. Each graph shows an individual's estimated social influence (y-axis) in relation to his own phenotypic expression (x-axis) for that same behavior (n = 142 individuals analyzed as both focal and partner). Strength, degree, and importance all have strong positive correlations, indicating that indirect effects can facilitate the spread of these behaviors through the social network. In contrast, the correlation for effort is not statistically significant. Data points and error bars are derived from the posterior medians and standard deviations, respectively. The regression fit lines account for posterior estimates of the error in both variables. Pearson's correlation coefficients are also given on the lower right of each panel.