Interaction of taste and place coding in the hippocampus

Running title: Taste and place coding in the hippocampus

Authors: Linnea E. Herzog[1], Leila May Pascual[2], Seneca J. Scott[2], Elon R. Mathieson[2], Donald

B. Katz[1,2*] and Shantanu P. Jadhav[1,2*]

[1]Graduate Program in Neuroscience, Brandeis University, Waltham, MA, 02453, USA

[2]Neuroscience Program, Department of Psychology, and Volen National Center for Complex

Systems, Brandeis University, Waltham, MA, 02453, USA

* These authors contributed equally to this work.

Correspondence: Donald B. Katz (dbkatz@brandeis.edu), Brandeis University, 415 South St.,

MS 062, Waltham, MA 02453

Shantanu P. Jadhav (shantanu@brandeis.edu), Brandeis University, 415 South St., MS 062,

Waltham, MA 02453

Number of Pages: 42

Number of Figures: 7   Number of Tables: 1

Number of words: Abstract (246 words); Introduction (645 words); Discussion (1479 words)

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgements: This work was supported by a Sloan Research Fellowship in

Neuroscience (Alfred P. Sloan Foundation) and Whitehall Foundation award to SPJ; NIH Grants

R01 DC006666 and R01 DC007703 to DBK; Training grant T90 DA032435 to LEH.

Abstract

An animal's survival depends on finding food, and the memory of food and contexts are often

linked. Given that the hippocampus is required for spatial and contextual memory, it is

reasonable to expect related coding of space and food stimuli in hippocampal neurons.

However, relatively little is known about how the hippocampus responds to tastes, the most

central sensory property of food. In this study, we examined the taste-evoked responses and

spatial firing properties of single units in the dorsal CA1 hippocampal region as male rats

received a battery of taste stimuli differing in both chemical composition and palatability within a

specific spatial context. We identified a subset of hippocampal neurons that responded to

tastes, some of which were place cells. These taste and place responses had a distinct

interaction: taste-responsive cells tended to have less spatially specific firing fields, and place

cells only responded to tastes delivered inside their place field. Like neurons in the amygdala

and lateral hypothalamus, hippocampal neurons discriminated between tastes predominantly on

the basis of palatability, with taste-selectivity emerging concurrently with palatability-

relatedness; these responses did not reflect movement or arousal. However, hippocampal taste

responses emerged several hundred msec later than responses in other parts of the taste

system, suggesting that the hippocampus does not influence real-time taste decisions, instead

associating the hedonic value of tastes with a particular context. This incorporation of taste

responses into existing hippocampal maps could be one way that animals use past experience

to locate food sources.

Significance statement

Finding food is essential for animals' survival, and taste and context memory are often linked.

While hippocampal responses to space and contexts have been well characterized, little is

known about how the hippocampus responds to tastes. Here, we identified a subset of

hippocampal neurons that discriminated between tastes based on palatability. Cells with

stronger taste responses typically had weaker spatial responses, and taste responses were

confined to place cells' firing fields. Hippocampal taste responses emerged later than in other

parts of the taste system, suggesting that the hippocampus does not influence taste decisions,

but rather, associates the hedonic value of tastes consumed within a particular context. This

could be one way that animals use past experience to locate food sources.

Introduction

The hippocampus is essential for spatial learning and memory, and is thought to provide a

cognitive map of animals' experience. The central data for this view come from studies of place

cells that respond to specific locations as animals explore their environments (O'Keefe and

Nadel, 1978; Moser et al., 2008).

Given that one of the most obvious uses for such a mental map is to aid in the finding of

food, it is surprising how little is known about how the hippocampus processes taste, the most

central sensory property of food. It is reasonable to expect that taste information reaches the

hippocampus; although not traditionally considered to be part of the taste system, anatomical

studies show that the hippocampus receives projections, either directly or indirectly through the

entorhinal cortex, from several brain regions in which taste information is processed, including

the gustatory cortex (GC), orbitofrontal cortex, and amygdala (Suzuki and Amaral, 1994; von

Bohlen und Halbach and Albrecht, 2002). Functional imaging studies in humans also indicate

that the hippocampal formation is active during taste ingestion and discrimination (Zald et al.,

1998; Haase et al., 2009; Spetter et al., 2010), and rodent lesion studies suggest that the

hippocampus plays a role in taste learning (Reilly et al., 1993; Stone et al., 2005;

Chinnakkaruppan et al., 2014).

A great deal is known about taste responses in other parts of the taste system: in cortex,

these responses evolve dynamically, reflecting taste presence, identity and palatability in

distinct epochs preceding the decision to consume or expel a given taste (Katz et al., 2001;

Sadacca et al., 2012; Sadacca et al., 2016). However, it remains unclear if or how hippocampal

taste responses co-exist and interact with representations of space. While hippocampal neurons

are known to respond to tastes in a context-dependent manner (Ho et al., 2011), no studies to

date have directly measured single unit responses to tastes in the hippocampus alongside

spatial firing properties, or examined the dynamics of these responses.

Hippocampal place cells are certainly capable of encoding non-spatial information such

as odors (Wood et al., 1999), visual cues (Fried et al., 1997), textures (Shapiro et al., 1997),

tones (Moita et al., 2003) and time (Kraus et al., 2013). Place cells typically respond to these

stimuli by modulating their firing rate ("rate remapping," Leutgeb et al., 2004; Allen et al., 2012)

or firing location ("global remapping," Leutgeb et al., 2005; Fyhn et al., 2007). A new cognitive

map can also be formed based on the parameters of a behaviorally relevant non-spatial

stimulus (Kraus et al., 2013; Aronov et al., 2017). The difficulty inherent in dissociating spatial

from non-spatial influences in behaving rodents (O'Keefe, 1999), however, has led some

researchers to propose that seeming responses to non-spatial stimuli may simply reflect arousal

triggered by the onset of the stimulus, rather than true sensory responses (Shan et al., 2016).

To establish that non-spatial responses are genuine, it is necessary to show that spatially tuned

neurons can discriminate between sensory stimuli.

Here, we did just this, recording single-unit activity in the dorsal CA1 region of awake

rats while exposing them to four taste solutions. We identified subsets of place cells and

interneurons that discriminated between tastes based predominantly on palatability; this pattern

was consistent with those observed in basolateral amygdala (BLA; Fontanini et al., 2009) and

lateral hypothalamus (LH; Li et al., 2013), although hippocampal taste dynamics evolved much

more slowly. Neurons classified as taste-responsive place cells responded exclusively to tastes

delivered within their place field, and tended to have lower spatial selectivity than non-taste-

responsive place cells. Together, these results establish that hippocampal responses to sensory

stimuli do not simply reflect changes in arousal state, and can encode sensory parameters

relevant for behavior. Further, they suggest that hippocampal taste responses may be used to

form value-related associations between tastes and contexts, which can facilitate using past

experience to locate food sources.

Materials and Methods

Animals and surgery

Five adult (450-550 g) male Long-Evans rats (Charles River Laboratories, RRID:

RGD_2308852) were used as subjects in this study. Rats were kept on a 12 h light/dark cycle,

with all sessions taking place around the same time during the light period. All surgical and

experimental procedures were conducted in accordance with the National Institutes of Health

guidelines and approved by the Brandeis University Institutional Animal Care and Use

Committee.

After several weeks of habituation to daily handling, animals were chronically implanted

with a microdrive array consisting of 25-30 independently moveable tetrodes in the right dorsal

hippocampal region CA1 (-3.6 mm AP, 2.2 mm ML), and an intra oral cannula (IOC). Each IOC

consisted of a polyethylene tube inserted beneath the temporalis muscle and terminating

anterolateral to the first maxillary molar, allowing for the precise delivery of taste solutions onto

the rat's tongue (Grill and Norgren, 1978; Travers and Norgren, 1986; Katz et al., 2001).

Following recovery from the implantation surgery (~7-8 days), rats were water-deprived

to 85-90% of their *ad libitum* weight to ensure taste consumption during the recording sessions.

At ~14 d after implantation, animals were habituated for at least 3 days to the behavioral

chamber, sleep box, and the delivery of taste solutions through the IOC; place cells were

expected to be stable after habituation (Thompson and Best, 1990; Agnihotri et al., 2004).

Following habituation, we performed daily recording sessions in which rats were exposed to

pseudo-randomized sequences of four standard taste stimuli (see **Figure 1A**, **1B** and *Passive*

taste administration paradigm). Following the conclusion of experiments, we made electrolytic

lesions through each electrode tip to mark recording locations. Brains were sectioned into 50

μm slices and stained with cresyl violet to confirm electrode placement in the hippocampal cell

layer (see **Figure 1C**).

<u>Passive taste administration paradigm</u>

Each recording session typically lasted between 2 and 3 hours, and consisted of three sessions

in a ~30 x 35 x 40 cm Plexiglass behavioral chamber (J. Green, Charles River Maker Lab)

interleaved with four 15-20 minute sleep sessions in a ~30 x 30 x 40 cm black box (rest box).

The first and last sessions in the behavioral chamber consisted of 15-20 minute periods in which

animals were habituated to the behavioral chamber in the absence of tastes. During the middle

experimental session (depicted in **Figure 1A**), rats received a pseudo-randomized sequence of

four standard taste stimuli [sweet: 4 mM saccharin (S); salty: 100 mM sodium chloride (N);

neutral: distilled water (W); and bitter: 5 mM quinine hydrochloride (Q)] that varied in hedonic

value and fell within the range of concentrations typically used by other groups (3-20 mM

saccharin, 10-300 mM sodium chloride, and 1-10 mM quinine; for review, see Frank and Brown,

2003; Kobayakawa et al., 2005; Accolla and Carleton, 2008; Geran and Travers, 2009; Rosen

et al., 2010; Chen et al., 2011; MacDonald et al., 2012; Li and Lemon, 2015; Sadacca et al.,

2016). Taste solutions were delivered directly onto the tongue in ~40 μL aliquots via four

polyamide tubes inserted into the IOC, with a separate tube for each solution to prevent the

mixing of tastes. Rats received 50 pseudo-randomized repeats of each of the four taste stimuli,

for a total of 200 taste deliveries. This number of deliveries allowed us to obtain sufficient

numbers of in-field and out-of-field trials for each taste stimulus (see *In-field vs. out-of-field*

analysis) before rats became satiated (Fontanini and Katz, 2005). This requisite number of trials

per stimulus is much greater than what is typically used in studies of taste coding in awake

rodents (10-30 trials, see Katz et al., 2001; Li et al., 2013; Baez-Santiago et al., 2016; Li et al.,

2016), and therefore necessitated limiting the number of stimuli (e.g., sour taste) required for

these comparisons (Moita et al., 2003). The interval between taste deliveries was randomized to

13-17 seconds, which is sufficiently long enough to prevent mixture effects or contamination by

previous deliveries, even without rinses factored into the experimental paradigm, since awake

rats have previously been shown to engage in what amounts to a constant saliva rinse

(Fontanini and Katz, 2006). The total amount of fluid delivered in each ~50 minute taste

administration period was 8 mL, after which animals had access to an additional 15-20 mL of

water in their home cage.

Electrophysiology

Electrophysiological recordings were conducted using a SpikeGadgets system (Tang et al.,

2017). Spikes were sampled at 30 kHz and bandpass filtered between 600 Hz and 6 kHz. Local

field potentials (LFPs) were sampled at 1.5 kHz and bandpass filtered between 0.5 and 400 Hz.

During recording sessions, the animal's position and speed were recorded using an overhead

monochrome CCD camera (30 fps) and tracked by LEDs affixed to the headstage.

Over ~14 d following surgery, tetrodes were gradually advanced to the CA1 hippocampal

cell layer, as identified by characteristic EEG patterns (sharp-wave ripples, or SWRs; theta

rhythm) as previously described (Jadhav et al., 2012; Jadhav et al., 2016; Tang et al., 2017).

Tetrodes were readjusted after each day's recordings. Each animal had one hippocampal

reference tetrode in corpus callosum, which was also referenced to a ground screw installed in

the skull overlying cerebellum.

Single units were isolated offline based on peak amplitude and principal components

(Matclust, M.P. Karlsson). Only well-isolated units with stable waveforms that fired at least 100

spikes per session were included in our analysis. As is typical for studies of hippocampal CA1

neurons (Fox and Ranck, 1981; Jadhav et al., 2016; Tang et al., 2017), putative interneurons

(*Int*) were identified on the basis of firing rate (> 8.5 Hz) and spike width (< 0.35 ms) parameters

(**Figure 1D**). All other isolated units were classified as putative pyramidal cells (*Pyr*). We

emphasize that these distinctions are putative, as definitive classifications of cell types cannot

be determined from extracellular recordings alone. We isolated a total of 482 neurons from five

rats, conducted across nine experiments. **Table 1** shows the distribution of cells across all five

animals.

SWR detection

SWRs were detected as previously described (Jadhav et al., 2016; Tang et al., 2017) using the

ripple-band (150-250 Hz) filtering of LFPs from multiple tetrodes. A Hilbert transform was used

to determine the envelope of band-passed LFPs, and events that exceeded a threshold (mean

+ 3 SD) were detected. SWR events were defined as the times around initially detected events

when the envelope exceeded the mean. SWR periods were excluded from place field analysis,

similar to previous studies (Jadhav et al., 2016; Tang et al., 2017).

Palatability/preference data

Palatability can be defined as the relative hedonic value of tastes; this characteristic is related,

but not identical to reward, as palatability is a continuously-varying property of taste that is

easily manipulated by experience (Stone et al., 2005; Katz and Sadacca, 2011;

Chinnakkaruppan et al., 2014). Taste palatability was assessed using a brief-access task (BAT,

Davis Rig Gustometer, Med Associates; for details, see Sadacca et al., 2016) in a separate

cohort of adult male rats (n = 7) that underwent the same water restriction protocol as the rats

used in the recording experiment. Consumption data were averaged across two testing days for

each animal. The palatability rank order determined by the brief-access test (S > N > W > Q,

see **Figure 6C**) matches what has been observed in numerous studies across a broad range of

stimulus delivery methods and assessment techniques (Travers and Norgren, 1986; Breslin et

al., 1992; Clarke and Ossenkopp, 1998; Fontanini and Katz, 2006; Sadacca et al., 2016).

Experimental design and statistical analysis

Spatial maps:

To characterize the spatial firing properties of neurons, two-dimensional occupancy-normalized

firing rate maps (**Figures 2**, **4, 5**) were made using 0.5 cm square bins and smoothed with a 2D

Gaussian (σ = 3 cm; Tang et al., 2017). Data from taste delivery (500 ms before to 2500 ms

after) and SWR periods (see *SWR detection and modulation*) were excluded from spatial map

analysis. Peak rates for each cell were defined as the maximum firing rate across all spatial bins

in the spatial map. Each cell's place field was defined as the largest cluster of neighboring

spatial bins which had firing rates >= 20% of the peak rate; place field sizes were then

calculated by multiplying the number of bins by the bin size (Brun et al., 2002). Unpaired t-tests

were used to determine whether the mean firing field size differed significantly between taste-

responsive and non-taste-responsive neurons of each cell type (**Figure 4B**; pyramidal cells: n =

65 taste-responsive cells, n = 376 non-taste-responsive cells; interneurons: n = 31 taste-

responsive cells, n = 10 non-taste-responsive cells).

Spatial specificity was determined by calculating the spatial information content, or

amount of information that a single spike conveys about the animal's location in bits/spike using

the formula:

Spatial information content $= \sum P_i(R_i/R)log_2(R_i/R),$

where i is the bin number, P_i is the probability of occupancy for bin i, R_i is the mean firing rate for

bin i, and R is the overall mean firing rate of the cell (Skaggs et al., 1993).

Unpaired t-tests were used to determine whether the average spatial information content

differed significantly between taste-responsive and non-taste-responsive neurons of each cell

type (**Figure 4C**; pyramidal cells: n = 65 taste-responsive cells, n = 376 non-taste-responsive

cells; interneurons: n = 31 taste-responsive cells, n = 10 non-taste-responsive cells).

In-field vs. out-of-field analysis:

To analyze how place cells responded to tastes delivered inside or outside of their place fields

(**Figure 5**), only pyramidal cells exhibiting place-specific activity (n = 395 cells, defined as

neurons whose peak rate exceeded 1 Hz and spatial information content exceeded 0.2

bits/spike, similar to Moita et al., 2003) were considered. Only place cells that contained at least

ten in-field and out-of-field deliveries of each taste were included in this in-field vs. out-of-field

analysis (n = 26 taste-responsive cells, n = 153 non-taste-responsive cells). A one-way ANOVA

was used to assess differences between the mean in-field and out-of-field taste response

magnitude (calculated using η^2, see *Taste selectivity*) of taste-responsive and non-taste-

responsive cells (**Figure 5C**). To ensure that comparable numbers of trials of each taste were

delivered in-field, a one-way ANOVA was used to compare the mean number of in-field trials for

each of the four tastes. An unpaired t-test was also used to compare the total number of in-field

trials for taste-responsive and non-taste-responsive cells. To determine whether numbers of

taste-responsive cells were underreported due to inter-trial variability, we repeated our initial

analyses of taste responsiveness (see *Taste response properties*) using only in-field trials of

place cells that met the in-field vs. out-of-field analysis criteria (n = 179 cells). A chi-square test

was used to evaluate whether this approach resulted in comparable numbers of presence-,

identity- and palatability-responsive cells.

Taste response properties:

The pseudo-randomized taste delivery paradigm used to characterize hippocampal responses

to tastes is described above (see *Passive taste administration paradigm*). Taste responses were

characterized separately for each of the 482 isolated neurons, focusing on the 2500 ms of

spiking activity following each taste delivery, a time period that includes previously identified

taste-related responses, but precedes swallowing behaviors that remove tastes from the tongue

and make neural responses difficult to interpret (Travers and Norgren, 1986; Katz et al., 2001).

We analyzed a set of response properties ranging from general to specific, as have been

identified in other parts of the taste system, including the GC (Katz et al., 2001; Sadacca et al.,

2012), BLA (Fontanini et al., 2009; Piette et al., 2012), and LH (Li et al., 2013). Neurons were

classified as "taste-responsive" (see **Table 1** for summary) if they exhibited responses to taste

presence, identity and/or palatability, as described below. All other neurons were classified as

"non-taste-responsive." All statistical tests were performed in MATLAB (MathWorks, Natick, MA,

RRID: SCR_001622) and evaluated at a level of α = 0.05 unless otherwise specified, with a

Bonferroni correction applied for multiple comparisons.

First, non-specific responses to taste <u>presence</u> (**Figure 6B**, light gray lines), which are

common across all four types of taste delivery and thought to originate from somatosensory

responses detecting a taste on the tongue, were determined by assessing whether evoked

responses differed significantly from the baseline firing rate in responses collated across all 200

taste delivery trials (Katz et al., 2001). The significance of the difference was first established

using the main effect for time in a two-way, mixed-effect ANOVA (taste [saccharin, NaCl, water,

quinine] x time [successive 500 ms bins of firing rate]).

Next, responses to taste <u>identity</u> (**Figure 6B**, dark gray lines), in which at least one taste

can be discriminated from the others, were assessed by determining if the evoked responses to

the four tastes (this time, collated across the 50 deliveries of each unique taste) differed from

each other. We employed a similar strategy as the one used to evaluate taste responsiveness,

except in this case, the main effect for taste was considered.

Finally, responses to taste <u>palatability</u> (**Figure 6D**), which reflected the relative hedonic

value of tastes as assessed in the BAT (see **Figure 6C** and *Palatability/preference data*) were

computed using a Spearman rank-order correlation (r_s) between the evoked response and the

palatability of the associated taste. Specifically, neurons whose evoked firing rates matched the

ranking of taste preference (S > N > W > Q) in increasing or decreasing order had higher

palatability index scores.

Population responses to tastes:

For all taste-responsive cells (n = 96, see *Taste response properties* for classification criteria),

the mean evoked firing rate across trials was determined (window size, 500 ms; step size, 50

ms; span, 0-2500 ms after taste delivery; Sadacca et al., 2012; Li et al., 2013) and Z-scored

separately for each taste (**Figure 3A**). Responses to all four tastes were sorted by the timing of

their peak firing to saccharin.

Taste selectivity:

The magnitude of taste responsiveness for each cell was quantified using eta-squared (η^2), a

standard measure of ANOVA effect sizes that describes the proportion of variance in a

dependent variable explained by each factor:

309             $\eta^2 = SS_{(factor)}/SS_{(total)}$,

where SS is the sum of squares (Maier et al., 2015). In our analysis, we used the summed SS

of the two main factors (time + taste) to calculate η^2. A chi-square goodness-of-fit test was

used to determine if η^2 values followed a normal distribution (**Figure 3B**). The Pearson

correlation (R) between spatial information content and η^2 was computed separately for place

cells (n = 395 cells) and interneurons (n = 41 cells). As described above (see *In-field vs. out-of-*

field analysis), a one-way ANOVA was used to assess differences in η^2 for the in-field and out-

of-field regions of taste-responsive (n = 26 cells) and non-taste-responsive (n = 153 cells) place

cells that fit our analysis criteria (**Figure 5C**).

Taste response dynamics:

To determine the timing of presence-, identity- and palatability-related responses in single

neurons, Student's t-tests were conducted on successive time windows of each neuron's

evoked response (window size, 500 ms; step size, 50 ms; span, 0-2500 ms after taste delivery;

Sadacca et al., 2012; Li et al., 2013). To determine which taste(s) each of the identity- and

palatability-responsive neurons (n = 40) preferentially responded to, the taste-evoked and

baseline (-500 to 0 ms before taste delivery) firing rates were compared using successive t-

tests. A chi-square goodness-of-fit test was used to determine whether cells responded to one

taste, or to a particular number of tastes, more than the others.

To analyze taste-related dynamics on a population-wide level, we constructed a

histogram showing what percentage of the total 482 recorded neurons exhibited responses to

taste presence, identity and palatability at each time point following stimulus delivery (**Figure

7A**). To investigate the timing of different aspects of the taste experience present in

hippocampal responses, we compared response onset times of presence- and identity-related

firing (**Figure 7B**) as well as identity- and palatability-related firing (**Figure 7C**) in the subset of

cells that exhibited both (n = 19 and 14 cells, respectively). Principal component analysis (PCA,

see Briggman et al., 2005; Harvey et al., 2012) was conducted on the pooled subset of 36

identity-responsive cells to determine when discriminative firing emerged in the population

response following taste delivery, with significance assessed at the $\alpha = 0.01$ level comparing the

neural data to 10,000 instances of firing-rate-shuffled controls (**Figure 7D**).

Speed and position controls:

To ensure that hippocampal responses to tastes were not actually caused by overall differences

in movement following taste delivery or in response to different tastes, we used a one-way

ANOVA to compare the average speed and distance traveled during the pre- vs. post-taste

period (2.5 seconds before or after taste delivery, segmented into 500 ms bins with a 50 ms

step size) across all tastes (n = 1800 total trials across 9 sessions), as well as separately for

each of the four tastes (n = 450 trials of each taste across 9 sessions). To confirm that tastes

were delivered in different spatial locations in our paradigm, we used a one-way ANOVA to

compare the mean number of taste trials in each spatial quadrant of the behavioral chamber.

Results

Hippocampal place cells and interneurons respond to tastes.

We examined taste responses in a total of 482 CA1 neurons recorded across nine sessions in

five rats (mean ± SEM: 53.6 ± 5.34 neurons/session) that received a battery of four standard

tastes *via* IOC (**Figure 1A**). Tastes were delivered in random order and timing as rats explored

the behavioral chamber, leading to a varied distribution of taste delivery locations, as

exemplified in **Figure 1B** (mean number of trials across sessions, quadrant 1: 46.4 ± 6.57 trials,

quadrant 2: 55.2 ± 14.58 trials, quadrant 3: 64.2 ± 12.16 trials, quadrant 4: 34.1 ± 5.34 trials;

one-way ANOVA, $p = 0.23$). Histology confirmed that the majority of our tetrodes were located

intermediately along the proximodistal axis of dorsal CA1 (**Figure 1C**; Henriksen et al., 2010).

Isolated single neurons were classified as either putative pyramidal cells (91.5%, 441/482) or

interneurons (8.5%, 41/482) on the basis of baseline firing rates and action potential shape

(**Figure 1D**).

366   In total, 395 of the 441 pyramidal neurons were classified as place cells using standard

analysis of the spatial specificity of firing rate responses (see Materials and Methods, Moita et

al., 2003). The spatial firing maps of four representative place cells and interneurons (all of

which were computed with taste delivery periods omitted from the analysis) are shown in the top

row of **Figure 2**. As expected (O'Keefe and Dostrovsky, 1971; O'Keefe and Nadel, 1978), only

the pyramidal cells had place fields (**Figure 2A, 2B**)—interneurons (**Figure 2C**, **2D**) typically

exhibited high spontaneous firing rates regardless of the rat's position.

A cell was considered "taste-responsive" if significant firing rate modulations were

evoked by taste presence, identity, and/or palatability (see **Figure 6** for more details). In total,

96/482 (19.9%) cells were classified as taste-responsive, which is similar to the proportion

reported in the only previous study to assess taste responses in individual hippocampal neurons

(Ho et al., 2011). We found taste-responsive and unresponsive units on tetrodes across the

proximodistal axis of dorsal CA1 (n = 50/60 tetrodes with taste-responsive units). **Table 1** shows

the distribution of taste-responsive cells across animals.

The raster plots and peri-stimulus time histograms (PSTHs) for an example taste-

responsive place cell and interneuron are depicted in **Figure 2B** and **2D**. The place cell in

Figure 2B responded to NaCl from 1550-1750 ms following taste delivery, while the interneuron

in **Figure 2D** responded to NaCl, water and quinine from 1200-2500 ms following taste delivery

(light gray lines). In contrast, the raster plots and PSTHs for non-taste-responsive cells (**Figure**

2A, **2C**) show no differences in evoked activity from baseline or between tastes.

The Z-scored firing rates (**Figure 3A**) reveal the diversity of taste responses within the

population of taste-responsive cells (n = 96 cells). When Z-scored firing rates are ordered by the

timing of each cell's peak excitatory response to saccharin (S, **Figure 3A**), these responses tile

the entirety of the taste delivery period. The breakdown of this sequence for NaCl (N), water

(W), and quinine (Q) when the same cell order is conserved illustrates that taste-responsive

cells respond preferentially and dynamically to different tastes. **Figure 3B** reveals that the

distribution of taste response magnitudes as quantified by eta-squared (η^2, a standard measure

of ANOVA effect size; see Maier et al., 2015) does not follow a normal distribution (number of

histogram bins = 50, bin width = 2e-04; chi-square goodness-of-fit test, χ^2 = 57.5, p = 3.3e-14).

Instead, the majority of neurons (n = 386 cells) were non-taste-responsive and clustered around

low η^2 values, while a smaller proportion of neurons (n = 96 cells) were taste-responsive and

had higher η^2 values (mean η^2, taste-responsive cells: 0.024 ± 1.6e-04, non-taste-responsive

cells: 0.010 ± 1.3e-05).

Since hippocampal activity is affected by animals' location and movement, one possible

explanation of these results is that different tastes have different impacts on animals' motor

behavior, and that therefore any perceived "taste"-evoked responses can simply result from

changes in the animal's speed or position (O'Keefe, 1999; Shan et al., 2016). To control for this

possibility, we assessed differences in rats' pre- and post-taste speed and position, both overall

and between each of the four tastes. We found no differences in the average speed (before

taste delivery: 1.07 ± 0.031 cm/s, after taste delivery: 1.12 ± 0.029 cm/s; one-way ANOVA, p =

0.29) or distance traveled (before taste delivery: 1.57 ± 0.050 cm, after taste delivery: 1.59 ±

0.045 cm; one-way ANOVA, p = 0.74) in the 2.5 seconds preceding and following taste

deliveries; the same was true when trials were split up by taste identity (post-pre taste delivery

speed, saccharin: -0.081 ± 0.063 cm/s, NaCl: 0.037 ± 0.061 cm/s, quinine: 0.11 ± 0.061 cm/s,

water: 0.12 ± 0.070 cm/s; one-way ANOVA, p = 0.10; post-pre taste delivery distance:

saccharin: -0.0026 ± 0.11 cm, NaCl: -0.029 ± 0.11 cm, quinine: 0.093 ± 0.093 cm, water: 0.027

± 0.11 cm; one-way ANOVA, p = 0.86). Therefore, it is unlikely that hippocampal responses to

tastes were caused by changes in animals' position or locomotion; rather, they reflected sensory

responses to some aspect of the taste experience itself.

Taste responses are gated by the spatial firing properties of hippocampal neurons.

We found that 14.7% of place cells (n = 58/395 cells) had significant responses to tastes; a far-

higher percentage of interneurons (75.6%; n = 31/41 cells) were taste-responsive (**Figure 2**).

The significance of this larger likelihood of taste-responsiveness amongst spatially diffuse

interneurons than in spatially-specific place cells (chi-square test, χ^2 = 84.87, p = 3.19e-20)

suggests that taste responsiveness depends on the spatial firing properties of hippocampal

neurons. To further investigate the relationship between place- and taste-specific firing, we

compared the place field size (Brun et al., 2002), spatial information content (Skaggs et al.,

1993), and taste response magnitudes (Maier et al., 2015) of taste-responsive and non-taste-

responsive hippocampal neurons.

Figure 4A depicts the firing fields of 12 example non-taste-responsive and taste-

responsive place cells and interneurons, all of which were computed with taste delivery periods

(extending from 500 ms before to 2500 ms after taste delivery) omitted from the analysis. As

expected, place cells had a smaller firing field size (mean field size, place cells: 525.1 ± 18.1

cm^2, interneurons: 988.7 ± 46.4 cm^2; unpaired t-test, p = 1.4e-14) and higher spatial information

content than interneurons (mean spatial information content, place cells: 1.3 ± 0.034 bits/spike,

interneurons: 0.12 ± 0.034 bits/spike; unpaired t-test, p = 2.6e-25; higher values = smaller, more

concentrated regions of enhanced firing). Therefore, taste responses (which were found

predominately in interneurons) were associated with less spatially selective firing.

This same pattern was found to hold even within each cell type, however: cells with

stronger taste-evoked responses tended to have larger place fields (**Figure 4B**) and exhibit

weaker spatial responses (**Figure 4C**) in analyses restricted to place cells (mean firing field

size, taste-responsive place cells: 626.0 ± 46.1 cm^2, non-taste-responsive place cells: 507.7 ±

19.6 cm^2; unpaired t-test, p = 0.021; mean spatial information content, taste-responsive place

cells: 0.91 ± 0.050 bits/spike, non-taste-responsive place cells: 1.37 ± 0.038 bits/spike; unpaired

t-test, p = 1.03 e-06) as well as interneurons (mean firing field size, taste-responsive

interneurons: 1066.7 ± 49.9 cm^2, non-taste-responsive interneurons: 746.8 ± 69.7 cm^2; unpaired

t-test, p = 0.002; mean spatial information content, taste-responsive interneurons: 0.065 ± 0.040

bits/spike, non-taste-responsive interneurons: 0.29 ± 0.015 bits/spike; unpaired t-test, p =

0.0027). There was a negative correlation between the spatial information content and

magnitude of taste responsiveness (η^2) within each cell type (place cells: Pearson correlation,

R = -0.18, p = 3.27e-04; interneurons: Pearson correlation, R = -0.58, p = 5.95e-05), confirming

that hippocampal neurons that respond strongly to taste delivery tend to have more diffuse firing

in space.

The above analysis implies that while place cells tended to exhibit fewer and lower-

magnitude taste responses than interneurons, a subset of place cells exhibited taste-specific

firing (n = 58/395 cells; example in **Figure 2B**). Thus it is important to ask how place and taste

responses interact when an animal receives familiar tastes in a specific spatial context: can

place cells acquire sensory responses regardless of location, or are responses to tastes gated

by spatial firing, as suggested for other sensory modalities (Moita et al., 2003; Shan et al.,

2016)? To investigate this question, we compared the specificity of taste responses inside and

outside each place cell's firing field.

Only place cells (n = 58 taste-responsive cells, n = 337 non-taste-responsive cells) were

considered for in-field vs. out-of-field analysis. We defined a cell's place field as the largest

number of adjacent spatial bins which had firing rates >= 20% of the peak firing rate (Brun et al.,

2002). To ensure sufficient sampling of taste responses for statistical comparisons, only cells

that contained at least ten in-field and out-of-field trials of each taste stimulus were included in

our analysis (n = 26 taste-responsive cells, n = 153 non-taste-responsive cells). Taste response

magnitude (η^2) was then determined separately for trials taking place within and outside each

cell's place field.

The top row of **Figure 5** shows the spatial firing maps of representative taste-responsive

(**Figure 5A**) and non-taste-responsive (**Figure 5B**) place cells, with in-field and out-of-field trials

indicated in pink. The distribution of taste trials in these cells was representative of the entire

population analyzed, with comparable numbers of taste trials delivered in-field (number of in-

field trials, taste-responsive cells, saccharin (S): 25.1 ± 1.5 trials, NaCl (N): 25.5 ± 1.5 trials,

water (W): 24.4 ± 1.4 trials, quinine (Q): 24.8 ± 1.5 trials; one-way ANOVA, p = 0.96; number of

in-field trials, non-taste-responsive cells, saccharin: 23.4 ± 0.7 trials, NaCl: 24.1 ± 0.7 trials,

water: 23.3 ± 0.7 trials, quinine: 23.8 ± 0.7 trials; one-way ANOVA, p = 0.89). The total number

of in-field trials was also comparable across groups (number of in-field trials, taste-responsive

cells: 99.7 ± 5.7 trials, non-taste-responsive cells: 94.6 ± 2.7 trials; unpaired t-test, p = 0.46).

Repeating our initial analyses of taste responsiveness using only the in-field trials of units that

met our analysis criteria (n = 179 cells) resulted in comparable numbers of taste-responsive

cells (n = 17 cells using in-field trials, n = 26 cells using all trials; chi-square test, χ^2 = 1.69, p =

0.19), indicating that the total number of taste-responsive neurons was not underestimated due

to inter-trial variability, and cells previously classified as non-taste-responsive did not typically

contain any regions of taste-responsiveness, either in- or out-of-field (see example in **Figure**

5B).

For the taste-responsive cell (**Figure 5A**), virtually all responses occurred within the

cell's place field (left panel), with very little taste-evoked firing out of field (right panel). On the

other hand, no taste-evoked responses were observed in- or out-of-field for the non-taste-

responsive cell (**Figure 5B**). This trend was representative of the entire population of place cells

(**Figure 5C**): the in-field region of taste-responsive cells had a higher mean eta-squared value

than the out-of-field region, or either region of non-taste-responsive cells (mean eta-squared

values, taste-responsive cells, in-field: 0.036 ± 0.0042; taste-responsive cells, out-of-field: 0.023

± 0.0027; non-taste-responsive cells, in-field: 0.025 ± 0.0013; non-taste-responsive cells, out-of-

field: 0.0208 ± 0.0011; 1-way ANOVA, p = 3e-05). Together, these results indicate that

hippocampal taste responses are gated by the spatial firing properties of place cells—a finding

that is consistent with previous studies investigating tone-evoked sensory responses during

auditory fear conditioning (Moita et al., 2003; Shan et al., 2016).

Hippocampal taste-specific responses reflect taste palatability, at a relatively long delay.

Previous work has shown that taste-specific firing in GC neurons evolves through three stages:

following an initial, nonspecific response to taste presence, a discriminative response conveys

information about taste identity starting at approximately 200 ms after stimulus administration;

after approximately 500 ms, responses then change to reflect palatability, specifically

anticipating an animal's decision to consume or expel a particular taste between 600 and 1600

ms after taste delivery (Katz et al., 2001; Piette et al., 2012; Sadacca et al., 2012; Maier and

Katz, 2013; Li et al., 2016; Sadacca et al., 2016). Brainstem taste responses in the parabrachial

nucleus (PbN) are similarly organized (Baez-Santiago et al., 2016). Other nodes of the taste

CNS, however, such as the BLA and LH, appear instead to respond primarily and immediately

to the hedonic value of tastes, regardless of identity (Nishijo et al., 1998; Fontanini et al., 2009;

Li et al., 2013). To determine which components are present in hippocampal taste responses

(and when), we performed analyses similar to those brought to bear on firing in these other

structures.

Many hippocampal neurons responded nonspecifically to taste presence, providing

information that could allow for the detection of tastants on the tongue (**Figure 6A**). In a subset

of these cells, responses were more discriminative, providing information about taste identity

and/or palatability. Two such neurons are shown in **Figure 6B**—one of which (place cell, left)

rapidly developed a response primarily to quinine, and one of which (interneuron, right)

produced a longer-latency response that differentiated each of the four tastes, and which

notably involved a sudden change of firing rate to saccharin. As illustrated previously (**Figure**

3A), we observed a relatively uniform distribution among taste-discriminative cells (n = 40) of

which taste(s) cells responded to (number of cells with significant evoked responses to

saccharin: 20 cells, NaCl: 29 cells, water: 26 cells, quinine: 25 cells; chi-square goodness-of-fit

test, χ^2 = 1.7, p = 0.64). Similar to the interneuron shown in **Figure 6B**, most taste-discriminative

cells responded to more than one taste (number of cells responding to one taste: 11 cells, two

tastes: 4 cells, three tastes: 19 cells, four tastes: 6 cells; chi-square goodness-of-fit test, χ^2 =

13.4, p = 0.0038).

536   Closer examination revealed that the patterning of both responses in **Figure 6B**

reflected taste palatability across the entirety of the periods of taste-specific firing (**Figure 6D**).

Responses to taste palatability were assessed, as is typical in studies of taste temporal coding

(Li et al., 2013; Baez-Santiago et al., 2016; Sadacca et al., 2016), in terms of the correlation

between neuronal firing rates and the order of taste preference, which was assayed in a brief-

access task (Li et al., 2013; Monk et al., 2014; Sadacca et al., 2016) run on a separate cohort of

experimental rats (**Figure 6C**). The observed order of taste preference (S > N > W > Q) shown

in **Figure 6C** is consistent with that observed across a broad range of stimulus delivery methods

and assessment techniques (Travers and Norgren, 1986; Breslin et al., 1992; Clarke and

Ossenkopp, 1998; Fontanini and Katz, 2006; Sadacca et al., 2016).

547   **Figure 6D** reveals that palatability correlations for the example neurons shown in **Figure**

6B developed as the taste-specific responses themselves developed: the place cell's responses

(**Figure 6B**, left) were significantly correlated with taste palatability between 600 and 2100 ms

(**Figure 6D**, left), while the interneuron's responses (**Figure 6B**, right) were palatability-related

between 2000 and 2500 ms (**Figure 6D**, right; compare these periods with the dark gray line in

Figure 6B, which marks the period of significantly taste-specific firing).

Again, the examples shown in **Figure 6** suggest that, like responses observed in limbic

structures (i.e., BLA and LH; Fontanini et al., 2009; Li et al., 2013) but unlike those in the main

taste axis (i.e., GC and PbN; Sadacca et al., 2012; Baez-Santiago et al., 2016), hippocampal

taste responses do not go through a period of "pure" taste specificity prior to becoming

palatability-related. These appearances were borne out in an analysis of the entire neural

dataset. **Figure 7A** shows, similar to what has been observed in all other parts of the taste

system (Katz et al., 2001; Sadacca et al., 2012; Baez-Santiago et al., 2016; Fontanini et al;

2009; Li et al., 2013), that totally non-specific responses to taste presence emerged first in

hippocampus, followed by responses to taste identity and palatability. However, both taste

specificity and palatability-relatedness appeared in hippocampal taste responses at similarly

long latencies (average onset, presence: 1032.5 ± 55.73 ms; identity: 1443.1 ± 108.07 ms;

palatability: 1797.2 ± 118.51 ms).

Direct within-neuron comparisons strongly supported the group analysis. Presence-

related responses reliably arose before identity-related responses in cells that responded to

both properties (n = 19 cells, paired t-test, p = 7.6e-05), as also evidenced by the cloud of points

above the unity line (**Figure 7B**; regression slope: 0.079, p = 0.79). A plot of the onset latency of

identity- and palatability-related responses in cells where both properties were present,

meanwhile, revealed tight clustering around the unity line (**Figure 7C**; n = 14 cells; regression

slope: 0.92, p = 8.7e-06) with no significant differences between onset times (paired t-test, p =

0.83), suggesting that these properties arose simultaneously in single-unit responses.

Finally, we performed PCA by pooling responses of identity-responsive cells (n = 36

cells) to visualize the onset of taste-specific firing in the population as a whole. This analysis

suggests that tastes are discriminated based on palatability, as shown by significant encoding of

palatability rank (here, in reverse order as shown in **Figure 6C**) by the principal components

starting at ~1.4 s after stimulus delivery (**Figure 7D**). It is important to note that this analysis is

not representative of every cell: we identified a small but significant contingent of exclusively

identity-coding neurons (n = 15/96 taste-responsive cells; see **Figure 6A**, and **Figure 2B** for

example); it's worth noting, however, that even many of the neurons that responded strongly

and specifically to single tastes did so to either the most rewarding or aversive tastants in our

battery (see **Figure 6B** and **D** for examples).

We therefore conclude that hippocampal "taste codes" do not contain the purely identity-

related component found in gustatory brainstem and cortex; rather, taste selectivity emerges at

the same time as palatability-relatedness, and as a whole, the taste-reactive hippocampal

population delivers information on palatability without a prior epoch of less structured taste-

specificity. In this regard, hippocampal responses are similar to those observed in other non-

cortical parts of the taste system, such as the BLA (Fontanini et al., 2009) and LH (Li et al.,

2013); notably, however, palatability coding appears in hippocampus much later than it appears

in these other limbic structures—a difference that likely has strong implications for the potential

roles of the hippocampus in taste (see Discussion below).

Discussion

Our findings suggest that place and taste responses can co-exist within the same hippocampal

neurons, and that these response modalities influence one another. Taste-responsive cells

tended to have less spatially specific firing fields (**Figures 2**, **4**). On the other hand, place cells

that responded to tastes did so in a spatially specific manner (**Figure 5**), with responses only

occurring within that cell's place field. Hippocampal neurons discriminated between tastes at

relatively long latencies and predominantly on the basis of palatability (**Figures 6**, **7**), confirming

that these responses can encode sensory parameters, and do not simply reflect changes in

animals' movement or attentive state, as has been suggested (O'Keefe, 1999; Shan et al.,

2016). These results stand alongside similar studies of audition (Moita et al., 2003) and olfaction

(Wood et al., 1999), and further show that spatially tuned hippocampal neurons can discriminate

between behaviorally relevant sensory stimuli. Our observations add to an expanding view of

the hippocampal cognitive map as a representation that encompasses both spatial and non-

spatial aspects of an animal's environment (Shapiro et al., 1997; Eichenbaum et al., 1999;

Wood et al., 1999; Moita et al., 2003; Kraus et al., 2013; Aronov et al., 2017).

In total, about 20% of recorded hippocampal cells in our study were classified as taste-

responsive (**Figure 2**), which is comparable to the proportion reported in the only previous study

to assess taste responses in individual hippocampal neurons (Ho et al., 2011). Our usage of

IOCs for enhanced stimulus control allowed us to build upon Ho and colleagues' findings,

uncovering new understanding of the structure, content, and dynamics of hippocampal taste

responses. Similar to what is seen elsewhere in the taste system, including the GC (Katz et al.,

2001), BLA (Fontanini et al., 2009), LH (Li et al., 2013), and PbN (Baez-Santiago et al., 2016),

we observed larger-than-chance numbers of CA1 neurons that responded discriminatively to

taste stimuli (**Figure 6A**), with response specificity evolving over time (**Figure 7**). This taste-

distinctiveness, along with the history of taste research that has been conducted in rodents

using IOCs, further underscores that hippocampal taste responses are truly gustatory, and do

not arise from stress or fear responses caused by the delivery of solutions via IOC (Travers and

Norgren, 1986; Katz et al., 2001; Sadacca et al., 2016).

Unlike what has been observed in GC (Katz et al., 2001; Sadacca et al., 2016) and PbN

(Baez-Santiago et al., 2016), we found little evidence of pure identity coding in the hippocampus

in this passive administration paradigm. Instead, hippocampal neurons distinguish between

tastes based on palatability (**Figure 7**), similar to other taste system limbic structures such as

the BLA (Fontanini et al., 2009) and LH (Li et al., 2013). However, palatability-related

hippocampal coding emerged much later than in BLA or LH, and likely after the time (although

more direct measurements must be taken to ascertain this) that animals make decisions about

palatability-related orofacial behaviors (Li et al., 2016; Sadacca et al., 2016). These results

support the idea that the hippocampus does not contribute to an animal's decision to consume

or expel a given taste. Rather, it responds to the hedonic value of tastes consumed within a

particular context, forming associations between place and reward that may be relayed

downstream via hippocampal projections to the ventral tegmental area and other neural reward

centers (Lisman et al., 2011). This could serve as a means of relating place to taste in terms of

its inherent reward value, allowing animals to use past experience to locate food sources.

While spatial learning is indisputably considered to be a hippocampal-dependent

process (Morris, 1984; Moser et al., 2008), the role of the hippocampus in non-spatial taste

learning is less clear-cut. Some forms of taste learning, such as social transmission of food

preferences (Bunsey and Eichenbaum, 1995; Countryman et al., 2005) are considered to be

hippocampal-dependent; other paradigms (e.g., conditioned taste aversion) undergo variable

effects after hippocampal lesions (Yamamoto et al., 1995; Stone et al., 2005; Chinnakkaruppan

et al., 2014). Future studies in which individual neurons are recorded during taste learning,

which have been informative when focused on other nodes of the taste system (Grossman et

al., 2008; Lavi et al., 2018), may help to decipher how the hippocampus encodes tastes and

contexts to guide future food choices.

Our study provides the first direct evidence that hippocampal taste responses are almost

entirely gated by the neurons' spatial firing properties. More specifically, taste responses are

more prevalent in interneurons than place cells (**Figure 2**), associated with broader spatial

responsiveness regardless of cell type (**Figure 4**), and limited to cells' spatial firing fields

(**Figure 5**). Our finding that place cells only respond to tastes delivered within their place field is

consistent with previous studies (Moita et al., 2003; Shan et al., 2016), indicating that taste

responses can best be understood as a rate code overlaid on existing representations of space.

Since place fields can be modulated by food reward (Dupret et al., 2010; Allen et al., 2012), it

seems likely that taste responses could arise as a consequence of place cell remapping.

However, we could not address this question in the current study, since rats did not explore the

behavioral chamber in the absence of tastes, making it difficult to calculate place fields. We

specifically chose a smaller open field environment for these experiments because it allowed

adequate coverage of place fields as well as multiple taste stimuli, allowing us to study the real-

time interaction between taste and place. Future studies that incorporate place-specific taste

delivery in a larger or more structured environment, such as a linear track, will be able to

explore whether taste experience can modify animals' hippocampal representation of

environments through rate or global remapping, as has been shown for other sensory modalities

(Moita et al., 2004; Fyhn et al., 2007; Zhang and Manahan-Vaughan, 2015).

Whatever the relationship between spatial and gustatory firing, more hippocampal

interneurons—by their very nature, non-place cells—responded to non-spatial stimuli than place

cells (**Figure 2**). This result is consistent with studies that measured hippocampal responses to

auditory (Moita et al., 2003) or olfactory (Deshmukh and Bhalla, 2003) stimuli. This finding may

reflect the intrinsic properties of each cell type: place cells have lower mean firing rates and

rarely respond outside of their place field, while interneurons exhibit high firing rates regardless

of location (see **Figure 2**), making it much easier to obtain statistical significance in the latter.

Another (though not mutually exclusive) possibility is that interneurons could be influencing

place cell activity. While place cells' responses to space have been well-characterized (O'Keefe

and Nadel, 1978; Moser et al., 2008), recent work suggests that interneurons also contribute to

hippocampal representations of space (Wilent and Nitz, 2007), and disinhibit place cell firing

through location-specific decreases in activity (Hangya et al., 2010; Royer et al., 2012). Since

interneurons' projections are primarily internal to the hippocampal network (Freund and Buzsáki,

1996; Kullman, 2011), these responses may play a role in shaping activity within the network,

such as place fields (**Figure 4**) or location-specific responses to tastes (**Figure 5**). However,

how non-spatial information is transmitted within hippocampal microcircuits remains an open

question, one that may be investigated in future studies by determining the effect of cell-type-

specific inhibition on hippocampal taste responses.

There is strong evidence that the behavioral relevance of sensory stimuli within a task

influences what proportion of hippocampal neurons respond to non-spatial cues. In our study,

rats passively received tastes via IOC, a paradigm that requires no learning, other than

associating tastes with a context for the first time. The total proportion of taste-responsive

neurons in our study (~20%, **Figure 2**) is comparable to the proportion of tone-responsive cells

(16%) found by one study analyzing the auditory evoked responses of hippocampal neurons

(Moita et al., 2003); however, this proportion increased to 52% following auditory fear

conditioning. Similarly, ~40% of hippocampal neurons exhibited non-spatial firing during an

odor-guided non-match-to-sample task that required working memory of spatial and non-spatial

factors (Wood et al., 1999). More pronounced changes in hippocampal responsiveness are

observed when reward contingencies are entirely dependent on discriminating between non-

spatial cues, such as in one study where rats learned to manipulate a joystick to modulate a

tone within a target frequency range (Aronov et al., 2017). During this task, about 40% of

hippocampal neurons responded to specific tone frequencies, compared to only 2% during the

passive playback of tones. In our study, 8.3% (40/482) of CA1 neurons discriminated between

tastes (**Figure 6A**), with an additional 11.6% (56/482) of cells responding nonspecifically to

taste presence. These results suggest that the hippocampus forms a flexible map of spatial and

non-spatial stimuli based on current behavioral demands. For the case of taste, gustatory

responses in spatially tuned hippocampal neurons may allow animals to form value-related

associations between tastes and contexts, aiding in the finding of food. Future work will assess

how taste experience affects this ongoing mental map.

References

Accolla R, Carleton A (2008) Internal body state influences topographical plasticity of sensory

representations in the rat gustatory cortex. Proc Natl Acad Sci 105: 4010-4015.

Agnihotri NT, Hawkins RD, Kandel ER, Kentros C (2004). The long-term stability of new

hippocampal place fields requires new protein synthesis. Proc Natl Acad Sci 101(10):

3656-3661.

Allen K, Rawlins JN, Bannerman DM, Csicsvari J (2012) Hippocampal place cells can encode

multiple trial-dependent features through rate remapping. J Neurosci 32:14752-14766.

Aronov D, Nevers R, Tank DW (2017) Mapping of a non-spatial dimension by the hippocampal-

entorhinal circuit. Nature 543: 719-722.

Baez-Santiago MA, Reid EE, Moran A, Maier JX, Marrero-Garcia Y, Katz DB (2016) Dynamic

taste responses of parabrachial pontine neurons in awake rats. J Neurophysiol 115:

1314-1323.

Breslin PA, Spector AC, Grill HJ (1992) A quantitative comparison of taste reactivity behaviors

to sucrose before and after lithium chloride pairings: a unidimensional account of

palatability. Behav Neurosci 106: 820-836.

Briggman KL, Abarbanel HD, Kristan WB, Jr. (2005) Optical imaging of neuronal populations

during decision-making. Science 307: 896-901.

Brun VH, Otnass MK, Molden S, Steffenach HA, Witter MP, Moser MB, Moser EI (2002) Place

cells and place recognition maintained by direct entorhinal-hippocampal circuitry.

Science 296: 2243-2246.

Bunsey M, Eichenbaum H (1995) Selective damage to the hippocampal region blocks long-term

retention of a natural and nonspatial stimulus-stimulus association. Hippocampus 5: 546-

556.

Chen X, Gabitto M, Peng Y, Ryba NJ, Zuker CS (2011) A gustotopic map of taste qualities in

the mammalian brain. Science 333: 1262-1266.

Chinnakkaruppan A, Wintzer ME, McHugh TJ, Rosenblum K (2014) Differential contribution of

hippocampal subfields to components of associative taste learning. J Neurosci 34:

11007-11015.

Clarke SN, Ossenkopp KP (1998) Taste reactivity responses in rats: influence of sex and the

estrous cycle. Am J Physiol 274: R718-724.

Countryman RA, Kaban NL, Colombo PJ (2005) Hippocampal c-fos is necessary for long-term

memory of a socially transmitted food preference. Neurobiol Learn Mem 84: 175-183.

Deshmukh SS, Bhalla US (2003) Representation of odor habituation and timing in the

hippocampus. J Neurosci 23: 1903-1915.

Dupret D, O'Neill J, Pleydell-Bouverie B, Csicsvari J (2010) The reorganization and reactivation

of hippocampal maps predict spatial memory performance. Nat Neurosci 13: 995-1002.

Eichenbaum H, Dudchenko P, Wood E, Shapiro M, Tanila H (1999) The hippocampus, memory,

and place cells: is it spatial memory or a memory space? Neuron 23: 209-226.

Fontanini A, Katz DB (2005) 7 to 12 Hz activity in rat gustatory cortex reflects disengagement

from a fluid self-administration task. J Neurophysiol 93: 2832-2840.

Fontanini A, Katz DB (2006) State-dependent modulation of time-varying gustatory responses. J

Neurophysiol 96: 3183-3193.

Fontanini A, Grossman SE, Figueroa JA, Katz DB (2009) Distinct subtypes of basolateral

amygdala taste neurons reflect palatability and reward. J Neurosci 29: 2486-2495.

Fox SE, Ranck JB (1981) Electrophysiological characteristics of hippocampal complex-spike

cells and theta cells. Exp Brain Res 41: 399-410.

Frank LM, Brown EN (2003) Persistent activity and memory in the entorhinal cortex. Trends

Neurosci 26: 400-401.

Fried I, MacDonald KA, Wilson CL (1997) Single neuron activity in human hippocampus and

amygdala during recognition of faces and objects. Neuron 18: 753-765.

Freund TF, Buzsáki G (1996) Interneurons of the hippocampus. Hippocampus 6: 347-470.

Fyhn M, Hafting T, Treves A, Moser MB, Moser EI (2007) Hippocampal remapping and grid

realignment in entorhinal cortex. Nature 446: 190-194.

Geran LC, Travers SP (2009) Bitter-responsive gustatory neurons in the rat parabrachial

nucleus. J Neurophysiol 101: 1598-1612.

Grill HJ, Norgren R (1978) The taste reactivity test. I. Mimetic responses to gustatory stimuli in

neurologically normal rats. Brain Res 143: 263-279.

Grossman SE, Fontanini A, Wieskopf JS, Katz DB (2008) Learning-related plasticity of temporal

coding in simultaneously recorded amygdala-cortical ensembles. J Neurosci 28: 2864-

2873.

Haase L, Cerf-Ducastel B, Murphy C (2009) Cortical activation in response to pure taste stimuli

during the physiological states of hunger and satiety. Neuroimage 44: 1008-1021.

Hangya B, Li Y, Muller RU, Czurko A (2010) Complementary spatial firing in place cell-

interneuron pairs. J Physiol 588: 4165-4175.

Harvey CD, Coen P, Tank DW (2012) Choice-specific sequences in parietal cortex during a

virtual-navigation decision task. Nature 484: 62-68.

Henriksen EJ, Colgin LL, Barnes CA, Witter MP, Moser MB, Moser EI (2010) Spatial

representation along the proximodistal axis of CA1. Neuron 68: 127-137.

Ho AS, Hori E, Nguyen PH, Urakawa S, Kondoh T, Torii K, Ono T, Nishijo H (2011)

Hippocampal neuronal responses during signaled licking of gustatory stimuli in different

contexts. Hippocampus 21: 502-519.

Jadhav SP, Kemere C, German PW, Frank LM (2012) Awake hippocampal sharp-wave ripples

support spatial memory. Science 336: 1454-1458.

Jadhav SP, Rothschild G, Roumis DK, Frank LM (2016) Coordinated Excitation and Inhibition of

Prefrontal Ensembles during Awake Hippocampal Sharp-Wave Ripple Events. Neuron

90: 113-127.

Katz DB, Simon SA, Nicolelis MA (2001) Dynamic and multimodal responses of gustatory

cortical neurons in awake rats. J Neurosci 21: 4478-4489.

Kobayakawa T, Wakita M, Saito S, Gotow N, Sakai N, Ogawa H (2005) Location of the primary

gustatory area in humans and its properties, studied by magnetoencephalography.

Chem Senses 30 Suppl 1: i226-227.

Kraus BJ, Robinson RJ, 2nd, White JA, Eichenbaum H, Hasselmo ME (2013) Hippocampal

"time cells": time versus path integration. Neuron 78: 1090-1101.

Kullman DM (2011) Interneuron networks in the hippocampus. Curr Opin Neurobiol 21(5): 709-

716.

Lavi K, Jacobson GA, Rosenblum K, Luthi A (2018) Encoding of Conditioned Taste Aversion in

Cortico-Amygdala Circuits. Cell Reports 24: 278-283.

Leutgeb S, Leutgeb JK, Treves A, Moser MB, Moser EI (2004) Distinct ensemble codes in

hippocampal areas CA3 and CA1. Science 305: 1295-1298.

Leutgeb S, Leutgeb JK, Barnes CA, Moser EI, McNaughton BL, Moser M-B (2005) Independent

Codes for Spatial and Episodic Memory in Hippocampal Neuronal Ensembles. Science

309: 619-623.

Li J, Lemon CH (2015) Influence of stimulus and oral adaptation temperature on gustatory

responses in central taste-sensitive neurons. J Neurophysiol 113(7): 2700-2712.

Li JX, Yoshida T, Monk KJ, Katz DB (2013) Lateral hypothalamus contains two types of

palatability-related taste responses with distinct dynamics. J Neurosci 33: 9462-9473.

Li JX, Maier JX, Reid EE, Katz DB (2016) Sensory Cortical Activity Is Related to the Selection of

a Rhythmic Motor Action Pattern. J Neurosci 36: 5596-5607.

Lisman J, Grace AA, Duzel E (2011) A neoHebbian framework for episodic memory; role of

dopamine-dependent late LTP. Trends Neurosci 34(10): 536-547.

Maier JX, Katz DB (2013) Neural dynamics in response to binary taste mixtures. J Neurophysiol

109: 2108-2117.

Maier JX, Blankenship ML, Li JX, Katz DB (2015) A Multisensory Network for Olfactory

Processing. Curr Biol 25: 2642-2650.

Moita MA, Rosis S, Zhou Y, LeDoux JE, Blair HT (2003) Hippocampal place cells acquire

location-specific responses to the conditioned stimulus during auditory fear conditioning.

Neuron 37: 485-497.

Moita MA, Rosis S, Zhou Y, LeDoux JE, Blair HT (2004) Putting fear in its place: remapping of

hippocampal place cells during fear conditioning. J Neurosci 24: 7015-7023.

Monk KJ, Rubin BD, Keene JC, Katz DB (2014) Licking microstructure reveals rapid attenuation

of neophobia. Chem Senses 39: 203-213.

Moran A, Katz DB (2014) Sensory cortical population dynamics uniquely track behavior across

learning and extinction. J Neurosci 34: 1248-1257.

Morris R (1984) Developments of a water-maze procedure for studying spatial learning in the

rat. J Neurosci Methods 11: 47-60.

Moser EI, Kropff E, Moser MB (2008) Place Cells, Grid Cells, and the Brain's Spatial

Representation System. Annu Rev Neurosci 31: 69-89.

Nishijo H, Uwano T, Tamura R, Ono T (1998) Gustatory and multimodal neuronal responses in

the amygdala during licking and discrimination of sensory stimuli in awake rats. J

Neurophysiol 79: 21-36.

O'Keefe J (1999) Do hippocampal pyramidal cells signal non-spatial as well as spatial

information? Hippocampus 9: 352-364.

O'Keefe J, Dostrovsky J (1971) The hippocampus as a spatial map. Preliminary evidence from

unit activity in the freely-moving rat. Brain Res 34: 171-175.

O'Keefe J, Nadel L (1978) The hippocampus as a cognitive map. London: Oxford University

Press.

Piette CE, Baez-Santiago MA, Reid EE, Katz DB, Moran A (2012) Inactivation of basolateral

amygdala specifically eliminates palatability-related information in cortical sensory

responses. J Neurosci 32: 9981-9991.

Reilly S, Harley C, Revusky S (1993) Ibotenate lesions of the hippocampus enhance latent

inhibition in conditioned taste aversion and increase resistance to extinction in

conditioned taste preference. Behav Neurosci 107: 996-1004.

Rosen AM, Roussin AT, Di Lorenzo PM (2010) Water as an independent taste modality. Front

Neurosci 15(4): 175.

Roux L, Hu B, Eichler R, Stark E, Buzsáki G (2017) Sharp wave ripples during learning stabilize

the hippocampal spatial map. Nat Neurosci 261: 1055.

Sadacca BF, Katz DB (2011) Taste. In: Neurobiology of Sensation and Reward (Gottfriend JA,

ed), pp 127-141. Boca Raton, FL: CRC Press/Taylor & Francis.

Sadacca BF, Rothwax JT, Katz DB (2012) Sodium concentration coding gives way to evaluative

coding in cortex and amygdala. J Neurosci 32: 9999-10011.

Sadacca BF, Mukherjee N, Vladusich T, Li JX, Katz DB, Miller P (2016) The Behavioral

Relevance of Cortical Neural Ensemble Responses Emerges Suddenly. J Neurosci 36:

655-669.

Shan KQ, Lubenov EV, Papadopoulou M, Siapas AG (2016) Spatial tuning and brain state

account for dorsal hippocampal CA1 activity in a non-spatial learning task. eLife 5.

Shapiro ML, Tanila H, Eichenbaum H (1997) Cues that hippocampal place cells encode:

dynamic and hierarchical representation of local and distal stimuli. Hippocampus 7: 624-

642.

Skaggs WE, McNaughton BL, Gothard K, Markus E (1993) An information theoretic approach to

deciphering the hippocampal code. In: Advanced in Neural Information Processing

Systems (Hanson S, Cowan JD, Giles CL, eds), pp 1030-1037. San Mateo, CA: Morgan

Kaufmann Publishers.

Spetter MS, Smeets PA, de Graaf C, Viergever MA (2010) Representation of sweet and salty

866   taste intensity in the brain. Chem Senses 35: 831-840.

867   Stone ME, Grimes BS, Katz DB (2005) Hippocampal inactivation enhances taste learning.

868   Learn Mem 12: 579-586.

869   Suzuki WA, Amaral DG (1994) Perirhinal and parahippocampal cortices of the macaque

870   monkey: cortical afferents. J Comp Neurol 350: 497-533.

871   Tang W, Shin JD, Frank LM, Jadhav SP (2017) Hippocampal-prefrontal reactivation during

872   learning is stronger in awake compared with sleep states. J Neurosci 37: 11789-11805.

873   Thompson LT, Best PJ (1990) Long-term stability of the place-field activity of single units

874   recorded from the dorsal hippocampus of freely behaving rats. Brain Res 509(2): 299-

875   308.

876   Travers JB, Norgren R (1986) Electromyographic analysis of the ingestion and rejection of sapid

877   stimuli in the rat. Behav Neurosci 100: 544-555.

878   Tsao A, Moser MB, Moser EI (2013) Traces of experience in the lateral entorhinal cortex. Curr

879   Biol 23: 399-405.

880   van de Ven GM, Trouche S, McNamara CG, Allen K, Dupret D (2016) Hippocampal offline

881   reactivation consolidates recently formed cell assembly patterns during sharp wave-

882   ripples. Neuron 92: 968-974.

883   von Bohlen und Halbach O, Albrecht D (2002) Reciprocal connections of the hippocampal area

884   CA1, the lateral nucleus of the amygdala and cortical areas in a combined horizontal

885   slice preparation. J Neurosci Res 44: 91-100.

886   Wilent WB, Nitz DA (2007) Discrete place fields of hippocampal formation interneurons. J

887   Neurophysiol 97: 4152-4161.

888   Wood ER, Dudchenko PA, Eichenbaum H (1999) The global record of memory in hippocampal

889   neuronal activity. Nature 397: 613-616.

Yamamoto T, Fujimoto Y, Shimura T, Sakai N (1995) Conditioned taste aversion in rats with

excitotoxic brain lesions. Neurosci Res 22: 31-49.

Zald DH, Lee JT, Fluegel KW, Pardo JV (1998) Aversive gustatory stimulation activates limbic

circuits in humans. Brain 121 (Pt 6): 1143-1154.

Zhang S, Manahan-Vaughan D (2015) Spatial olfactory learning contributes to place field

formation in the hippocampus. Cereb Cortex 25: 423-432.

Legends

Figure 1. Experimental design and electrophysiology.

A, A portion of the timeline of an example taste delivery experiment. Colored bars indicate

individual deliveries of taste stimuli: green (S, 4 mM saccharin), yellow (N, 100 mM sodium

chloride), blue (W, distilled water) and red (Q, 5 mM quinine hydrochloride). Taste deliveries

occurred at a randomized timing of 13-17 s, with the taste identity randomized for each trial. *B*,

Example session showing all 200 taste delivery locations (colored symbols) overlaid on top of

the rat's position in the behavioral chamber (gray circles) during one recording experiment. *C*,

Histological verification of tetrode locations in intermediate dorsal CA1. Dotted lines indicate the

extent of recording sites across all five animals. *D*, Classification of putative interneurons (Int,

gray crosses) from pyramidal cells (Pyr, black circles) based on spike width (> 8.5 Hz) and firing

rate (< 0.35 ms) parameters. The top panel depicts average waveform shapes for putative

interneurons (gray) and pyramidal cells (black).

Table 1. Cell distribution across animals.

Summary of the number of taste-responsive and total CA1 cells recorded from each animal.

Only the cells meeting the inclusion criteria (see Materials and Methods) are reported. Putative

pyramidal cells (Pyr) and interneurons (Int) were identified on the basis of firing rate and spike

width parameters. Neurons were classified as "taste-responsive" if they exhibited responses to

taste presence, identity and/or palatability.

Figure 2. Subsets of hippocampal place cells and interneurons respond to tastes.

A-D, Top panels, Example spatial firing maps of two place cells (left) and interneurons (right),

calculated with taste delivery periods (500 ms before to 2500 ms after taste delivery) omitted

from the analysis. Numbers on the bottom right of each plot denote peak spatial firing rate (FR)

in Hz. Middle panels, Raster plots of the cells in **A-D** responding to each of the four tastes, with

trials aligned to the time of taste delivery (S, green line, 4 mM saccharin; N, yellow line, 100 mM

sodium chloride; W, blue line, distilled water; Q, red line, 5 mM quinine hydrochloride) and black

dots indicating when spikes occurred during each trial. Light gray boxes indicate the taste(s) for

which there was a significant evoked response (*p < 0.05, t-tests on successive time windows).

Bottom panels, Taste-evoked responses of each of the above neurons. Each colored trace

represents the mean firing rate to one of the four tastes, smoothed with a 1D Gaussian filter

(σ = 5 ms). Light gray lines indicate the periods of significant taste responsiveness (*p < 0.05, t-

tests on successive time windows) for the place cell in **B** and interneuron in **D**.

Figure 3. Diversity among the population of taste-responsive cells.

A, Z-scored firing rates of the 96 taste-responsive cells, ordered by the timing of their peak

response to saccharin (S). Note the dissimilarity of responses when Z-scored firing rates to

NaCl (N), water (W), or quinine (Q) are depicted using the same order, indicating that different

cells respond preferentially and dynamically to each of the four administered tastes. **B**,

Histogram depicting the non-normal distribution of taste response magnitudes (quantified using

effect size, η^2) within the entire population of neurons (n = 482 cells; chi-square goodness-of-fit

test, χ^2 = 57.5, ***p = 3.3e-14), as well as the average η^2 value for taste-responsive (n = 386

cells; mean η^2 = 0.024 ± 1.6e-04) and non-taste-responsive cells (n = 96 cells; mean η^2 =

0.010 ± 1.3e-05) within this distribution.

Figure 4. Taste-responsive hippocampal neurons exhibit weaker spatial selectivity than non-

taste-responsive hippocampal neurons.

A, Example spatial firing maps of twelve non-taste-responsive (left) and taste-responsive (right)

place cells (Place, top row) and interneurons (Int, bottom row). Note that taste-responsive cells

tend to exhibit more diffuse spatial firing. All firing maps were computed with taste delivery

periods (500 ms before to 2500 ms after taste delivery) omitted from the analysis. Numbers on

the bottom right of each plot denote peak spatial firing rate (FR) in Hz. **B**, Mean firing field size

for non-taste-response (white bars) and taste-responsive (gray bars) place cells and

interneurons. For each cell type, taste-responsive neurons had larger firing fields than non-

taste-responsive neurons (place cells: n = 337 non-taste-responsive cells, n = 58 taste-

responsive cells; unpaired t-test, *p = 0.021; interneurons: n = 10 non-taste-responsive cells, n

= 31 taste-responsive cells; unpaired t-test, **p = 0.002). **C**, Mean spatial information content for

non-taste-responsive (white bars) and taste-responsive (gray bars) place cells and interneurons.

Within each cell type, taste-responsive neurons had a lower spatial information content than

non-taste-responsive neurons (place cells, unpaired t-test, ***p = 1.03e-06; interneurons,

unpaired t-test, **p = 0.0027).

Figure 5. Place cells respond to tastes delivered within their place field.

A-B, Example in-field and out-of-field responses for a taste-responsive (**A**) and non-taste-

responsive (**B**) place cell. Top panels show the spatial firing maps for each cell, with place field

boundaries (defined as largest cluster of neighboring spatial bins which had firing rates >= 20%

of the peak rate) indicated by black lines. The colored symbols represent locations of individual

taste deliveries (green: S, 4 mM saccharin; yellow: N, 100 mM sodium chloride; blue: W,

distilled water; red: Q, 5 mM quinine hydrochloride), with either in-field (left) or out-of-field (right)

trials highlighted in pink. All firing maps were computed with taste delivery periods (500 ms

before to 2500 ms after taste delivery) omitted from the analysis. Numbers on the bottom right

of each plot denote peak spatial firing rate (FR) in Hz. Middle panels depict raster plots of each

cell's in-field (left) and out-of-field (right) responses to each of the four tastes. Black dots

indicate spike times during each individual trial, aligned to the time of taste delivery (S, green

line, 4 mM saccharin; N, yellow line, 100 mM sodium chloride; W, blue line, distilled water; Q,

red line, 5 mM quinine hydrochloride). Bottom panels show the evoked responses to taste

deliveries taking place in-field and out-of-field for each cell. Each colored trace represents the

mean firing rate to one of the four tastes, smoothed with a 1D Gaussian filter (σ = 5 ms). Note

that taste responses are only found within the taste-responsive cell's place field (left panel of **A**).

C, Mean magnitude of taste responsiveness (eta-squared, or η^2) for the in-field and out-of-field

regions of taste-responsive and non-taste-responsive place cells that fit our criteria (>= 10 trials

in- and out-of-field; n = 26 taste-responsive cells, n = 153 non-taste-responsive cells). The mean

η^2 value for the in-field region of taste-responsive place cells (striped gray bar) was higher than

that of the in-field region of non-taste-responsive cells (striped white bar), the out-of-field region

of taste-responsive cells (gray bar), or the out-of-field region of non-taste-responsive (white bar)

cells (1-way ANOVA, ***p = 3e-05), indicating that place cells only respond to tastes delivered

within their place field.

Figure 6. Example hippocampal responses to different elements of the taste experience.

A, Summary of the number of taste-responsive cells (n = 96/482 cells) that responded to taste

presence (Pres, n = 77 cells), identity (ID, n = 36 cells), and/or palatability (Pal, n = 18 cells). **B**,

Example raster plots and PSTHs from a taste-responsive place cell (left) and interneuron (right).

Black dots in the raster plots (top four panels) represent spike times during each trial, aligned to

taste delivery time (S, green line, 4 mM saccharin; N, yellow line, 100 mM sodium chloride; W,

blue line, distilled water; Q, red line, 5 mM quinine hydrochloride). Each colored trace in the

PSTHs (bottom panels) represents the mean firing rate to one of the four tastes, smoothed with

a 1D Gaussian filter (σ = 5 ms). Light gray lines indicate periods of significant responses to

taste presence, while dark gray lines indicate periods of significant responses to taste identity

(*p < 0.05, t-tests on successive time windows). **C**, Relative palatability of the four taste stimuli

as determined by a brief-access task (BAT). Palatability rank is determined by the average

number of licks per 15 s of exposure to the given taste. **D**, Rank-order correlation (r_s) between

the taste-evoked firing rates and palatability rank (S > N > W > Q) for the place cell (left) and

interneuron (right) depicted in **B**. Black lines indicate periods of significant palatability-

relatedness (*p < 0.05, t-tests on successive time windows). Note the similarity between the

timing of palatability- and identity-related responses (dark gray lines in **B**).

1003

Figure 7. Hippocampal neurons discriminate between tastes based on palatability.

A, Histogram showing the percentage of hippocampal neurons that responded significantly to

taste presence (Pres, light gray line), identity (ID, dark gray line) or palatability (Pal, black line)

at each time point relative to taste delivery. Nonspecific responses to taste presence emerged

before responses to taste identity or palatability, which occurred at similarly long latencies. *B*,

For the 19 cells that responded significantly to taste presence and identity, the onset of the

identity-related response (y-axis) is plotted against the onset of the presence-related response

(x-axis). Similar to what is seen in **A**, single-neuron responses to taste presence typically

preceded responses to taste identity, as evidenced by the cloud of points above the unity line

(black dashed line). *C*, For the 14 cells that responded significantly to taste identity and

palatability, the onset of the palatability-related response (y-axis) is plotted against the onset of

the identity-related response (x-axis). Responses to identity and palatability tended to emerge

simultaneously in single units, as evidenced by tight clustering around the unity line (regression

slope = 0.92, ***p = 8.68e-06). *D*, PCA of identity-responsive cells (n = 36 cells). Each colored

line depicts the PC of pooled responses to each of the four tastants (green: S, 4 mM saccharin;

yellow: N, 100 mM sodium chloride; blue: W, distilled water; red: Q, 5 mM quinine

hydrochloride) over time. Palatability-relatedness emerges at the same time as taste selectivity,

as shown by significant encoding of palatability rank (S > N > W > Q, in reverse order here; **p

< 0.01, comparison to firing-rate-shuffled controls) by the PCs starting at ~1.4 s after stimulus

delivery (black line).

1024

1025 **Table 1.**

1026 Cell distribution across animals

1027

Animal	CA1 cells					
	All	**Pyr**	**Int**	**Taste responsive**	**Pyr**	**Int**
EM5	162	155	7	24	18	6
LH36	41	39	2	7	5	2
EM6	136	118	18	52	35	17
LP1	91	79	12	11	6	5
LH42	52	50	2	2	1	1
Total	482	441	41	96	65	31

1028

Figure 2

A

B

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7