Altered Gene Regulatory Networks are Associated with the Transition from C_3 to Crassulacean Acid Metabolism in *Erycina* (Oncidiinae: Orchidaceae)

Karolina Heyduk (karohey@gmail.com)[1*], Michelle Hwang (Mxhwang@gmail.com)[1], Victor A. Albert (vaalbert@buffalo.edu)[2,3], Katia Silvera (katia.silvera@ucr.edu)[4, 5], Tianying Lan (tianying@buffalo.edu)[2], Kimberly M. Farr (kmfarr@buffalo.edu)2, Tien-Hao Chang (cardiandra@gmail.com)[2], Ming-Tsair Chan (mbmtchan@gate.sinica.edu.t)[6], Klaus Winter (winterk@si.edu)[5], and Jim Leebens-Mack (jleebensmack@uga.edu)[1]

– Department of Plant Biology, University of Georgia, Athens, GA, USA
– Department of Biological Sciences, University at Buffalo, Buffalo, NY, USA
– School of Biological Sciences, Nanyang Technological University, Singapore
– Department of Botany and Plant Sciences, University of California-Riverside, Riverside, CA, USA
– Smithsonian Tropical Research Institute, Panama City, Panama
– Agricultural Biotechnology Research Center, Academia Sinica, Nankang, Taipei, Taiwan, ROC

Correspondence:
Dr. Karolina Heyduk
karohey@gmail.com

Abstract

Crassulacean acid metabolism (CAM) photosynthesis is a modification of the core C_3 photosynthetic pathway that improves the ability of plants to assimilate carbon in water-limited environments. CAM plants fix CO_2 mostly at night, when transpiration rates are low. All of the CAM pathway genes exist in ancestral C_3 species, but the timing and magnitude of expression are greatly altered between C_3 and CAM species. Understanding these regulatory changes is key to elucidating the mechanism by which CAM evolved from C_3. Here we use two closely related species in the Orchidaceae, *Erycina pusilla* (CAM) and *Erycina crista-galli* (C_3), to conduct comparative transcriptomic analyses across multiple time points. Clustering of genes with expression variation across the diel cycle revealed some canonical CAM pathway genes similarly expressed in both species, regardless of photosynthetic pathway. However, gene network construction indicated that 149 gene families had significant differences in network connectivity and were further explored for these functional enrichments. Genes involved in light sensing and ABA signaling were some of the most differently connected genes between the C_3 and CAM *Erycina* species, in agreement with the contrasting diel patterns of stomatal conductance in C_3 and CAM plants. Our results suggest changes to transcriptional cascades are important for the transition from C_3 to CAM photosynthesis in *Erycina*.

Keywords: RNA-seq, transcriptomics, photosynthesis, time-course, gene network

Introduction

      Crassulacean acid metabolism (CAM) is a carbon concentrating mechanism that evolved multiple times in response to CO_2 limitation caused by water stress. In C_3 species, stomata remain open during the day to assimilate atmospheric CO_2, but water limitation can force stomata to close, resulting in impaired CO_2 fixation at the expense of growth. When water stress is prolonged, stomatal closure in C_3 plants can become debilitating. CAM species circumvent prolonged stomatal closure by opening stomata at night and fix CO_2 nocturnally, when evapotranspiration rates are on average lower. CO_2 is temporarily stored as malic acid in the vacuoles until day time, when stomata close and malic acid is moved back into the cytosol for decarboxylation. The resulting increase of CO_2 levels near ribuslose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO) results in highly efficient CO_2 reduction via C_3 photosynthesis CAM is associated with a number of anatomical, physiological and genetic change, including alterations to leaf anatomy (Nelson and Sage, 2008; Zambrano et al., 2014), stomatal opening at night, and tight regulation of metabolic genes within day/night cycles. Despite the complexity of these evolutionary novelties, CAM plants are found in a wide range of plant families, including eudicot species in the Euphorbiaceae (Horn et al., 2014) and Caryophyllales (Guralnick et al., 1984; Moore et al., 2017; Winter and Holtum, 2011) and monocot lineages in Agavoideae (Abraham et al., 2016; Heyduk et al., 2016), Orchidaceae (Silvera et al., 2009, 2010), and Bromeliaceae (Crayn et al., 2004).

      The CAM pathway is well-described biochemically (Holtum et al., 2005) and contemporary genomics approaches are beginning to shed light on the genetic basis of CAM (Abraham et al., 2016; Cushman et al., 2008; Dever et al., 2015) (Fig. 1). As CO_2 enters the chloroplast-containing cells as night, it is initially converted to HCO_3^- facilitated by a carbonic anhydrase (CA). HCO_3^- is then fixed by phosphoenolpyruvate carboxylase (PEPC) using phosphoenolpyruvate (PEP) as the substrate. Carboxylation of PEP results in oxaloacetate (OAA), which is subsequently converted to malic acid by malate dehydrogenase (MDH). Malic acid is then moved into the vacuole for storage. The vacuolar transporter of malic acid is not known for certain, although previous studies have pointed to aluminum-activated malate transporters (ALMT) as a candidate (Kovermann et al., 2007; Yang et al., 2017). During the day, the malic acid is released from the vacuoles either via a passive process or through as-yet undescribed transporter. The malic acid is then decarboxylated to CO_2 and PEP using two decarboxylation pathways: NAD and/or NADP malic enzymes together with pyruvate, phosphate dikinase (PPDK), or MDH and phosphoenolpyruvate carboxykinase (PEPCK)

      While the roles of canonical CAM enzymes are considered novel in CAM species, they are all present in C_3 ancestral species as well. As a result, the evolution of CAM likely involved alterations to gene copies, including changes to protein sequences and regulatory motifs. For example, PEPC in C_3 species can play several roles depending on tissue type and developmental stage, including providing carbon backbones to the citric acid cycle and providing malate for cellular pH balance (Aubry et al., 2011; Winter et al., 2015), but its core function as a carboxylating enzyme remains unchanged in C_3 and CAM species. Studies of molecular evolution of PEPC in CAM species have largely determined that there are CAM-specific copies of the enzyme that are differentially expressed in C_3 and CAM species (Gehrig et al., 2001; Lepiniec et al., 1994; Ming et al., 2015; Silvera et al., 2014). In some cases, these CAM-specific PEPC gene copies have been shown to share sequence similarity across closely related but

independently derived CAM taxa (Christin et al., 2014; Silvera et al., 2014). Additionally,
genomic screens across many canonical CAM gene promoters have revealed an enrichment of
circadian clock motifs (Ming et al., 2015), implicating alterations to transcription factor binding
sites during CAM evolution. Because of the strong influence of the internal circadian clock on
CAM (Hartwell, 2005), we might expect that genes involved in the CAM regulatory pathway
should be controlled by a co-expressed circadian master regulator (Borland et al., 1999; Nimmo,
2000; Taybi et al., 2000b).
99
Elucidation of regulatory changes requires comparative analysis between closely related
C_3 and CAM species, and this can be accomplished through RNA-Seq analyses. One of the
largest plant families with multiple origins of CAM is the Orchidaceae; known for floral
diversity and inhabiting a broad range of habitats, the evolution of CAM in predominantly
epiphytic lineages may have also contributed to orchid diversity (Silvera et al., 2009). Epiphytic
species constitute more than 70% of the Orchidaceae (Chase et al., 2015; Gravendeel et al.,
2004), and many exhibit different degrees of CAM (Silvera et al., 2005). In a large proportion of
these, CAM is weakly expressed relative to C_3. Weakly expressed CAM may represent an
evolutionary end point, or may be an important intermediate step on the evolutionary path
between constitutive C_3 and constitutive CAM. Because many genera within the Orchidaceae
include both C_3 and weak/strong CAM species, the orchids are an attractive family to study the
evolution of CAM photosynthesis. The subtribe Oncidiinae is one of the most diverse subtribes
within Orchidaceae and it is part of a large epiphytic subfamily (Epidendroideae) in which CAM
may have facilitated the expansion into the epiphytic habitat (Silvera et al., 2009). Despite the
prevalence of CAM within the subtribe, the genus *Erycina* is particularly interesting because it
has both CAM and C_3 species. *Erycina pusilla* is a fast-growing CAM species with
transformation capability and has the potential to be a model species for studying CAM
photosynthesis in monocots (Lee et al., 2015). Comparative investigations of *E. pusilla* and its C_3
relative, *E. crista-galli*, can therefore offer valuable insight into studying the evolution and
regulation of CAM photosynthesis in the Orchidaceae. Through comparative, time-course RNA-
Seq analysis of *E. pusilla* and *E. crista-galli*, we aim to understand 1) the changes in expression
of core CAM genes between C_3 and CAM *Erycina* species and 2) which regulatory changes are
required for the evolution of CAM.
123
Materials and Methods
Plant growth and RNA-Seq tissue collection
Erycina pusilla (L.) N.H.Williams & M.W.Chase (CAM) seedlings were cultivated on
solid PSYP medium comprising 2 g/L Hyponex No. 1, 2 g/L tryptone, 20 g/L sucrose, 0.1 g/L
citric acid, and 1 g/L active charcoal in flasks. The pH of the medium is adjusted to 5.4 before
autoclaving and gelling with 3 g/L Phytagel. Plants were grown in 12-hour day and 12-hour
night conditions over three independent dates in a growth chamber at the University at Buffalo,
with temperatures set to 22-25C and lights on at 6 am for a 12 hour photoperiod. Light intensity
was between 95-110 μmol m^{-2} s^{-1}. Leaf samples for RNA-sequencing were collected every 4
hours directly from plants grown on sealed flasks for the first two experiments (January and
February 2015, Set 1 and Set 2 respectively) and every 2 hours from the final experiment
(October 2015, Set 3), where both medium-sized and large plants were collected. Because
Erycina species are considered miniatures and are therefore relatively small for destructive leaf

sampling, we use individual genets as biological replicates at each time point. Leaf samples were
flash frozen in liquid N_2 and stored at -80°C.

Erycina crista-galli (Rchb.f.) N.H.Williams & M.W,Chase (C_3) plants were wild-
collected from Peña Blanca, District of Capira, Republic of Panama at 858m above sea level,
then grown and propagated in a commercial orchid greenhouse in Bajo Bonito, District of
Capira, Republic of Panama. Plants were fertilized once a week week alternatively with a 20-20-
20 or 16-32-16 N-P-K fertilizer. Similarly sized and aged plants were moved into an
environmental growth chamber at the Smithsonian Tropical Research Institute laboratories
(Panama City, Panama) in April 2016, where they were allowed to acclimate for 48 hours to the
following conditions: 12 hour light/dark cycle (lights on 6 a.m.), 25°C/22°C day/night
temperatures, 60% humidity, and a light intensity of 30 μmol m^{-2} s^{-1}, which is similar to the light
intensity this species would experience naturally. Biological replicates (consisting of entire
shoots without root tissue) were sampled every 4 hours over a 24-hour period, starting at ZT0
(lights on, 6 a.m.) with 4 replicates per time point. Tissue was flash frozen in liquid nitrogen and
stored as described above for *E. pusilla*.

RNA was isolated from leaf tissue of both *Erycina* species using the RNeasy Plant Mini
Kit (Qiagen). RNA samples were subsequently quantified via Nanodrop and checked for
integrity with a Bioanalyzer v2100. RNA libraries were constructed using the Kapa mRNA
stranded kit with a combinatorial barcoding scheme (Glenn et al., 2016). Libraries were
sequenced on an Illumina NextSeq500 with PE75 reads, pooling 30-32 samples per run. A
summary of the data can be found in Supplemental Table 1.

Gas exchange
Individual shoots (leaves emerging from a common base) of *E. pusilla* and *E. crista-galli*
species were individually sealed within a CQP 130 porometer gas-exchange cuvette (Walz,
Effeltrich, Germany), located inside an environmental chamber (Environmental Growth
Chambers, OH, USA) operating on 12h light (6AM to 6PM) at 28°C, and 12h dark (6PM to
6AM) cycle at 22°C. Light intensity inside the chamber was 230 μmol m^{-2} s^{-1}. Plants were
watered 3-4 times daily and humidity inside the chamber was maintained at near 60%.
Continuous net CO_2 exchange was measured for each plant for up to 8 day/night cycles with data
points obtained every 4 minutes. For *E. pusilla*, water was withheld from one of the three plants
measured (drought stress) between the fifth and the eighth day; regarding *E. crista-galli*, water
was withheld from the third to the fifth day for one plant of the three measured. Data is presented
for all 3 replicates of each of the two *Erycina* species in Supplemental Figure 1.

Titratable Acidity
Erycina crista-galli plants were too small for both transcriptomic and titratable acidity
analysis, therefore leaf tissue from this species was collected only for RNA-Seq. Leaf samples
from *E. pusilla* were collected from the same plant and at the same time such that half the plant
was used for RNA-sequencing and the other half was used for titratable acidity assays. Leaves
from mature plants were collected every 4 hours, flash frozen in liquid N_2, weighed, and boiled
in 20% ethanol and deionized water. Titratable acidity was measured as the amount of 0.002M
NaOH required to neutralize the extract to a pH of 7. Because leaf tissue was limited for *E.*
crista-galli plants, we conducted titrations on three plants that were not sampled for RNAseq to

confirm their status as C_3. Samples were collected in greenhouse conditions at dawn and dusk
with three replicates at each time. Titratable acidity was measured as for *E. pusilla* but using
0.001M KOH.
186
Transcriptome assembly
An initial *de novo* transcriptome was assembled from Sets 1 and 2 for *E. pusilla*
sequences and from all samples sequenced for *E. crista-galli* using Trinity v2.0.6 (Haas et al.,
2013). Reads were cleaned using Trimmomatic (Bolger et al., 2014) as implemented in Trinity,
and assemblies were made on *in-silico* normalized reads. An initial evaluation of read mapping
results from *E. pusilla* Sets 1, 2, and 3 showed a large degree of variation among replicates; to
reduce this variation, we used only reads from Sets 1 and 2, as well as medium sized plants from
Set 3. All reads for *E. crista-galli* were included in the analysis. These data were further reduced
to include only 4 replicates per time points for a total of 24 samples. Replicates were chosen
randomly. Read mapping and abundance estimation for transcripts was conducted separately in
each species using RSEM v1.3.0 (Li and Dewey, 2011) and Kallisto (Bray et al., 2016).
Transcripts with a transcripts per kilobase million mapped (TPM) < 2 were removed and the
reads were re-mapped to the filtered assemblies.
200
Ortholog circumscription and isoform filtering
To determine gene family circumscription and annotation, all transcripts were sorted into
14 orthogroup gene families from the genomes of the following: *Amborella trichopoda, Ananas*
comosus, Arabidopsis thaliana, Asparagus officinalis, Brachypodium distachyon, Carica
papaya, Dendrobium catenatum, *Elaeis guineensis*, *Musa acuminata*, *Oryza sativa*,
Phalaenopsis equestris, *Solanum lycopersicum* , *Sorghum bicolor, Spirodella polyrrhiza* , *Vitis*
vinifera , and *Zostera marina*. Assembled transcripts were first used to query the genome
database using blastx and sorted to gene families (orthogroups) based on best BLAST hit. The
gene families were annotated by *Arabidopsis* members, using TAIR 10 (www.arabidopsis.org)
classifications. Transcripts were retained only if they 1) had a length less than the maximum
sequence of a gene family based on the sequenced genomes and 2) had a length no less than 50%
of the minimum sequence length based on sequenced genome members of that gene family.
213
Trinity produces both gene components and subsidiary isoforms, which may represent
true alternative splice isoforms or allelic or paralogous sequence variants. To mitigate dilution of
read mappings to multiple isoforms, we instead used gene components (hereafter referred to as
transcripts) for all further analyses (including gene level read mapping from RSEM). For gene
tree estimation we took the longest isoform per component per orthogroup, using our
minimum/maximum orthogroup filtered data set. Scripts for orthogroup sorting and filtering can
be found at www.github.com/kheyduk/Erycina.
221
Time-dependent clustering
To incorporate time into our clustering analysis, we used R software package maSigPro
v1.46.0 (Conesa et al., 2006; Nueda et al., 2014), which analyzes expression data for patterns
across time by fitting each gene's expression pattern to a polynomial using stepwise regression.
Cross-normalized read counts and a negative binomial distribution for the generalized linear
models were used. For each transcript, maSigPro estimated up to a 4[th] degree polynomial and
tested the fit via ANOVA. Transcripts that had significantly time-structured expression

(Benjamini & Hochberg adjusted p < 0.05) were retained while all others were removed from
further analysis. Additionally, any genes considered overly influential based on DFBETAS
diagnostic (how much an observation affects the estimate of a regression coefficient) (Belsley et
al., 1980) were also removed. In total, 1,515 transcripts from *E. pusilla* and 505 transcripts from
E. crista-galli were removed as influential genes.

The remaining transcripts that did show time-dependent expression (n=7,066 in *E. pusilla*
and n=7,127 in *E. crista-galli*) were clustered by fuzzy clustering based on similarity in
expression profiles. An optimal fuzzifier *m*, a parameter that determines how much clusters can
overlap, was calculated in the Mfuzz package (Kumar and E Futschik, 2007) of R for each
species (*m*=1.09 for both *E. pusilla* and *E. crista-galli*). The number of groups *k* for each species
was determined by choosing a value which minimizes the within-group variance (Supplemental
Figure 2); a *k* of 6 was used for both *E. pusilla* and *E. crista-galli*. Z-scores of normalized counts
were calculated for each gene in each cluster, as well as a median cluster expression, for each
species separately.

Gene trees and expression
Gene trees were estimated for phosphoenolpyruvate carboxylase (PEPC) and its kinase
(PPCK) by first aligning nucleotide sequences from *Erycina* transcripts and their associated gene
family members from the sequenced genomes using PASTA (Mirarab et al., 2014), then
estimating trees using RAxML (Stamatakis, 2006). Gene expression for genes of interest was
plotted based on averaged transcripts per million mapped (TPM) for each replicate.

Network analysis
To identify regulatory candidates possibly involved in CAM and examine the
relationships between genes within clusters, we used the ARACNe-AP v1.4 (Lachmann et al.,
2016) algorithm to create networks of co-expressed transcripts from both species separately.
Briefly, the algorithm randomly samples gene pairs and uses an adaptive partitioning approach to
infer a pairwise mutual information (MI) statistic, or measure of statistical dependence, between
them. This process is repeated iteratively for a specified number of bootstraps, while at each step
removing indirect interactions. A final network is built based on the consensus of all bootstrap
runs. Although ARACNe provides an option to specify transcription factors to generate a
directed network by only considering interactions with a transcription factor source, we chose to
generate an undirected gene co-expression network of genes that were significantly time-
structured based on our maSigPro analysis, using 100 bootstrap replicates in ARACNe.

We imported network data into Cytoscape (Shannon et al., 2003) to generate
visualizations and calculate network statistics. Nodes were color coded by their cluster
membership and scaled to represent number of connections. Network statistics were exported
and further analyzed at the orthogroup level. We calculated which orthogroups had the largest
average difference in connectivity between the two species. We first calculated the average
connectivity (number of directed edges, output from Cytoscape Network Analysis) for each
orthogroup per species, then normalized these via Z-scores and subtracted the Z-score of the
orthogroup in *E. pusilla* from that in *E. crista-galli*. Outliers – those orthogroups with the largest
difference between species in connectivity – had Z-score differences above and below the upper
and lower quantiles (Supplemental Figure 3).

275
The outlier orthogroups with large changes to connectivity between species
(Supplemental Table 3) were explored for genes of interest. The largest difference in
connectivity was found in a E3 ubiquitin ligase gene family shown to play a role in ABA
signaling, with the *Arabidopsis* homolog known as ring finger of seed longevity1 (RSL1). To
explore differences in network connections of RSL1 between the two species, we employed the
diffusion algorithm (Carlin et al., 2017) in Cytoscape which finds strongly interactive nodes to a
target of interest. For both species, we found the diffusion network for the RSL1 gene. Only a
single gene copy of RSL1 was time-structured in *E. crista-galli*, but *E. pusilla* had two copies
found in the ARACNe network. One gene copy had only a single connection to any other gene in
the network and was not analyzed further. The other copy in *E. pusilla*, which had 72 directed
connections, was used as the center of the diffusion network. Diffusion networks were compared
for orthogroup content between species using a hypergeometric test. Gene Ontology (GO) terms
were compared for the two RSL1 subnetworks and checked for enrichment using hypergeometric
test (using all GO terms found in either ARACNe network as the universe), correcting for
multiple testing with Bonferroni-Holm significance correction.
291
Results
Gas exchange patterns and titratable acidity
Gas exchange data collected continuously showed net nighttime CO_2 uptake in CAM *E.*
pusilla under both well-watered conditions and while drought stressed (Fig. 2a). C_3 *Erycina*
crista-galli displayed net CO_2 uptake during the light period only. There was no net uptake of
CO_2 at night. Nonetheless, under drought stress, a slight decrease in respiratory loss of CO_2 at
night may indicate low levels of CAM cycling. Titration data collected from the same plants and
at the time of RNA-sampling in *E. pusilla* confirms CAM function in the plants used for gene
expression analysis, with a significant increase in leaf titratable acids occurring towards the end
of the dark period (Fig. 2A, 6AM), and a reduction in total acids during the day period. Although
the C_3 *E. crista-galli* had higher overall levels of leaf acids, there was no significant diurnal
fluctuation (Fig. 2b).
304
Clustering of genes with time-structured expression profiles
After filtering by minimum/maximum length each transcript's orthogroup, 23,596 and
26,437 genes were retained in *E. pusilla* and *E. crista-galli*, respectively. Both species had a
similar number of genes that were significantly time-structured (~7,000) according to maSigPro.
Each species had best fit to k=6 clusters, with three clusters showing nighttime biased expression
and three with daytime bias (Fig. 3). Expression of PEPC, the initial carboxylating enzyme in the
CO_2 fixation pathway at night, increased in expression just before the onset of darkness in the
CAM species *E. pusilla* (Fig. 4A). In contrast, there was a low, but significant, time-structured
expression pattern of PEPC in the C_3 species *E. crista-galli* (Fig. 4B). The dedicated kinase,
PPCK, which phosphorylates PEPC and allows it to function in the presence of malate, likewise
showed a strong nocturnal increase in expression in the CAM species, with similar levels of
expression in the C_3 species (Fig. 4B).
317
Network Comparisons
The network for C_3 *E. crista-galli* had 119,338 directed connections between 4,828
nodes, whereas the CAM *E. pusilla* network was notably less connected, with only 76,071

connections between 4,591 nodes. Although the number of genes in each network was similar,
overall connectivity of *E. pusilla* is easily seen in both the fewer number of connections as well
as the mean number of connections (34.3 in *E. pusilla* vs. 50.3 in *E. crista-galli*). As expected,
genes from the same co-expressed cluster (Fig. 3) were grouped within the larger ARACNe
network for each species (Fig. 5A,C).
326
Comparison of network connectivity of the time-structured genes found 149 outlier
transcripts that had large species differences in the number of connected directed edges
(Additional file 4); these were largely skewed toward increased connectivity in *E. crista-galli*
(n=90). Annotations of these outliers revealed a number of genes involved in stomatal
opening/closing and ABA signaling. GO term enrichment indicates that outliers that skew
toward more connectivity in C_3 *E. crista-galli* were enriched for vacuolar and tonoplast
membrane proteins and potassium and calcium transport. Genes that were more connected in
CAM *E. pusilla* were enriched for genes involved in aldehyde dehydrogenase activity, among
other functions (Supplemental Table 4).
336
A E3 ubiquitin ligase also known as ring finger of seed longevity1 (RSL1) had the
greatest difference in connectivity between the two species. RSL1 has been shown to be a
negative regulator of ABA signaling (Bueso et al., 2014) and was chosen as a center node for
comparison between the two species. The diffusion algorithm used to create subnetworks
defaults to producing a subnetwork that is 10% of the total nodes in the larger network; as a
result, both species subnetworks were roughly the same size, containing about 400 genes.
However, the connectivity of those subnetworks differed greatly (Fig. 5B,D); the C_3 *E. crista-*
galli RSL1 subnetwork contained 30,392 connections, whereas the CAM *E. pusilla* network had
only 9,244. The subnetworks differed in their gene content as well. There contained 429 and 427
orthogroups in *E. pusilla* and *E. crista-galli,* respectively, but only 57 orthogroups were shared
between the two (this was not significantly under-enriched via a hypergeometric test (p=1)).
While all of the genes in the *E. crista-galli* RSL1 subnetwork were in night-biased expression
clusters (485), *E. pusilla* had more subnetwork genes in day-biased clusters (298) than in night
biased ones (162). GO term enrichment indicates both subnetworks are enriched for chloroplast,
chloroplast stroma, and photosynthesis (Supplemental Table 5).
352
While both subnetworks were centered on RSL1 and generally were enriched for similar
types of genes involved in chloroplast functions and photosynthesis, there were substantial
differences in gene content between the networks. The focal gene RSL1 acts as a master negative
regulator of ABA signaling pathway by targeting pyrabactin resistance 1 (PYR1) and PYR-like
(PYL) ABA receptors for degradation. More generally its role in protein ubiquitination is
relatively unknown. RSL1 was the third most connected node in the C_3 *E. crista-galli*
subnetwork (419 directed connections; most connected node had 423 connections). In CAM *E.*
pusilla, RSL1 was 173[rd] out of 460 genes in connectivity. *E. pusilla* had a number of ABA
responsive genes in its RSL1 subnetwork that *E. crista-galli* did not, including protein
phosphatase 2C (PP2C), a gene encoding a member of the Snf1-related kinase family, plus a
homolog of *ABA Overly Sensitive 5*. Recent work has shown that ABA responses in stomata
including *PYR/PYL* and downstream genes are responsible for constitutive stomatal aperture, as
well as stomatal responses to drought stress (Gonzalez-Guzman et al., 2012).
366

Erycina crista-galli, on the other hand, had a number of light sensing and circadian clock
genes that CAM *E. pusilla* did not have in the RSL1 subnetwork, including *lov kelch protein 2*
(LKP2/ZTL gene family), *time for coffee* (*TIC*), *phytochrome B/D* (*PHYB*), and *protein*
phosphatase 2A (*PP2A*). LKP2/ZTL and PHYB are both involved in light sensing (blue and
red/far-red, respectively). Specifically, *LKP2/ZTL* genes are thought to regulate light induced
protein degradation via their function as E3 ligases (Demarsy and Fankhauser, 2009; Ito et al.,
2012), and *ztl* mutants in *Arabidopsis* show a prolonged clock period under constant light due to
the lack of degradation of clock components via ZTL (Más et al., 2003; Somers et al., 2000). TIC
has been found to be responsible for the amplitude of the circadian clock but is not thought to be
directly involved in light signaling (Hall et al., 2003). TIC is also implicated in daytime
transcriptional induction via its association with the central circadian oscillator late elongated
hypocotyl (LHY) (Ding et al., 2007). Finally, PP2A is a member of a large family of plant
phosphoprotein phosphatases (PPP) with several cellular roles (Uhrig et al., 2013). PP2A
specifically has been implicated in brassinosteroid signaling, light signaling via
dephosphorylation of phototropin2 (Tseng and Briggs, 2010), flowering time control (Kim et al.,
2002), as well as the induction of CAM under certain abiotic stresses (Cushman and Bohnert,
1999).

Although the C_3 *E. crista-galli* RSL1 subnetwork contained circadian and light sensing
transcripts that were absent in the *E. pusilla* RSL1 subnetwork, these transcripts largely did not
have different expression patterns between the two species with the exception of *PHYB* (Fig.
6B). The contrasting gene content of the RSL1 subnetwork suggests light sensing and circadian
regulatory cascades comprise large differences between C_3 and CAM species, rather than levels
of gene expression, which were quite similar.

Discussion
Shared gene expression patterns
While traditionally it was thought that canonical CAM genes should have large
differences in timing and magnitude of expression between C_3 and CAM species, recent work
has highlighted that between closely related C_3 and CAM species, that may not always be the
case (Heyduk et al., 2018b). In *Erycina*, it appears that a similar pattern holds. Both the CAM
and C_3 species have time-structured expression of PEPC albeit at very different expression
levels. While this alone says little about PEPC's function in both species, the high nocturnal
expression of PEPC's dedicated kinase, PPCK, in both *E. pusilla* and *E. crista-galli* suggests that
PEPC is being phosphorylated in both species and therefore has the need to function in the
presence of malate.
It is worth noting that all C_3 species have the genes involved in the CAM cycle. Many,
including PEPC, function in anaplerotic reactions of the TCA cycle. PEPC has also been shown
to have a role in malate production for osmotic regulation of stomatal aperture and in CO_2
fixation in guard cells of tobacco (Asai et al., 2000; Daloso et al., 2015). Because most RNA-seq
or gene expression studies to date in C_3 species sample during the day, understanding how
common nocturnal *PEPC* expression is across flowering plants will require more nighttime gene
expression studies in C_3 species, especially in lineages closely related to CAM species. While
nearly all the genes in the CAM CO_2 fixation pathway have known functions in C_3 species,
PPCK is a notable exception. Phosphorylation of PEPC by PPCK in CAM and C_4 species is
well-described (Jiao and Chollet, 1989; Nimmo et al., 1986; Taybi et al., 2000a); malate and

other organic acids act as negative regulators of PEPC, but phosphorylation of PEPC renders it
immune to these negative effects. Thus, phosphorylated PEPC via PPCK is required for high
levels of CAM and C_4 malic acid production. In C_3 species, however, there is no clear need for
heavily phosphorylated PEPC, especially at night, making the expression of *PEPC* and *PPCK* in
C_3 *E. crista-galli* intriguing (though see Sullivan et al., 2004 and Fukayama et al., 2006 for work
on nocturnal *PPCK* expression in soybean and rice, respectively).
419
Alterations in regulatory pathways between C_3 and CAM orchids
Network analysis of co-expressed genes and subsequent comparisons between species
can give insights not only into the changes in expression, but also the degree to which a given
gene changes in connectivity between species. Extensive connectivity for a gene has long
thought to be a signal of a "hub" or master regulatory gene – one that cannot experience large
changes in timing or magnitude of expression without significant perturbations to the entire
network. A major assumption of co-expression networks is that they rely on mRNA as an
accurate predictor of downstream processes; while this is not always the case, recent work
showed that although mRNA and protein networks differed in their gene content, they
overlapped in gene ontologies and were predictive of pathway regulation in maize (Walley et al.,
2016). Additionally, it has been shown that a correlation exists between connectivity of a gene
within a network and the evolutionary conservation of the gene's sequence across a number of
flowering plant species (Masalia et al., 2017). It is therefore somewhat surprising to observe that
as many as 149 gene families have large changes in connectivity between the two closely related
Erycina species.
435
The 149 outlier gene families in *Erycina* were enriched for functions in protein
degradation via phosphorylation and ubiquitination. While typically differences in phenotype are
considered the result of changes in the abundance of gene products, our data highlight the
importance of considering protein degradation as well. The differences in connectivity of protein
degradation pathway genes were unbiased between species – in other words, genes involved in
protein degradation have increased connectivity in both species. More interestingly, genes that
had increased connectivity in C_3 *E. crista-galli* were enriched for GO terms involved in
potassium and chloride channels and membrane proteins associated with chloroplasts and
vacuoles. Greater connectivity of such genes in the C_3 species indicates an increased reliance on
ion and metabolite fluxes. In stomatal guard cells, which make up a smaller portion of the whole-
leaf transcriptome, these fluxes directly affect stomatal aperture and may play a role in
alternative regulation of stomatal opening in C_3 and CAM species.
448
Regulatory changes in ABA, light, and clock perception
Stomatal opening in CAM species has been vastly understudied, despite the opportunities
it presents for understanding a fundamental biological process. In C_3 species, stomatal opening is
thought to be regulated by blue and red light inputs, whereas stomatal closing is driven by efflux
of potassium cations. It remains largely unknown how stomata sense darkness, but experimental
data have suggested a large role of CO_2 concentrations on stomatal aperture (Cockburn et al.,
1979). Draw down of CO_2 concentrations at night in the intercellular airspace would result in
stomatal opening, whereas high concentrations of CO_2 from decarboxylation during the day may
promote stomatal closure. CO_2 concentrations undoubtedly play some role in the inverted
stomatal aperture of CAM species, but more contemporary genomic work has implicated

additional levels of regulation in CAM stomata (Abraham et al., 2016; Cushman and Bohnert,
1997; Wai et al., 2017). For example, ABA may be one signaling molecule for nocturnal
stomatal closure (Desikan et al., 2004; Gonzalez-Guzman et al., 2012; Merilo et al., 2013). Gene
expression results coupled with network analysis in *Erycina* indicate that both ABA signaling
and light sensing are likely to be altered. Indeed, the gene with the largest difference in
connectivity between species is *RSL1*, which encodes a E3 ubiquitin ligase known to function in
stomatal response to ABA. Expression of RSL1 is higher in CAM *E. pusilla* and is clustered with
day-biased genes (Fig. 6A), whereas in C_3 *E. crista-galli* expression is about half that in *E.*
pusilla and slightly increases in expression during the dark period. These expression patterns are
consistent with stomatal regulation between C_3 and CAM species. In *E. crista-galli* nighttime
ABA may play a larger role in drought-induced or nighttime stomatal closure than in the CAM
E. pusilla, where the nocturnal stomatal closure driven by ABA signaling must be repressed to
allow for nighttime CO_2 acquisition. It is also unknown what causes daytime stomatal closure in
CAM species; while high intracellular CO_2 concentrations may play a role, so might ABA
signaling. *RSL1*'s high connectivity to other genes in the subnetwork of *E. crista-galli* suggests
that alterations to regulatory networks are also important for fully functional CAM.
475
Stomata, while highly responsive to ABA, are also strongly regulated by light inputs. The
gene encoding phytochrome B (phyB), a photoreceptor that both regulates transcriptional
responses to red and far-red light as well as entrains the circadian clock (Goosey et al., 1997;
Más et al., 2000; Ni et al., 1999), was differentially regulated and expressed in the two *Erycina*
species (Fig. 6B). While both species had copies of *phyB* that showed time-structured expression,
only *E. crista-galli* had a copy that was in the same network as *RSL1* and had many connections
to other genes in the same network. A single copy of *phyB* was time-structured in CAM *E.*
pusilla, and the expression level relative to copies in C_3 *E. crista-galli* was quite low. Instead, the
constitutively expressed copy of *phyB* in *E. pusilla* had the highest expression, but what it's role
might be given constant expression across time is unclear. The stark difference in both
connectivity and expression levels of *phyB* in *E. pusilla* and *E. crista-galli* suggests that light-
induced transcriptional regulation has a greater role in the C_3 species, and that phyB mediated
transcriptional regulation in CAM species may be significantly reduced. Additionally, the
presence of *PP2A* in the C_3 *E. crista-galli* RSL1 subnetwork, but not in *E. pusilla,* further
indicates that light induced responses are differentially regulated (although expression was
similar between the two species, Fig. 6D). PP2A is, among many other tasks, responsible for
dephosphorylation of photoropin 2, which subsequently promotes stomatal opening (Tseng and
Briggs, 2010; Uhrig et al., 2013).
494
Differences in light input sensing and signaling between C_3 and CAM species is not
surprising, but relatively little work has focused on this aspect of CAM biology. Previous work
assessed light responses in the facultative CAM species *Portulacaria afra* (Lee and Assmann,
1992) and *Mesembryanthemum crystallinum* (Tallman et al., 1997) and showed that both species
had reduced stomatal response to blue light signals when relying on the CAM cycle for carbon
fixation (though see Ceusters et al., 2014). Additionally, *M. crystallinum* had reduced guard cell
zeaxanthin production during the day in the CAM state compared to the C_3 state. The reduction
in zeaxanthin in the CAM state was shown to be a result of the downregulation of the pathway,
rather than an aberration in guard cell chloroplasts. Changes to regulation and expression of

phyB in *Erycina* further support significantly altered light-induced pathways in CAM species
relative to C_3.
506
TIME FOR COFFEE (*TIC*) also stood out in its altered connectivity between the C_3 and
CAM species. In C_3 *E. crista-galli*, TIC was present in the RSL1 diffusion network whereas it
was not for *E. pusilla*. TIC has been shown to be involved in maintaining the amplitude of the
circadian clock in *Arabidopsis* (Ding et al., 2007; Hall et al., 2003), as well as regulating
metabolic homeostasis and response to environmental cues (Sanchez-Villarreal et al., 2013). *tic*
mutants in *Arabidopsis* showed large phenotypic effects ranging from late flowering to
anatomical abnormalities. The *tic* mutants also showed extreme tolerance to drought, likely due
to increased amounts of osmolytes such as proline and myo-inositol, as well as an accumulation
of starch. As a result of the pleiotropic effects of TIC, altered network status of TIC in CAM *E.*
pusilla compared to C3 *E. crista-galli* likely results a complex alteration in phenotype. In
general, the mechanism that link the circadian clock and CAM photosynthesis are unknown
(Boxall et al., 2005), and research to uncover circadian regulation of CAM is limited to
transcriptomic studies. Gene expression comparisons between CAM *Agave* and C_3 *Arabidopsis*
revealed changes to the timing of expression of *REVEILLE 1,* a clock output gene that integrates
the circadian network to metabolic activity (Yin et al., 2018). While it's possible that changes to
the timing of *REVEILLE 1* are required for CAM evolution, expression differences may also be a
result of lineage-specific changes to expression unrelated to CAM. Comparisons of closely
related C3 and CAM species of *Erycina* suggest that alteration of transcriptional cascades from
circadian oscillators may play a role in the evolution of CAM, rather than large scale changes to
the timing or abundance of expression, but additional work to link clock outputs to the CAM
phenotype are necessary.
528
Implications for the evolution of CAM in Oncidiinae
Both CAM and C_4 photosynthesis have evolved multiple times across the flowering plant
phylogeny, suggesting that the evolution of these complex traits cannot be insurmountably
difficult. Recent physiological work demonstrates that the evolution of anatomical traits required
for CAM or C_4 often predates the emergence of strong, constitutive carbon concentrating
mechanisms (Christin et al., 2013; Heyduk et al., 2016). Other transcriptomic work has shown
that closely related C_3 and CAM species share the expression of canonical CAM genes,
especially *PEPC* and *PPCK* as seen here in *Erycina* (Heyduk et al., 2018b). It has even been
suggested that many C_3 plants already have the nocturnal CAM cycle in place for fixation of
respired CO_2 and generation of amino acids (Bräutigam et al., 2017), but this alone is unlikely to
entirely explain the repeated and relatively frequent emergence of CAM on the angiosperm
phylogeny.
541
Indeed it appears that gene expression alone would not facilitate the large-scale transition
from C_3 to CAM. Recent comparative work across multiple C_3 and C_4 transcriptomes highlighted
the recurrent co-option of highly expressed genes from C_3 species into C_4 (Moreno-Villena et al.,
2018). The initial co-option of highly expressed gene copies happened early in the evolutionary
trajectory between C_3 and C_4, and later steps included the refinement of C_3 enzymes, including
kinetic and tissue specificity. It is quite likely that a similar model of evolution holds for CAM,
in that C_3 relatives of CAM lineages have been shown to have similar expression patterns of
canonical CAM genes (though, it is worth noting, they are not typically highly expressed in the

C$_3$ species). In *Erycina*, both C$_3$ and CAM species share similar expression profiles for *PEPC*
and *PPCK*, with the latter having nearly identical levels of expression in both taxa. It is possible
that low levels of nocturnal CO$_2$ fixation via PEPC evolved in the ancestor of both species for
non-photosynthetic reasons; for example, tobacco leaves (Kunitake et al., 1959) and cotton
ovules (Dhindsa et al., 1975) show various levels of carbon concentration, and fixation of
cytosolic CO$_2$ via PEPC is required to replenish the citric acid cycle (Aubry et al., 2011).
Because these pathways already exist, slight upregulation of some components in a shared
ancestor may have enabled the origins of CAM in certain lineages (Bräutigam et al., 2017).
558
However, gene expression alone clearly does not make a species CAM, and further
refinement is necessary. Refinement of CAM may take the form of improving secondary
metabolic processes or genomic characteristics that allow for strong and constitutive CAM to
exist. For example, fine tuning of carbohydrate turnover is necessary for CAM function (Borland
et al., 2016; Ceusters et al., 2014). Experiments placing *Mesembryanthemum crystallinum* in
CO$_2$-free air at night resulted in a dampened CAM cycle (Dodd et al., 2003) and comparative
RNA-seq in facultative CAM and C$_3$/CAM comparisons has shown increased reliance on
carbohydrate breakdown as CAM function increases (Brilhaus et al., 2016; Heyduk et al.,
2018a). Circadian regulation of CAM genes in pineapple was shown to be the result of promoters
that induce evening expression (Ming et al., 2015), and genes that link the clock to metabolic
outputs had shifts in expression phasing between *Agave* (CAM) and *Arabidopsis* (C$_3$) (Yin et al.,
2018). Variation in timing and magnitude of expression of various light sensing and clock genes
has been described not only here in *Erycina*, but also in *Agave* (Abraham et al., 2016). In all, the
growing evidence suggests that the secondary processes that make a CAM plant, including
stomatal regulation, light sensing and downstream transcriptional responses, and carbohydrate
metabolism feedbacks, undergo fine-tuning along the evolutionary trajectory between C$_3$ and
CAM. Further work that characterizes closely related C$_3$ and CAM species has the potential to
greatly advance our understanding of the integration of nighttime CO$_2$ acquisition with more
complex regulatory pathways.
578
579 **Data Availability**
580 Raw sequence reads are available on NCBI Short Read Archive, under the BioProject
581 PRJNA483943. All scripts used that are not part of existing programs are available at
582 www.github.com/kheyduk.
583

Acknowledgements
The authors would like to sincerely thank Orquideas Tropicales in Panama for generously
donating *Erycina crista-galli* plants. Additional gratitude goes to Saravanaraj Ayyampalayam for
assistance with data processing, Jeremy Ray for assistance with RNA preparation, and Rishi
Masalia for giving feedback on figures. This work was supported by NSF grants to J.L.M, V.A,
and K.S (DEB #1442199 and 1442190) and the by Smithsonian Tropical Research Institute.

Author Contributions Statement
JLM and VAA conceived and led the project; KS and KW collected, phenotyped plants, and
conducted continuous gas exchange measurements, and KS additionally sampled *E. crista-galli*
for RNA-seq; VAA and TL grew and sampled *Erycina pusilla* individuals for RNA and
physiology; KH and MH sequenced and analyzed the data and prepared the manuscript; JLM

oversaw general experimental design, and all authors contributed to the final version of this
manuscript.
598
599 **Conflict of Interest**
600 The authors declare no personal, professional, or financial conflicts of interest.
601
602 **References**
603

Abraham, P. E., Yin, H., Borland, A. M., Weighill, D., Lim, S. D., De Paoli, H. C., et al. (2016).
Transcript, protein and metabolite temporal dynamics in the CAM plant Agave. *Nat. Plants*
2, 16178. doi:10.1038/nplants.2016.178.
Asai, N., Nakajima, N., Tamaoki, M., Kamada, H., and Kondo, N. (2000). Role of Malate
Synthesis Mediated by Phosphoenolpyruvate Carboxylase in Guard Cells in the Regulation
of Stomatal Movement. *Plant Cell Physiol.* 41, 10–15. doi:10.1093/pcp/41.1.10.
Aubry, S., Brown, N. J., and Hibberd, J. M. (2011). The role of proteins in C3 plants prior to
their recruitment into the C4 pathway. *J. Exp. Bot.* 62, 3049–3059. doi:10.1093/jxb/err012.
Belsley, D. A., Kuuh, E., and E., W. R. (1980). *Wiley Series in Probability and Statistics*.
doi:10.1002/9780470316559.oth1.
Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30, 2114–20. doi:10.1093/bioinformatics/btu170.
Borland, A. M., Guo, H.-B., Yang, X., and Cushman, J. C. (2016). Orchestration of carbohydrate
processing for crassulacean acid metabolism. *Curr. Opin. Plant Biol.* 31, 118–124.
doi:10.1016/j.pbi.2016.04.001.
Borland, A. M., Hartwell, J., Jenkins, G. I., Wilkins, M. B., Nimmo, H. G., Borland, A. M., et al.
(1999). Metabolite Control Overrides Circadian Regulation of Phosphoenolpyruvate
Carboxylase Kinase and CO□ Fixation in Crassulacean Acid Metabolism. *Plant Physiol.*
121, 889–896.
Boxall, S. F., Foster, J. M., Bohnert, H. J., Cushman, J. C., Nimmo, H. G., and Hartwell, J.
(2005). Conservation and Divergence of Circadian Clock Operation in a Stress-Inducible
Crassulacean Acid Metabolism Species Reveals Clock Compensation against Stress.
PLANT Physiol. 137, 969–982. doi:10.1104/pp.104.054577.
Bräutigam, A., Schlüter, U., Eisenhut, M., and Gowik, U. (2017). On the Evolutionary Origin of
CAM Photosynthesis. *Plant Physiol.* 174, 473–477. doi:10.1104/pp.17.00195.
Bray, N. L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-
seq quantification. *Nat. Biotechnol.* 34, 525–527. doi:10.1038/nbt.3519.
Brilhaus, D., Bräutigam, A., Mettler-Altmann, T., Winter, K., and Weber, A. P. M. (2016).
Reversible Burst of Transcriptional Changes during Induction of Crassulacean Acid
Metabolism in Talinum triangulare. *Plant Physiol.* 170, 102–22. doi:10.1104/pp.15.01076.
Bueso, E., Rodriguez, L., Lorenzo-Orts, L., Gonzalez-Guzman, M., Sayas, E., Muñoz-Bertomeu,
J., et al. (2014). The single-subunit RING-type E3 ubiquitin ligase RSL1 targets PYL4 and
PYR1 ABA receptors in plasma membrane to modulate abscisic acid signaling. *Plant J.* 80,
1057–1071. doi:10.1111/tpj.12708.
Carlin, D. E., Demchak, B., Pratt, D., Sage, E., and Ideker, T. (2017). Network propagation in
the cytoscape cyberinfrastructure. *PLOS Comput. Biol.* 13, e1005598.
doi:10.1371/journal.pcbi.1005598.
Ceusters, J., Borland, A. M., Taybi, T., Frans, M., Godts, C., and De Proft, M. P. (2014). Light

quality modulates metabolic synchronization over the diel phases of crassulacean acid
metabolism. *J. Exp. Bot.* 65, 3705–3714. doi:10.1093/jxb/eru185.

Chase, M. W., Cameron, K. M., Freudenstein, J. V., Pridgeon, A. M., Salazar, G., van den Berg,
C., et al. (2015). An updated classification of Orchidaceae. *Bot. J. Linn. Soc.* 177, 151–174.
doi:10.1111/boj.12234.

Christin, P.-A., Arakaki, M., Osborne, C. P., Bräutigam, A., Sage, R. F., Hibberd, J. M., et al.
(2014). Shared origins of a key enzyme during the evolution of C4 and CAM metabolism. *J.*
Exp. Bot. 65, 3609–3621. doi:10.1093/jxb/eru087.

Christin, P.-A., Osborne, C. P., Chatelet, D. S., Columbus, J. T., Besnard, G., Hodkinson, T. R.,
et al. (2013). Anatomical enablers and the evolution of C4 photosynthesis in grasses. *Proc.*
Natl. Acad. Sci. U. S. A. 110, 1381–6. doi:10.1073/pnas.1216777110.

Cockburn, W., Ting, I. P., and Sternberg, L. O. (1979). Relationships between Stomatal Behavior
and Internal Carbon Dioxide Concentration in Crassulacean Acid Metabolism Plants. *Plant*
Physiol. 63, 1029–32. doi:10.1104/PP.63.6.1029.

Conesa, A., Nueda, M. J., Ferrer, A., and Talón, M. (2006). maSigPro: a method to identify
significantly differential expression profiles in time-course microarray experiments.
Bioinformatics 22, 1096–1102. doi:10.1093/bioinformatics/btl056.

Crayn, D. M., Winter, K., and Smith, J. A. C. (2004). Multiple origins of crassulacean acid
metabolism and the epiphytic habit in the Neotropical family Bromeliaceae. *Proc. Natl.*
Acad. Sci. U. S. A. 101, 3703–8. doi:10.1073/pnas.0400366101.

Cushman, J. C., and Bohnert, H. J. (1997). Molecular Genetics of Crassulacean Acid
Metabolism. *Plant Physiol.* 113, 667–676. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/15012212.

Cushman, J. C., and Bohnert, H. J. (1999). CRASSULACEAN ACID METABOLISM:
Molecular Genetics. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 50, 305–332.
doi:10.1146/annurev.arplant.50.1.305.

Cushman, J. C., Tillett, R. L., Wood, J. A., Branco, J. M., and Schlauch, K. A. (2008). Large-
scale mRNA expression profiling in the common ice plant, Mesembryanthemum
crystallinum, performing C3 photosynthesis and Crassulacean acid metabolism (CAM). *J.*
Exp. Bot. 59, 1875–94. doi:10.1093/jxb/ern008.

Daloso, D. M., Antunes, W. C., Pinheiro, D. P., Waquim, J. P., AraÚJo, W. L., Loureiro, M. E.,
et al. (2015). Tobacco guard cells fix CO_2 by both Rubisco and PEPcase while sucrose acts
as a substrate during light-induced stomatal opening. *Plant. Cell Environ.* 38, 2353–2371.
doi:10.1111/pce.12555.

Demarsy, E., and Fankhauser, C. (2009). Higher plants use LOV to perceive blue light. *Curr.*
Opin. Plant Biol. 12, 69–74. doi:10.1016/J.PBI.2008.09.002.

Desikan, R., Cheung, M.-K., Clarke, A., Golding, S., Sagi, M., Fluhr, R., et al. (2004). Hydrogen
peroxide is a common signal for darkness- and ABA-induced stomatal closure in *Pisum*
sativum. *Funct. Plant Biol.* 31, 913. doi:10.1071/FP04035.

Dever, L. V, Boxall, S. F., Kneřová, J., and Hartwell, J. (2015). Transgenic perturbation of the
decarboxylation phase of Crassulacean acid metabolism alters physiology and metabolism
but has only a small effect on growth. *Plant Physiol.* 167, 44–59.
doi:10.1104/pp.114.251827.

Dhindsa, R. S., Beasley, C. A., and Ting, I. P. (1975). Osmoregulation in Cotton Fiber:
Accumulation of Potassium and Malate during Growth. *Plant Physiol.* 56, 394–8. Available
at: http://www.ncbi.nlm.nih.gov/pubmed/16659311 [Accessed September 21, 2018].

Ding, Z., Millar, A. J., Davis, A. M., and Davis, S. J. (2007). TIME FOR COFFEE encodes a
nuclear regulator in the Arabidopsis thaliana circadian clock. *Plant Cell* 19, 1522–36.
doi:10.1105/tpc.106.047241.
Dodd, A. N., Griffiths, H., Taybi, T., Cushman, J. C., and Borland, A. M. (2003). Integrating diel
starch metabolism with the circadian and environmental regulation of Crassulacean acid
metabolism in Mesembryanthemum crystallinum. *Planta* 216, 789–797.
doi:10.1007/s00425-002-0930-2.
Fukayama, H., Tamai, T., Taniguchi, Y., Sullivan, S., Miyao, M., and Nimmo, H. G. (2006).
Characterization and functional analysis of phospho *enol* pyruvate carboxylase kinase genes
in rice. *Plant J.* 47, 258–268. doi:10.1111/j.1365-313X.2006.02779.x.
Gehrig, H., Heute, V., and Kluge, M. (2001). New partial sequences of phosphoenolpyruvate
carboxylase as molecular phylogenetic markers. *Mol. Phylogenet. Evol.* 20, 262–274.
doi:10.1006/mpev.2001.0973.
Glenn, T. C., Nilsen, R., Kieran, T. J., Finger, J. W., Pierson, T. W., Bentley, K. E., et al. (2016).
Adapterama I: Universal stubs and primers for thousands of dual-indexed Illumina libraries
(iTru & iNext). *bioRxiv*. Available at:
http://biorxiv.org/content/early/2016/06/15/049114 [Accessed May 31, 2017].
Gonzalez-Guzman, M., Pizzio, G. A., Antoni, R., Vera-Sirera, F., Merilo, E., Bassel, G. W., et
al. (2012). Arabidopsis PYR/PYL/RCAR receptors play a major role in quantitative
regulation of stomatal aperture and transcriptional response to abscisic acid. *Plant Cell* 24,
2483–96. doi:10.1105/tpc.112.098574.
Goosey, L., Palecanda, L., and Sharrock, R. A. (1997). Differential patterns of expression of the
Arabidopsis PHYB, PHYD, and PHYE phytochrome genes. *Plant Physiol.* 115, 959–69.
doi:10.1104/PP.115.3.959.
Gravendeel, B., Smithson, A., Slik, F. J. W., and Schuiteman, A. (2004). Epiphytism and
pollinator specialization: drivers for orchid diversity? *Philos. Trans. R. Soc. Lond. B. Biol.*
Sci. 359, 1523–1535. doi:10.1098/rstb.2004.1529.
Guralnick, L. J., Rorabaugh, P. A., and Hanscom, Z. (1984). Seasonal Shifts of Photosynthesis in
Portulacaria afra (L.) Jacq. *Plant Physiol.* 76, 643–646. doi:10.1104/pp.76.3.643.
Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Philip, D., Bowden, J., et al. (2013).
De novo transcript sequence reconstruction from RNA-Seq: reference generation and
analysis with Trinity. Nature Protocols doi:10.1038/nprot.2013.084.De.
Hall, A., Bastow, R. M., Davis, S. J., Hanano, S., McWatters, H. G., Hibberd, V., et al. (2003).
The TIME FOR COFFEE gene maintains the amplitude and timing of Arabidopsis
circadian clocks. *Plant Cell* 15, 2719–29. doi:10.1105/tpc.013730.
Hartwell, J. (2005). The co-ordination of central plant metabolism by the circadian clock.
Biochem. Soc. Trans. 33, 945–8. doi:10.1042/BST20050945.
Heyduk, K., McKain, M. R., Lalani, F., and Leebens-Mack, J. (2016). Evolution of CAM
anatomy predates the origins of Crassulacean acid metabolism in the Agavoideae
(Asparagaceae). *Mol. Phylogenet. Evol.* 105, 102–113.
Heyduk, K., Ray, J. N., Ayyampalayam, S., and Leebens-Mack, J. (2018a). Shifts in gene
expression profiles are associated with weak and strong Crassulacean acid metabolism. *Am.*
J. Bot. doi:10.1002/ajb2.1017.
Heyduk, K., Ray, J. N., Ayyampalayam, S., Moledina, N., Borland, A., Harding, S., et al.
(2018b). Nocturnal expression of Crassulacean acid metabolism (CAM) genes predates the
origin of CAM in the genus Yucca. *bioRxiv*, 371344. doi:10.1101/371344.

Holtum, J. A. M., Smith, J. A. C., and Neuhaus, H. E. (2005). Intracellular transport and
pathways of carbon flow in plants with crassulacean acid metabolism. *Funct. Plant Biol.* 32,
429. doi:10.1071/FP04189.

Horn, J. W., Xi, Z., Riina, R., Peirson, J. A., Yang, Y., Dorsey, B. L., et al. (2014). Evolutionary
bursts in Euphorbia (Euphorbiaceae) are linked with photosynthetic pathway. *Evolution (N.
Y).* 68, 3485–3504. doi:10.1111/evo.12534.

Ito, S., Song, Y. H., and Imaizumi, T. (2012). LOV Domain-Containing F-Box Proteins: Light-
Dependent Protein Degradation Modules in Arabidopsis. *Mol. Plant* 5, 573–582.
doi:10.1093/MP/SSS013.

Jiao, J.-A., and Chollet, R. (1989). Regulatory seryl-phosphorylation of C4 phosphoenolpyruvate
carboxylase by a soluble protein kinase from maize leaves. *Arch. Biochem. Biophys.* 269,
526–535. doi:10.1016/0003-9861(89)90136-7.

Kim, D.-H., Kang, J.-G., Yang, S.-S., Chung, K.-S., Song, P.-S., and Park, C.-M. (2002). A
phytochrome-associated protein phosphatase 2A modulates light signals in flowering time
control in Arabidopsis. *Plant Cell* 14, 3043–56. doi:10.1105/TPC.005306.

Kovermann, P., Meyer, S., Hörtensteiner, S., Picco, C., Scholz-Starke, J., Ravera, S., et al.
(2007). The Arabidopsis vacuolar malate channel is a member of the ALMT family. *Plant
J.* 52, 1169–1180. doi:10.1111/j.1365-313X.2007.03367.x.

Kumar, L., and E Futschik, M. (2007). Mfuzz: a software package for soft clustering of
microarray data. *Bioinformation* 2, 5–7. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/18084642 [Accessed September 20, 2016].

Kunitake, G., Stitt, C., and Saltman, P. (1959). Dark Fixation of CO(2) by Tobacco Leaves.
Plant Physiol. 34, 123–7. doi:10.1104/PP.34.2.123.

Lachmann, A., Giorgi, F. M., Lopez, G., and Califano, A. (2016). ARACNe-AP: Gene network
reverse engineering through adaptive partitioning inference of mutual information.
Bioinformatics 32, 2233–2235. doi:10.1093/bioinformatics/btw216.

Lee, D. M., and Assmann, S. M. (1992). Stomatal responses to light in the facultative
Crassulacean acid metabolism species, Portulacaria afra. *Physiol. Plant.* 85, 35–42.
doi:10.1111/j.1399-3054.1992.tb05260.x.

Lee, S. H., Li, C. W., Liau, C. H., Chang, P. Y., Liao, L. J., Lin, C. S., et al. (2015).
Establishment of an Agrobacterium-mediated genetic transformation procedure for the
experimental model orchid Erycina pusilla. *Plant Cell. Tissue Organ Cult.* 120, 211–220.
doi:10.1007/s11240-014-0596-z.

Lepiniec, L., Vidal, J., Chollet, R., Gadal, P., and Cretin, C. (1994). Phosphoenolpyruvate
carboxylase: structure, regulation and evolution. *Plant Sci.* 99, 111–124. doi:10.1016/0168-
9452(94)90168-6.

Li, B., and Dewey, C. N. (2011). RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* 12, 323. doi:10.1186/1471-2105-
12-323.

Más, P., Devlin, P. F., Panda, S., and Kay, S. A. (2000). Functional interaction of phytochrome B
and cryptochrome 2. *Nature* 408, 207–211. doi:10.1038/35041583.

Más, P., Kim, W.-Y., Somers, D. E., and Kay, S. A. (2003). Targeted degradation of TOC1 by
ZTL modulates circadian function in Arabidopsis thaliana. *Nature* 426, 567–570.
doi:10.1038/nature02163.

Masalia, R. R., Bewick, A. J., Burke, J. M., Irizarry, R., Bertranpetit, J., and Laayouni, H.
(2017). Connectivity in gene coexpression networks negatively correlates with rates of

molecular evolution in flowering plants. *PLoS One* 12, e0182289. doi:10.1371/journal.pone.0182289.

Merilo, E., Laanemets, K., Hu, H., Xue, S., Jakobson, L., Tulva, I., et al. (2013). PYR/RCAR receptors contribute to ozone-, reduced air humidity-, darkness-, and CO2-induced stomatal regulation. *Plant Physiol.* 162, 1652–68. doi:10.1104/pp.113.220608.

Ming, R., VanBuren, R., Wai, C. M., Tang, H., Schatz, M. C., Bowers, J. E., et al. (2015). The pineapple genome and the evolution of CAM photosynthesis. *Nat. Genet.* 47, 1435–42. doi:10.1038/ng.3435.

Mirarab, S., Nguyen, N., and Warnow, T. (2014). "PASTA: Ultra-Large Multiple Sequence Alignment," in *Research in Computational Molecular Biology: 18th Annual International Conference, RECOMB 2014, Pittsburgh, PA, USA, April 2-5, 2014, Proceedings*, ed. R. Sharan (Cham: Springer International Publishing), 177–191. doi:10.1007/978-3-319-05269-4_15.

Moore, A., de Vos, J. M., Hancock, L. P., Goolsby, E., and Edwards, E. J. (2017). Targeted Enrichment of Large Gene Families for Phylogenetic Inference: Phylogeny and Molecular Evolution of Photosynthesis Genes in the Portullugo (Caryophyllales). *bioRxiv*. Available at: http://biorxiv.org/content/early/2017/06/04/145995 [Accessed June 19, 2017].

Moreno-Villena, J. J., Dunning, L. T., Osborne, C. P., and Christin, P.-A. (2018). Highly Expressed Genes Are Preferentially Co-Opted for C4 Photosynthesis. *Mol. Biol. Evol.* 35, 94–106. doi:10.1093/molbev/msx269.

Nelson, E. A., and Sage, R. F. (2008). Functional constraints of CAM leaf anatomy: tight cell packing is associated with increased CAM function across a gradient of CAM expression. *J. Exp. Bot.* 59, 1841–50. doi:10.1093/jxb/erm346.

Ni, M., Tepperman, J. M., and Quail, P. H. (1999). Binding of phytochrome B to its nuclear signalling partner PIF3 is reversibly induced by light. *Nature* 400, 781–784. doi:10.1038/23500.

Nimmo, G. A., Nimmo, H. G., Hamilton, I. D., Fewson, C. A., and Wilkins, M. B. (1986). Purification of the phosphorylated night form and dephosphorylated day form of phosphoenolpyruvate carboxylase from Bryophyllum fedtschenkoi. *Biochem. J.* 239, 213–20. doi:10.1042/BJ2390213.

Nimmo, H. G. (2000). The regulation of phosphoenolpyruvate carboxylase in CAM plants. *Trends Plant Sci.* 5, 75–80. doi:10.1016/S1360-1385(99)01543-5.

Nueda, M. J., Tarazona, S., and Conesa, A. (2014). Next maSigPro: updating maSigPro bioconductor package for RNA-seq time series. *Bioinformatics* 30, 2598–2602. doi:10.1093/bioinformatics/btu333.

Sanchez-Villarreal, A., Shin, J., Bujdoso, N., Obata, T., Neumann, U., Du, S.-X., et al. (2013). *TIME FOR COFFEE* is an essential component in the maintenance of metabolic homeostasis in *Arabidopsis thaliana*. *Plant J.* 76, n/a-n/a. doi:10.1111/tpj.12292.

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., et al. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* 13, 2498–504. doi:10.1101/gr.1239303.

Silvera, K., Santiago, L. S., Cushman, J. C., and Winter, K. (2009). Crassulacean Acid Metabolism and Epiphytism Linked to Adaptive Radiations in the Orchidaceae. *Plant Physiol.* 149. Available at: http://www.plantphysiol.org/content/149/4/1838.short [Accessed June 21, 2017].

Silvera, K., Santiago, L. S., Cushman, J. C., and Winter, K. (2010). The incidence of

crassulacean acid metabolism in Orchidaceae derived from carbon isotope ratios: a checklist
of the flora of Panama and Costa Rica. *Bot. J. Linn. Soc.* 163, 194–222. doi:10.1111/j.1095-
8339.2010.01058.x.

Silvera, K., Santiago, L. S., and Winter, K. (2005). Distribution of crassulacean acid metabolism
in orchids of Panama: evidence of selection for weak and strong modes. *Funct. Plant Biol.*
32, 397. doi:10.1071/FP04179.

Silvera, K., Winter, K., Rodriguez, B. L., Albion, R. L., and Cushman, J. C. (2014). Multiple
isoforms of phosphoenolpyruvate carboxylase in the Orchidaceae (subtribe Oncidiinae):
implications for the evolution of crassulacean acid metabolism. *J. Exp. Bot.* 65, 3623–36.
doi:10.1093/jxb/eru234.

Somers, D. E., Schultz, T. F., Milnamow, M., and Kay, S. A. (2000). ZEITLUPE Encodes a
Novel Clock-Associated PAS Protein from Arabidopsis. *Cell* 101, 319–329.
doi:10.1016/S0092-8674(00)80841-7.

Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with
thousands of taxa and mixed models. *Bioinformatics* 22, 2688–90.
doi:10.1093/bioinformatics/btl446.

Sullivan, S., Jenkins, G. I., and Nimmo, H. G. (2004). Roots, Cycles and Leaves. Expression of
the Phosphoenolpyruvate Carboxylase Kinase Gene Family in Soybean. *Plant Physiol.* 135.
Available at: http://www.plantphysiol.org/content/135/4/2078.full [Accessed September 14,
2017].

Tallman, G., Zhu, J., Mawson, B. T., Amodeo, G., Nouhi, Z., Levy, K., et al. (1997). Induction
of CAM in Mesembryanthemum crystallinum Abolishes the Stomatal Response to Blue
Light and Light-Dependent Zeaxanthin Formation in Guard Cell Chloroplasts. *Plant Cell*
Physiol. 38, 236–242. doi:10.1093/oxfordjournals.pcp.a029158.

Taybi, T., Patil, S., Chollet, R., and Cushman, J. C. (2000a). A minimal serine/threonine protein
kinase circadianly regulates phosphoenolpyruvate carboxylase activity in crassulacean acid
metabolism-induced leaves of the common ice plant. *Plant Physiol.* 123, 1471–82.
Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=59103&tool=pmcentrez&render
type=abstract [Accessed October 10, 2015].

Taybi, T., Patil, S., Chollet, R., Cushman, J. C., Taybi, T., Patil, S., et al. (2000b). A Minimal
Serine / Threonine Protein Kinase Circadianly Regulates Phosphoenolpyruvate Carboxylase
Activity in Crassulacean Acid Metabolism-Induced Leaves of the Common Plant. *Plant*
Physiol. 123, 1471–1481.

Tseng, T.-S., and Briggs, W. R. (2010). The Arabidopsis rcn1-1 mutation impairs
dephosphorylation of Phot2, resulting in enhanced blue light responses. *Plant Cell* 22, 392–
402. doi:10.1105/tpc.109.066423.

Uhrig, R. G., Labandera, A.-M., and Moorhead, G. B. (2013). Arabidopsis PPP family of
serine/threonine protein phosphatases: many targets but few engines. *Trends Plant Sci.* 18,
505–513. doi:10.1016/J.TPLANTS.2013.05.004.

Wai, C. M., VanBuren, R., Zhang, J., Huang, L., Miao, W., Edger, P. P., et al. (2017). Temporal
and spatial transcriptomic and microRNA dynamics of CAM photosynthesis in pineapple.
Plant J. doi:10.1111/tpj.13630.

Walley, J. W., Sartor, R. C., Shen, Z., Schmitz, R. J., Wu, K. J., Urich, M. A., et al. (2016).
Integration of omic networks in a developmental atlas of maize. *Science* 353, 814–8.
doi:10.1126/science.aag1125.

Winter, K., and Holtum, J. A. M. (2011). Induction and reversal of crassulacean acid metabolism
in *Calandrinia polyandra*: effects of soil moisture and nutrients. *Funct. Plant Biol.* 38, 576–
582. Available at: https://doi.org/10.1071/FP11028.
Winter, K., Holtum, J. A. M., and Smith, J. A. C. (2015). Crassulacean acid metabolism: a
continuous or discrete trait? *New Phytol.* doi:10.1111/nph.13446.
Yang, X., Hu, R., Yin, H., Jenkins, J., Shu, S., Tang, H., et al. (2017). The Kalanchoë genome
provides insights into convergent evolution and building blocks of crassulacean acid
metabolism. *Nat. Commun.* 8. doi:10.1038/s41467-017-01491-7.
Yin, H., Guo, H.-B., Weston, D. J., Borland, A. M., Ranjan, P., Abraham, P. E., et al. (2018).
Diel rewiring and positive selection of ancient plant proteins enabled evolution of CAM
photosynthesis in Agave. *BMC Genomics* 19, 588. doi:10.1186/s12864-018-4964-7.
Zambrano, V. A. B., Lawson, T., Olmos, E., Fernández-García, N., and Borland, A. M. (2014).
Leaf anatomical traits which accommodate the facultative engagement of crassulacean acid
metabolism in tropical trees of the genus Clusia. *J. Exp. Bot.* 65, 3513–23.
doi:10.1093/jxb/eru022.

887
888
889
Figure 1 - A simplified diagram of the Crassaluceuan acid metabolism (CAM) pathway under day
and night conditions. ALMT9 – aluminum activated malate transporter; CA – carbonic
anhydrase; MDH – malate dehydrogenase; OAA – oxaloacetate; ME – malic enzyme (NAD or
NADP); PEPC – phosphoenolypyruvate; PEPCK – PEP carboxykinase; PPCK – PEPC kinase;
PPDK – pyruvate, phosphate dikinase.

Figure 2 – Gas exchange and titratable acidity for A) *Erycina pusilla* and B) *Erycina crista-*
galli. Gas exchange is shown for a single plant; replicate plant gas exchange plots can be found
in Supplemental Figure 1. Drought induction is indicated with a red arrow. Titrations are shown
for dawn and dusk; full titratable acidity values are in Supplemental Table 2.

Figure 3 – Expression z-scores for each gene in each cluster for both A) *E. pusilla* and B) *E.*
crista-galli, with median expression shown in the black line. Clusters represent co-expressed
genes within each species' time-structured transcripts. Cool colors (blue and purple) are clusters
with nighttime biased expression, whereas warmer colors (orange and yellow) are clusters whose
transcripts increase in expression during the day.

Figure 4 – Expression of A) phosphoenolpyruvate carboxylase (PEPC) and B) PEPC kinase in
E. pusilla (squares, blue tones) and *E. crista-galli* (circles, purple tones). Different shades of
color represent different assembled gene copies, and only transcripts that were found to be
significantly time-structured are shown. Average TPM and standard deviation, scaled to the
maximum mean TPM value across all copies and species, is plotted for all 6 time points, with the
grey background indicating nighttime samples.

Figure 5 – Network constructions of A) *E. pusilla* and C) *E. crista-galli* based on ARACNe
network analysis, and the ring finger seed longevity1 (RSL1) subnetworks for B) *E. pusilla* and
D) *E. crista-galli* calculated via the diffusion algorithm in Cytoscape. Colors correspond to

Figure 3 cluster colors – cooler colors (blues in A and B and purples in C and D) are clusters that
have an increase in expression at night and warmer colors (oranges in A and B and yellows in C
and D) are clusters that have increases in expression during the day. Dots represent genes, scaled
by the number of edges (maximum number of directed edges=435).

Figure 6 - Expression of A) ring finger of seed longevity 1 (RSL1), B) phytochrome B/D
(PHYB) C) time for coffee (TIC), and D) protein phosphatase 2A in *E. pusilla* (squares) and *E.*
crista-galli (circles). Points are colored by which category of cluster they belong to: blues and
purples are genes with increases in expression in the dark in *E. pusilla* and *E. crista-galli*,
respectively, while oranges and yellows are genes with increases in daytime expression in *E.*
pusilla and *E. crista-galli*, respectively (see Fig. 3). Time-structured genes are marked with an
asterisk, while those that were not significantly time-structured are shown in greys. Number of
directed edges per gene is shown if they belonged to the RSL1-subnetwork. Average TPM and
standard deviation, scaled to the maximum mean TPM value across all copies and species, is
plotted for all 6 time points, with the grey background indicating nighttime samples.

A

Cluster 2 (1482) Cluster 4 (1231) Cluster 5 (863)

Cluster 1 (1035) Cluster 3 (1070) Cluster 6 (1385)

Erycina pusilla (CAM)

B

Cluster 5 (1124) Cluster 6 (871) Cluster 3 (1135)

Cluster 2 (989) Cluster 1 (1310) Cluster 4 (1100)

Erycina crista-galli (C$_3$)