Differences in protein dosage underlie nongenetic differences in traits

Nikolay Burnaevskiy[1], Bryan Sands[1], Soo Yun[1], Patricia M Tedesco[2], Thomas E Johnson[2], Matt

Kaeberlein[1], Roger Brent[3,4], Alexander Mendenhall[1,4*]

[1] Department of Pathology, University of Washington, Seattle, Washington

[2] Department of Integrative Physiology, University of Colorado, Boulder, Colorado

[3] Division of Basic Sciences, Fred Hutchinson Cancer Research Center, Seattle, Washington

[4] Co-Senior Authors

[*] To whom correspondence should be addressed: alexworm@uw.edu

ABSTRACT

Genes and environments do not entirely explain variation in biological traits. Discrete and complex

traits manifest distinctly in isogenic organisms in homogenous environments. One cause of

variation in traits is variation in the expression of genes. We do not understand how variation in

gene expression arises among isogenic individuals, nor do we fully understand the

consequences. Here, for the first time in a metazoan, we quantified sources of cell-to-cell variation

in gene expression to understand how it arises in the cells of *Caenorhabditis elegans*. We adapted

an experimental approach and analytical framework we used previously in *Saccharomyces*

cerevisiae. We measured cell-to-cell variation in gene expression attributable to: 1) stochastic

molecular events in transcription and translation (a.k.a. intrinsic noise) and/or allele access, 2)

pathway-specific signaling noise, and 3) general differences in protein dosage resulting from

global differences in protein production or turnover. Surprisingly, and in contrast with single celled

organisms, in *C. elegans* cells, general differences in protein dosage constitute the major source

of variation. Cells with high protein dosage came from animals with high protein dosage. These

differences in protein dosage were consequential. Animals that made more reporter proteins had

discrete traits consistent with higher dosage of unrelated neomycin resistance and Ras gain-of-

function proteins, consistent with global variation in protein dosage. Our results suggest that the

natural axis of protein dosage underlies nongenetic, nonenvironmental differences in complex

and discrete traits, including penetrance and expressivity of hypomorphic and hypermorphic

alleles. They also raise the possibility that differences in penetrance caused by differences in

protein dosage might be responsible for some of the missing heritability in genome wide

association studies.

INTRODUCTION

Genes and environments are not the sole determinants of discreet or complex traits[1]. That

is, complex and discreet traits vary among isogenic organisms existing in the same homogeneous

environment. In the early 20th century, Timoféeff-Ressovsky[2] and Romaschoff[3], observed that,

in isogenic populations of *Drosophila funebris*, some mutations affected only some individuals

(*incomplete penetrance* of a mutation's phenotype, a discreet trait), and the severity of the

phenotype varied among affected individuals (*expressivity*)[4]. Later, scientists characterized

numerous differences in traits among isogenic unicellular organisms in the same environment,

including differences in the lifespan of single yeast cells[5] (a complex trait). Similarly, researchers

showed that individuals in isogenic populations of *C. elegans* in homogeneous environments

manifest different lifespans[6-8] (a complex trait) and had incomplete penetrance of mutant alleles[9]

(discreet traits).

Nongenetic differences in the manifestation of traits are not stochastic; they can be

predicted by measured variables. Consider the, perhaps, seemingly peculiar, series of reports

wherein differences in the measured expression level of a single gene correlates with and/or

predicts differences in discreet and complex traits, in systems ranging from yeast to human stem

cells. For example, there can be a nine-fold difference in the expression level of a pheromone

reporter gene among individual yeast cells; cells that express more reporter gene increase in

volume more rapidly[10]. In *C. elegans*, we have shown that high expression levels of a single gene,

a GFP-based reporter of the molecular chaperone encoded by *hsp-16.2*, predicted increased

lifespan and thermotolerance[7,8] (complex traits). The same reporter gene can also predict

differences in the penetrance of hypomorphic mutations that cause defects in larval development[9]

(discreet traits). Similarly, mammalian stem cells that express high levels of a cell surface protein,

SCA-1, are more likely to differentiate into myeloid cells[11]. In these experiments where we and

others measure a reporter gene and then see some other trait covary, there must be some

underlying causative gene or genes that covary with the measured gene. In cases like the

measurement of endogenous SCA-1, the measured gene may be causative; however, this is not

the case for the freely diffusing GFP reporter protein in *C. elegans*.

The molecular basis for variation in gene expression phenotypes (such as *hsp-16.2*

reporter expression[12-14]) and their relation to variation in more complex traits they predict (such

as lifespan[7,8]) is not well understood. In *C. elegans*, *hsp-16.2* reporters are expressed most

abundantly in the 20 cells of the intestine, which form distinct segments (rings; Figure 1a)[12,14].

Genetic experiments suggest that the physiology of the worm intestine affects lifespan. For

example, intestinal expression of the forkhead transcription factor homolog of FOXO3, *daf-16*[15],

is sufficient to extend lifespan. Similarly, the expression level of a number of reporter genes of

seemingly disparate function correlated with lifespan when measured from intestine cells[16]. Yet

we do not entirely know how differences in reporter gene expression arise, nor why they are

consequential.

Ultimately, differences in traits must be caused by differences in the expression of some

causative gene or genes. Thus, it is important to understand how nongenetic, nonenvironmental

differences in gene expression arise. One way to understand how variation in gene expression

arises is to directly observe and quantify differences in gene expression using fluorescent reporter

genes. The ability to directly observe the fluorescent signals in living cells allowed us[10] and

others[17-19] to isolate and study different sources of variation in gene expression in single celled

microbes.

To better understand variation in gene expression and its relationship to discreet and

complex traits such as mutation penetrance and lifespan, we developed means to precisely

quantify differences in reporter protein expression in intestine cells[14]. We measured animal-to-

animal and cell-to-cell variation in the expression of a set of *hsp-16.2* based reporter genes. Our

experiments showed, perhaps surprisingly, that differences in site of reporter integration, copy

number, identity of the fluorophore, or sequence of 3'UTR did not affect variation in gene

expression[14]. Consistent with this observation, we also found that different *hsp-16.2* reporters

have identical abilities to predict lifespan and heat tolerance[8]. Moreover, high or low expression

of these reporters defined a whole-organism physiological state; that is, bright intestine cells came

from animals composed of bright cells, and dim intestine cells came from animals composed of

relatively dim cells[14].

Here, to better understand the relationship between the gene expression phenotype and

other traits, we built on previous work in *S. cerevisiae* to dissect and quantify different sources of

gene expression variation[10]. This is the first time these sources of variation have been quantified

in a metazoan. The results were surprising.

RESULTS

Here we adapted an approach we used in yeast[10], wherein we compared the outputs of

two differently colored (different fluorescent proteins) versions of the same reporter gene

expressed from two identical loci on homologous chromosomes (Type I experiments) or the

outputs of two different reporter genes (Type II experiments), shown in Figure 1. Measuring the

pairs of reporter genes in these two different kinds of experiments allows us to quantify the amount

of cell-to-cell variation in gene expression attributable to three distinct, experimentally tractable

bins, each with distinct underlying molecular causes. Figure 1 shows how we can use this reporter

gene measurement scheme to quantify differences in intrinsic noise[17,18] ($\eta^2(\gamma)$), signaling through

different pathways[10] ($\eta^2(P)$), or general protein expression capacity[10] ($\eta^2(G)$). A complete

description of the analytical framework is provided in Supplemental Materials Section 2: Analytical

Framework.

Type I experiments allow us to measure how much cell-to-cell variation there is in

apparently stochastic to-promoter or to-transcript binding events, or allele access. Type I

experiments reveal how much cell-to-cell variation there is in allele expression/access as

uncorrelated variation, often referred to as intrinsic noise ($\eta^2(\gamma)$); Figure 1 shows an example of a

scatterplot for the same cell type (same lineage/fate) measured from many different animals from

one of these experiments. If there were a lot of uncorrelated variation (a cloud like appearance of

points) it would mean that there were significant cell-to-cell differences in to-promoter binding, to-

transcript binding, or allele access. If there was little uncorrelated variation (alignment of points

along the major axis; Figure 1), it would mean that most cells would have had access to both

alleles. It would also mean that there were few differences attributable to probabilistic differences

in biochemical binding events (to-promoter/to-transcript) that tend to dominate when there are

small numbers of molecules involved[18]; we would say that intrinsic noise was low.

In Type I experiments for *hsp-16.2*, and for all the other reporter genes we measured,

intrinsic noise was low. The expression level of one allele accounted for 90% or more of the

expression of the other allele in animals. Figure 2 shows the average intrinsic noise levels for a

few genes we measured. Intrinsic noise was a minor component of the cell-to-cell variation. We

show individual Type I scatterplots and bar graphs organized by cell type and experiment in

Supplemental Figures 1&2.

Type II experiments compare the expression levels from two distinct genes to tell us about

cell-to-cell differences in signaling through different pathways ($\eta^2(P)$). In the scatterplot from a

Type II experiment shown in Figure 1, the uncorrelated variation (dispersion across the major

axis) is a measure of cell-to-cell differences in signaling through distinct pathways. We subtract

the average intrinsic noise ($\eta^2(\gamma)$) for the gene pairs we are measuring, so we know that the

remaining dispersion is due to signaling/pathway noise ($\eta^2(P)$) (shown in Figure 1). Type II

experiments also quantify how much cell-to-cell variation is attributable to differences in protein

expression capacity ($\eta^2(G)$), which we define as the ability to express and maintain proteins. In

the scatter plot of a Type II experiment on the bottom right of Figure 1, the correlated variation

(dispersion along the major axis) is a measure of ($\eta^2(G)$).

In Type II experiments, we found that cell fate (which ring a cell was in) was the primary

determinant of a rigid cell-fate specific ratiometric setpoint for any given gene pair, shown in

scatterplots of cells grouped together and by ring in Supplementary Figure 3. Thus, we grouped

cells we measured by the ring from which they came to prevent artificial inflation of ($\eta^2(P)$). We

found that there was little cell-to-cell (and thus, animal-to-animal) variation in signaling through

distinct pathways to activate distinct genes. Figure 3 shows that there was little signaling noise

($\eta^2(P)$). Figure 3 also shows that we found that most of the variation in gene expression was

attributable to cell-to-cell (and thus, animal-to-animal) differences in protein expression capacity

($\eta^2(G)$). These intestine cells express more protein from other, distinctly regulated genes, even if

they are on other chromosomes, or fused to other genes (Emerin::GFP; Figure 3e). Figure 3f

shows ($\eta^2(G)$) visually; two different types of animals expressing two distinct sets of reporter

genes from Type II experiments are arranged from dimmest to brightest. Note that the patterns of

expression are maintained but that the overall expression level of both reporters is different

between animals; to observe this phenomenon, there must be little intrinsic noise ($\eta^2(P)$) and little

signaling noise ($\eta^2(\gamma)$). We show individual Type II scatterplots and bar graphs organized by cell

type and experiment in Supplemental Figures 4, 5 & 6.

To corroborate the finding that protein expression capacity is a major mechanism of cell-

to-cell (and thus, animal-to-animal) variation, we conducted two additional types of experiments

where we sorted animals on the expression level of a fluorescent reporter gene (approximately

top or bottom 5-10% of animals), and measured phenotypes indicative of the activity of particular

non-fluorescent genes. First, we sorted larval (L1) animals on GFP expressed from a site on

chromosome I, controlled by the promoter of another ubiquitously expressed gene, *eft-3*; these

animals also expressed the neomycin resistance gene (Neo^R) on a different chromosome

(chromosome II) under control of the *rps-27* promoter. If general protein expression capacity (G)

was a major mechanism of cell-to-cell (and animal-to-animal) variation, then animals that have

more GFP should have more NeoR and resist extreme neomycin concentrations to a greater

extent. Figure 4A shows that we found this to be the case; brighter green animals can grow to

adulthood in a high neomycin environment better than dimmer animals.

Next, we used the *hsp-16.2* lifespan biomarker gene to sort animals expressing an

incompletely penetrant, dominant Ras gain of function mutation that caused animals to develop

between 0-4 hypodermal neoplasias[20]. If animals that made less biomarker had less protein

expression capacity, we would expect them to have lower penetrance and expressivity for this

gain of function mutation. In fact, Figure 4B shows that animals that make less of the biomarker

have a lower penetrance; brighter animals have more hypodermal neoplasias and a higher

probability of developing them. Thus, animals seem to vary in effective gene dosage for multiple

different genes. Supplemental Figure 7 shows images of animals in which the nuclear envelope

of the somatic cells, including the neoplastic hypodermal cells, is delineated by a green

fluorescent protein fusion to Emerin, a nuclear envelope component.

DISCUSSION

Here, for the first time, we quantified different sources of cell-to-cell variation in gene

expression in a metazoan. Work was facilitated by development of precise microscopic methods

that enabled accurate measurements of gene expression in individual *C. elegans* intestine cells[14].

Our experiments allowed us to independently quantify three possible sources of variation in gene

expression. We found that the major contribution to cell-to-cell variation among isogenic animals

in the same environment was correlated variation in the expression of reporter genes (protein

expression capacity; G; Figures 1&3). Our previous work identified whole animal differences in

reporter gene expression and suggested that these differences defined a whole organism

physiological state[14]. Our work here shows that these differences are caused by differences in

general protein expression capacity (quantified by a variable, G), which define differences in

protein dosage in cells throughout the animal. Our experiments here showed that animals with

higher G showed greater penetrance of two single gene traits: antibiotic resistance conferred by

a Neo^R transgene, and developmental hypodermal neoplasia caused by a gain of function in the

sole *C. elegans* Ras gene (*let-60*[20]).

The fact that, in *C. elegans*, differences in G constitute the major contribution to organism-

to-organism variation in gene expression, was somewhat surprising. In *S. cerevisiae*, differences

in G are also important for variation, but differences in *P*, the ability to send signal to activate

different promoters, are of similar magnitude[10]. Our findings do not exclude the possibility that

variation in other cell types at other points in the life of an organism might be dominated by other

sources of variation in gene expression. In this vein, it will be interesting when future researchers

quantify the different contributions to the increased cell-to-cell variation in gene expression

observed in aged mammalian cardiomyocytes[21].

Work in simple metazoans like *C. elegans*, whose cells are observable in intact animals

via light microscopy, is an essential complement to work in mammalian cell culture. The work in

simple metazoans allows us to observe and understand types of cell-to-cell variation, such as

variation in G for the whole intestine tissue (and effectively whole animals), which might not be

observed in cultures of cells lacking the proper milieu and control signals. For example, in *C.*

elegans, intestinal proteostasis is known to be controlled by insulin signaling and neurons[22,23],

and variation in general protein dosage, G, might stem from differences in neuronal and/or insulin

signals[24].

Our findings on animal-to-animal differences in protein dosage cause us to revisit previous

results on a complex trait, lifespan. Earlier, we found that young adult *C. elegans* with greater

expression of a *hsp-16.2* reporter ($P_{hsp16.2}::gfp$) lived longer and were more tolerant to stress[7,8].

We asserted that high reporter gene expression in these animals defined a physiological state

that protected the animals from the "rigors of living". The *hsp-16.2* gene encodes a small

molecular chaperone, and it seemed possible that overexpression of *hsp-16.2* and other

chaperones[13] might account for the longevity phenotype[25,26], perhaps by protecting cells from the

consequences of damage due to accumulated misfolded proteins. However, Sanchez-Blanco

and Kim identified a number of reporter genes that predict lifespans, and not all of them were

chaperones (for example, superoxide dismutase, *sod-3*). Significantly, in these experiments

lifespan prediction power increased when the investigators quantified correlated expression of

any two reporters (thus measuring the variable we call G).

Our finding that that individual animals (and the cells that comprise them) show general

differences in protein dosage may help explain incomplete penetrance. Burga et al. showed that

animals with increased expression of a *daf-21* reporter gene (P_{daf-21}*::mCherry*) had decreased

penetrance of a hypomorphic *tbx-9* mutation[27]. Since *daf-21* encodes the Hsp90 chaperone, it

seemed possible that the diminished penetrance of the *tbx-9* mutant phenotype might be caused

by increased chaperone activity that increased the activity of the hypomorphic TBX-9 protein.

Consistent with this idea, Casaneuva et al. showed that animals with chaperone levels increased

by three different experimental means exhibited decreased penetrance of a few different

hypomorphic mutant alleles. These observations are also consistent with the idea that animals

that make more chaperone reporters have higher doses of most of the proteome. Furthermore,

their experiments suggest that only mutant proteins with some residual function could benefit from

the physiological state defined by increased chaperone reporter expression level. For example,

penetrance of the hypomorphic *lin-31(n1053)* allele (a point mutation changing a single amino

acid) showed differential penetrance in animals with different measured expression levels of a

hsp-16.2 reporter gene, but the *lin-31(gk569)* null deletion (presumed null, first three of four total

exons deleted) mutant did not[9]; this is consistent with the idea that high G animals that make

more of a hypomorphic mutant protein get some increased functional capacity from increased

dosage of that partially functional protein, but animals that make more of a null/truncated protein
do not.

Thus, taken with previous work, our results suggest a complementary interpretation for
the relationship between reporter gene expression, longevity (complex trait), and incomplete
penetrance (discreet trait). We suggest that: 1) the physiological state first defined operationally
by high expression of a $P_{hsp16.2}$::GFP reporter is a global increase in protein dosage defined here
(increased G), 2) that this state may cause increased longevity by its general increase in dosage
of proteins (in addition to chaperones), and 3) that changes in penetrance of particular
hypomorphic and hypermorphic mutations may be due to changes in the dosage of these mutant
proteins.

We do not understand the molecular origins of organism-to-organism differences in protein
dosage. In principle, animals might differ in their ability to transcribe DNA into RNA, translate
mRNA, or in the rates at which mRNA and proteins are degraded. Regarding how differences in
mRNA production may come to be, in mammals, expression of c-Myc proteins globally increases
transcription [28,29]. Thus, in *C. elegans*, differences in G could be due to differential expression of
genes such as *mml-1*, the (sole) Myc family gene[30], which might lead to global differences in
transcription. For instance, measurements of single mRNAs suggest that incomplete penetrance
of *skn-1* hypomorphic mutants (which fail to form intestines) in some animals might be due to high
expression of mRNA for *end-1*[31]. It is possible that global differences in mRNA caused differences
in *end-1*. Alternatively, increased chaperone abundance in some animals[13], might lead to global
increases in translation, for example by increasing the protein folding efficiency of the nascent
protein associated complex[32].

However, we are tempted to speculate that a major contribution to animal-to-animal
differences in G might be differences in the rate of protein turnover. Activation the *daf-2/daf-16*
insulin signaling system decreases general protein turnover[33-35]. We showed recently that

activation of this system increases mean expression of the *hsp-16.2* biomarker gene[24], consistent

with the idea that decreased protein turnover conveyed by the *daf-2(e1370)* mutation would lead

to increased expression level. In the same report, we also showed that the AFD sensory neuron

pair and insulin signaling controlled the animal-to-animal variation in reporter gene expression[24].

Taken together with this report, our prior investigation suggests that variation in G might be

controlled by sensory neurons and insulin signaling (see working model in Supplemental Figure

8). Perhaps significantly, Sanchez-Blanco and Kim reported that the ability of a lifespan biomarker

(P_{sod-3}::*GFP*) to predict lifespan depended on an intact insulin signaling system.

265   In animals ranging from humans to flies to worms, the manifestation of traits conferred by

specific alleles is often incomplete. Because selection acts on phenotype, incomplete penetrance

has long been recognized as a factor affecting the rate of evolution[36]. Our results suggest that in

other multicellular organisms, global differences in protein dosage might account for some

organism-to-organism phenotypic difference in discreet and complex traits, such as lifespan

(Figure 5). Variation in allelic penetrance bedevils detection of gene-trait associations in human

genome wide association studies[37] (GWAS). Differences in the phenotypic penetrance of alleles

contributes to the phenomenon called "missing heritability[38]; specifically, the summed

contributions of loci identified in GWAS studies cannot account for all of the heritability in the

disease or trait. Missing heritability might have multiple causes including, for example, epistatic

gene-gene interactions whose effect (on fitness) can now be comprehensively surveyed and

quantified (in *S. cerevisiae*[39]). We suggest that differences in protein dosages causing cells and

clones of cells to manifest different phenotypes and exist in different physiological states[7,40,41]

might also account for some of the missing heritability of traits.

Acknowledgements: Some strains were provided by the CGC, which is funded by NIH Office of

Research Infrastructure Programs (P40 OD010440). We thank the other laboratories and

scientists that have contributed to the understanding of cell-to-cell variation in gene expression

not referenced in this manuscript; a complete literature review on non-genetic, non-environmental

biological variation was beyond the scope of this study. We thank George Martin for thoughtful

discussions about the manuscript.

Funding: Funding was provided by the National Institutes of Health, National Institute on Aging

R01AG039025 to TEJ, National Institutes of Health, National Institute of General Medical

Sciences R01GM97479 to RB, the National Institutes of Health National Institute on Aging K99/

R00AG045341 to AM, P50 AG005136 to MK, a training grant from the National Institute on Aging,

T32AG000057, to support NB, and Pilot grant from the Nathan Shock Center for Excellence in

the Basic Biology of Aging to AM (NIA Grant P30AG013280 to MK).

Author Contributions: AM, TEJ and RB designed the study. NB, AM and RB analyzed and

interpreted the results. AM, RB, TEJ and MK provided funding. BS, PT, AM and SY generated

DNA and transgenic nematode strains. AM and NB conducted the imaging experiments and

performed image cytometry. AM, PT and BS conducted the flow sorting and phenotyping

experiments. NB and AM analyzed the data. AM, RB and NB wrote the initial manuscript. RB,

MK, TEJ, PT, BS, AM and NB reviewed and revised the manuscript.

Figure 1. Experimental design and analytical framework. **a.** Schematic view of the *C. elegans* body. Intestine is colored pink. The twenty cells of the intestine are organized into nine segments, called rings. **b.** Overview of reporter gene measurement scheme. In the left panel, two differently colored copies of the same gene respond to the same signaling system (Type I experiments), or, in the right panel two differently colored reporters controlled by two different promoters respond to two distinct signaling systems (Type II experiments)**. c.** Data plotting schematic. We imaged intestine cells in rings one through four (the cells that fit in a single field of view with a 40x objective) in animals on day two of adulthood, at the point when lifespan can be predicted from $P_{hsp-16.2}$*::gfp* expression levels. We then plotted expression levels of reporter pairs expressed in the intestine cells, grouped by cell type; the same cell types (e.g., cells in ring three; int3 cells) from different animals are plotted. Left scatterplot shows typical results of type I experiment. Since both genes receive the signals from the same upstream regulators and share the same downstream expression machinery, uncorrelated variation of expression results only from stochastic noise of transcription/translation or variable allele access $\eta^2(\gamma)$. Right scatterplot shows typical results of type II experiment. In the case of two different genes, uncorrelated variation of expression can result from stochastic noise of transcription/translation or variable allele access $\eta^2(\gamma)$ and variation in activation of particular signaling pathways $\eta^2(P)$. Correlated variation results from variation in shared gene expression machinery $\eta^2(G)$. **d.** A simplified version of the analytical framework derived from Colman-Lerner et al 2005 is shown; it depicts the three experimentally tractable bins into which we can attribute cell-to-cell variation in gene expression. Each bin has different underlying molecular causes. A complete, mathematically detailed description of the analytical framework is shown in Supplemental Materials Section 2.

Figure 2

Figure 2. Type I experiments reveal low intrinsic noise. a. An example of the animal producing fluorescent reporters (mEGFP and mCherry) from two identical promoters ($P_{hsp-16.2}$). We did not observe patchwork of green and red cells. Cells were nearly uniformly yellow, indicating low intrinsic noise. **b.** Scatterplot of $P_{hsp-16.2}$::GFP and $P_{hsp-16.2}$::mCherry expression in cells in ring 3 of the intestine from ten animals examined in one experiment. **c.** Average correlated and uncorrelated variations for $P_{hsp-16.2}$, P_{eft-3}, P_{vit-2} and P_{daf-21} in the intestine cells in rings 1-4. In type I experiment uncorrelated variation arises from stochastic noise of transcription/translation or variable allele access – $\eta^2(\gamma)$; correlated variation is a combined result of variation in gene expression capacity $\eta^2(G)$ and variation in pathway activation $\eta^2(P)$. * indicates statistical significance $P < 0.05$ by t-test.

Figure 3. Variation in protein expression capacity, η²(G), is the major source of inter-individual variation in expression. **a**. An example of the animal producing fluorescent reporters (mEGFP and mCherry) from two different promoters ($P_{hsp-16.2}$ and $P_{vit-.2}$). **b**. Scatterplot of $P_{hsp-16.2}$::GFP and P_{vit-2}::mCherry expression in the intestine cells in ring 3. **c**. Average stochastic noise η²(γ), variation in pathway activation η²(P) and variation in protein expression capacity η²(G) for $P_{hsp-16.2}$ promoter in cells in the intestine rings 1-4. * indicated statistical significance $p<0.05$ analyzed by one-way ANOVA with post-hoc Tukey's HSD test. **d**. Average stochastic noise η²(γ), variation in pathway activation η²(P) and variation in protein expression capacity η²(G) for P_{vit-2}, P_{eft-3}, P_{daf-21} in the intestine cells in rings 1-4. * indicated statistical significance $p<0.05$ analyzed by one-way ANOVA with post-hoc Tukey's test. **e**. Average correlated and uncorrelated variations for different promoters couples in the intestine cells in rings 1-4. Uncorrelated variation combines stochastic noise of transcription/translation or variable allele access – η²(γ) and variation in pathway activation η²(P). Correlated variation results from variation in protein expression capacity η²(G). * indicates statistical significance $p<0.05$ by t-test. **f**. G is a whole organism property. Multiple pairs of reporter genes covary; animals are generally brighter or dimmer for any pair of reporter genes we measure. Shown are individual animals that vary in expression level for both P_{mtl-2}::GFP and P_{daf-21}::mCherry (top panel), or, P_{eft-3}::BFP and P_{vit-2}::mCherry (bottom panel).

Figure 4 Differences in gene expression capacity are associated with differences in penetrance and drug resistance. a. Drug resistance conferred by Neo^R expressed from P_{rps-27} promoter correlates with expression of P_{eft-3}::GFP reporter. P_{eft-3}::GFP reporter does not predict ability to grow in the absence of neomycin (striped bars). High P_{eft-3}::EGFP expression predicts a significant difference in the ability to grow to adulthood on a high neomycin concentration (solid bars; $P < 0.05$, Paired t-test). See Tables S1 and S2 for numerical data. **b**. Penetrance of Ras/let-60 gain of function mutation *let-60(n1046)* correlates with the ability to express $P_{hsp-16.2}$::GFP reporter. Chi Square for distributions of bright or dim vs unselected $P < 0.0001$. Difference in penetrance between bright and dim $P < 0.05$, Paired t-test. See Tables S3a-c for numerical data.

418 **Figure 5 Phenotypic expression of allelic forms of Gene A, Gene B, and Gene C. a.** An
419 individual with average G and wild-type phenotypes. **b.** An individual with low G and penetrance
420 of a hypormorphic phenotype for the allele of Gene C. **c.** An individual with high G and penetrance
421 of a hypermorphic phenotype for the allele of Gene B.

Supplemental Figure 1. Type I experiments supplemental scatterplots.

Scatterplots of two identical promoters with different fluorescent protein outputs expressed from homologous chromosomes (type I experiment). Far left scatterplots show all cells measured in a given experiment. Scatterplots on the right show expression of reporters in the cells from particular intestine rings.

 Supplemental Figure 2. Type I experiments supplemental bargraphs.

Correlated and uncorrelated variation for $P_{hsp\text{-}16.2}$, $P_{eft\text{-}3}$, $P_{vit\text{-}2}$ and $P_{daf\text{-}21}$ based reporter genes in
intestine cells in rings 1-4. In type I experiment uncorrelated variation arises from stochastic noise
of transcription/translation or variable allele access – $\eta^2(\gamma)$; correlated variation is a combined
result of variation in gene expression capacity $\eta^2(G)$ and variation in pathway activation $\eta^2(P)$.

<u>Supplemental Figure 3. Evidence for splitting cells by fate for analysis in type II</u>
<u>experiments.</u>

Scatterplots of expression of two reporters from the different reporter genes, P_{hsp-17} and P_{mtl-2},
grouped by ring (one through four) or combined in one plot (right panel). Cell fate determines
ratiometric setpoint for expression of two distinct genes. When cells are not split by fate,
correlation of expression is quite low. Correlation of expression is much higher when cells are
grouped by fate.

 Supplemental Figure 4. Type II experiments supplemental scatterplots.

445 Scatterplots of expression of two distinct reporter genes (type II experiment). Far left scatterplots
446 show all cells measured in a given experiment. Scatterplots on the right show expression of
447 reporters in the cells from particular intestine rings.

  Supplemental Figure 5. Type II experiments supplemental bargraphs.

456   Stochastic noise $\eta^2(\gamma)$, variation in pathway activation $\eta^2(P)$ and variation in gene expression
457   capacity $\eta^2(G)$ for $P_{hsp-16.2}$, P_{vit-2}, P_{eft-3} and P_{daf-21} for cells in intestine rings 1-4.

Supplemental Figure 6. Type II experiments supplemental bargraphs (no promoter

splitting)

462   Correlated and uncorrelated variation for different gene pairs in cells in intestine rings 1-4.
463   Uncorrelated variation combines stochastic noise of transcription/translation or variable allele
464   access – $\eta^2(\gamma)$, and variation in pathway activation $\eta^2(P)$. Correlated variation results from
variation in gene expression capacity $\eta^2(G)$.

Supplemental Figure 7. Penetrance and Expressivity of Ras Gain of Function Mutation in
C. elegans.

Animals express EMR-1::GFP to mark cell nuclei. Wild type and _let-60(n1046)_ animals are shown
in the top two panels. Some _let-60_ mutants do not develop extra pseudo-vulvae/neoplasias
(impenetrant, compare to wild type). Other _let-60_ animals exhibit variable number of hypodermal
neoplasias.

wild-type

let-60(n1046); impenetrant

let-60(n1046); penetrant, 2 hypodermal neoplasias

let-60(n1046); penetrant, 2 hypodermal neoplasias

let-60(n1046); penetrant, 2 hypodermal neoplasias

let-60(n1046); penetrant, 3 hypodermal neoplasias

Supplementary Figure 8. Working Model for how Differences in G Might Arise. a) Genetic
experiments from Mendenhall et. al. [24] show that interindividual variation in the *hsp-16.2* reporter
gene that is a biomarker for mutation penetrance and lifespan arises from differences in insulin
signaling and depolarization of the AFD neuron pair. Thus, animals with low insulin signaling and
high neuronal depolarization would have the highest expression of the *hsp-16.2*
lifespan/penetrance biomarker and have the longest lifespans, highest penetrance of
hypermorphic phenotypes and lowest penetrance of hypomorphic phenotypes. **b)** Diagram of the
possible molecular differences in cells that might underlie global differences in protein dosage.
Solid arrows are directly supported by experiments in *C. elegans* using *hsp-16.2* reporter genes,
and arrows with question marks are possibilities supported by less direct experimental evidence
(see discussion). Here we currently favor the interpretation that decreased protein turnover, and
not protein production, results in higher protein dosage, at least in part because we showed
animals with high biomarker expression transcribe less of a AAATPase ribosome export homolog,
mac-1[13], consistent with the idea that brighter animals are not making more protein. Furthermore,
isogenic animals with lower insulin signaling, measured by the duration of DAF-16 nuclear
translocation (which would also have higher *hsp-16.2* biomarker expression[24]), produce fewer
progeny[9], also consistent with the idea that animals with high biomarker expression do not
produce more protein, but instead, maintain existing protein longer, resulting in higher effective
protein dosages.

Table S1 Growth to gravid adulthood after 72 hours development on regular NGM

Experiment	Bright adulthood	Bright total	Bright % adulthood	Dim adulthood	Dim total	Dim % adulthood	Unselected adulthood	Unselected total	Unselected % adulthood
1							37	45	82%
2									
3	46	48	96%	46	48	96%	48	49	98%
4									
5	38	39	97%	38	39	97%	36	44	82%

Table S2 Growth to gravid adulthood after 96 hours on high Neomycin concentration NGM

Experiment	Bright neo adulthood	Bright neo total	Bright neo % adulthood	Dim neo adulthood	Dim neo total	Dim neo % adulthood	Unselected neo adulthood	Unselected neo total	Unselected neo % adulthood
1	15	48	31%	2	43	5%	1	45	2%
2	26	46	57%	1	42	2%			
3	25	49	51%	15	51	29%	9	47	19%
4	25	48	52%	15	43	35%			
5	6	37	16%	2	36	6%	1	40	3%

Table S3a Number of hypodermal neoplasias

Experiment 1

# Growths	Dim		Unselected		Bright	
	# Animals	% Animals	# Animals	% Animals	# Animals	% Animals
0	10	23%	2	3%	0	0%
1	19	44%	15	21%	7	19%
2	11	26%	25	35%	15	42%
3	3	7%	24	34%	12	33%
4	0	0%	5	7%	2	6%
Total # Animals	43		71		36	

Table S3b Number of hypodermal neoplasias

Experiment 2

# Growths	Dim		Unselected		Bright	
	# Animals	% Animals	# Animals	% Animals	# Animals	% Animals
0	27	29%	31	16%	6	7%
1	30	32%	66	34%	21	26%
2	30	32%	79	40%	41	51%
3	7	7%	16	8%	11	14%
4	0	0%	5	3%	2	2%
Total # Animals	94		197		81	

Table S3c Number of hypodermal neoplasias

Experiment 3

# Growths	Dim		Unselected		Bright	
	# Animals	% Animals	# Animals	% Animals	# Animals	% Animals
0	42	22%	45	18%	18	11%
1	62	32%	79	32%	33	21%
2	68	36%	95	38%	73	46%
3	19	10%	24	10%	31	19%
4	0	0%	5	2%	4	3%
Total # Animals	191		248		159	

Supplemental Material Section 1: Materials and Methods

<u>Genome Engineering and Creation of *C. elegans* Strains</u>

We generated strains for these studies using MosSCI transgenesis[42]. We also used some

transgenes from prior studies that were multicopy; RBW2 contains a biolistically integrated

transgene (*emr-1::gfp)* and was described previously[14]. TJ2741 contains a multicopy version of

$P_{hsp-16.2}$*::gfp*, *zIS2735* [$P_{hsp-16.2}$*::egfp::T$_{unc-54}$*] V, which shows the same physiological worm to worm

variation as the several other $P_{hsp-16.2}$ reporters[14]. We used this particular version because it was

genetically compatible with the *let-60(n1046)* IV mutation and enabled our sorter to detect signal

in the heatshock-pathway-attenuating *let-60(n1046)* background. We made the DNA constructs

fusing promoter sequences to the coding sequence of different fluorescent proteins (XFPs) and

the terminator of *unc-54* by using yeast gap repair[43]. All reporter constructs carried the 5'UTR (the

upstream regulatory sequences including the promoter) of the selected genes (i.e., $P_{hsp-16.2}$) fused

to an XFP coding sequence (*megfp*, *mcherry*, *mtagbfp2*, *mneptune*) and the 3'UTR of *unc-54*. We

integrated fluorescent reporters into Chromosome II in the parent strain RBW6699 (an outcrossed

version of EG6699 without an extrachromosomal array; *ttTi5605 II; unc-119(ed9)*), unless

otherwise noted. To construct DNA for transgenes, we introduced reporter sequences into a

vector that targeted the DNA to be inserted at the chromosome II MosSCI transposon at site

ttTi5605. We then injected each construct at 50ng/µL into EG6699 animals and recovered single

copy insertions that we size and site validated by PCR. Thus, through transformation and/or

standard genetic crosses, we created the following strains:

RBW2661 (*hutSi2661[P$_{daf-21}$::megfp::T$_{unc-54}$ + Cbr-unc-119(+)]* II), RBW2642 (*hutSi2642[P$_{daf-}$*

$_{21}$::mcherry::T$_{unc-54}$ + Cbr-unc-119(+)] II), RBW2601 (*hutSi2601[P$_{hsp-16.2}$::megfp::T$_{unc-54}$ + Cbr-unc-*

119(+)] II), RBW2561 (*hutSi2561[P$_{hsp-16.2}$::mcherry::T$_{unc-54}$ + Cbr-unc-119(+)]* II), RBW2621

(*hutSi2621[P$_{vit-2}$::megfp::T$_{unc-54}$ + Cbr-unc-119(+)]* II), RBW2581 (*hutSi2581[P$_{vit-2}$::mCherry::T$_{unc-54}$*

+ Cbr-unc-119(+)] II), RBW2531 (*hutSi2531[P$_{mtl-2}$::mcherry::T$_{unc-54}$ + Cbr-unc-119(+)]* II), ARM1

(*let-60(n1046)*) IV; *jjIs699[P$_{lmn-1}$::emr-1::gfp::T$_{unc-54}$ + unc-119(+)]* ?, not II), ARM6 (*wamSi6[P$_{eft-3}$::mtagbfp2::T$_{unc-54}$ + Cbr-unc-119(+)]* II), ARM5 (*wamSi5[P$_{eft-3p}$::mneptune::T$_{unc-54}$ + Cbr-unc-119(+)]* II), RBW3211 (*hutSi3211[P$_{hsp-17}$::megfp:T$_{unc-54}$ + Cbr-unc-119(+)]* II), TJ2741 ((*zIs2735[P$_{hsp-16.2}$::gfp::T$_{unc-54}$]*) V; *let-60(n1046)* IV), RBW99 (*oxSi259[P$_{eft-3}$::megfp::T$_{unc-54}$ + Cbr-unc-119(+)]* I; *oxTi179[P$_{rps-27}$::Neo^{R}::T$_{unc-54}$ + unc-18(+)]* II); RBW2 (*zSi3002[P$_{hsp-16.2}$::mcherry::T$_{unc-54}$ + Cbr-unc-119(+)]* II; *jjIs699[P$_{lmn-1}$::emr-1::gfp::T$_{unc-54}$ + unc-119(+)]*?)

Table S4: List of crosses to make different F1 animals for Type I & II Experiments
P_{daf-21}::*gfp* (RBW2661) x P_{daf-21}::*mcherry* (RBW2642)
P_{daf-21}::*gfp* (RBW2661) x P_{vit-2}::*mcherry* (RBW2581)
$P_{hsp-16.2}$::*gfp* (RBW2601) x P_{daf-21}::*mcherry* (RBW2642)
$P_{hsp-16.2}$::*gfp* (RBW2601) x $P_{hsp-16.2}$::*mcherry* (RBW2561)
$P_{hsp-16.2}$::*gfp* (RBW2601) x P_{vit-2}::*mcherry* (RBW2581)
P_{vit-2}::*gfp* (RBW2621) x P_{vit-2}::*mcherry* (RBW2581)
P_{mtl-2}::*mcherry* (RBW2531) x P_{daf-21}::*gfp* (RBW2661)
P_{mtl-2}::*mcherry* (RBW2531) x $P_{hsp-16.2}$::*gfp* (RBW2601)
P_{mtl-2}::*mcherry* (RBW2531) x P_{vit-2}::*gfp* (RBW2621)
P_{eft-3}::*mtagbfp2* (ARM6) x P_{daf-21}::*mcherry* (RBW2642)
P_{eft-3}::*mtagbfp2* (ARM6) x $P_{hsp-16.2}$::*mcherry* (RBW2561)
P_{eft-3}::*mtagbfp2* (ARM6) x P_{mtl-2}::*mcherry* (RBW2531)
P_{eft-3}::*mtagbfp2* (ARM6) x P_{vit-2}::*mcherry* (RBW2581)
P_{eft-3}::*mtagbfp2* (ARM6) x P_{eft-3}::*mneptune* (ARM5)
P_{hsp-17}::*gfp* (RBW3211) x P_{daf-21}::*mcherry* (RBW2642)
P_{hsp-17}::*gfp* (RBW3211) x $P_{hsp-16.2}$::*mcherry* (RBW2561)
P_{hsp-17}::*gfp* (RBW3211) x P_{mtl-2}::*mcherry* (RBW2531)
P_{hsp-17}::*gfp* (RBW3211) x P_{vit-2}::*mcherry* (RBW2581)
P_{hsp-17}::*gfp* (RBW3211) x P_{eft-3}::*mneptune* (ARM5)
$P_{hsp-16.2}$::*mcherry* (RBW2561) x *emr-1::gfp* (stable RBW2)

 Culture conditions

We maintained animals on NGM plates seeded with live OP50 *E. coli* at 20°[44]. We ensured that

the stocks were not starved for at least two generations prior to use for experimentation. Single-

color reporter animals (see above) were mated to produce two-color F1 progeny for microscopy

analysis. As a result we analyzed expression of reporters driven by identical or different promoters

integrated at the identical loci on both copies of chromosome II. The only exception is RBW2

strain that carried a transcriptional reporter ($P_{hsp-16.2}$::*mCherry*) and *emr-1::gfp* fusion transgene

on different chromosomes and was maintained as a stable strain. Animals that carried $P_{hsp-16.2}$ –

driven reporters were heat shocked on the first day of adulthood by exposure to 35°C for one hour

on a solid NGM plate. All reporter strains were analyzed on the second day of adulthood. We

analyzed crosses listed in Table S4.

Mounting animals for microscopy

We prepared animals for microscopy as previously described in[14]. Briefly, we prepared a 1%

agarose pad (about 1-2 cm^2) on a standard glass microscope slide. We used worm anesthesia

solution containing 0.2% tricaine and 0.02% tetramisole (Sigma, Inc., St. Louis) in M9. We

carefully picked animals off their NGM plates and into the drop of anesthesia on the pad. We

allowed the worms to swim off the pick into the drop. After that, we covered animals with a glass

coverslip. We waited until worms ceased most active movements (about 15-25 minutes), and

placed the prepared slide on the slide holder mounted on the microscope stage.

Microscopic image acquisition

We acquired images of animals expressing multiple fluorescent proteins in a similar fashion to

Mendenhall et al, 2015[14]. Briefly, we used a 40X 1.2 NA water immersion objective and acquired

a z stack of images of each animal in one field of view. Thus, we were able to capture intestine

cells in rings one through four of the intestine. We used minimal laser power to prevent

photobleaching and heat damage to the cells. We adjusted gain and laser power to ensure that

the reporter signals fit within the dynamic range of the photomultiplier tubes and did not saturate

them. We only imaged animals lying on their left side to prevent signal loss attributable to imaging

through the gonad. We digitally rotated the field of view to arrange the animals into diagonal

orientation to fit maximal number of intestine cells in a field of view. We then acquired sequential

z-slices to span most of the intestine cells' depth. We acquired an optical slice every two

micrometers from the proximal starting point at the objective-proximal side of the intestine, and

continued imaging until we acquired signal from intestine cell nuclei in the first four rings. We set

optical slice thickness to two micrometers (all photons from plus or minus two micrometers of the

image plane). Imaging of each animal typically took two minutes at 1024x1024 resolution. Each

pixel value was the average of four samples. We imaged approximately ten animals per

experiment.

<u>Image Cytometry</u>

We first determined the orientation of the animals in images and then identified individual cells as

described previously, in Mendenhall et al, 2015[14]. As the intestine is anchored anteriorly and

posteriorly we used the known positions of the anteriormost cells to determine the identity of

adjacent cells. Once we identified a cell, we measured signal within an equatorial slice of the cell's

nucleus, as a proxy for the whole cell, as previously described; the nuclear signal is nearly

perfectly correlated with the cytoplasmic contents. We used the ImageJ software as well as

custom built Nuclear Quantification Support Plugin for nucleus segmentation and signal

quantification, we call C Entmoot (Alexander Seewald, Seewald Solutions, Inc., Vienna). The

algorithm uses a meeting of decision trees (an Ent Moot – a meeting of tree-beings; from J. R. R.

Tolkien's *The Lord of the Rings*) used to delineate a nuclear boundary based on the drop in signal

intensity, trained on user delineated images (Seewald et al, manuscript in preparation). The

values of the nuclei from binucleate cells were averaged.

<u>Sorting Animals in Flow</u>

For an extensive description of our flow cytometry methods, please see Mendenhall et al, 2015[14].

Also bear in mind that these experiments require the growth of large amounts of animals for

sorting – more so than for simply measuring the distribution of a population. Briefly, for the animals

bearing the *let-60* mutation to be sorted on $P_{hsp-16.2}$*::EGFP* expression level, we grew a

hypochlorite-synchronized population of animals to gravid adulthood, as in [8,14]. For animals we

would sort during the L1 diapause on P_{eft-3}*::GFP* expression level, we performed a hypochlorite

synchronization and allowed the animals to hatch and enter the L1 diapause in a 3-5mm deep

pool of S-basal swirling at approximately 30 rpms on an unseeded 10cm NGM petri dish. At the

start of worm flow sorting, we first flowed a sample of the population of animals to determine the

distribution of values in order to select the animals at the extremes of the distribution of fluorescent

reporter gene signal; we measured between 2-300 animals to get an estimate of the distribution

of the population. We then selected animals at the top or bottom 5-10%, or selected all animals,

and sorted between 50-250 animals per group (bright dim, unselected). Actual numbers of

animals examined were often less due to loss of animals from escape onto the side of the plate.

We hid the identity of each group from the person scoring to prevent bias in scoring. Scoring

proceeded immediately after sorting for $P_{hsp-16.2}$*::EGFP* expression levels for adult animals bearing

the *let-60(n1046)* mutation. To score, we counted the number of distinct growths emanating from

the ventral hypodermis. A distinct growth was considered a protrusion from the soma

unconnected to any other apparent protrusion; we verified that these protrusions were caused by

the growth of extra cells using strain ARM1, shown in Supplemental Figure 7. The *let-60* gain of

function mutation decreases the expression level of the heatshock response, necessitating the

use of the multicopy insertion to ensure detection of GFP signal in this genetic background. We

showed previously that the single copy and multicopy reporters have identical lifespan prediction

capabilities and the same amount of worm-to-worm variation in biomarker expression. For larvae

expressing Neo[R], we used a high magnification stereoscope and scored animals for development

to gravid adulthood (at least one fertilized embryo in the uterus) after 96 hours of development on

food, after being sorted on $P_{eft-3}::GFP$ expression while in the L1 diapause (72 hours for

experiments on regular NGM).

<u>Statistical Analysis</u>

We used Sigma Stat (Systat Software, Inc., San Jose) for statistical analyses of the bright and

dim animals we sorted. We first determined if measurements comprising each dataset were

normally distributed, then, depending on the results of those tests, used appropriate parametric

or non-parametric statistics to determine if there was significant difference in any measured

parameters. Details of specific tests are shown in figure legends. Additional details regarding

grouping of different types of measurements for calculations of η^2(G, P, or γ) and statistical

analyses of different variation bins see Supplementary Materials Section 2: Analytical Framework.

Supplemental Material Section 2: Analytical Framework

S2.1 Analytical Framework: Defining Pathway Output and Expression Capacity

For quantitative analysis we adapted the analytical framework developed previously by Coleman-Lerner et al and briefly described below. We considered the production of fluorescent proteins (e.g. GFP and mCherry) in individual cells to be a measure of activities of the promoters that regulate their expression. Furthermore, to understand cell-to-cell and animal-to-animal differences in activities of various promoters we considered the amount of fluorescent protein produced to be the product of two subsystems: "pathway" and "expression".

"Pathways" is the first subsystem. For instance, for a heat shock inducible promoter $P_{hsp-16.2}$, the input to pathway is the activation of transcription factors, e.g. HSF-1, that bind to *hsp-16.2* promoter, and the output is the activation of the inducible promoter that drives the fluorescent protein. Pathway output depends on the summed activity of upstream, DNA-bound transcription factors.

For a given cell *i,* pathway output is proportional to P_i, the time-averaged of the level of pathway activation. We separate P_i as $P_i = L_i + \lambda_i$ where L_i is the expectation value of P_i for a given cell type in a cohort of isogenic age-matched animals, and λ_i is the stochastic fluctuation term for that cell. We refer to L_i as the pathway power. The pathway "power" is a function of the activities of upstream signaling molecules and transcription factors that lead to activation of the promoter. We refer to cell-to-cell differences in P_i as "variation in pathway power", and we describe differences that result from the stochastic fluctuation term λ_i as "transmission noise". Thus, we can treat the output of the cell as being decomposed into an expectation value, which depends on the number and activity of the molecules that comprise the pathway, and a stochastic fluctuation that occurs because of the inherent randomness of the activity of the pathway, as well as its composition over the time of the experiment. Different cells may have different pathway capacities, and therefore, can have different values of L_i.

The second subsystem consists of the sequence of events from gene transcription through

protein translation. We call this subsystem "*expression*". It includes transcriptional initiation,

elongation, mRNA maturation, nuclear export, and mRNA translation and degradation (of protein

and mRNA). We use our data to measure cell-to-cell variation in expression, but we cannot isolate

the contribution of each part of the expression machinery to the overall variation.

The output of the expression subsystem is the total amount of the fluorescent reporter

protein, and the input is proportional to the level of promoter activity (the output of the pathway

subsystem). For analytical purposes, we assume here that the expression per unit of input is

independent of the level of input. For cell *i,* we describe the expression per unit input as the

variable E_i, where E_i is given by the sum of G_i and γ_i. The quantity G_i is the expectation value of

the expression per unit input for cell *i,* and this quantity may vary from cell to cell. We refer to G_i

as the "expression capacity" for cell *i*, and we refer to the cell-to-cell differences in G_i as "variation

in expression capacity". The quantity γ_i is the stochastic fluctuation that occurred in cell *i*, and we

refer to differences that result from this stochastic fluctuation term as "expression noise."

Using this model, we described the total amount of GFP, $y_{i,}$ in cell *i* as

673
$$y_i = (L_i + \lambda_i) \text{ x } (G_i + \gamma_i) \text{ x } \Delta\text{T} \tag{1}$$

where ΔT is the time of reporter protein production. For heat inducible promoter we can consider

ΔT as the time since the heat shock. For constitutive promoters like *vit-2* we cannot precisely

measure what fraction of the observed reporter protein pool was produced in the same time

frame. Thus, we started this work under "steady state" assumption that reporter protein synthesis

and degradation are balanced in young animals. In that case reporter proteins level directly mirror

activity of the promoters and expression capacity. Highly correlated expression of the examined

promoters suggests that this assumption is applicable and acceptable.

S2.2 Extracting Pathway and Expression Capacity Information from the Data

Variance and Covariance

We define the variance of a given quantity *x* for a population of cells as

684
$$\sigma^2 = \frac{1}{N}\sum(x_i - \bar{x})^2 \tag{2}$$

685
$$= \overline{(x_i - \bar{x})} \tag{3}$$

686 where *N* is the number of cells in the population. Since, we examined homologous cells of *C.*

687 *elegans* nematodes, this number is equal to the number of animals in the population. We use the

688 overbar symbol ($\bar{x}$) to represent the population average. Similarly we define the covariance,

689 *Cov(x,y)*, of two quantities *x* and *y* as

690
$$Cov(x,y) = \frac{1}{N}\sum(x_i - \bar{x})(y_i - \bar{y}) \tag{4}$$

691
$$= \overline{(x_i - \bar{x})(y_i - \bar{y})} \tag{5}$$

693 The covariance will be non-zero if x and y are correlated (or anti-correlated).

694 The correlation coefficient, ρ(x,y), is the covariance scaled by the standard deviations.

695 Specifically

697
$$\rho(x,y) = \frac{Cov(x,y)}{\sigma(x)\sigma(y)} \tag{6}$$

S2.3 Calculating Variance and Covariance of G**, γ, L, and** λ

As discussed above we assumed that in standard conditions, the average pathway output (*P*) for

a given cell has a pathway power *L* associated with it, along with a stochastic term, λ. Thus, we

have $P=L+\lambda$. Similarly, the expression subsystem of the cell (E) contains a capacity term G and

a stochastic term γ.

Thus, the amount of fluorescent reporter protein (y_i) produced for a given cell i is the

product of the terms

$$y_i = (L_i + \lambda_i) \times (G_i + \gamma_i) \times \Delta T \tag{7}$$

$$= (L_i G_i + L_i \gamma_i + \lambda_i G_i + \lambda_i \gamma_i) \times \Delta T \tag{8}$$

Since the expectations of the stochastic fluctuation terms are zero, and since their fluctuations

are uncorrelated to other terms, we expect that the population average of y_i reduces to the

population average of the quantity $L_i G_i \Delta T$.

To calculate the average of y, we re-write the product $L_i \times G_i$ in terms of population averages and

deviations from the average. For these two terms, we use a capital delta (Δ) to represent a

deviation from a population average. We get

$$L_i = \bar{L} + \Delta L_i \tag{9}$$

$$G_i = \bar{G} + \Delta G_i \tag{10}$$

718 We then calculate the population average of y to be

$$\bar{y} = \frac{\Delta}{T}\Sigma(L_i \times G_i) \tag{11}$$

$$= (\bar{L} \times \bar{G} + Cov(L, G))\Delta T \tag{12}$$

721 where we used equation (4) for the definition of covariance and the fact that $\Delta G=\Delta L=0$. Thus the

722 average number of GFP molecules is the product of L and G plus an extra term to account for

723 their correlation.

724   To calculate the variance on y, we re-write equation (7) in terms of equations (9) and (10). We

725   get

726
$$y_i = (\bar{L} + \Delta L_i + \lambda_i)\Delta T \times (\bar{G} + \Delta G_i + \gamma_i) = \qquad (13)$$

727
$$= \left(1 \frac{\Delta L_i}{\bar{L}} + \frac{\lambda_i}{\bar{L}}\right) \times \left(1 + \frac{\Delta G_i}{\bar{G}} + \frac{\gamma_i}{\bar{G}}\right) \times \bar{G} \times \bar{L} \times \Delta T \qquad (14)$$

728
$$= \left(1 + \frac{\Delta L_i}{\bar{L}} + \frac{\lambda_i}{\bar{L}}\right) \times \left(1 + \frac{\Delta G_i}{\bar{G}} + \frac{\gamma_i}{\bar{G}}\right) \times \bar{y}(1 - \frac{Cov(L,G)\Delta T}{\bar{y}}) \qquad (15)$$

729
$$\approx y \times (1 + \frac{\Delta L_i}{\bar{L}} + \frac{\lambda_i}{\bar{L}} + \frac{\Delta G_i}{\bar{G}} + \frac{\gamma_i}{\bar{G}}) \qquad (16)$$

where we used equation (12) to make the substitution for $\bar{L} \times \bar{G} \times \Delta T$ in going from equation (14)

to (15); and where we have dropped, in going from equation (15) to (16) all higher order terms of

fractional deviations from the means. These higher order terms are various products of $\frac{\Delta L_i}{\bar{L}}, \frac{\lambda_i}{\bar{L}}, \frac{\Delta G_i}{\bar{G}}$

, and $\frac{\gamma_i}{\bar{G}}$. We assume that each of the higher order terms is small relative to the lower order terms

that are retained in equation (16), and therefore we have neglected them. The term $\frac{Cov(L,G)\Delta T}{\bar{y}}$ is

second order, and has also been neglected. Discussion of the magnitude of the error introduced

by this approximation is provided in Coleman-Lerner et al.

We use equation (16) to calculate the variance on the number of GFP molecules using the

definitions of variance and covariance above. We get

742
$$\frac{\sigma^2(y)}{\bar{y}^2} = \frac{\sigma^2(L)}{\bar{L}^2} + \frac{\sigma^2(\lambda)}{\bar{L}^2} + \frac{\sigma^2(G)}{\bar{G}^2} + \frac{\sigma^2(\gamma)}{\bar{G}^2} + 2\rho(L,G)\frac{\sigma(L)}{\bar{L}}\frac{\sigma(G)}{\bar{G}} \qquad (17)$$

where $\rho(L,G)$ is the correlation coefficient between L and G.

The correlation coefficient is always between -1 and 1, and a value of 0 corresponds to no

correlation. There may be no correlation between L and G, but, for example, one possibility that

would lead to a correlation is that cells with higher expression capacity may have stronger or

weaker pathway output, perhaps because they have higher amounts of positive regulators or

negative regulators of the pathway, respectively. This would show up as a positive or negative

value for $\rho(L,G)$, respectively.

We do not expect that the stochastic fluctuations λ or γ are correlated to any other variables,

and thus the terms $\rho(L,\lambda)$, $\rho(L,\gamma)$, $\rho(\gamma,G)$, $\rho(\lambda,G)$, and $\rho(\lambda,\gamma)$ were omitted from equation (17).

All the variances in equation (17) are expressed as fractions of the means squared. We

define the variables

755
$$\eta(L) \equiv \sigma(L)/\bar{L} \qquad (18)$$

756
$$\eta(\lambda) \equiv \sigma(\lambda)/\bar{L} \qquad (19)$$

757
$$\eta(G) \equiv \sigma(G)/\bar{G} \qquad (20)$$

758
$$\eta(\gamma) \equiv \sigma(\gamma)/\bar{G} \qquad (21)$$

The quantity η is simply the width of different distributions expressed as a fraction of their means.

For the stochastic variables λ and γ, we will refer to $\eta(\lambda)$ and $\eta(\gamma)$ as the "noise" associated with

those quantities, and for the non-stochastic variables L and G, we will refer to $\eta(L)$ and $\eta(G)$ as

the "variation."

We quantified cell-to-cell variation in system output using "normalized variance" ($\eta^2 = \sigma^2/\mu^2$)

rather than "noise strength" (σ^2/μ), a measure others have used to describe deviations from purely

stochastic Poisson-type biological processes. We used normalized variance for a number of

reasons. First, we found that most of the cell-to-cell differences in system behavior we reported

are not due to stochastic differences in signal transmission or gene expression, as described in

the main text. Second, use of η^2 allowed examination of different amounts of variation in terms of

the fraction of the mean. Third, and most important, because η^2 is unitless, it allowed direct

comparison of different measurements, for example, from different fluorescent proteins, and it

allowed the definition of total variation as the sum of individual sources of variation plus additional

terms to account for correlations.

We re-write equation (17) in terms of η as

$$\eta^2(y) = \eta^2(P) + \eta^2(G) + \eta^2(\gamma) + 2\rho(L,G)\eta(L)\eta(G)$$

(22)

We have written $\eta^2(P)$ for $\eta^2(L)+\eta^2(\lambda)$ since the data will not be able to distinguish the cell-to-cell

variation in pathway output from the noise or stochastic fluctuations. The term $\eta^2(P)$ is the variation

in average pathway output per unit time, which, because pathway output is given by $P\Delta T$ and ΔT

is the same for every cell, is identical to variation in pathway output.

To separate $\eta(G)$ from $\eta(P)$ and $\eta(\gamma)$ in the data from the two-promoter, two-color experiments,

discussed further below, we introduce the quantity

$$Z(GFP, mCherry) = \frac{\eta^2(GFP) + \eta^2(mCherry)}{2} -$$

$$\eta(GFP)\eta(mCherry)\rho(GFP, mCherry)$$

$$= \frac{\eta^2(GFP) + \eta^2(mCherry)}{2} - \frac{Cov(GFP, mCherry)}{\overline{GFP} \times \overline{mCherry}} \qquad (23)$$

The quantity Z is the average variance divided by the mean square of the two fluorescence

signals GFP and mCherry with the correlated part subtracted out. Thus, Z is a measure of the

uncorrelated part of the GFP vs mCherry scatter plot. Depending of the type of the experiment

(two identical promoters or two different promoters) Z encompasses distinct sources of variation.

S2.4 Two Color Variants Driven by the Same Promoter and Gene Expression Noise

We calculate the correlation coefficient between GFP and mCherry reporter proteins for the

case that both genes are expressed in the same cell and with the same promoter as

$$\rho(GFP, mCherry) \equiv \frac{1}{\sigma(GFP)\sigma(mCherry)} \frac{1}{N} \sum_i \Delta(GFP_i) \Delta(mCherry_i)$$

(24)

$$= \frac{1}{\eta(GFP)\eta(mCherry)} \frac{1}{N} \sum_i \left(\frac{\Delta L_i}{\bar{L}} + \frac{\lambda_i}{\bar{L}} + \frac{\Delta G_i}{\bar{G}} + \frac{\gamma_{GFP,i}}{\bar{G}}\right)\left(\frac{\Delta L_i}{\bar{L}} + \frac{\lambda_i}{\bar{L}} + \frac{\Delta G_i}{\bar{G}} + \frac{\gamma_{mCherry,i}}{\bar{G}}\right)$$

(25)

where we have used equation (16) for the deviations from the mean, $\Delta(GFP_i)$, and an analogous

equation for $\Delta(mCherry_i)$. Since the GFP gene has the same promoter as mCherry, ΔL_i is the

same for both color variants. The quantity λ_i is the same since stochastic fluctuations in pathway

output occur upstream of the promoters (with the exception of the binding of the transcription

factors to each individual copy of the reporter genes, see the end of this section), and the quantity

ΔG_i is the same since the proteins are being expressed in the same cell. The stochastic

fluctuations in gene expression, γ_i, however, are different for the two color variants and we write

them as $\gamma_{GFP,i}$ and $\gamma_{mCherry,i}$ for GFP and mCherry respectively.

The uncorrelated terms drop out of equation (25) when we perform the population average.

We do not expect the stochastic fluctuations to be correlated with any other terms, and equation

(25) becomes

$$\rho(GFP, mCherry) = \frac{\eta^2(P) + \eta^2(G) + 2\rho(L,G)\eta(L)\eta(G)}{\eta(GFP)\,\eta(mCherry)} \qquad (26)$$

Where $\eta^2(P) = \eta^2(L) + \eta^2(\lambda)$ as discussed above.

The quantity $Z(GFP, mCherry)$ as defined in equation (23), gives the contribution of the

uncorrelated part of the expression of the two color variants to the average η^2 of the two colors.

Only the gene expression noise is uncorrelated, and we get

$$Z(GFP, mCherry) = \frac{\eta^2(\gamma_{GFP}) + \eta^2(\gamma_{mCherry})}{2} \qquad (27)$$

$Z(GFP, mCherry)$ is the average of gene expression noise of the two promoters. Since the same

promoters are driving the GFP and mCherry genes, we expect equal levels of mRNA for the two

variants on average, and therefore the gene expression noise is the same for the GFP and

mCherry signal. We get $\eta^2(\gamma_{GFP}) = \eta^2(\gamma_{mCherry}) = \eta^2(\gamma)$ and therefore, from equation (27),

$Z(GFP, mCherry) = \eta^2(\gamma)$. Thus, equation (23) gives direct measurement of gene expression

noise γ when applied to an experiment with two identical promoters:

$$\eta^2(\gamma) = \frac{\eta^2(GFP) + \eta^2(mCherry)}{2} - \frac{Cov(GFP, mCherry)}{\overline{GFP} \times \overline{mCherry}} \qquad (28)$$

<u>We used this equation (28) for practical calculations of gene expression noise of individual</u>

<u>promoters.</u>

We note that for the experiment described above, GFP and mCherry were not driven by

the same promoter, but rather by identical *copies* of the same promoter. Therefore, stochasticity

in the binding of transcription factors to the DNA and in the subsequent activation of transcription

contributes to the uncorrelated part of the total variation. In our *ad hoc* model, these molecular

steps are part of the pathway subsystem, and therefore fluctuations in this step should contribute

to the transmission noise, $\eta^2(\lambda)$. However, due to experimental limitations of the type of

experiment we have just described, the noise caused by these molecular steps is, instead,

included in the measure of gene expression noise $\eta^2(\gamma)$.

S2.5 Two Color Variants Driven by Different Promoters and Pathway Variation

For this case, the calculation of *ρ(GFP,mCherry)* is the same as in equation (25) above except

that the pathway activity is different for the two color variants since different promoters are driving

the GFP and mCherry genes. If we assume that the two terms *L* and *λ* are uncorrelated for the

two promoters then we get

$$\rho(GFP, mCherry) = \frac{\eta^2(G) + \rho(L_{GFP}, G)\eta(L_{GFP})\eta(G) + \rho(L_{mCherry}, G)\eta(L_{mCherry})\eta(G)}{\eta(GFP)\,\eta(mCherry)}$$ (29)

where we have written *L_{GFP}* and *L_{mCherry}* for capacities of the pathways that lead to the activation

of the GFP and mCherry gene respectively. The uncorrelated part of the average η^2 will now

include a contribution from the pathway variation; and the quantity *Z(gfp, mCherry)* includes this

extra contribution. We get

$$Z(GFP, mCherry) = \frac{\eta^2(P_{GFP}) + \eta^2(P_{mCherry})}{2} + \frac{\eta^2(\gamma_{GFP}) + \eta^2(\gamma_{mCherry})}{2}$$ (30)

where $\eta^2(\lambda_{GFP})$ is the transmission noise for the pathway that leads to GFP activation and $\eta^2(P_{GFP})$

$= \eta^2(L_{GFP}) + \eta^2(\lambda_{GFP})$, and similarly for $\eta^2(\lambda_{mCherry})$ and $\eta^2(P_{mCherry})$.

Thus, when uncorrelated expression variation for two independent promoters includes an average

variation in pathways activation and gene expression variations for each promoter. The latter can

be calculated from type I experiments. Knowing expression variations $\eta^2(\gamma)$ for each promoter

and uncorrelated variation Z from equation (23) allows to calculate of pathway variations $\eta^2(P)$.

To split contribution of variation of each pathway we need to perform three experiments with

pairwise analysis of three promoters and then solve a system linear equations to calculate $\eta^2(P)$

of each pathway.

Below we give an example with expression of three promoters, *vit-2*, eft-3, daf-21, in cells Int3V

and Int3D.

$Z\left(P_{eft-3} :: BFP, P_{vit-2} :: mCherry\right) = \dfrac{\eta^2(P_{eft-3})+\eta^2(P_{vit-2})}{2} + \dfrac{\eta^2\left(\gamma_{P_{eft-3::BFP}}\right)+\eta^2\left(\gamma_{P_{vit-2::mCherry}}\right)}{2}$

(31)

$Z\left(P_{eft-3} :: BFP, P_{daf-21} :: mCherry\right) = \dfrac{\eta^2(P_{eft-3})+\eta^2(P_{daf-21})}{2} + \dfrac{\eta^2\left(\gamma_{P_{eft-3::BFP}}\right)+\eta^2\left(\gamma_{P_{daf-21::mCherry}}\right)}{2}$

(32)

$Z\left(P_{daf-21} :: GFP, P_{vit-2} :: mCherry\right) = \dfrac{\eta^2(P_{daf-21})+\eta^2(P_{vit-2})}{2} + \dfrac{\eta^2\left(\gamma_{P_{daf-21::GFP}}\right)+\eta^2\left(\gamma_{P_{vit-2::mCherry}}\right)}{2}$

(33)

Average gene expression noise of two promoters, e.g. $\dfrac{\eta^2\left(\gamma_{P_{eft-3::BFP}}\right)+\eta^2\left(\gamma_{P_{vit-2::mCherry}}\right)}{2}$ is

calculated from type I experiments and equation (28). It is therefore a known value for each cell

type and promoter. We calculate quantity Z directly from equation (23). Quantity Z and average

expression noise therefore give a particular number of each of the equations above. We can

rewrite them in more suitable form to calculate pathways variations

880
$$\frac{\eta^2(P_{eft-3}) + \eta^2(P_{vit-2})}{2} = A_1$$

881
(34)

882
$$\frac{\eta^2(P_{eft-3}) + \eta^2(P_{daf-21})}{2} = A_2$$

883
(35)

884
$$\frac{\eta^2(P_{daf-21}) + \eta^2(P_{vit-2})}{2} = A_3$$

885
(36)

886 Where A_1, A_2 and A_3 are known values. To calculate pathway variation for a particular promoter,

887 e.g. eft-3, we will sum up (34) and (35) and subtract (36):

888
$$\frac{\eta^2(P_{eft-3}) + \eta^2(P_{vit-2})}{2} + \frac{\eta^2(P_{eft-3}) + \eta^2(P_{daf-21})}{2} - \frac{\eta^2(P_{daf-21}) + \eta^2(P_{vit-2})}{2} = A_1 + A_2 - A_3$$

889
(37)

890
$$\frac{\eta^2(P_{eft-3}) + \eta^2(P_{vit-2}) + \eta^2(P_{eft-3}) + \eta^2(P_{daf-21}) - \eta^2(P_{daf-21}) - (P_{vit-2})}{2} = A_1 + A_2 - A_3$$

891
(38)

892
$$\frac{2\eta^2(P_{eft-3})}{2} = A_1 + A_2 - A_3 \qquad (39)$$

893
$$\eta^2(P_{eft-3}) = A_1 + A_2 - A_3 \qquad (40)$$

Thus, we have got one estimate of $\eta^2(P_{eft-3})$. To obtain it we used results of the experiments

involving P_{eft-3}, P_{vit-2} and P_{daf-21} reporters in pairwise manner. We say that these 3 pairs of

experiments form a 'triangle'. However, we could choose another triangle consisting of

experiments with P_{eft-3}, P_{vit-2} and $P_{hsp-16.2}$ reporters, or a triangle of P_{eft-3}, $P_{hsp-16.2}$ and P_{daf-21}

reporters. To get a better estimate of the true $\eta^2(P_{eft-3})$ value we have calculated it from all three

possible triangles and averaged. We did the same for other reporters too. The resulting $\eta^2(P)$

values are shown in Figure 3 bar graphs.

Similarly, we can calculate pathway variation for other promoters for which we have

measured their expression noise. In case when we did not measure expression noise, e.g. hsp-

17 promoter, we did calculate pathway variation for these promoters as well. In these cases we

have limited our analysis with comparison of total correlated variation of two promoters to their

uncorrelated variation. In all cases we have observed that correlated variation dominates over

uncorrelated term:

$$\frac{Cov(P_{hsp-17}::GFP, P_{eft-3}::mNeptune)}{\overline{P_{hsp-17}::GFP} \times P_{eft-3}::mNeptune} \gg Z(P_{hsp-17}::GFP, P_{eft-3}::mNeptune)$$

References:

1   Kirkwood, T. B. & Finch, C. E. *Chance, Development, and Aging.*, (Oxford University Press, 2000).
2   Timofeeff-Ressovsky, N. W. Über den Einfluss des Genotypus auf das phanotypen Auftreten
eines einzelnes. *Gens. J. Psychol. Neurol.* **31**, 305-310 (1925).
3   Romaschoff, D. D. Uber die Variabilität in der Manifestierung eines erblichen Merkmales
(Abdomen abnormalis) bei Drosophila funebris. *F. J. Psychol. Neurol.* **31**, 323-325 (1925).
4   Laubichler, M. D. & Sarkar, S. in *Mutaliflg Concepts. Evolving Disciplines: Genetics, Medicine and*
Society (eds L.S. Parker & R.A. Ankeny) Ch. 4, 63-85 (Kluwer Academic Publishers, 2002).
5   Mortimer, R. K. & Johnston, J. R. Life span of individual yeast cells. *Nature* **183**, 1751-1752
(1959).
6   Kirkwood, T. B. & Finch, C. E. Ageing: the old worm turns more slowly. *Nature* **419**, 794-795
(2002).
7   Rea, S. L., Wu, D., Cypser, J. R., Vaupel, J. W. & Johnson, T. E. A stress-sensitive reporter predicts
longevity in isogenic populations of Caenorhabditis elegans. *Nature genetics* **37**, 894-898 (2005).
8   Mendenhall, A. R. *et al.* Expression of a single-copy hsp-16.2 reporter predicts life span. *The*
journals of gerontology **67**, 726-733 (2012).
9   Casanueva, M. O., Burga, A. & Lehner, B. Fitness trade-offs and environmentally induced
mutation buffering in isogenic C. elegans. *Science* **335**, 82-85 (2011).
10   Colman-Lerner, A. *et al.* Regulated cell-to-cell variation in a cell-fate decision system. *Nature*
437, 699-706 (2005).
11   Chang, H. H., Hemberg, M., Barahona, M., Ingber, D. E. & Huang, S. Transcriptome-wide noise
controls lineage choice in mammalian progenitor cells. *Nature* **453**, 544-547 (2008).
12   Seewald, A. K., Cypser, J., Mendenhall, A. & Johnson, T. Quantifying phenotypic variation in
isogenic Caenorhabditis elegans expressing Phsp-16.2::gfp by clustering 2D expression patterns.
PloS one **5**, e11426 (2010).
13   Cypser, J. R. *et al.* Predicting longevity in C. elegans: fertility, mobility and gene expression.
Mechanisms of ageing and development **134**, 291-297, doi:10.1016/j.mad.2013.02.003 (2013).
14   Mendenhall, A. R., Tedesco, P. M., Sands, B., Johnson, T. E. & Brent, R. Single Cell Quantification
of Reporter Gene Expression in Live Adult Caenorhabditis elegans Reveals Reproducible Cell-
Specific Expression Patterns and Underlying Biological Variation. *PloS one* **10**, e0124289,
doi:10.1371/journal.pone.0124289 (2015).
15   Libina, N., Berman, J. R. & Kenyon, C. Tissue-specific activities of C. elegans DAF-16 in the
regulation of lifespan. *Cell* **115**, 489-502 (2003).
16   Sanchez-Blanco, A. & Kim, S. K. Variable pathogenicity determines individual lifespan in
Caenorhabditis elegans. *PLoS genetics* **7**, e1002047 (2011).
17   Elowitz, M. B., Levine, A. J., Siggia, E. D. & Swain, P. S. Stochastic gene expression in a single cell.
Science **297**, 1183-1186, doi:10.1126/science.1070919 (2002).
18   Swain, P. S., Elowitz, M. B. & Siggia, E. D. Intrinsic and extrinsic contributions to stochasticity in
gene expression. *Proc Natl Acad Sci U S A* **99**, 12795-12800, doi:10.1073/pnas.162041399
(2002).
19   Raser, J. M. & O'Shea, E. K. Control of stochasticity in eukaryotic gene expression. *Science* **304**,
1811-1814 (2004).
20   Han, M. & Sternberg, P. W. let-60, a gene that specifies cell fates during C. elegans vulval
induction, encodes a ras protein. *Cell* **63**, 921-931 (1990).
21   Bahar, R. *et al.* Increased cell-to-cell variation in gene expression in ageing mouse heart. *Nature*
441, 1011-1014 (2006).

22 Prahlad, V. & Morimoto, R. I. Neuronal circuitry regulates the response of Caenorhabditis
elegans to misfolded proteins. *Proc Natl Acad Sci U S A* **108**, 14204-14209,
doi:10.1073/pnas.1106557108 (2011).
23 Prahlad, V., Cornelius, T. & Morimoto, R. I. Regulation of the cellular heat shock response in
Caenorhabditis elegans by thermosensory neurons. *Science* **320**, 811-814 (2008).
24 Mendenhall, A., Crane, M. M., Tedesco, P. M., Johnson, T. E. & Brent, R. Caenorhabditis elegans
Genes Affecting Interindividual Variation in Life-span Biomarker Gene Expression. *The journals*
of gerontology, doi:10.1093/gerona/glw349 (2017).
25 Walker, G. A. & Lithgow, G. J. Lifespan extension in C. elegans by a molecular chaperone
dependent upon insulin-like signals. *Aging cell* **2**, 131-139 (2003).
26 Baird, N. A. *et al.* HSF-1-mediated cytoskeletal integrity determines thermotolerance and life
span. *Science* **346**, 360-363, doi:10.1126/science.1253168 (2014).
27 Burga, A., Casanueva, M. O. & Lehner, B. Predicting mutation outcome from early stochastic
variation in genetic interaction partners. *Nature* **480**, 250-253 (2011).
28 Lin, C. Y. *et al.* Transcriptional amplification in tumor cells with elevated c-Myc. *Cell* **151**, 56-67,
doi:10.1016/j.cell.2012.08.026 (2012).
29 Nie, Z. *et al.* c-Myc is a universal amplifier of expressed genes in lymphocytes and embryonic
stem cells. *Cell* **151**, 68-79, doi:10.1016/j.cell.2012.08.033 (2012).
30 Pickett, C. L., Breen, K. T. & Ayer, D. E. A C. elegans Myc-like network cooperates with
semaphorin and Wnt signaling pathways to control cell migration. *Developmental biology* **310**,
226-239, doi:10.1016/j.ydbio.2007.07.034 (2007).
31 Raj, A., Rifkin, S. A., Andersen, E. & van Oudenaarden, A. Variability in gene expression underlies
incomplete penetrance. *Nature* **463**, 913-918 (2010).
32 Kirstein-Miles, J., Scior, A., Deuerling, E. & Morimoto, R. I. The nascent polypeptide-associated
complex is a key regulator of proteostasis. *The EMBO journal* **32**, 1451-1468,
doi:10.1038/emboj.2013.87 (2013).
33 Dhondt, I. *et al.* FOXO/DAF-16 Activation Slows Down Turnover of the Majority of Proteins in C.
elegans. *Cell reports* **16**, 3028-3040, doi:10.1016/j.celrep.2016.07.088 (2016).
34 Depuydt, G., Shanmugam, N., Rasulova, M., Dhondt, I. & Braeckman, B. P. Increased Protein
Stability and Decreased Protein Turnover in the Caenorhabditis elegans Ins/IGF-1 daf-2 Mutant.
The journals of gerontology **71**, 1553-1559, doi:10.1093/gerona/glv221 (2016).
35 Visscher, M. *et al.* Proteome-wide Changes in Protein Turnover Rates in C. elegans Models of
Longevity and Age-Related Disease. *Cell reports* **16**, 3041-3051,
doi:10.1016/j.celrep.2016.08.025 (2016).
36 Schmalhausen, I. *Factors of evolution; the theory of stabilizing selection*. (Blakiston, 1949).
37 Hirschhorn, J. N. & Daly, M. J. Genome-wide association studies for common diseases and
complex traits. *Nat Rev Genet* **6**, 95-108, doi:10.1038/nrg1521 (2005).
38 Manolio, T. A. *et al.* Finding the missing heritability of complex diseases. *Nature* **461**, 747-753,
doi:10.1038/nature08494 (2009).
39 Baryshnikova, A., Costanzo, M., Myers, C. L., Andrews, B. & Boone, C. Genetic interaction
networks: toward an understanding of heritability. *Annu Rev Genomics Hum Genet* **14**, 111-133,
doi:10.1146/annurev-genom-082509-141730 (2013).
40 Mendenhall, A., Driscoll, M. & Brent, R. Using measures of single-cell physiology and
physiological state to understand organismic aging. *Aging cell* **15**, 4-13, doi:10.1111/acel.12424
(2016).
41 Lehner, B. Genotype to phenotype: lessons from model organisms for human genetics. *Nat Rev*
Genet **14**, 168-178, doi:10.1038/nrg3404 (2013).

42   Frokjaer-Jensen, C., Davis, M. W., Ailion, M. & Jorgensen, E. M. Improved Mos1-mediated
transgenesis in C. elegans. *Nature methods* **9**, 117-118 (2012).
43   Sands, B. *et al.* A toolkit for DNA assembly, genome engineering and multicolor imaging for C.
elegans. *Translational Medicine of Aging* **X**, x-xx In Press, doi:In Press (2018).
44   Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71-94 (1974).

1007