1 Using camouflage for conservation: colour change in

2 juvenile European lobster

3 Sara Mynott[1]*, Carly Daniels[2], Stephen Widdicombe[3], Martin Stevens[1]

4 [1] Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Penryn, UK; [2] The

5 National Lobster Hatchery, Padstow, UK; [3]Plymouth Marine Laboratory, Plymouth, UK

6 *Corresponding author: s.mynott@exeter.ac.uk

7 Abstract

Changes in coloration enable animals to refine their camouflage to match different visual

backgrounds, responding to spatial and temporal changes in the environment. Such

plasticity provides ecological benefits and could be exploited to support conservation or

stock enhancement efforts. One application could be to ensure that hatchery-reared animals,

reared to stock wild populations, are appropriately matched to their environment on release.

Following crashes in wild European lobster (*Homarus gammarus*) populations, hatcheries

were established across Europe to restock or enhance local lobster stocks by rearing

juveniles through their most vulnerable stages, then releasing them into the wild. However,

little consideration has yet been given to their camouflage and the implications of matching

individuals to the appearance of their release site. This study assesses to what extent

juvenile lobsters can change appearance to match their background and whether hatchery

practices can be altered to enhance lobster camouflage and potentially survivorship in the

wild. We test this by switching individuals between black or white backgrounds in the

laboratory and monitoring their coloration over time. We show that juvenile lobsters are

capable of changes in luminance (perceived lightness) to better match their background over

2-3 weeks. These changes potentially correspond to improved camouflage, based on a

model of predator (fish, *Pollachius pollachius*) vision. Over a longer period (5 weeks),

lobsters maintained on either black or white backgrounds converged on the same darker coloration, likely due to ontogenetic (developmental) changes. Overall, our results indicate that hatcheries could rear lobsters on backgrounds that better match the habitat into which they will be released, but such manipulations should be considered in the context of ontogenetic changes and the timing of release. We recommend rearing individuals on darker, more natural-coloured substrates throughout their early benthic phase in order to maximise the benefits of camouflage on release into the wild. This could potentially increase juvenile survival during the critical days immediately after their release. The findings highlight the potential benefits of using camouflage and colour change in stocking programmes and aquaculture alike.

Keywords

Camouflage, European lobster, nature-based solutions, plasticity, predation, stocking, ontogenetic, colour change

Introduction

Background matching is one of the most widely used anti-predator defence strategies in nature (Stevens & Merilaita, 2011), with many species using colours and patterns to match their surroundings and avoid detection and predation. In a wide range of taxa, both terrestrial and aquatic, camouflage can be achieved through plastic changes in appearance (Stuart-Fox & Moussalli 2011; Duarte et al. 2017). This enables animals to respond to either fast and unpredictable changes in the visual environment through rapid (seconds and minutes) colour change, or slower and more predictable environmental change with gradual (hours, days, and weeks) appearance changes (Caro et al. 2016). One of the most widely studied groups, particularly in terms of the mechanisms and functions of colour change, are the crustaceans (Abramowitz, 1937; Duarte, Flores, & Stevens, 2017). Many crustacean species employ camouflage to conceal themselves from predators and several have been shown to

change colour to better match their background, including crabs (Hultgren & Stachowicz,

2008; Russell & Dierssen, 2015), prawns (Duarte et al. 2017; Brown 1935) and isopods

(Kleinholz, 1937; Oguro, 1962), leading to phenotype-environment matches (Stevens et al.,

2015; Todd, Oh, Loke, & Ladle, 2012). It is likely that many juvenile crustaceans are

phenotypically plastic with the ability to match the environment in which they settle (Palma &

Steneck, 2001; Todd, Qiu, & Chong, 2009). The benefits of understanding camouflage and

how it works have long been realised in applied areas such as the military and art and

design (Behrens, 2002; Talas, Baddeley, & Cuthill, 2017), and colour change applications

are growing in other fields, such as biomimicry (Isapour & Lattuada, 2018) and animal

welfare (Rotllant et al., 2003). However, an important application, seldom considered, is how

an understanding of camouflage and colour change may be harnessed for conservation. For

example, how colour change for camouflage could be harnessed in captive breeding and

stocking programmes to improve post release survival and therefore stocking success.

Stocking, including stock enhancement and restocking, is used around the world to meet

conservation needs and future seafood demands (Born, Immink, & Bartley, 2004; Kitada,

2018; Okuzawa et al., 2008). Such programs improve and sustain capture fisheries *via* the

release of cultured juveniles into the wild. There are some 180 cultivated marine species

worldwide (Kitada, 2018). These are reared in captivity (when they are most vulnerable to

predation) before release into the wild (at a larger, more resilient size) in order to overcome

challenges in recruitment and restore spawning stock biomass (Bell, Bartley, Lorenzen, &

Loneragan, 2006). While extensive research has been carried out to ensure the viability of

released individuals, little consideration has been given to their ecology on release (Molony,

Lenanton, Jackson, & Norriss, 2003), in particular to the development of appropriate anti-

predator defence behaviours. Unlike their wild counterparts, hatchery-reared juveniles are

naïve to predators, which often puts them at greater risk (Olla & Davis, 1989). The lack of

habitat enrichment limits shelter-seeking behaviour in released European lobster, *Homarus*

gammarus and those reared alone are slow to seek shelter (Aspaas et al., 2016), making

them more vulnerable to predation (Mercer et al., 2002). While training individuals alongside conspecifics can mitigate this (Agnalt, Grefsrud, Farestveit, & Jørstad, 2017), European lobsters are often reared individually to prevent agonistic interactions between individuals (Carlberg, Olst, & Ford, 2009). With this vulnerability to predation in mind, it is important to maximise anti-predator defences prior to release.

European lobsters have attracted significant attention for restocking. Their populations collapsed during the 1970s (van der Meeren, 2005) and several hatcheries have been established across Europe to help restock the natural population – for the benefit of both conservation and fisheries. Restocking is achieved in hatchery aquaria by rearing larvae through their planktonic and early benthic phases, keeping them in captivity during a time when, in the wild, they are most vulnerable to predation (van der Meeren, 2000, 2005). Clawed lobster stocking programs use variable approaches, releasing lobsters from stage IV onwards into suitable sites to mature in their natural environment. Release strategies consider available shelter, as well as physical and oceanographic conditions required by European lobster (Galparsoro, Borja, Bald, Liria, & Chust, 2009). However, the visual components of the habitat (colour, pattern) have been neglected to date.

Juvenile European lobsters have seldom been observed in the wild, but those reared in hatchery aquaria show considerable individual variation in coloration with few resembling wild-caught adults (Fig. 1). This presents a potential problem on release into the wild, as individuals are likely to be conspicuous to predators if poorly matched to their surroundings (Stevens & Merilaita, 2011). Predation rates are highest in the first 24 hours following release (van der Meeren, 2000), making this period a critical point in the restocking program. Given that there is considerable variation in the colour of juvenile lobsters, knowing whether they can adapt to match the habitat, and whether aquarium colour can be altered to enhance habitat matching, should help to enhance their survival on release. To date, no work has tested the capacity for this species to change colour and research on lobster coloration is limited to the influence of feed and genetics on pigmentation in the American lobster, *H.*

americanus (Lim, Sakurai, Sugihara, & Kittaka, 1997; Tlusty & Hyland, 2005). Given the

prevalence of background matching in crustaceans (Umbers et al. 2014; Duarte et al 2017;

Caro 2018), it seems reasonable that juvenile lobsters could be capable of changing colour

for the purposes of camouflage.

By placing hatchery-reared lobsters on artificial backgrounds and monitoring their coloration

over time, we were able to test the ability of juvenile lobsters to change brightness in

response to their surroundings. This study quantifies the capacity of lobsters to match their

background using a model of fish vision (pollack, *Pollachius pollachius*), allowing us to

assess coloration from the predator's perspective. Ultimately, this paper aims to assess

whether altered hatchery practices can be used to improve lobster camouflage, and by

implication survival, following release. We used a 35-day-long laboratory experiment to test

the hypothesis that lobsters will change their coloration to better match their background

over time, and that this will result in improvements in camouflage, when modelled to the

visual systems of relevant predators.

Figure 1 Variation in European lobster coloration (stage IV juveniles following collection from

the National Lobster Hatchery (NLH), Padstow, UK). Individuals here are housed within an

Aquahive® (Shellfish Hatchery Systems Ltd, Orkney, UK), a compartmentalised, stacked

system used to rear juvenile European lobsters in captivity.

Materials and methods

Husbandry

A total of 80 juvenile lobsters (Stage IV, approximately 1 month old) were sourced from the

National Lobster Hatchery (NLH) in Padstow (UK) and transported to the aquarium facility of

the University of Exeter (Penryn Campus, UK). Lobsters were transported in an Aquahive®

tray (Fig. 1), enclosed within a heavy-duty plastic bag containing seawater (32 +/- 2 ‰

salinity) and pure oxygen. This was secured within a 50 x 50 x 100 cm cool box to stabilise

temperature during transport. Experimental glass tanks were set up to mimic hatchery

rearing conditions as closely as possible. Water was supplied using uPVC push-fit pipe

drilled with 1.5 mm diameter holes to allow aerated water to flow into each container. Before

being recirculated, water was filtered (Classic 350 filter; Eheim GmbH & Co., Deizisau,

Germany) and run through a heating system (DC300 Aquarium Chiller; D-D The Aquarium

Solution Ltd., Ilford, UK) in order to maintain tank temperature (18-19 °C). Each tank was

filled to 125 L with artificial seawater. Saltwater was made up to 32 +/- 2 ‰ salinity using

Instant Ocean Salt (Instant Ocean, Blacksburg, Virginia) and dechlorinated water. Tanks

were topped up with freshwater to compensate for evaporation during the course of the

experiment. Tanks were prepared 48 hours before lobster arrival to allow them to reach a

stable temperature.

Lobsters were housed individually (to prevent harm from aggressive interactions) in

containers made from square uPVC gutter pipe (65 mm by 65 mm) cut to 60 mm lengths

and covered with a mesh base to allow water through (Fig. S1). All containers were fixed to

corner braces and suspended within the tank above waterproof paper corresponding to the

experimental treatment (a black or white background). Each juvenile was fed one formulated

pellet (1.5 mm diameter – formula undisclosed) daily, in accordance with the NLH feeding

regime. Any uneaten food was removed the following day. Tanks were cleaned and half the

water was changed twice weekly to limit the build up of bacteria and algae in the tanks. The

light regime was set to 12 hours of light and 12 hours of darkness, with lights on from 07:30

to 19:30. Where handling was required, lobsters were pipetted between containers using a

modified turkey baster, following the approach used by NLH.

Experiment protocol

To determine the capacity of lobsters for background matching, juvenile lobsters were

randomly assigned to either a black or a white compartment for a 2.5 or 5-week period.

Individuals were photographed to determine their initial luminance (lightness as perceived by

a particular predator), and then photographed at various intervals (described below) to

monitor changes in coloration over time. 80 individuals were used in the study; 40 of which were used to determine short (3 hours) and medium (2.5 weeks) term changes in coloration, and 40 of which were used to determine changes in luminance over the longer term (5 weeks). The allocation of individuals to each treatment is described in Table 1.

Initial photographs of all lobsters were taken 24 hours after introduction to the tank to prevent undue stress on individuals following transport. Further photos were taken after 2.5 weeks to quantify colour change over the medium-term. At this point, 40 individuals (20 of those on a white background and 20 of those on a black background) were transferred to the alternative treatment (black to white and *vice versa*) to assess plasticity in colour change over a further 2.5 weeks as well as any short-term changes in coloration. The switched group were photographed after 3 hours and 2.5 weeks. The group that remained on their original background were photographed after 2.5 and 5 weeks to assess longer-term changes in coloration. Lobsters were photographed in water within a 10 mm deep PTFE chamber, under diffuse lighting conditions to minimise stress during data collection. Lobsters that moulted on the same day that photography was scheduled were not photographed to prevent damage to the new exoskeleton during handling.

Table 1 Experimental design showing the treatments and recording intervals used in each experiment, and the number of juvenile European lobsters allocated to each treatment. Note that the same individuals were used in both the short- and medium-term experiments.

Experiment	Aim	Black	White	Photography intervals
Short-term	Quantify short-term change	n = 20	n = 20	0 and 3 hours
Medium-term	Quantify plasticity & medium-term change	n = 20	n = 20	0 and 2.5 weeks
Long-term	Monitor ontogenetic & longer-term change	n = 20	n = 20	0, 2.5 and 5 weeks

Quantifying colour change

Both colour change and camouflage was quantified with respect to predator vision (Stevens, Párraga, Cuthill, Partridge, & Troscianko, 2007; Stevens, Stoddard, & Higham, 2009; Troscianko & Stevens, 2015). Photos were taken using a Nikon D7000 SLR, fitted with a 60 mm quartz lens (Coastal Optics). A 400-700 nm, Baader Venus U filter was fixed in front of the lens, allowing relevant wavelengths to be recorded. All photos were taken under simulated daylight conditions, achieved using an arc lamp (Ventronic) equipped with a daylight 65 bulb. All lobsters were photographed against a white background in a clear PTFE chamber filled to 1 cm deep with artificial seawater. A translucent PTFE diffuser was placed between the photography chamber and the light source to ensure lighting was even. Two grey standards (7 % and 93 % reflectance) were used in every photo to account for any variation in illumination over time. The camera white balance was set to manual and the aperture was kept constant between photos. All photographs were taken during daylight hours to account for any potential coloration with day-night cycles, as has been observed in other crustaceans (Keeble & Gamble, 1904; Powell, 1962; Stevens, Rong, & Todd, 2013).

Image analysis

To establish how lobsters would be perceived by predators, the images were mapped to fish vision using an established polynomial mapping technique (Troscianko & Stevens, 2015), which yields predicted cone catch data that are highly accurate compared to data obtained *via* reflectance spectrometry (Pike, 2011; Stevens & Cuthill, 2006; Troscianko & Stevens, 2015). Images were analysed in ImageJ (National Institute of Health, NIH) using the Multispectral Image Calibration and Analysis Toolbox (Troscianko & Stevens, 2015). The longwave channel was used to calculate luminance (lightness according to a specific visual system) as potentially perceived by a dichromatic predatory fish, using spectral sensitivity data from the pollack, *Pollachius pollachius* (Shand, Partridge, Archer, Potts, & Lythgoe, 1988). Average luminance was calculated for each lobster by selecting a rectangular region

of interest (ROI) that covered as much of the carapace as possible while excluding the white

patterning observed at the edges of the cephalothorax. This patterning develops with age

and was excluded from the ROI so that plasticity in coloration could be quantified.

To assess the lobsters' level of camouflage, a widely-implemented predator discrimination

model, based on receptor-noise limited discrimination, was used (Siddiqi, Cronin, Loew,

Vorobyev, & Summers, 2004; Vorobyev & Osorio, 1998). This calculates Just Noticeable

Differences (JNDs) between one object (the lobster) and another (the background) to predict

how easily the two can be distinguished, according to a particular visual system (here,

pollack). Luminance JNDs were calculated using a modified (log) version of the Vorobyev

and Osorio model for luminance discrimination (Siddiqi et al., 2004; Vorobyev & Osorio,

1998), using a Weber fraction of 0.05, which is thought to be appropriate for many fish

(Olsson, Lind, Kelber, & Simmons, 2018).

Statistical analysis

All statistical analyses were carried out using R version 3.31 (R Core Team 2016). Linear

mixed effects models were used to assess the effect of time, background colour and their

interaction on luminance, and to assess the effect of time on camouflage (expressed in

JNDs). Lobster ID was included as a random effect in all models in order for any potential

temporal autocorrelation to be accounted for. For all models, the following approach was

used: linear mixed models were fitted by restricted maximum likelihood (REML), with

Kenward-Roger approximations to degrees of freedom using the lme4 package (Bates,

Maechler, Bolker, & Walker, 2015). The minimum adequate model was determined by

successively removing non-significant terms, starting with the highest order terms in the

model. Analysis of variance (ANOVA) was used to determine which model best explained

the variation in the response. Assumptions of normality were supported for all datasets,

which were checked through visual inspections of quantile-quantile plots, residual

distributions and residual vs fitted values plots. In addition, Welch two sample t-tests were

used to evaluate differences in mean luminance change over the experimental period.

Results

Short-term change

Juvenile lobsters show no significant response to their background in the short-term; the only

significant term was the intercept (GLM: t = 20.46, d.f. = 42, p < 0.001)), resulting in a null

model (see Table S1 for model output). After 3 hours of exposure (Fig. S2), there was no

significant interaction between time and background (ANOVA: Chi-sq = 3.07, d.f. = 1, p =

0.080), no significant effect of background on coloration (ANOVA: Chi-sq = 0.02, d.f. = 1, p =

0.889) and no significant effect of time (ANOVA: Chi-sq = 2.40, d.f. = 1, p = 0.121), resulting

in a null model.

Medium-term change

During the first 2.5 weeks in the laboratory, lobsters were observed to darken over time

(GLM: t = 16.53, d.f. = 39, p <0.001), (Fig. 2a), with individuals on a darker background as

dark as those on a light one, on average (t-test: t = 1.06, d.f. = 37, p = 0.298), (Fig. 2b).

However, neither the interaction between time and background (ANOVA: Chi-sq = 1.16, d.f.

= 1, p = 0.282) nor background as an independent variable (ANOVA: Chi-sq = 1.32, d.f. = 1,

p = 0.250) had a significant effect on lobster coloration. Model parameters are detailed in

Table 2.

253 (a)

(b)

(c)

(d)

255 **Figure 2** Medium-term luminance change observed in juvenile lobsters placed on a black

256 (dark grey) and white (light grey) background. Panels (a) and (b) show the initial luminance

257 change; panels (c) and (d) show the plastic change following transfer to the alternative

background treatment (those initially on black were transferred to white and *vice versa*). Panels (a) and (c) show the variation in luminance across the experimental population, where central lines are medians, boxes are interquartile ranges and whiskers are 95% quartiles. Panels (b) and (d) show the mean luminance change observed following exposure to the background treatments, together with standard errors. Luminance is presented according to pollack vision. Model parameters are detailed in Table 2.

Table 2 Parameter estimates from the minimum adequate model describing the initial (a) and plastic (b) changes in juvenile lobster coloration for individuals allocated to a black or white background over the medium term. Initial luminance change describes the change in luminance when placed on the first background (black or white), plastic luminance change describes the change in luminance when placed on the second background (black to white and *vice versa*). Linear mixed models were fitted by restricted maximum likelihood (REML) using the lme4 package (Bates et al., 2015). The Kenward-Roger approximation for degrees of freedom was used to determine p-values. Lobster ID and length were included as random effects.

(a) Luminance change: medium-term (initial)

Source	Estimate	SE	d.f.	t	p
Intercept	0.1700	0.0049	67	34.55	<0.001
Time	-0.0892	0.0054	39	16.53	<0.001
Model formula	lmer(Luminance ~ Time + (1 | ID) + (1 | Length)				

(b) Luminance change: medium-term (plastic)

Source	Estimate	SE	d.f.	t	p
Intercept	0.0882	0.0051	66	17.29	<0.001
BackgroundWhite	-0.0150	0.0072	66	2.07	0.042
Time	-0.0003	0.0004	31	0.74	0.463
BackgroundWhite	0.0012	0.0005	31	2.33	0.023
Model formula	lmer(Luminance ~ Background + Time + Background:Time + (1 | ID) + (1 | Length))				

Despite initially darkening over time (Fig. 2a,b), lobsters showed some plasticity in their ability to change luminance when switched to the alternative background type, with a significant interaction between time and background (GLM: t = 2.33, d.f. = 31, p = 0.026),

(Fig. 2c). Those on a black background darkened whereas those on a light background became lighter, on average (t-test: t = 2.45, d.f. = 32, p = 0.020), (Fig. 2d). The full model containing the interaction between time and background was significantly better than simpler alternatives (ANOVA: Chi-sq = 5.27, d.f. = 1, p = 0.022); model parameters are detailed in Table 2b.

The initial darkening observed in both treatments (Fig 2a,b) corresponds to a significant decrease in camouflage over time for lobsters on a white background (GLM: t = 10.14, d.f. = 19, p < 0.001), (Fig. 3a) and a significant increase in camouflage for those initially placed on a black background (GLM: t = 13.65, d.f. = 19, p < 0.001), (Fig. 3b). In both cases, the full model was a significantly better explainer for the variation in camouflage than simpler alternatives (ANOVA$_{white}$: Chi-sq = 40.68, d.f. = 1, p = <0.001; ANOVA$_{black}$: Chi-sq = 62.55, d.f. = 1, p = <0.001).

Plastic changes in coloration (Fig. 2c,d) corresponded to a small but significant increase in camouflage over time for those on a white background (GLM: t = 2.45, d.f. = 36, p = 0.020), (Fig. 3c), but no change in camouflage for those on a black background (GLM: t = 13.38, d.f. = 18, p < 0.001), (Fig. 3d), with no significant change in background matching over time (ANOVA: Chi-sq = 0.46, d.f. = 1, p = 0.499), likely because individuals were already quite dark (Fig. 3). Individuals allocated to a black background were already well matched to their surroundings at the start of the plastic trial (JND close to one, Fig. 3d). Model parameters are detailed in Table 3 (see Table S2 for mean JNDs).

Table 3 Parameter estimates from the minimum adequate models describing the initial (a, b)

and plastic (c, d) changes in juvenile lobster camouflage according to Just Noticeable

Differences (JNDs) individuals allocated to a white background (a, c) and black background

(b, d) over the medium term. Estimates are for individuals that were placed on either a black

or a white background, then switched to the alternative background type (i.e. plastic

camouflage change). Linear mixed models were fitted by restricted maximum likelihood

(REML) using the lme4 package (Bates et al., 2015). The Kenward-Roger approximation for

degrees of freedom was used to determine p-values. Lobster ID was included as a random

effect.

(a) Camouflage: medium-term, white background (initial)

Source	Estimate	SE	d.f.	t	p
Intercept	12.2531	0.1593	31	76.93	<0.001
Time	1.6705	0.1647	19	10.14	<0.001
Model formula	lmer(JND ~ Time + (1 | ID))				

(b) Camouflage: medium-term, black background (initial)

Source	Estimate	SE	d.f.	t	p
Intercept	2.933	0.1147	26	25.69	<0.001
Time	-1.898	0.1391	19	13.65	<0.001
Model formula	lmer(JND ~ Time + (1 | ID))				

(c) Camouflage: medium-term, white background (plastic)

Source	Estimate	SE	d.f.	t	p
Intercept	14.22	0.0866	36	164.17	<0.001
Time	-0.0171	0.0070	36	2.45	0.020
Model formula	lmer(JND ~ Time + (1 | ID))				

(d) Camouflage: medium-term, black background (plastic)

Source	Estimate	SE	d.f.	t	p
Intercept	1.335	0.1147	18	13.38	<0.001
Model formula	lmer(JND ~ Time + (1 | ID))				

(a) Time (days)

(b) Time (days)

(c) Time (days)

(d) Time (days)

(e) (f)

Figure 3 Variation in juvenile lobster camouflage, according to pollack vision, over time.

Figures on the left and right are for individuals allocated to a white (light grey) or black (dark

grey) background, respectively. The change in detectability according to Just Noticeable

Differences (JNDs) is shown for both initial changes in coloration (a, b) and for individuals

placed on the alternative background (c, d). Note that a decline in JND corresponds to an

increase in camouflage. Central lines are medians, boxes are interquartile ranges and

whiskers are 95% quartiles. Both (e) and (f) are example individuals, showing level of

lightness/darkness attained after 18 days on their respective backgrounds. Model

parameters are detailed in Table 3.

Long-term change

Lobsters that were not presented with a new background darkened significantly throughout

their time in the laboratory (GLM: t = 10.35, d.f. = 107, p < 0.001), (Fig. 4). However, neither

the interaction between time and background (ANOVA: Chi-sq = 0.04, d.f. = 1, p = 0.834)

nor background as an independent variable (ANOVA: Chi-sq = 2.43, d.f. = 1, p = 0.119) had

a significant effect on luminance. Model parameters are detailed in Table 4. Lobsters

darkened by the same extent over a 35-day period, regardless of their background (t-test: t =

0.06, d.f. = 31, p = 0.956), (Fig. 4 insert). This darkening resulted in an increase in

detectability for those on a light background (GLM: t = 8.38, d.f. = 34, p <0.001), (Fig. 5a)

and decrease in detectability for those on a black one (GLM: t = 6.34, d.f. = 56, p < 0.001),

(Fig. 5b). Model parameters are summarised in Table 5.

Table 4 Parameter estimates from the minimum adequate model describing the change in

juvenile lobster luminance (lightness according to pollack vision) over a 35-day period in the

laboratory. Linear mixed models were fitted by restricted maximum likelihood (REML) using

the lme4 package (Bates et al., 2015). The Kenward-Roger approximation for degrees of

freedom was used to determine p-values. Lobster ID and length were included as random

effects.

Luminance change: long-term

Source	Estimate	SE	d.f.	t	p
Intercept	0.1470	0.0051	90	28.87	<0.001
Time	-0.0023	0.0002	107	10.35	<0.001
Model formula	lmer(Luminance ~ Time + (1 | ID) + (1 | Length))				

Figure 4 Long-term luminance change in juvenile lobsters according to pollack vision.

Boxplots show the variation across the experimental population at each time point, where

central lines are medians, boxes are interquartile ranges and whiskers are 95% quartiles.

Insert shows the mean change in luminance observed for each background treatment from

day 0 to day 35. Dark boxes and points correspond to individuals on a black background;

light boxes and points correspond to individuals on a white background. Error bars in insert

show standard errors. Error bars in insert show standard errors.

(a)

(b)

Figure 5 Change in juvenile lobster camouflage over the long-term. The change in

detectability according to Just Noticeable Differences (JNDs) is shown for (a) juvenile

lobsters placed on a white background (light grey) and (b) for those on black (dark grey).

Central lines are medians, boxes are interquartile ranges and whiskers are 95% quartiles.

JNDs are calculated according to pollack vision. Note that a decline in JND corresponds to

an increase in camouflage. Model parameters are detailed in Table 5.

Table 5 Parameter estimates from the minimum adequate models describing the changes in

camouflage according to Just Noticeable Differences (JNDs) for individuals allocated to a

white background (a) and black background (b) over the long term. JNDs are according to

pollack vision. Linear mixed models were fitted by restricted maximum likelihood (REML)

using the lme4 package (Bates et al., 2015). The Kenward-Roger approximation for degrees

of freedom was used to determine p-values. Lobster ID was included as a random effect.

(a) Camouflage: long-term, white background

Source	Estimate	SE	d.f.	t	p
Intercept	12.68	0.1265	41	100.3	<0.001
Time	0.0432	0.0052	34	8.38	<0.001
Model formula	lmer(JND ~ Time + (1 | ID))				

(b) Camouflage: long-term, black background

Source	Estimate	SE	d.f.	t	p
Intercept	2.410	0.1489	56	16.19	<0.001
Time	-0.0444	0.0067	56	6.64	<0.001
Model formula	lmer(JND ~ Time + (1 | ID))				

Discussion

Juvenile European lobsters show no significant change in coloration in response to their

background in the short-term (3 hours, Fig. S2). The absence of rapid camouflage in this

species explains, in part, why mortality due to predation is so great in the first 24 hours

following release into the wild (van der Meeren, 2000). Given that individuals do not change

their coloration rapidly, if they are reared on substrates that are not representative of the

natural environment, it is likely that they will stand out in the wild, making them an easy

target for predators (van der Meeren, 2000). When transferred to a new substrate,

individuals became lighter on a white background and darker on a black one over 2-3 weeks

(Fig. 2c,d, Fig. 3). This response shows that juvenile European lobsters are capable of some

degree of plastic background matching and, over time, have the potential to become

increasingly difficult to discern from their surroundings (Fig. 3). However, some of the

changes in discriminability are small (less than 1 JND for those on a white background), so

may not always be perceptible to predator visual systems. More effective changes in

camouflage may be achieved or through the use of more natural substrates during rearing or

with earlier exposure to the background. While black and white features like pebbles occur in

the environment, the backgrounds used here were artificial and not fully representative of the

range of substrates that would be encountered in nature. It is entirely possible that when

faced with more naturalistic substrates that lobsters show a greater degree of plasticity and

matching. Greater background matching could also occur if individuals are exposed to

different substrates as soon as they settle from the plankton, when, in accordance with their

life history, there would be the greatest need for habitat matching. These gradual changes in

response to the background (Fig. 2c,d) are likely to correspond to improvements in survival

(Troscianko et al. 2016; Duarte et al. 2017), but because colour change in European lobster

is relatively slow, this capacity to adapt will only benefit hatchery-reared juveniles if they are

pre-conditioned for the habitat they are released into.

In both the medium- and long-term experiment, individuals darken initially, with most of that

darkening occurring within the first 2.5 weeks (Fig. 2, Fig. 4). Individuals that remained on

the same background became darker over a longer time period (5 weeks), regardless of

their background treatment (Fig. 5). Given that darkening occurs over the longer-term, it

likely reflects an ontogenetic change in coloration (Booth, 1990). Ontogenetic changes in

coloration may reflect a number of drivers, including relaxed selection pressure on coloration

(e.g. fewer predators and less need for camouflage), movement into new habitats with age,

or changes in camouflage strategy (Booth 1990; Todd et al. 2009; Wilson et al. 2007; Duarte

et al. 2017). In many animals, predation risk changes with age as individuals gain a size

refuge from predation. The risk is greatest for younger, smaller individuals and lessens as

individuals grow and develop weapons, (such as chelae) for defence, or a more robust body

morphology (Anderson, Spadaro, Baeza, & Behringer, 2013; Todd et al., 2009). As such,

there may be stronger selective pressure for effective anti-predator defences, like

camouflage, in early life stages (Duarte et al. 2017). Ontogenetic changes in coloration can

also coincide with changes in habitat use throughout development (Hultgren & Stachowicz,

2010; Todd et al., 2009), as different colours and patterns may provide better camouflage in

different habitats. Lobsters are initially pelagic in their early life stages and while in the

brightly light surface oceans it pays to be lightly coloured or even translucent, when juveniles

reach the deeper, darker seabed they will need to be correspondingly dark, so it stands to

reason that settlement could act as a trigger for darkening. Long-term darkening is slightly,

and temporarily, offset by the plastic response to their background in the medium-term, with

those on a white background becoming less dark than those on black (Fig. 4). This indicates

that rearing benthic stages in white containers in the hatchery could impede individuals from

darkening as they would in a more natural (darker) habitat.

Planktonic lobster larvae can disperse over a large area (Huserbråten et al., 2013), with

single populations extending up to 230 kilometres (Ellis, Hodgson, Daniels, Collins, &

Griffiths, 2017). Given this extensive larval dispersal, plasticity in coloration (in order to

match the local environment after settling) has clear advantages. The ability to change

coloration to match their surroundings means juvenile European lobsters have the capacity

for adaptive camouflage, a strategy that could afford them protection from predators in a

variable environment (Stevens et al., 2015; Todd, Briers, Ladle, & Middleton, 2006) and

across the range of habitats into which they might settle. This has implications for the

survival of hatchery-reared juveniles post-release: ecological drivers of colour change could

be harnessed to provide nature-based solutions for stocking. If reared in an environment that

resembles their release site, individuals may be more likely to evade predator detection

when released into the wild. Furthermore, the ability to fine-tune their brightness to match a

specific habitat may have added benefits for individuals as they age, given the site-fidelity

exhibited by European lobsters (Bannister & Addison, 1998; Bannister, Addison, & Lovewell,

1994). However, in the longer-term, ontogenetic changes are likely to override such

plasticity, as adult European lobsters are much less variable than juveniles. Consequently,

the timing of rearing individuals on different backgrounds and the timing of release needs

careful thought in order to maximise any applied benefits of camouflage for survival.

These findings also have implications for shellfish product coloration, particularly given

recent advances in the potential for aquaculture of this species (Halswell, Daniels, &

Johanning, 2016). Individuals with darker, more striking coloration often attract a higher price

(Parisenti et al. 2011a). If the plasticity in coloration seen here exists in later developmental

stages, then responses to background colour and brightness could be harnessed to produce

higher value shellfish for market. While some individuals in our study converged on the same

brightness in the longer-term (Fig. 4), on other backgrounds and with differences in diet

more plasticity may be possible. These considerations apply not only to lobster, but also

other crustacean aquaculture, as many commercial species are capable of colour change,

including giant tiger prawns, *Penaeus monodon*, and Pacific white shrimp, *Litopenaeus*

vannamei, and are more valuable when exhibiting a particular colour (Parisenti et al. 2011b;

Wade et al. 2015). Understanding and applying camouflage offers significant advantages to

stocking and aquaculture programs, as individuals reared in environments that appear

natural are likely to be more natural in colour. Natural coloration will likely afford individuals

reared for conservation and stock enhancement with protection from predators on release.

Releasing individuals with effective anti-predator defences is vital to ensure the success of

stocking programs. Similarly, as either more natural or darker coloration can enhance

product value, aquaculturists stand to benefit from modifying rearing environments to

promote such traits, particularly as the economic viability of aquaculture ventures depends

on product value (Parisenti, Beirão, Tramonte, et al., 2011; Wade et al., 2015).

This study demonstrates the capacity of early benthic phase lobster to match their

background, and shows they are capable of changing brightness, which is a strong

contributor to visual detectability. These findings have direct relevance to restocking

programmes, which aim to maximise the survival of captive-reared lobster on release.

Building on this, further work should test the capacity of early benthic phase larvae to match

the colour of ecologically relevant substrates, such as sand, mud, cobbles and seaweed

(Botero & Atema, 1982; Linnane et al., 2001; Linnane, Mazzoni, & Mercer, 2000).

Understanding the implications of hue as well as brightness on lobster coloration will greatly

inform our understanding of phenotypic plasticity in this species, and determine the extent to

which rearing environments can be modified to increase the chances of survival in the

natural environment. We recommend that hatchery-reared lobsters are reared on

backgrounds that match their release site prior to release into the wild, and that precise

timings should consider when plasticity is highest during development. Given the high

predation rate experienced immediately following release, such measures may improve the

survivorship of released juveniles. Other conservation projects and aquaculture programmes

may reap similar benefits from careful selection of artificial substrates and altering the colour

of rearing environments to enhance camouflage and, hence, survival.

Authors' contributions

476   SM designed the study, carried out data collection and analysis, completed statistical

477   analysis, and drafted the manuscript. CD participated in study design and development of

478   hatchery-relevant protocols and helped draft the manuscript. SW and MS participated in

479   study design and helped to draft and refine the manuscript.

Acknowledgements

481   Thanks to National Lobster Hatchery staff for their support throughout this project, particular

482   thanks to Charlie Ellis for his help initiating the work. SM was supported by a Natural

483   Environment Research Council Studentship within the GW4+ Doctoral Training Partnership

484   [NE/L002434/1].

Data accessibility

486   Data will be archived in the Dryad Digital Repository on publication of this manuscript.

References

Abramowitz, A. A. (1937). The chromatophorotropic hormone of the crustacea: standardization, properties and physiology of the eye-stalk glands. *The Biological Bulletin*, *72*(3). doi:doi.org/10.2307/1537694

Agnalt, A. L., Grefsrud, E. S., Farestveit, E., & Jørstad, K. E. (2017). Training camp—A way to improve survival in European lobster juveniles? *Fisheries Research*, *186*, 531–537. doi:10.1016/j.fishres.2016.09.021

Anderson, J. R., Spadaro, A. J., Baeza, J. A., & Behringer, D. C. (2013). Ontogenetic shifts in resource allocation: colour change and allometric growth of defensive and reproductive structures in the Caribbean spiny lobster *Panulirus argus*. *Biological Journal of the Linnean Society*, *108*(1), 87–98. doi:10.1111/j.1095-8312.2012.01998.x

Aspaas, S., Grefsrud, E. S., Fernö, A., Jensen, K. H., Trengereid, H., & Agnalt, A.-L. (2016). An enriched environment promotes shelter-seeking behaviour and survival of hatchery-produced juvenile European lobster (*Homarus gammarus*). *PLOS ONE*, *11*(8), e0159807. doi:10.1371/journal.pone.0159807

Bannister, R. C. A., & Addison, J. T. (1998). Enhancing lobster stocks: a review of recent European methods, results, and future prospects. *Bulletin of Marine Science*, *19*, 369–387.

Bannister, R. C. A., Addison, J. T., & Lovewell, S. R. J. (1994). Growth, movement, recapture rate and survival of hatchery-reared lobsters (*Homarus gammarus* (Linnaeus, 1758)) released into the wild on the English east coast. *Crustaceana*, *67*(672), 156–172.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48. doi:10.18637/jss.v067.i01

Behrens, R. A. (2002). *False Colors: Art, Design, and Modern Camouflage*. Bobolink Books.

doi:10.1353/mod.2003.0035

Bell, J. D., Bartley, D. M., Lorenzen, K., & Loneragan, N. R. (2006). Restocking and stock

enhancement of coastal fisheries: potential, problems and progress. *Fisheries*

Research, *80*(1), 1–8. doi:10.1016/J.FISHRES.2006.03.008

Booth, C. L. (1990). Evolutionary significance of ontogenetic colour change in animals.

Biological Journal of the Linnean Society, *40*(2), 125–163. doi:10.1111/j.1095-

8312.1990.tb01973.x

Born, A. F., Immink, A. J., & Bartley, D. M. (2004). Marine and coastal stocking: global status

and information needs. In D. M. Bartley & K. M. Leber (Eds.), *Marine Ranching* (429th

ed., pp. 1–18). FAO Fisheries Technical Paper.

Botero, L., & Atema, J. (1982). Behavior and substrate selection during larval settling in the

lobster *Homarus Americanus*. *Journal of Crustacean Biology*, *2*(1), 59–69.

doi:10.2307/1548113

Brown, F. A. (1935). Color changes in Palaemonetes. *Journal of Morphology*, *57*(2), 317–

333. doi:10.1002/jmor.1050570202

Carlberg, J. M., Olst, J. C., & Ford, R. F. (2009). Potential for the communal rearing of the

nephropid lobsters (*Homarus spp.*). *Proceedings of the World Mariculture Society*,

10(1–4), 840–853. doi:10.1111/j.1749-7345.1979.tb00082.x

Caro, T. (2018). The functional significance of coloration in crabs. *Biological Journal of the*

Linnean Society, *124*(1), 1–10. doi:10.1093/biolinnean/bly021

Duarte, R. C., Flores, A. A. V., & Stevens, M. (2017). Camouflage through colour change:

mechanisms, adaptive value and ecological significance. *Philosophical Transactions of*

the Royal Society of London B: Biological Sciences, *372*(1724).

doi:10.1098/rstb.2016.0342

Duarte, R. C., Flores, A. A. V., Vinagre, C., & Leal, M. C. (2017). Habitat-dependent niche

partitioning between colour morphs of the algal-dwelling shrimp *Hippolyte*

obliquimanus. *Marine Biology*, *164*(11), 215. doi:10.1007/s00227-017-3247-1

Ellis, C., Hodgson, D., Daniels, C., Collins, M., & Griffiths, A. (2017). Population genetic

structure in European lobsters: implications for connectivity, diversity and hatchery

stocking. *Marine Ecology Progress Series*, *563*, 123–137. doi:10.3354/meps11957

Galparsoro, I., Borja, Á., Bald, J., Liria, P., & Chust, G. (2009). Predicting suitable habitat for

the European lobster (*Homarus gammarus*), on the Basque continental shelf (Bay of

Biscay), using Ecological-Niche Factor Analysis. *Ecological Modelling*, *220*(4), 556–

567. doi:10.1016/J.ECOLMODEL.2008.11.003

Halswell, P., Daniels, C. L., & Johanning, L. (2016). Sea based container culture (SBCC)

hydrodynamic design assessment for European lobsters (*Homarus gammarus*).

Aquacultural Engineering, *74*, 157–173. doi:10.1016/J.AQUAENG.2016.08.003

Hultgren, K. M., & Stachowicz, J. J. (2008). Alternative camouflage strategies mediate

predation risk among closely related co-occurring kelp crabs. *Oecologia*, *155*(3), 519–

528. doi:10.1007/s00442-007-0926-5

Hultgren, K. M., & Stachowicz, J. J. (2010). Size-related habitat shifts facilitated by positive

preference induction in a marine kelp crab. *Behavioral Ecology*, *21*(2), 329–336.

doi:10.1093/beheco/arp192

Huserbråten, M. B. O., Moland, E., Knutsen, H., Olsen, E. M., André, C., & Stenseth, N. C.

(2013). Conservation, spillover and gene flow within a network of Northern European

Marine Protected Areas. *PLoS ONE*, *8*(9), e73388. doi:10.1371/journal.pone.0073388

Isapour, G., & Lattuada, M. (2018). Bioinspired stimuli-responsive color-changing systems.

Advanced Materials, *30*(19), 1707069. doi:10.1002/adma.201707069

Kitada, S. (2018). Economic, ecological and genetic impacts of marine stock enhancement
and sea ranching: A systematic review. *Fish and Fisheries*, *19*(3), 511–532.
doi:10.1111/faf.12271

Kleinholz, L. H. (1937). Studies in the pigmentary system of Crustacea I. Color changes and
diurnal rhythm in *Ligia baudiniana*. *Biological Bulletin*, *72*(1), 24–36.

Lim, B. K., Sakurai, N., Sugihara, T., & Kittaka, J. (1997). Survival and growth of the
American lobster *Homarus americanus* fed formulated feeds. *Bulletin of Marine
Science*, *61*(1), 159–163.

Linnane, A., Ball, B., Mercer, J. P., Browne, R., Meeren, G. van der, Ringvold, H., …
Munday, B. (2001). Searching for the early benthic phase (EBP) of the European
lobster: a trans-European study of cobble fauna. *Hydrobiologia*, *465*(1/3), 63–72.
doi:10.1023/A:1014547618888

Linnane, A., Mazzoni, D., & Mercer, J. P. (2000). A long-term mesocosm study on the
settlement and survival of juvenile European lobster *Homarus gammarus* L. in four
natural substrata. *Journal of Experimental Marine Biology and Ecology*, *249*(249), 51–
64. doi:doi.org/10.1016/S0022-0981(00)00190-8

Mercer, J. P., Bannister, R. C. A., van der Meeren, G. I., Debuse, V., Mazzoni, D., Lovewell,
S., … Ball, B. (2002). An overview of the LEAR (Lobster Ecology and Recruitment)
project: results of field and experimental studies on the juvenile ecology of *Homarus
gammarus* in cobble. *Marine and Freshwater Research*, *52*(8), 1291.
doi:10.1071/MF01216

Molony, B. W., Lenanton, R., Jackson, G., & Norriss, J. (2003). Stock enhancement as a
fisheries management tool. *Reviews in Fish Biology and Fisheries*, *13*(4), 409–432.
doi:10.1007/s11160-004-1886-z

584 Oguro, C. (1962). On the physiology of melanophores of the marine isopod, *Idotea japonica*.

585  III. The role of the eyes in background response. *Crustaceana*, *4*, 85–92.

586  doi:10.2307/20102466

587 Okuzawa, K., Maliao, R. J., Quinitio, E. T., Buen-Ursua, S. M. A., Lebata, M. J. H. L.,

588  Gallardo, W. G., … Primavera, J. H. (2008). Stock enhancement of threatened species

589  in Southeast Asia. *Reviews in Fisheries Science*, *16*(1–3), 394–402.

590  doi:10.1080/10641260701678496

591 Olla, B. L., & Davis, M. W. (1989). The role of learning and stress in predator avoidance of

592  hatchery-reared coho salmon (*Oncorhynchus kisutch*) juveniles. *Aquaculture*, *76*(3–4),

593  209–214. doi:10.1016/0044-8486(89)90075-6

594 Olsson, P., Lind, O., Kelber, A., & Simmons, L. (2018). Chromatic and achromatic vision:

595  parameter choice and limitations for reliable model predictions. *Behavioral Ecology*,

596  *29*(2), 273–282. doi:10.1093/beheco/arx133

597 Palma, A. T., & Steneck, R. S. (2001). Does variable coloration in juvenile marine crabs

598  reduce risk of visual predation? *Ecology*, *82*(10), 2961–2967. doi:10.1890/0012-

599  9658(2001)082[2961:DVCIJM]2.0.CO;2

600 Parisenti, J., Beirão, L. H., Tramonte, V. L. C. G., Ourique, F., da Silveira Brito, C. C., &

601  Moreira, C. C. (2011). Preference ranking of colour in raw and cooked shrimps.

602  *International Journal of Food Science & Technology*, *46*(12), 2558–2561.

603  doi:10.1111/j.1365-2621.2011.02781.x

604 Parisenti, J., Luiz, Beirão, H., Mouriño, J. L., Vieira, F. do N., … Maraschim, M. (2011).

605  Effect of background colour on shrimp pigmentation. *Boletim Do Instituto de Pesca Sao*

606  *Paulo*, *37*(372), 177–182.

607 Pike, T. W. (2011). Using digital cameras to investigate animal colouration: estimating

608  sensor sensitivity functions. *Behavioral Ecology and Sociobiology*, *65*(4), 849–858.

doi:10.1007/s00265-010-1097-7

Rotllant, J., Tort, L., Montero, D., Pavlidis, M., Martinez, M., Wendelaar Bonga, S. E., &

Balm, P. H. M. (2003). Background colour influence on the stress response in cultured

red porgy *Pagrus pagrus*. *Aquaculture*, *223*(1–4), 129–139. doi:10.1016/S0044-

8486(03)00157-1

Russell, B. J., & Dierssen, H. M. (2015). Use of hyperspectral imagery to assess cryptic

color matching in sargassum associated crabs. *PLOS ONE*, *10*(9), e0136260.

doi:10.1371/journal.pone.0136260

Shand, J., Partridge, J. C., Archer, S. N., Potts, G. W., & Lythgoe, J. N. (1988). Spectral

absorbance changes in the violet/blue sensitive cones of the juvenile pollack,

Pollachius pollachius. *Journal of Comparative Physiology A*, *163*, 699–703.

doi:10.1007/BF00603854

Siddiqi, A., Cronin, T. W., Loew, E. R., Vorobyev, M., & Summers, K. (2004). Interspecific

and intraspecific views of color signals in the strawberry poison frog *Dendrobates*

pumilio. *The Journal of Experimental Biology*, *207*, 2471–2485. doi:10.1242/jeb.01047

Stevens, M., Broderick, A. C., Godley, B. J., Lown, A. E., Troscianko, J., Weber, N., &

Weber, S. B. (2015). Phenotype–environment matching in sand fleas. *Biology Letters*,

11(8), 20150494. doi:10.1098/rsbl.2015.0494

Stevens, M., & Cuthill, I. C. (2006). Disruptive coloration, crypsis and edge detection in early

visual processing. *Proceedings of the Royal Society B: Biological Sciences*, *273*, 2141–

2147. doi:10.1098/rspb.2006.3556

Stevens, M., Lown, A. E., & Wood, L. E. (2014). Color change and camouflage in juvenile

shore crabs *Carcinus maenas*. *Frontiers in Ecology and Evolution*, *2*, 1–14.

doi:10.3389/fevo.2014.00014

Stevens, M., & Merilaita, S. (2011). *Animal camouflage mechanmisms and function*.

Cambridge University Press.

Stevens, M., Párraga, C. A., Cuthill, I. C., Partridge, J. C., & Troscianko, T. S. (2007). Using

digital photography to study animal coloration. *Biological Journal of the Linnean*

Society, *90*, 211–237. doi:10.1111/j.1095-8312.2007.00725.x

Stevens, M., Stoddard, M. C., & Higham, J. P. (2009). Studying primate color: towards visual

system-dependent methods. *International Journal of Primatology*, *30*(6), 893–917.

doi:10.1007/s10764-009-9356-z

Stuart-Fox, D., & Moussalli, A. (2011). Camouflage in colour-changing animals. In S. M.

Martin Stevens (Ed.), *Animal Camouflage: Mechanisms and Function*. Cambridge:

Cambridge University Press.

Talas, L., Baddeley, R. J., & Cuthill, I. C. (2017). Cultural evolution of military camouflage.

Philosophical Transactions of the Royal Society of London. Series B, Biological

Sciences, *372*(1724), 20160351. doi:10.1098/rstb.2016.0351

Team, R. C. (2016). R: A language and environment for statistical computing. Vienna,

Austria: R Foundation for Statistical Computing.

Tlusty, M., & Hyland, C. (2005). Astaxanthin deposition in the cuticle of juvenile American

lobster (*Homarus americanus*): implications for phenotypic and genotypic coloration.

Marine Biology, *147*(1), 113–119. doi:10.1007/s00227-005-1558-0

Todd, P. A., Briers, R. A., Ladle, R. J., & Middleton, F. (2006). Phenotype-environment

matching in the shore crab (*Carcinus maenas*). *Marine Biology*, *148*(6), 1357–1367.

doi:10.1007/s00227-005-0159-2

Todd, P. A., Oh, J., Loke, L. H. L., & Ladle, R. J. (2012). Multi-scale phenotype-substrate

matching: Evidence from shore crabs (*Carcinus maenas* L.). *Ecological Complexity*, *12*,

58–62. doi:10.1016/j.ecocom.2012.09.005

Todd, P. A., Qiu, W., & Chong, K. Y. (2009). Ontogenetic shifts in carapace patterning

and/or colouration in intertidal and subtidal brachyuran crabs. *Raffles Bulletin of*

Zoology, *57*(2), 543–550.

Troscianko, J., & Stevens, M. (2015). Image calibration and analysis toolbox - a free

software suite for objectively measuring reflectance, colour and pattern. *Methods in*

Ecology and Evolution, *6*, 1320–1331. doi:10.1111/2041-210X.12439

Troscianko, J., Wilson-Aggarwal, J., Stevens, M., & Spottiswoode, C. N. (2016). Camouflage

predicts survival in ground-nesting birds. *Scientific Reports*, *6*(1), 19966.

doi:10.1038/srep19966

Umbers, K. D. L., Fabricant, S. A., Gawryszewski, F. M., Seago, A. E., & Herberstein, M. E.

(2014). Reversible colour change in Arthropoda. *Biological Reviews*, *89*(4), 820–848.

doi:10.1111/brv.12079

van der Meeren, G. I. (2000). Predation on hatchery-reared lobsters released in the wild.

Canadian Journal of Fisheries and Aquatic Sciences, *57*, 1794–1803.

van der Meeren, G. I. (2005). Potential of ecological studies to improve survival of cultivated

and released European lobsters, *Homarus gammarus*. *New Zealand Journal of Marine*

and Freshwater Research, *39*(2), 399–424. doi:10.1080/00288330.2005.9517321

Vorobyev, M., & Osorio, D. (1998). Receptor noise as a determinant of colour thresholds.

Proceedings of The Royal Society B, *265*(1394), 351–358. doi:10.1098/rspb.1998.0302

Wade, N. M., Budd, A., Irvin, S., & Glencross, B. D. (2015). The combined effects of diet,

environment and genetics on pigmentation in the giant tiger prawn, *Penaeus monodon*.

Aquaculture, *449*, 78–86. doi:10.1016/J.AQUACULTURE.2015.01.023

Wilson, D., Heinsohn, R., & Endler, J. A. (2007). The adaptive significance of ontogenetic

colour change in a tropical python. *Biology Letters*, *3*(1), 40–3.

doi:10.1098/RSBL.2006.0574