SHORT COMMUNICATION

Do Cells use Passwords in Cell-State Transitions? Is Cell Signaling sometimes Encrypted?

Alex Root [1,*]

[1] Molecular Biology Program, Memorial Sloan Kettering Cancer Center, New York, NY, USA

AR: Alex Root 0000-0003-1206-7010

* Correspondence: Alex Root

Molecular Biology Program

Memorial Sloan Kettering Cancer Center

New York, NY, USA

Tel 646-286-3612

Email alex.root.research@gmail.com

Abstract

Organisms must maintain proper regulation including defense and healing. Life-threatening problems may be caused by pathogens or by a multicellular organism's own cells through cancer or auto-immune disorders. Life evolved solutions to these problems that can be conceptualized through the lens of information security, which is a well-developed field in computer science. Here I argue that taking an information security view of cells is not merely semantics, but useful to explain features of signaling, regulation, and defense. An information security perspective also offers a conduit for cross-fertilization of advanced ideas from computer science, and the potential for biology to inform computer science. First, I consider whether cells use passwords, i.e., initiation sequences that are required for subsequent signals to have effects, by analyzing the concept of pioneer transcription factors in chromatin regulation and cellular reprogramming. Second, I consider whether cells may encrypt signal transduction cascades. Encryption could benefit cells by making it more difficult for pathogens or oncogenes to hijack cell networks. By using numerous molecules cells may gain a security advantage in particular against viruses, whose genome sizes are typically under selection pressure. I provide a simple conceptual argument for how cells may peform encryption through post-translational modifications, complex formation, and chromatin accessibility. I invoke information theory to provide a criterion of an entropy spike to assess whether a signaling cascade has encryption-like features. I discuss how the frequently invoked concept of context-dependency may over-simplify more advanced features of cell signaling networks, such as encryption. Therefore, by considering that biochemical networks may be even more complex than commonly realized we may be better able to understand defenses against pathogens and pathologies.

Keywords: cell signaling; reprogramming; host-pathogen interactions; information security; biochemical passwords; biochemical encryption; post-translational modification; chromatin state; context-dependency

Main

Biochemical networks transmit, receive, and process information resulting in decision making concerning growth, defense, differentiation, migration, apoptosis, metabolism, and other processes[1,2]. Groundbreaking studies over the last several decades have elucidated properties of signaling and regulatory networks, including scale-freeness, robustness, fragility, noise-filtering,

bistability, controllability, ultrasensitivity, signal dissipation, amplification, memory, modularity, feedfoward and other motifs, which are reviewed by Krakauer and colleagues[2], Uda and Kuroda[3], Mousavian and colleagues[4,5], Walterman and Klipp[6], Azeloglu and Iyengar[1], and Antebi and colleagues[7]. Biochemical networks can become dysfunctional through genetic mutation, chemical injury, infection, or other processes, that might achieve varying degrees of control over the network[8]. Here, I consider these processes through the lens of information security, which as far as I can determine is not common. This is in contrast to telecommunications, where cybersecurity is of paramount importance[9]. In an elegant and trenchant examination of theoretical biology, Krakauer and colleagues argue "before we can look for patterns, we often need to know what kinds of patterns to look for, which requires some fragments of theory to begin with[10]."

By explicitly incorporating information security concepts into thinking about biological systems, several outcomes are possible in general: (1) distinctions without differences, that is, we are merely rephrasing familiar concepts of immunity and regulation in terms of information security, which adds no value; (2) cross-disciplinary fertilization occurs as information security concepts are imported into biological theory; (3) new information security knowledge arises from examination of biological systems. Recent studies on network controllability provide one framework for examining security in biochemical networks[11-15]. Here, a different perspective is taken to analyze whether cells may use passwords and encrypt their signaling cascades.

Immune systems and biological security

The evolution of immune systems and self-defense against injury and mutation are major innovations in the history of life on earth[16-18]. By total volume, life on earth has its largest habitat in the deep ocean with an abundance of bacteriophages, seemingly saturating the environment[19]. Indeed, parasitism has been estimated to have evolved independently several hundred times. Single-celled and multi-cellular organisms evolved a wide-variety of defense systems, often dichotomized into innate and adaptive systems[20]. These systems can be conceptualized more generally to include protective mechanisms against both external and internal damage. The

connection between external and internal injury is seen in the study of viruses, which led to

insights in cancer biology and the discovery of oncogenes[17]. Organisms developed the ability to

recognize self from non-self and destroy xenobiotic material. However, not all foreign genetic

material is completely destroyed, because it can increase fitness, e.g., antibiotic resistance

plasmids[20,21]. On the intracellular level, bacterial defense mechanisms include blocking receptor

binding (surface modification), genome injection (superinfection exclusion), viral replication

(restriction modification, CRISPR-Cas, and prokaryotic Argonaute), and abortive infection

(programmed cell death)[21]. Similar mechanisms exist in eukaryotic cells, including, RIG-like

receptor proteins that recognize RNA[16], xenophagy[22], advanced intracellular nucleic acid

recognition systems and other cell-autonomous mechanisms[23]. In plants, sophisticated DICERs

defend against retroviruses[24]. Similarly, pathogens use a variety of mechanisms to co-opt, hijack,

and counteract host defenses[25-28]. Mutations leading to oncogenes reprogram signaling

networks[29]. All of these attacks and counter-attacks involve changes in signaling and regulatory

networks, and therefore, changes in information. Intriguingly, the ciliates *Oxytricha* and

Stylonychia have scrambled or encrypted genomes that undergo massive re-arrangement during

development into functional genes[30]. Presumably, this evolved as a defense mechanism against

pathogens and provides examples that cells encrypt information.

Information security in computer science

Information security has been important for millennia, with the Caesar substitution cipher being a

known early example[31]. (The cipher works by shifting each letter of the English alphabet by 3, i.e.,

A->C, B->D,...,X->A[31]). Within several decades of the development of computers, security against

computer viruses became an important concern. Computer viruses achieved notoriety in 1987

when the Brain, Lehigh, and April Fool viruses came to worldwide attention[32]. Hackers achieved

infamy and also contributed to the advancement of information technology[33]. Information security

depends on the use of passwords for system access and encryption[34]. Development of secure

encryption systems, e.g., the RSA asymmetric public key cryptography, was an essential

innovation in the history of the internet[34] and must evolve to meet new threats[9]. Steganography is

an altogether different approach that conceals the existence of information, e.g., writing with

invisible ink, and appears to have had played less importance in the history of information

technology than cryptography[34]. Attacks on encrypted systems can involve interception,

modification, fabrication, or interruption of information[34]. There has been considerable work in

adapting biomolecules for use in information security in human telecommunications using

biosteganography[35] where information is invisible and molecular cryptography, where synthetic

biology is used to re-engineer molecules to decode and encode information[36]. Despite obvious

parallels to computer science, less explicit attention appears to have been paid to theoretical

descriptions of cells in terms of information security.

Information systems in cells

Organisms have a variety of sophisticated information systems. They encode information through

the nucleic acid sequences, which utilize complementary base pairing of two strands to provide

built-in error correction, which is a type of backup and repair system. At the proteome level, cells

can modify amino acids with a few hundred post-translational modifications in various

combinations, that give rise to numerous proteoforms[37], which form components of signaling and

regulatory networks. Somatic recombination in immunoglobulins and T-cell receptors can vastly

increase protein variants in certain cell types[38]. Interactions of these macromolecules form

networks that store and transmit information[6]. There is a context specificity to many signaling

pathways, including TGF-beta and AKT, which means that cells respond differently to pathway

activation depending on the cell type[39,40]. Many intracellular signaling pathways do not match one

receptor to a single ligand, but instead use multiple receptors and ligands that interact

combinatorially[41], or use combinations of numerous nuclear-receptor cofactors to regulate

activity[42]. Therefore, genetic, epigenetic, transcriptomic and proteomic variation gives rise to a

large repertoire of interacting components. These mechanisms are present in complex

multicellular organisms, where advanced regulation is needed to control differentiation[43] and also

in bacteria for complex behaviors, such as quorum sensing[2].

Cancer often involves rewiring cellular networks by oncogenes and therefore, in some sense,

these represent alterations in information transmission and compromised security[29,44]. Cancer

cells can transition because of the effects of microRNAs and oncogenes to different states with

distinct metabolism[45]. Similarly, viruses can substantially rewire signaling and regulatory networks

to hijack cellular machinery for viral benefit[46]. In the early days of cancer research, similarities

between the two systems caused the scientific community to think that viruses cause cancer, and

studies into viral biology provided insights into cancer[17,47]. Both pathogenic and pathological

processes can involve hijacking cellular networks.

In multicellular organisms, combinations of histone modifications give rise to varying

chromosomal accessibility and epigenetic states, which are read, written, and erased[48,49]. This

epigenetic regulation is capable of encoding memory at the single-cell level[50]. Redundancy and

correlation among epigenetic marks, transcription factors, and co-regulators provides a way to

specify cell states, off which there are more than 250 distinct cell types in adult humans[51]. For

example, the identity of a ligand can be encoded as pulsatile (DLL1-Notch1) or sustained (DLL4-

Notch1) to induce opposite cell fates. In the adult human body, several hundred distinct cell types

exist in cell states, some of which can be transitioned from one state to the next using

combinations of growth factors and transcription factors[52-54]. The language used to describe these

cellular properties (code, encode, read, write, memory, erase, rewire) suggests that they may be

information systems, but requires careful thought to address *how* it is that they are information

systems.

Do cells use passwords before transitioning between cell states?

Password authorization systems allow access based upon entry of a somewhat random code and

achieve security by creating a large search space. Passwords are also an initiation sequence of

signals without which a system will not respond to subsequent signals. In multicellular organisms,

cells develop into types that can be conceptualized as stable states. In order to change from one

state to another, that is, to differentiate or de-differentitate, the epigenome of cells changes[55].

Organization of chromatin into highly compact, inaccessible regions, and open, accessible

regions is a form of information security because some genes are "locked" and therefore, cannot

be transcribed. Chromatin is frequently characterized as being in "open" and "closed" states that

must be unlocked. This appears to be a potential case where cells use passwords because the

right combination of growth factors and transcription factors are necessary for reprogramming to

occur. The initial sequence is mediated by so-called pioneer transcription factors. There are

multiple algorithms to predict combinations of transcription factors to reprogram human cells from

one type to another with the number of successful conversions being relatively low[52]. The

systems work by engineering overexpression of transcription factors, rather than as it happens

normally in development through extracellular signaling molecules that signal to transcription

factors to achieve the rewiring. To my knowledge, no one has attempted to predict upstream

combinations of signals, e.g., growth factors, hormones, adhesion contacts, etc. that would trigger

the right combinations of transcription factors. Sampattavanich and colleagues demonstrated that

FOXO3 dynamics can code for different growth factors and their concentrations, which are under

combinatorial control of ERK and AKT pathways[56]. Typically, engineering the over-expression of

transcription factors does not introduce dynamics, that is, the over-expression simply turns on to

high concentration.

If cell reprogramming requires an initiation sequence of transcription factors to unlock the

epigenetic code then conceptualizing this as a password is appropriate. A password

conceptualization is distinct from simply requiring a series of events because of its defensive

purpose and strategy of employing a seemingly random draw from a large search space. If a set

of transcription factors are active during the entire reprogramming process, then they are not

performing an initiation sequence and therefore, not entering a password. Similarly, if only one

member of the combination can partially reprogram cells then it would seem inappropriate to

conceptualize the mechanism as a password. This conceptualization might predict that password-

length, i.e., the complexity of the reprogramming initiation is directly proportional the fitness cost

posed to the organism from the conversion. For example, because stem cells have greater

replicative potential, they might pose greater risk to develop into cancer and consequently,

require a more complex password for reprogramming. It is unclear to me whether there are

sufficient measurements to resolve whether there is a distinct initiation sequence during cell

reprogramming.

Do cells encrypt information in signaling cascades?

Cell signaling can be hijacked by pathogens or oncogenes; for example, the chrysanthemum

virus B p12 protein acts as a transcription factor to control cell size and proliferation for its

benefit[57]. A cell signaling system might be able to use an encryption-like system to block this sort

of attack. The way that it might work is to require different combinations of signals in different cell

types to control cell size and proliferation. What encryption systems do is to increase the entropy

of the signal by having machinery to encrypt and decrypt, e.g., rotors on an enigma machine. As

a general rule, there is evolutionary pressure on viruses to maintain relatively small genomes,

and less constraint on the sizes of non-viral genomes. This gives hosts a competitive advantage

over viruses to create more complex signaling systems by adding components, e.g.,

encrypters/decrypters, in the form of additional genes. It is also protection against deleterious

mutations because these are less likely to occur in multiple genes. The cost would seem to be

maintaining a larger genome.

One way encryption could work in cell signaling is through a system consisting of an input signal,

an encoder, an encrypter, a noisy signal transduction cascade, a decrypter, a decoder, and an

output signal. Hypothetically, an encrypter could consist of multiple post-translational

modifications on cell surface receptors and a decrypter could consist of transcription factor

complexes interacting with post-translationally modified histones, see figure 1A. For example,

suppose two different cell types A and B receive the same binary input signal, which they decode

identically to the same output signal, such as a cascade of gene expression to differentiate into a

another cell type. Suppose the phosphorylation patterns differ on the receptors, the transcription

factor complexes differ, and the chromatin state on the target genes differ, see figure 1. The

conventional explanation for these different signaling cascades is context-dependence, i.e.,

different signals in different cell types. Context-dependence is itself a useful form of information

security because a different attack may be required for each context, so that only a subset of an

organisms' tissues may be attacked at any one time. What is needed to distinguish context-

dependence from encryption is a consideration of what encryption does quantitatively, that is,

encryption causes an increase in entropy of the transmitted signal, see figure 1B. Figure 1A

depicts this as an increase in possible post-translational modification states going from the

encoder (receptor binding on/off) to encrypter (receptor dimerization partner with 3 on/off

phosphorylation sites (8 states)), where the inactive pathway randomly transmits the various

states. An attacker is therefore less likely to mimic the activation signal during the encrypted

chain, especially if the chain has dynamic behavior, such as periodic oscillations, see figure 1C.

However, vulnerabilities are still present before encryption and after decryption. This suggests

that a more secure biochemical system would combine signal reception, encoding, and

encryption into an integrated ligand-receptor-scaffold complex. Or it suggests that there may be

something about the biochemistry of the input and output signals that is harder for a pathogen or

pathology to mimic, e.g., an extracellular ligand when typically viruses infect intracellularly.

Conclusions

Evolutionary potential is vast and a complex interplay among environmental change, ecosystems,

speciation, niche diversification, extinctions, and innovation have shaped life on earth[58,59].

Considering how rapidly hacking, passwords, and encryption evolved in computers, it is natural to

ask whether they are used in nature by cells. This conceptual exploration suggests that cells may

use passwords to lock-in cell state, which must be unlocked through the right combination of

transcription factors. Open questions include if cells use passwords to initiate cell signaling

cascades, programmed cell death, neuron-to-neuron transmissions, or other areas. Also, is it the

case that password-length, i.e., the complexity of an initiation sequence is directly proportional

the fitness cost? When there is selection pressure due to co-evolution of pathogens are there

more complex initiation sequences, i.e., harder-to-crack passwords? Do these have greater

complexity in their molecular mechanisms? Another open question is do pathogens launch

attacks similar to those seen in computer science, e.g., denial-of-service attacks? What are

biological equivalents of two-factor authentication?

While I have not presented evidence for cell passwords, encryption, or other security measures, I

suggest that they may exist and provide fragments of theory and criteria that the community can

use to look for patterns that may demonstrate their existence. There are several examples

showing how to quantify Shannon entropy in biochemical systems and protein complexes,

reviewed by Waltermann and Klipp[6]. I also have not addressed the critical question of noise in

biological systems and measurements, which add considerable complexity to information

theoretic analysis of biological systems, which confounds the criterion of whether encryption is

present in a signaling cascade because both increase the entropy. If there is encryption and

decryption, identifying the molecular mechanisms by which it occurs could yield new and powerful

insights into signaling, pathogens, and pathologies.

Figures

Figure 1. How encryption might work in cell signaling. **A** Two different cell types encode and encrypt the same signal binary signal of ligand binding using different receptor dimerization pairs with 3 phosphorylation sites, allowing for 8 on/off states, only a subset of which encode the signal and the remainder are transmitted randomly when the receptor is inactive. The encrypted signal is relayed through a noisy channel where a complex of transcription factors and cofactors can also form 8 different association states, only a subset of which can interact with the chromatin state of the cell to produce an output response of expresed genes. **B** If an input signal is encrypted then its transmission will have a higher entropy. The decrypted output signal will have an entropy slightly higher than the input signal due to noise. The presence of error correction may make noise minimal. **C** The repressilator circuit can produce periodic oscillations in a biochemical system that in principle could function like a substitution cipher, i.e., A->A, A->B, A->C, A->A, etc.

Declarations

Author contributions

A.R. wrote the paper.

Acknowledgments

The author thanks H. Alexander Ebhardt for critical comments and discussion. The author also thanks two anonymous reviewers for critical comments and suggestions, including bringing to his attention the encrypted genomes of *Oxytricha* and *Stylonychia*.

Consent for publication

No humans were actively recruited to the work presented here.

Competing interests

The author declares that he has no competing interests.

Ethics approval and consent to participate

No humans were actively recruited to the work presented here.

Funding

U.S. National Cancer Institute (NCI) P30 Cancer Center Support Grant (CCSG) P30 CA008748 to A.R. The funding covers general support for the research center.

285 ***Publisher's Note***

286 Springer remains neutral with regard to jurisdictional claims in published maps and institutional

287 affiliations.

References

1.   Azeloglu, E.U. & Iyengar, R. Signaling networks: information flow, computation, and
decision making. *Cold Spring Harb Perspect Biol* **7**, a005934 (2015).
2.   Krakauer, D.C., Muller, L., Prohaska, S.J. & Stadler, P.F. Design specifications for
cellular regulation. *Theory Biosci* **135**, 231-240 (2016).
3.   Uda, S. & Kuroda, S. Analysis of cellular signal transduction from an information theoretic
approach. *Semin Cell Dev Biol* **51**, 24-31 (2016).
4.   Mousavian, Z., Kavousi, K. & Masoudi-Nejad, A. Information theory in systems biology.
Part I: Gene regulatory and metabolic networks. *Semin Cell Dev Biol* **51**, 3-13 (2016).
5.   Mousavian, Z., Diaz, J. & Masoudi-Nejad, A. Information theory in systems biology. Part
II: protein-protein interaction and signaling networks. *Semin Cell Dev Biol* **51**, 14-23
(2016).
6.   Waltermann, C. & Klipp, E. Information theory based approaches to cellular signaling.
Biochim Biophys Acta **1810**, 924-32 (2011).
7.   Antebi, Y.E., Nandagopal, N. & Elowitz, M.B. An operational view of intercellular signaling
pathways. *Curr Opin Syst Biol* **1**, 16-24 (2017).
8.   Vidal, M., Cusick, M.E. & Barabasi, A.L. Interactome Networks and Human Disease. *Cell*
144, 986-998 (2011).
9.   Bernstein, D.J. & Lange, T. Post-quantum cryptography. *Nature* **549**, 188-194 (2017).
10.   Krakauer, D.C. *et al.* The challenges and scope of theoretical biology. *J Theor Biol* **276**,
269-76 (2011).
11.   Liu, Y.Y., Slotine, J.J. & Barabasi, A.L. Controllability of complex networks. *Nature* **473**,
167-73 (2011).
12.   Gates, A.J. & Rocha, L.M. Control of complex networks requires both structure and
dynamics. *Sci Rep* **6**, 24456 (2016).
13.   Uhart, M., Flores, G. & Bustos, D.M. Controllability of protein-protein interaction
phosphorylation-based networks: Participation of the hub 14-3-3 protein family. *Sci Rep*
6, 26234 (2016).
14.   Li, A., Hu, Q., Liu, J. & Pan, Y. Resistance and Security Index of Networks: Structural
Information Perspective of Network Security. *Sci Rep* **6**, 26810 (2016).
15.   Albert, R., Jeong, H. & Barabasi, A.L. Error and attack tolerance of complex networks.
Nature **406**, 378-82 (2000).
16.   Mukherjee, K., Korithoski, B. & Kolaczkowski, B. Ancient origins of vertebrate-specific
innate antiviral immunity. *Mol Biol Evol* **31**, 140-53 (2014).
17.   Hanahan, D. & Weinberg, R.A. Hallmarks of cancer: the next generation. *Cell* **144**, 646-
74 (2011).
18.   Ameisen, J.C. On the origin, evolution, and nature of programmed cell death: a timeline
of four billion years. *Cell Death Differ* **9**, 367-93 (2002).
19.   Lara, E. *et al.* Unveiling the role and life strategies of viruses from the surface to the dark
ocean. *Sci Adv* **3**, e1602565 (2017).
20.   Sirisinha, S. Evolutionary insights into the origin of innate and adaptive immune systems:
different shades of grey. *Asian Pac J Allergy Immunol* **32**, 3-15 (2014).
21.   van Houte, S., Buckling, A. & Westra, E.R. Evolutionary Ecology of Prokaryotic Immune
Mechanisms. *Microbiol Mol Biol Rev* **80**, 745-63 (2016).
22.   Miller, C. & Celli, J. Avoidance and Subversion of Eukaryotic Homeostatic Autophagy
Mechanisms by Bacterial Pathogens. *J Mol Biol* **428**, 3387-98 (2016).
23.   Wu, J. & Chen, Z.J. Innate immune sensing and signaling of cytosolic nucleic acids. *Annu*
Rev Immunol **32**, 461-88 (2014).

24.   Ebhardt, H.A., Thi, E.P., Wang, M.B. & Unrau, P.J. Extensive 3' modification of plant
342   small RNAs is modulated by helper component-proteinase expression. *Proc Natl Acad*
343   *Sci U S A* **102**, 13398-403 (2005).

344   25.   Alto, N.M. & Orth, K. Subversion of cell signaling by pathogens. *Cold Spring Harb*
345   *Perspect Biol* **4**, a006114 (2012).

346   26.   Moffatt, J.H., Newton, P. & Newton, H.J. Coxiella burnetii: turning hostility into a home.
347   *Cell Microbiol* **17**, 621-31 (2015).

348   27.   Alcami, A. Viral mimicry of cytokines, chemokines and their receptors. *Nat Rev Immunol*
349   **3**, 36-50 (2003).

350   28.   Agol, V.I. & Gmyl, A.P. Viral security proteins: counteracting host defences. *Nat Rev*
351   *Microbiol* **8**, 867-78 (2010).

352   29.   Stuhlmiller, T.J., Earp, H.S. & Johnson, G.L. Adaptive reprogramming of the breast
353   cancer kinome. *Clin Pharmacol Ther* **95**, 413-5 (2014).

354   30.   Burns, J. *et al.* Recurring patterns among scrambled genes in the encrypted genome of
355   the ciliate Oxytricha trifallax. *J Theor Biol* **410**, 171-180 (2016).

356   31.   Gardner, M. *Codes, Ciphers, and Secret Writing*, (Dover Publications, Inc., New York,
357   NY, USA, 1972).

358   32.   Highland, H.J. History of Computer Viruses. *Computers & Security* **16**, 416-429 (1997).

359   33.   Levy, S. *Hackers*, (O'Reilly Media, Sebastopol, 2010).

360   34.   Tipton, H.F. & Krause, M. *Information Security Management Handbook*, 3280 (CRC
361   Press, Boca Raton, FL, 2007).

362   35.   Brunet, T.D. Aims and methods of biosteganography. *J Biotechnol* **226**, 56-64 (2016).

363   36.   Lustgarten, O., Motiei, L. & Margulies, D. User Authorization at the Molecular Scale.
364   *Chemphyschem* **18**, 1678-1687 (2017).

365   37.   Aebersold, R. *et al.* How many human proteoforms are there? *Nat Chem Biol* **14**, 206-
366   214 (2018).

367   38.   Hood, L., Kronenberg, M. & Hunkapiller, T. T cell antigen receptors and the
368   immunoglobulin supergene family. *Cell* **40**, 225-9 (1985).

369   39.   Feng, X.H. & Derynck, R. Specificity and versatility in tgf-beta signaling through Smads.
370   *Annu Rev Cell Dev Biol* **21**, 659-93 (2005).

371   40.   Toker, A. & Marmiroli, S. Signaling specificity in the Akt pathway in biology and disease.
372   *Adv Biol Regul* **55**, 28-38 (2014).

373   41.   Antebi, Y.E. *et al.* Combinatorial Signal Perception in the BMP Pathway. *Cell* **170**, 1184-
374   1196 e24 (2017).

375   42.   Broekema, M.F. *et al.* Profiling of 3696 Nuclear Receptor-Coregulator Interactions: A
376   Resource for Biological and Clinical Discovery. *Endocrinology* **159**, 2397-2407 (2018).

377   43.   Spickard, E.A., Joshi, P.M. & Rothman, J.H. The multipotency-to-commitment transition
378   in Caenorhabditis elegans-implications for reprogramming from cells to organs. *FEBS*
379   *Lett* **592**, 838-851 (2018).

380   44.   Drake, J.M. *et al.* Phosphoproteome Integration Reveals Patient-Specific Networks in
381   Prostate Cancer. *Cell* **166**, 1041-1054 (2016).

382   45.   Pinweha, P., Rattanapornsompong, K., Charoensawan, V. & Jitrapakdee, S. MicroRNAs
383   and oncogenic transcriptional regulatory networks controlling metabolic reprogramming in
384   cancers. *Comput Struct Biotechnol J* **14**, 223-33 (2016).

385   46.   McKinney, C. *et al.* Global reprogramming of the cellular translational landscape
386   facilitates cytomegalovirus replication. *Cell Rep* **6**, 9-17 (2014).

387   47.   Koonin, E.V., Dolja, V.V. & Krupovic, M. Origins and evolution of viruses of eukaryotes:
388   The ultimate modularity. *Virology* **479-480**, 2-25 (2015).

389   48.   Voigt, P. *et al.* Asymmetrically modified nucleosomes. *Cell* **151**, 181-93 (2012).

390   49.   Strahl, B.D. & Allis, C.D. The language of covalent histone modifications. *Nature* **403**, 41-
391   5 (2000).

392   50.   Bintu, L. *et al.* Dynamics of epigenetic regulation at the single-cell level. *Science* **351**,
393   720-4 (2016).

394   51.   Ahsendorf, T., Muller, F.J., Topkar, V., Gunawardena, J. & Eils, R. Transcription factors,
395   coregulators, and epigenetic marks are linearly correlated and highly redundant. *PLoS*
396   *One* **12**, e0186324 (2017).

52.   Kamaraj, U.S., Gough, J., Polo, J.M., Petretto, E. & Rackham, O.J. Computational
methods for direct cell conversion. *Cell Cycle* **15**, 3343-3354 (2016).
53.   Li, C. & Wang, J. Quantifying cell fate decisions for differentiation and reprogramming of
a human stem cell network: landscape and biological paths. *PLoS Comput Biol* **9**,
e1003165 (2013).
54.   Wang, J., Zhang, K., Xu, L. & Wang, E. Quantifying the Waddington landscape and
biological paths for development and differentiation. *Proc Natl Acad Sci U S A* **108**, 8257-
62 (2011).
55.   Graf, T. & Enver, T. Forcing cells to change lineages. *Nature* **462**, 587-94 (2009).
56.   Sampattavanich, S. *et al.* Encoding Growth Factor Identity in the Temporal Dynamics of
FOXO3 under the Combinatorial Control of ERK and AKT Kinases. *Cell Syst* (2018).
57.   Lukhovitskaya, N.I. *et al.* A viral transcription factor exhibits antiviral RNA silencing
suppression activity independent of its nuclear localization. *J Gen Virol* **95**, 2831-7
(2014).
58.   Knoll, A.H. & Nowak, M.A. The timetable of evolution. *Sci Adv* **3**, e1603076 (2017).
59.   Cazzolla Gatti, R., Fath, B., Hordijk, W., Kauffman, S. & Ulanowicz, R. Niche emergence
as an autocatalytic process in the evolution of ecosystems. *J Theor Biol* **454**, 110-117
(2018).

A

input signal

encoder

CELL TYPE 1

P
P
P
P

encrypter
8 phospho on/off states

MAPKKK

MAPKK

noisy channel

MAPK

decrypter
8 associations

output signal
gene expression {G}

input signal

encoder

CELL TYPE 2

P
P
P
P

encrypter
8 phospho on/off states

MAPKKK

MAPKK

noisy channel

MAPK

decrypter
8 associations

output signal
gene expression {G}

B

ENTROPY H

encryption
+
noise

decryption

signal

encrypted
transmission

decrypted
signal

C

Concentration

P1

P3 P2

time