Structure of the large extracellular loop of FtsX and its interaction with the

essential peptidoglycan hydrolase PcsB in *Streptococcus pneumoniae*

Britta E. Rued,[ab] Martín Alcorlo,[c] Katherine A. Edmonds,[b] Siseth Martínez-Caballero,[c]

Daniel Straume,[d] Yue Fu,[b]* Kevin E. Bruce,[a] Hongwei Wu,[b] Leiv Sigve Håvarstein,[d]

Juan A. Hermoso,[c,#] Malcolm E. Winkler,[a,#] and David P. Giedroc[b,#].

[a]Department of Biology, Indiana University Bloomington, Bloomington, IN, USA.

[b]Department of Chemistry, Indiana University Bloomington, Bloomington, IN, USA.

[c]Department of Crystallography and Structural Biology, Inst. Química-Física

Rocasolano, Consejo Superior de Investigaciones Científicas, Madrid, Spain.

[d]Department of Chemistry, Biotechnology and Food Science, Norwegian University of

Life Sciences, N-1432, Ås, Norway.

Running Head: FtsX$_{ECL1}$ structure and interaction with PcsB in *S. pneumoniae*

[#]Address correspondence to David P. Giedroc, giedroc@indiana.edu; Juan A.

Hermoso, xjuan@iqfr.csic.es; and Malcolm E. Winkler, winklerm@indiana.edu.

*Present address: Genentech, South San Francisco, California, USA.

B.E.R. and M.A. contributed equally to this work.

Abstract word count: 163 words

Text word count: 5,664 words

ABSTRACT

Streptococcus pneumoniae is a leading killer of infants and immunocompromised adults and has become increasingly resistant to major antibiotics. Therefore, the development of new antibiotic strategies is desperately needed. Targeting bacterial cell division is one such strategy, specifically targeting essential proteins for the synthesis and breakdown of peptidoglycan. One complex important to this process is FtsEX. FtsEX comprises an integral membrane protein (FtsX) and cytoplasmic ATPase (FtsE) that resembles an ATP-binding cassette (ABC) transporter. Here, we present NMR solution structural and crystallographic models of the large extracellular domain of FtsX, denoted ECL1. The structure of ECL1 reveals an upper extended β-hairpin and a lower α-helical lobe, each extending from a mixed α-β core. The helical lobe mediates a physical interaction with the peptidoglycan hydrolase PcsB, via the coiled-coil domain of PcsB (PcsB-CC). Characterization of *S. pneumoniae* D39 derived strains harboring mutations in the α-helical lobe shows that this subdomain is essential for cell viability and required for proper cell division of *S. pneumoniae*.

IMPORTANCE FtsX is a ubiquitous bacterial integral membrane protein involved in cell division that regulates the activity of peptidoglycan (PG) hydrolases. FtsX is representative of a large group of ABC3 superfamily proteins that function as "mechanotransmitters," proteins that relay signals from inside to the outside of the cell. Here we present a structural characterization of the large extracellular loop (ECL1) of FtsX from the human opportunistic pathogen *Streptococcus pneumoniae*. We show a direct interaction between the peptidoglycan hydrolase PcsB and FtsX, and

demonstrate that this interaction is essential for cell viability. As such, FtsX represents

an attractive, conserved target for the development of new classes of antibiotics.

INTRODUCTION

Streptococcus pneumoniae is a Gram-positive, opportunistic respiratory pathogen (1-3)

that has acquired antibiotic resistance worldwide (4-6). This ovococcal bacterium relies

on highly conserved cell wall machinery to divide and grow (7, 8). The cell wall is

primarily composed of peptidoglycan (PG), a macromolecule composed of repeating

subunits of N-acetylglucosamine and N-acetylmuramic acid linked by PG peptide side

chains (9, 10). Regulation of the synthesis and remodeling of PG is essential for

bacterial growth and viability, due to the turgor pressure bacterial cells must withstand

(10-12). One vital process for the synthesis of PG is the controlled insertion of new

strands of PG. This process requires timed cleavage of the old PG matrix to allow

incorporation of new nascent strands (13). PG hydrolases are the primary enzymes that

carry out PG cleavage and remodeling (14, 15). Thus, regulation of these hydrolases

and activation at specific times during the cell cycle is required for proper cell growth.

Specific protein complexes are utilized by bacterial cells to regulate these enzymes.

This work focuses on understanding the structure and function of one of these protein

complexes.

From *Mycobacterium tuberculosis* to *Caulobacter crescentus*, the ABC-

transporter-like protein complex FtsEX acts as a key regulator of PG hydrolysis and

divisome assembly (16-19). The proposed mechanism of FtsEX activation of PG

hydrolases is as follows. FtsE, upon sensing an unknown signal from inside the cell,

hydrolyzes ATP to ADP. Hydrolysis causes a conformational change transmitted

through the membrane via FtsX, an integral membrane protein with two extracellular

loops, denoted the large (ECL1) and small (ECL2) loops. These extracellular loops

interact with either the cell wall hydrolases or with effector proteins, which results in

activation of PG hydrolysis via an unknown mechanism (16, 18, 20-25). In *E. coli*, it has

been demonstrated that FtsX interacts with the effector protein EnvC to activate the PG

amidases AmiA and AmiB (24, 25). In addition, FtsX interacts with other division

proteins such as FtsA, where it regulates the polymerization of FtsA and recruitment of

downstream division proteins (26). In other organisms including *B. subtilis* and *M.*

tuberculosis, FtsEX also activates PG hydrolases (16, 23). Interestingly, FtsEX is non-

essential in rod-shaped bacteria such as *E. coli* and *B. subtilis* (23, 24, 26-28).

However, in *S. pneumoniae* FtsEX is absolutely essential (21) and depletion of FtsEX

results in cell rounding and cessation of growth (20, 21).

In the case of *S. pneumoniae*, genetic experiments suggest that both outward-

facing domains of FtsX, ECL1 and ECL2, interact with the essential PG hydrolase PcsB,

via its long coiled-coil (CC) domain (20, 21). However, there is little direct biochemical

evidence for this interaction. ECL1 and ECL2 are thought to allosterically activate the

catalytic activity of cysteine, histidine-dependent amidohydrolase/peptidase (CHAP)

domain of PcsB (20). The crystal structure of full-length PcsB including the CC domain,

an alanine-rich linker region, and the CHAP domain provides insight into the mechanism

of how this may occur (22). While the PcsB structure implies FtsEX activates PcsB by

displacing the catalytic domain from the CC domain, the exact nature of the FtsX::PcsB

interaction remains unknown.

In order to understand how FtsX activates PcsB, we determined the structure of

the large extracellular loop of FtsX (FtsX$_{ECL1}$) by both multidimensional NMR

spectroscopy and X-ray crystallography. FtsX$_{ECL1}$ harbors a conserved mixed α-β core

and a lower α-helical lobe extending from the core identified previously in *M.*

tuberculosis FtsX (16)*,* and an extended β-hairpin that is unique to *S. pneumoniae*

FtsX$_{ECL1}$. The N-terminal β1 and C-terminal β6 strands are adjacent in the core and

connect ECL1 to the TM1 and TM2 helices, respectively, in the membrane. PcsB-CC-

mediated chemical shift perturbations of ^{1}H-^{15}N HSQC spectra of FtsX$_{ECL1}$ reveal that

the helical lobe consisting of the α2 helix and the α2-β5 linker (residues 107-134) of

FtsX$_{ECL1}$ interacts with PcsB-CC. To determine if this interaction is required for FtsX

function in bacterial cells, we constructed a merodiploid strain that allows for conditional

expression of the mutant *ftsX*. We demonstrate that specific amino acid substitutions in

the FtsX-PcsB interface are lethal or cause pronounced morphological defects despite

the fact these FtsX$_{ECL1}$ mutant proteins are expressed at near wild-type levels. These

findings support the model that a direct physical interaction between FtsX and PcsB is

required for activation of PcsB PG hydrolytic activity.

RESULTS

The three-dimensional structure of FtsX$_{ECL1}$. The three-dimensional structure

of FtsX$_{ECL1}$ (residues 46-168) was solved by both NMR spectroscopy (Fig. 1A) and X-

ray crystallography (Fig. 1B). The folded structure (residues 57-166) reveals a central

core composed of a four-stranded antiparallel β-sheet (β1, β6, β4, β5) and two helices

(α1 and α3), an α-helical lobe (residues 107-135) harboring the α2 helix, and an

extended β-hairpin (β2, β3). The β-hairpin and helical lobes are connected to the central

core by hinges. Details for structural determination of FtsX$_{ECL1}$ by NMR are presented in

the methods section and structure statistics are summarized in Table 1. The solution

structure shows that while the central mixed α-β core adopts a well-defined

conformation, the two appended lobes are highly dynamic on multiple timescales (*vide*

infra), presenting a range of conformations among the 20 members of the FtsX$_{ECL1}$ NMR

structural ensemble (Fig. 1A).

Three different structures, with resolutions ranging from 2.0 to 2.3 Å, were solved

by X-ray crystallography, each in the presence of a different detergent. The presence of

detergents was critical as in their absence crystals diffracted at very low resolution (≤4

Å), suggesting significant mobility in some protein regions. Details for the

crystallographic determination are provided in the methods section and structure

statistics summarized in Table 2. In all cases, two protein molecules are present in the

asymmetric unit (Fig. S1), yielding a total of six independent structures and sufficient

electron density for both the protein structure and associated detergent molecules (Fig.

S2). Different conformations were observed for the β-hairpin and the helical lobe among

the six structures depending on the presence and identity of the bound detergent

molecule to FtsX$_{ECL1}$ (Fig. S1B). The structural variations observed in these

crystallographic structures, however, are less dramatic than those observed in the NMR

conformer bundle, obtained in the absence of ligand, although the regions with high

variability are similar.

In any case, the crystal structures suggest that changes in the protein backbone

and side chains of the helical lobe occur when a detergent ligand is bound (Fig. 1C).

These changes create a cavity in which the detergent molecules insert (Fig. 1D), with a

large part of the helical lobe affected by the interaction with detergent (from Q111 to

E127) (Fig. 1C). Residues Y112, W123, E127 and F126 are strongly perturbed upon

detergent ligand binding (Fig. 1C) with the hydroxyl moiety of Y112 interacting with

carboxylate of E127. Changes in W123 and F126 stabilize the hydrophobic region of

detergent 1. A similar interaction pattern is observed for detergents 2 and 3; additional

interactions are observed with N131 with detergents 2 and 3, which are characterized

by a larger hydrophilic head (maltose) (Fig. S1E-F). Although a physiological role for

detergent binding to the helical lobe is unknown, many of these same residues are

important for the interaction with PcsB (*vide infra*).

A structural comparison with *M. tuberculosis* FtsX$_{ECL1}$ (16) (PDB 4N8N) reveals

differences in both the overall structure (RMSD value of 2.2 Å) and in the appended

lobes of the core domain (Fig. 1F). The main differences between the mycobacterial

and pneumococcal FtsX$_{ECL1}$ domains are the presence of an extra helix (α1) and a

disulfide bond in *M. tuberculosis* ECL1 that are absent in pneumococcal ECL1, and an

extended β-hairpin (residues 71-87) unique to the pneumococcal ECL1 domain. It is

worth noting that in *M. tuberculosis* ECL1, the upper and lower lobes form a large

hydrophobic cleft with four exposed Phe residues (F61, F110, F113 and F122) and this

region was suggested as a strong candidate interaction surface between FtsX and RipC

(16). These phenylalanine residues are not conserved in the pneumococcal ECL1 (Fig.

1F-G) but the hydrophobic nature of this region is preserved (Y63, I118, W123 and

L133).

FtsX$_{ECL1}$ is dynamic in solution. We next performed additional NMR

experiments to explore the mobility of the FtsX$_{ECL1}$ domain, both to validate the

heterogeneity of the structural ensemble in solution, and to elucidate function. The $^1D_{HN}$

residual dipolar couplings (RDCs) obtained by weak alignment in Pf1 filamentous phage

correspond well to previously determined secondary structure elements (29), with

uniform values for the entire length of α1 and α2, as expected for straight helices. In

contrast, $^1D_{HN}$ values near 0 for the N-terminal tail, the very C-terminus, and the non-

helical part of the helical lobe between α2 and β5, are suggestive of significant

conformational disorder in solution (Fig. S3A). These regions of small or zero RDCs are

regions of very high RMSD in the NMR structure bundle (Fig. S3B). As anticipated, the

correlation between experimentally measured RDCs and predicted RDCs back-

calculated from the structures (30) is high, but only in the core subdomain, and match

poorly in the β-hairpin for most of the crystal structures (Fig. S3C-D).

We previously reported that the ^{15}N-{^{1}H} heteronuclear NOE (hNOE) is low or

negative at the termini, indicating that they are highly flexible in solution (29). The hNOE

is also smaller in the β2-β3 hairpin region, as well as in the C-terminal end of the α2

helix and the subsequent coiled region leading into β5 (29) (Fig. S3E). Mapping these

dynamics data onto a representative structure from the solution NMR ensemble shows

that these regions with fast-timescale dynamics correspond to the β-hairpin and the

helical lobe (Fig. S3F). Information on ps-ns fast-timescale motions extracted from the

R_2/R_1 ratio also reveals that α2-β5 linker is highly flexible, while the β1-β2 linker and β-

hairpin show an elevated R_2/R_1 ratio, specifically indicative of millisecond-timescale

slow conformational exchange (Fig. S3G-H). These findings were directly confirmed by

(Carr-Purcell-Meiboom-Gill) relaxation dispersion NMR spectroscopy (31) (Fig. S3I-J).

We conclude that the β-hairpin exhibits flexibility on both the sub-ns and ms timescales.

Interestingly, the β1-β2 linker also shows increased B-factors that are qualitatively

consistent with ms-timescale conformational exchange observed by NMR.

These complex motions observed in the solution dynamics experiments are

reflected in the heterogeneity of the NMR structure, with high $C\alpha$ RMSDs particularly in

the α2-β5 linker but also in the β-hairpin and β1-β2 linker, thus validating the

conformational spread in the ECL1 structure in solution (Fig. S3B). The dynamic nature

of the helical lobe is also reflected in the heterogeneity and B-factors of the crystal

structures (Fig. S3K-L). Full-length FtsX itself is likely a dimer *in vivo* (32), and one can

speculate that the flexible helical lobe and β-hairpin regions may contribute to

dimerization or to interactions with a binding partner such as PcsB.

The PcsB coiled-coil (PcsB-CC) domain interacts with FtsX$_{ECL1}$. The [1]H-[15]N

HSQC spectrum for ECL1 has excellent chemical shift dispersion and lends itself readily

to studies of protein-protein interactions (Fig. S4A). In contrast, full-length PcsB is 42

kDa and forms a dimer, and is thus challenging to study by NMR due to its size. We

therefore constructed truncation mutants of PcsB focusing on the coiled-coil domain

(PcsB-CC), thus limiting the molecular weight to 23-24 kDa. [15]N labeled PcsB-CC (47-

267) and PcsB-CC (47-254) are both characterized by [1]H-[15]N HSQC spectra with

limited [1]H signal dispersion (Fig. S6A), consistent with the high helical content of this

domain. Circular dichroism confirms that these proteins are primarily α-helical, in

agreement with the crystal structure (22), thus indicating that they are properly folded

(Fig. S6A) and can be used for ECL1 binding studies.

To determine if PcsB physically interacts directly with $FtsX_{ECL1}$, we titrated

unlabeled PcsB-CC (47-267) into ^{15}N labeled $FtsX_{ECL1}$ at molar ratios of 1:1, 2:1, 4:1,

and 6:1, respectively, and recorded ^1H-^{15}N HSQC spectra (Fig. S4A-B). Numerous

crosspeaks move in response to the addition of PcsB-CC to ^{15}N $FtsX_{ECL1}$. The largest

changes occur in the helical lobe of $FtsX_{ECL1}$ (Fig. 2A, S4-S5). In particular, residues

M119, W123, I125, F126, G128 exhibit the greatest changes in crosspeak position and

intensity in the 1:2 $FtsX_{ECL1}$ to PcsB-CC sample (Fig. 2A, Fig. S4C); when additional

PcsB-CC is added, these crosspeaks broaden beyond detection (Fig. S5B-C). A

reciprocal ^1H-^{15}N HSQC experiment with ^{15}N labeled PcsB-CC (47-254) further confirms

an interaction with unlabeled $FtsX_{ECL1}$ (Fig. S6B), as multiple crosspeaks shift upon the

addition of increasing $FtsX_{ECL1}$ (Fig. S6B). We measured the binding affinity of the

$FtsX_{ECL1}$- PcsB-CC complex using isothermal titration calorimetry (ITC), which reveals a

K_a of 3.0 x 10^4 M^{-1} (K_d ~ 34 µM) (Table 3, Fig. 2B). These data provide the first

biochemical evidence for a direct physical interaction between PcsB-CC and $FtsX_{ECL1}$

(Fig. 2, Fig. S4-S6).

The interaction region of $FtsX_{ECL1}$ with PcsB-CC is essential for cell growth

and proper morphology.   Having identified the interaction region between $FtsX_{ECL1}$

and PcsB-CC, we next sought to determine the degree to which this interaction

interface mapped by NMR spectroscopy contributes to pneumococcal viability. A

multiple sequence alignment of this region (residues 102-155) among bacterial species

in which FtsX has been studied and in related Streptococcal species (Fig. S7A) reveals

that amino acids in this region are either partially or completely conserved (Fig. S7A).

We therefore decided to target E109, Q111, L115, M119, W123, F126, and N131 singly

or in combination for substitution with alanine (Fig. S7A). Given the essentiality of the

$FtsX_{ECL1}$ and PcsB-CC interaction, we predicted that mutating these residues might be

lethal (20). To allow for the cross-in of potentially lethal point mutations, we employed

the Janus cassette method to insert point mutations at the native site of $ftsX$ into a strain

containing an ectopic copy of $ftsX^+$ under a zinc inducible promoter (33) (Fig. S7B). We

then transformed markerless mutant alleles of $ftsX$ in the presence of zinc. This allows

for expression of the wild-type $ftsX^+$ and mutant $ftsX$ simultaneously. As long as the

mutant $ftsX$ is not dominant negative, we could obtain a strain that expresses the wild-

type copy of $ftsX^+$ under zinc induction and mutant $ftsX$ only in the absence of zinc (Fig.

S7B).

Zinc toxicity has been observed to cause aberrant cell morphology and growth

inhibition in *S. pneumoniae*, when cells are not supplemented with manganese (34, 35).

To rule out any deleterious effects of the zinc/manganese (Zn/Mn) addition used to

induce $ftsX$ expression, we measured the growth of the parent and the FtsX merodiploid

strain in the presence of these metals. To verify that the addition of 0.45 mM $ZnCl_2$ and

0.045 mM $MnSO_4$ (Zn/Mn) did not cause growth or morphological defects, cells were

grown in the presence and absence of Zn/Mn and imaged at 3 h and 6 h into the growth

curve (Fig. S7C-D). Wild-type cells (D39 Δ*cps rpsL1*) had no morphological or growth

defects at these time points with or without the addition of Zn/Mn (Fig. S7C-E).

In contrast, the FtsX merodiploid strain (P_{zn}-*ftsX* Δ*ftsX*) had significant

morphological and growth defects at 3 or 6 h in the absence of Zn/Mn (Fig. S7C-E).

Cessation of growth and aberrant cell morphology was observed in 90% of cells at 3 h

and 95% of cells at 6 h growth (Fig. S7D). These cells are significantly shorter and

rounder than wild-type cells (Fig. S7E), and a large variability in their volumes was

observed (Fig. S7E), as previously found for a strain expressing $ftsX^+$ under a fucose-

inducible promoter (21). If the strain was grown in the presence of Zn/Mn, FtsX was

expressed and the strain has no growth or morphological defects (Fig. S7C-E). This

indicates the defects observed are solely due to the absence of FtsX.

We next constructed three classes of amino acid substitution or insertion mutants

(Table 4) in an effort to disrupt the $FtsX_{ECL1}$-PcsB-CC interaction defined by NMR

spectroscopy. These are designated Class I (single amino acid changes), Class II

(multiple amino acid changes), and Class III (insertion of a $(Gly_3Ser)_2$ linker) mutants.

Class I strains were made by introducing single amino acid substitutions in the

merodiploid strain and measuring growth or morphology defects (Fig. S8, Table 4).

Class I mutants targeted both the α2 helix and the loop region (residues 107-120, and

121-130, respectively) of the $FtsX_{ECL1}$ helical lobe (Table 4, Fig. 3A, S8A). Single amino

acid substitutions of FtsX (E109A, L115A, M119A, W123A, F126A, N131A, N131D)

exhibited morphological defects without Zn/Mn (Fig. S8C-D, Table 4), but did not induce

a measurable growth phenotype (Fig. S8B and Table 4). Expression of FtsX (L115A)

resulted in cell shape defects (aspect ratio, length, width, and volume) (Fig. S8C-D),

while expression of FtsX (M119A) only resulted in a change in cell volume (Fig. S8D).

These differences were not due to mis-expression of FtsX, as western blotting indicates

that all were expressed at or near wild-type levels (Fig. 4C). Two other single amino

acid substitutions (E109Q and Q111A) did not strongly affect growth, morphology, or

expression (Table 4, Fig. 4D).

In contrast to the somewhat modest physiological impact of Class I substitutions,

selected Class II mutants (Table 4) exhibited severe morphological and growth defects

(Fig. 3 and S9, Table 4). In strains targeting the α2 helix, FtsX(E109A, Q111A, L115A,

M119A), or the coil, FtsX(F126A, W123A, N131A), ≥98% of cells had severe growth

and morphology defects. The growth and morphology of these strains was similar to

ftsX depleted cells (Fig. 3B-D). Cells expressing these *ftsX* alleles became significantly

rounder and shorter (Fig. 3D), and growth was inhibited without zinc (Fig. 3B).

Importantly, in the presence of zinc the cells were indistinguishable from wild-type at 3 h

(Fig. 3D). At 6 h with zinc, these cells exhibited changes in width and volume, which

could be due to overexpression of FtsX at this time point or expression of wild-type and

mutant FtsX simultaneously (Fig. 3D). Western blotting confirmed that FtsX(E109A,

Q111A, L115A, M119A) and FtsX(F126A, W123A, N131A) were expressed in the

absence of zinc at 6 h post-depletion (Fig. 4A). The triple FtsX(F126A, W123A, N131A)

mutant migrated slightly higher on an SDS-PAGE gel compared to wild-type FtsX in the

absence of zinc, but was expressed (Fig. 4A). Taken together, these results reveal that

both the α2 helix and loop in the helical lobe of FtsX$_{ECL1}$ are important for FtsX function

in vivo and confirm the functional importance of the physical interaction of FtsX and

PcsB mapped by NMR.

Some of the Class II mutants we characterized had just two amino acid changes

in the α2 helix or the extended loop of the helical lobe (Table 4, Fig. S9). We observed

that FtsX (L115A, M119A) exhibited a strong growth and morphology defect in the

absence of zinc in 99% of cells at 6 h (Fig. S9B-C, Table 4), and these cells displayed a

decrease in length, width, and volume relative to wild-type cells (Fig. S9D). This mutant

was expressed in the absence of zinc at near wild-type levels (Fig. 4B). These data

confirm that the tandem L115A, M119A substitution disrupts FtsX function, even though

as individual single mutants they result only in slight morphological defects (Fig. S8B-D,

Fig. S9B-D, Table 4). Another double mutant FtsX(Q111A, L115C) induced the

formation of long chains and a "boxy" cell morphology (Fig. S9C, Table 4). This mutant

resulted in shorter cells with a significantly different aspect ratio from wild-type (Fig.

S9D), but this strain had no growth phenotype (Fig. S9B). In contrast, another double

substitution Class II mutant, *e.g.,* FtsX(E109A, N131A) (Table 4) exhibited no strong

morphology or growth defects.

Finally, Class III insertion mutants (Table 4) were constructed and used to

evaluate if other regions of FtsX$_{ECL1}$ were important for the FtsX$_{ECL1}$-PcsB interaction or

for FtsX function. We inserted a (Gly$_3$Ser)$_2$ flexible linker approximately where the

FtsX$_{ECL1}$ is predicted to enter (residue 51) or exit (residue 173) the membrane bilayer, or

in the β-hairpin, which exhibits significant conformational disorder over a range of

timescales (Fig. 1A). An insertion after residue 51 in FtsX$_{ECL1}$ was detrimental to both

growth and morphology (Table 4), and this insertion did not disrupt FtsX expression

(Fig. 4E). The insertion after residue 173 in FtsX$_{ECL1\ also}$ caused growth and morphology

defects, but this FtsX allele was not expressed in cells (Table 4, Fig. 4E). The β-hairpin

(Gly$_3$Ser)$_2$ insertion (Fig. 1B, Appendix) was introduced after amino acid 78 of FtsX$_{ECL1}$,

which corresponds to the tip of β-turn in the β-hairpin (Fig. 1B). This strain also

exhibited no growth defect, but these cells were significantly smaller, but only at the 3 h

time point (Appendix). We conclude that the β-hairpin does not play a major role in

FtsX-PcsB interaction, consistent with the NMR mapping experiments.

L115A/M119A FtsX$_{ECL1}$ is stably folded and unable to bind PcsB-CC. We

reasoned that if the defects observed in Class I and Class II mutants were due to the

disruption of the FtsX$_{ECL1}$-PcsB-CC interaction, this should impact the affinity of this

interaction as measured by ITC. We first characterized the L115A/M119A mutant by [1]H-

[15]N NMR and CD spectroscopies to confirm its structural integrity. The CD spectrum

resembled wild type, as did the [1]H-[15]N HSQC spectrum, with clear chemical shift

perturbations only among those resonances in the immediate vicinity of the double

substitution (Fig. S10B). Both pieces of data suggest a local rather than global

perturbation of α2-loop lobe in the FtsX$_{ECL1}$ structure upon introduction of the double

L115A/M119A substitution. As anticipated, titration of PcsB-CC (47-267) with

L115A/M119A FtsX$_{ECL1}$ reveals no detectable binding (no observable heat) (Fig. 2B,

Table 3) compared to wild-type FtsX$_{ECL1}$. These data confirm that the helical lobe of

FtsX$_{ECL1}$ interacts with PcsB-CC, and that this interaction is required for viability and

proper cell shape.

In contrast to the L115A/M119A double mutant, severely functionally

compromised representative triple mutant W123A, F126A, N131A FtsX$_{ECL1}$ and

quadruple mutant E109A, Q111A, L115A, M119A FtsX$_{ECL1}$ exhibited more pronounced

structural perturbations that nonetheless map only to the helical lobe. The triple mutant

was indistinguishable from the L115A/M119A and wild-type FtsX$_{ECL1}$ derivatives by CD

spectroscopy while the quadruple mutant exhibited less molar ellipticity, or secondary

structure (Fig. S10D). Inspection of their [1]H-[15]N HSQC spectra reveal that although the

core and β-hairpin domains essentially resemble wild-type, each exhibits considerable

perturbation of resonances throughout the entire helical lobe (Fig. S10A, S10C). Since

these mutants are functionally compromised, these structural findings strongly support

the conclusion that the structural integrity of the lower lobe is essential for the physical

interaction with PcsB, and the function of FtsX in pneumococcal cells.

DISCUSSION

This study presents a comprehensive analysis of the solution and x-ray

structures of the outward-facing large extracellular loop of FtsX (FtsX$_{ECL1}$) from *S.*

pneumoniae and defines a physical interaction site with the coiled-coil domain of

peptidoglycan hydrolase PcsB (PcsB-CC). Our FtsX$_{ECL1}$ structures reveal a globular fold

that while similar to the large extracellular loop of FtsX from *M. tuberculosis* (16) is

characterized by unique features. The upper β-hairpin distinguishes *S. pneumoniae*

FtsX$_{ECL1}$ from that of *M. tuberculosis*, and despite being characterized by significant

conformational dynamics on a range of timescales, is not required for the interaction of

S. pneumoniae FtsX$_{ECL1}$ with PcsB. The function of this domain is not well-defined by

our data, but could play a role in another process, *e.g.,* FtsX dimerization, interaction

with the small loop ECL2, or an interaction with another domain of PcsB. On the other

hand, the helical lobe of FtsX$_{ECL1}$, common to both *S. pneumoniae* and *M. tuberculosis*

FtsX structures, is vital for the interaction PcsB *in vitro*, and that this interface is

functionally important *in vivo*. Increasing numbers of Ala substitutions tested here

increasingly disrupt this interaction and ultimately cause dramatic growth and

morphology defects, indicating that the helical lobe of FtsX$_{ECL1}$ is essential for regulation

of PcsB during cell division. Interestingly, this region of $FtsX_{ECL1}$ corresponds with the

region shown to be important for the interaction of *M. tuberculosis* $FtsX_{ECD}$ with its PG

hydrolase, RipC (16). This suggests that the helical lobe could be a conserved

functional determinant for the interaction of FtsX with cognate hydrolases or adaptor

proteins across many species of bacteria.

We propose that the helical lobe of $FtsX_{ECL1}$ is important for the activation of

cognate hydrolase activity either directly or indirectly through their adaptors (Fig. 5). The

exact role of the second extracellular loop of FtsX ($FtsX_{ECL2}$) is unknown, but may also

regulate this process, as temperature-sensitive mutations in *pcsB* were found to be

suppressed by mutations in the coding region for $FtsX_{ECL2}$ (20). Previous work suggests

that FtsEX forms a dimer (32), as dimerization of the FtsE ATPase domain is likely a

necessary condition for ATP hydrolysis (36, 37) (Fig. 5A). After formation of the

complex, ATP hydrolysis by FtsE results in a conformational change in FtsX, releasing

PcsB from what we anticipate is an inhibited state (22) (Fig. 5B-C). This interaction is

mediated by the helical lobe of $FtsX_{ECL1}$, although the membrane, and possibly lipid

binding by $FtsX_{ECL1}$ and $FtsX_{ECL2}$ itself also may play a role as well. We propose that the

interaction of $FtsX_{ECL1}$ with the PcsB coiled-coil domain communicates release of the

PcsB CHAP domain from an inhibited state and thus is important for modulating PG

hydrolysis by PcsB (Fig. 5B-C).

Recently, the structure of *A. actinomycetemcomitans* MacB was reported and

suggested to be a structural paradigm for the ABC3 transporter superfamily that

includes FtsX (38). MacB was proposed to function as a mechanical pump to drive

enterotoxin transport through TolC in *E. coli* (38). Crow *et. al* found that MacB itself did

not transport enterotoxin but drove TolC to transport it instead, due to the lack of a

central cavity in the MacB structure (38). As such, they proposed that MacB as a model

for other "mechanotransmitters" belonging to this same ABC3 transporter superfamily.

While this proposed function of mechanotransmission may well characterize MacB and

FtsX, the overall structure of FtsX$_{ECL1}$ from *S. pneumoniae* is clearly distinct from the

periplasmic domain of MacB, thus revealing that MacB does not readily provide a

structural basis for understanding FtsX-dependent peptidoglycan hydrolases. Future

work using reconstituted FtsEX and PcsB complexes in membranes will allow for

understanding how this common mechanotransmission principle extends throughout the

ABC3 superfamily.

MATERIALS AND METHODS

NMR Spectroscopy. Spectra of ^{15}N- or ^{15}N^{13}C-labeled FtsX$_{ECL1}$ were recorded at 298 K

on Varian (Agilent) DDR 600 or 800 MHz spectrometers equipped with cryogenic

probes in the METACyt Biomolecular NMR Laboratory at Indiana University

Bloomington. NMR samples contained 50 mM potassium phosphate, pH 7.0, 50 mM

NaCl and 10% v/v D$_2$O, with 0.2 mM 4,4-dimethyl-4-silapentane-1-sulfonic acid (DSS)

for chemical shift referencing. Typical concentrations of FtsX$_{ECL1}$ for ^{15}N HSQC spectra

were 50 μM, and 400 μM for triple-resonance and dynamics experiments. $^1J_{HN}$ splittings

for residual dipolar couplings (RDCs) were measured using 2D IPAP [^{15}N, ^{1}H]-HSQC

spectra (39), recorded on an isotropic sample and on a sample aligned with 20 mg/mL

phage *Pf*1 (ASLA Biotech). $^1D_{HN}$ was calculated from $^1D_{HN} = {}^1J_{HN}$ (anisotropic) $- {}^1J_{HN}$

(isotropic). Aromatic sidechains were assigned using the HBCBCGCDHE and

HBCBCGCDCEHE experiments (40). For experiments detecting PcsB-CC binding, ^{15}N

FtsX$_{ECL1}$ was kept at 50 µM concentration, and ^{1}H-^{15}N HSQC spectra were recorded

with the following concentrations of PcsB-CC (47-267): 0, 50 µM, 100 µM, 200 µM, and

400 µM. nmrPipe, Sparky, CARA (http://cara.nmr.ch), CCPNMR, and NMRbox (41-44)

were used for data processing and analysis. Resonance assignments and dynamics

data are available in the BMRB under accession code 30523. These NMR data were

used to calculate and refine the solution structure of ECL1 (see *Supplemental Materials*

and Methods) with the ensemble of 20 lowest energy structures (see Table 1 for

structure statistics) deposited in the Protein Data Bank (accession code 6MK7).

X-Ray Crystallography. First crystallization screenings were performed by high-

throughput techniques in a NanoDrop robot and Innovadyne SD-2 microplates

(Innovadyne Technologies Inc.), screening PACT Suite and JCSG Suite (Qiagen),

JBScreen Classic 1-4 and 6 (Jena Bioscience) and Crystal Screen, Crystal Screen 2

and Index HT (Hampton Research). Positive conditions in which crystals grew were

optimized by sitting-drop vapor-diffusion method at 291 K by mixing 1 µL of protein

solution and 1 µL of precipitant solution, equilibrated against 150 µL of precipitant

solution in the reservoir chamber. The best crystals were obtained in a crystallization

condition containing 0.1 M sodium citrate pH=5.6, 0.2 M potassium-sodium tartrate and

2 M ammonium sulfate. These crystals were further optimized by cocrystallization in the

presence of detergents Dodecyltrimethylammonium chloride (detergent 1, 46 µM), n-

Undecyl-β-D-maltoside (detergent 2, 0.59 mM), and n-Decyl-β-D-maltoside (detergent

3, 1.8 mM). Crystals were cryo-protected in the precipitant solution supplemented with

25% (v/v) glycerol, prior to flash cooling at 100 K. Diffraction data were collected in

beamline XALOC at the ALBA synchrotron (CELLS-ALBA, Spain), using a Pilatus 6M

detector and a wavelength of 0.979257 Å. Crystals diffracted up to 2.0-2.3 Å resolution

and belonged to the P 4_3 2_1 2 space group. The collected datasets were processed with

XDS (45) and Aimless (46). Two FtsX$_{ECL1}$ molecules were found in the asymmetric unit,

yielding a Matthews coefficient of 2.49 Å^3/Da (47) and a solvent content of 50.73%.

Structure determination was performed by molecular replacement using the online

server Morda (http://www.biomexsolutions.co.uk/morda/). Refinement and manual

model building were performed with Phenix (48) and Coot (49), respectively. The data

presented translational non-crystallographic symmetry that were treated with Phenix

(48). The stereochemistry of the final model was checked by MolProbity (50). Data

collection and processing statistics are shown in Table 2. The atomic coordinates of

FtsX$_{ECL1}$ determined by co-crystallization in the presence of detergents 1, 2 and 3 have

been deposited in the Protein Data Bank with codes 6HE6, 6HEE and 6HFX,

respectively.

Bacterial strains, plasmids, and growth conditions. Bacterial strains and plasmids

used in this study are listed in Table S1. *S. pneumoniae* strains were derived from

IU1945, an unencapsulated derivative of serotype 2 *S. pneumoniae* strain D39 (51).

Strains were grown on trypticase soy agar II with 5% (vol/vol) defibrinated sheep blood

(TSAII-BA) plates or in Becton-Dickinson brain heart infusion (BHI) broth at 37°C in an

atmosphere of 5% CO_2. *E. coli* strains for protein expression were derived from

BL21(DE3) (NEB, C2527H). *E. coli* strains were grown in Luria-Bertani (LB) broth or in

M9 minimal media supplemented with $^{15}NH_4Cl$ at 37°C, with shaking at 150 rpm. When

required, tetracycline (0.25-2.5 µg/mL), kanamycin (250 µg/mL), spectinomycin (150

μg/mL), streptomycin (250 μg/mL), ampicillin (100 μg/mL) and/or isopropyl β-D-1-

thiogalactopyranoside (IPTG, 1 mM) were added to *S. pneumoniae* or *E. coli* culture

media. *S. pneumoniae* strains requiring zinc for expression of essential genes were

grown with 0.45 mM $ZnCl_2$ and 0.045 $MnSO_4$.

Growth curves and phase-contrast microscopy of strains. For physiological and

morphological analyses of strains, cells were inoculated from frozen glycerol stocks into

BHI broth, serially diluted, and incubated for 10-12 hours statically at 37°C in 5% CO2

overnight. If zinc was required for growth of cultures, 0.45 mM $ZnCl_2$ and 0.045 $MnSO_4$

was added to overnight tubes. The next day, cultures from OD_{620} ≈ 0.05 to 0.4 were

diluted into fresh BHI to OD_{620} ≈ 0.003 in 4 mL volumes, and two identical cultures for

each strain were prepared, one with 0.45 mM $ZnCl_2$/0.045 $MnSO_4$ and one without.

These cultures were grown under the same growth conditions as previously described

in *Materials and Methods*. Growth was monitored turbidimetrically every 45 min to 1

hour with a Genesys 2 spectrophotometer (Thermo Scientific). For microscopic

analyses, samples (1-2 μL) were taken at 3 hours and 6 hours and examined using a

Nikon E-400 epifluorescence phase-contrast microscope with a 100X Nikon Plan Apo

oil-immersion objective (numerical aperture, 1.40) connected to a CoolSNAP HQ2

charged-coupled device (CCD) camera (Photometrics). Images were processed using

NIS-Elements AR software (Nikon), and measurements and calculation of cell width,

length, volume, and aspect ratio were performed as described previously (52, 53).

Statistical significance was determined using GraphPad Prism (GraphPad Software,

Inc) by comparing values for cell width, length, volume, and aspect ratio measured for at

least 50 cells over two experimental replicates. To determine if values were significantly

different between strains and conditions, a Kruskal-Wallis (one-way ANOVA) with Dunn's multiple comparison post-test was used.

For additional materials and methods, please see Supplemental Materials and Methods.

ACKNOWLEDGEMENTS

We thank the members Winkler and Giedroc labs for their helpful discussion and insight, specifically Dr. Tiffany Tsui, Dr. Julia Martin, Melissa Lamanna and Dr. Hui Peng. We acknowledge the Indiana University Bloomington Department of Chemistry Mass Spectrometry Facility, specifically Dr. Jonathan Karty and Angela Hansen for their help with training and setting up mass spectrometry experiments and the MetaCyt Biomolecular NMR Laboratory. We also thank Dr. Daiana Capdevila for her help in acquiring and analyzing the isothermal titration calorimetry data, the Indiana University Biological Mass Spectrometry Facility and Dr. Giovanni Gonzales-Gutierrez in the Physical Biochemistry Instrumentation Facility at Indiana University Bloomington. We thank the staff from ALBA synchrotron facility for help during crystallographic data collection. This work was supported by NIH grants R01GM114315 and R01GM127715 to M.E.W., and NIH grant R35GM118157 to D.P.G. and by predoctoral Quantitative and Chemical Biology (QCB) NIH institutional training grant T32 GM109825 (to B.E.R.). The work in Spain was supported by grant from the Spanish Ministry of Science, Innovation and Universities BFU2017-90030-P to J.A.H.

REFERENCES

1.   Henriques-Normark B, Normark S. 2010. Commensal pathogens, with a focus on Streptococcus pneumoniae, and interactions with the human host. Exp Cell Res 316:1408-14.

2.   Kadioglu A, Weiser JN, Paton JC, Andrew PW. 2008. The role of Streptococcus pneumoniae virulence factors in host respiratory colonization and disease. Nat Rev Microbiol 6:288-301.

3.   Chao Y, Marks LR, Pettigrew MM, Hakansson AP. 2014. Streptococcus pneumoniae biofilm formation and dispersion during colonization and disease. Front Cell Infect Microbiol 4:194.

4.   Feldman C, Anderson R. 2014. Recent advances in our understanding of Streptococcus pneumoniae infection. F1000Prime Rep 6:82.

5.   CDC. 2013. Antibiotic resistance threats in the United States, 2013. http://www.cdc.gov/drugresistance/threat-report-2013/,

6.   Garcia-Bustos J, Tomasz A. 1990. A biological price of antibiotic resistance: major changes in the peptidoglycan structure of penicillin-resistant pneumococci. Proc Natl Acad Sci U S A 87:5415-9.

7.   Massidda O, Novakova L, Vollmer W. 2013. From models to pathogens: how much have we learned about Streptococcus pneumoniae cell division? Environ Microbiol 15:3133-57.

8.   Sham LT, Tsui HC, Land AD, Barendt SM, Winkler ME. 2012. Recent advances in pneumococcal peptidoglycan biosynthesis suggest new vaccine and antimicrobial targets. Curr Opin Microbiol 15:194-203.

9.   Egan AJ, Cleverley RM, Peters K, Lewis RJ, Vollmer W. 2017. Regulation of bacterial cell wall growth. FEBS J 284:851-867.

10.   Vollmer W, Blanot D, de Pedro MA. 2008. Peptidoglycan structure and architecture. FEMS Microbiol Rev 32:149-67.

11.   Rajagopal M, Walker S. 2017. Envelope Structures of Gram-Positive Bacteria. Curr Top Microbiol Immunol 404:1-44.

12.   Rojas ER, Huang KC. 2018. Regulation of microbial growth by turgor pressure. Curr Opin Microbiol 42:62-70.

13.   Holtje JV. 1998. Growth of the stress-bearing and shape-maintaining murein sacculus of Escherichia coli. Microbiol Mol Biol Rev 62:181-203.

14.   Holtje JV, Heidrich C. 2001. Enzymology of elongation and constriction of the murein sacculus of Escherichia coli. Biochimie 83:103-8.

15.   Alcorlo M, Martinez-Caballero S, Molina R, Hermoso JA. 2017. Carbohydrate recognition and lysis by bacterial peptidoglycan hydrolases. Curr Opin Struct Biol 44:87-100.

16.   Mavrici D, Marakalala MJ, Holton JM, Prigozhin DM, Gee CL, Zhang YJ, Rubin EJ, Alber T. 2014. Mycobacterium tuberculosis FtsX extracellular domain activates the peptidoglycan hydrolase, RipC. Proc Natl Acad Sci U S A doi:10.1073/pnas.1321812111.

17.   Margulieux KR, Liebov BK, Tirumala V, Singh A, Bushweller JH, Nakamoto RK, Hughes MA. 2017. Bacillus anthracis Peptidoglycan Integrity Is Disrupted by the Chemokine CXCL10 through the FtsE/X Complex. Front Microbiol 8:740.

18. Meier EL, Daitch AK, Yao Q, Bhargava A, Jensen GJ, Goley ED. 2017. FtsEX-
mediated regulation of the final stages of cell division reveals morphogenetic
plasticity in Caulobacter crescentus. PLoS Genet 13:e1006999.

19. Zielinska A, Billini M, Moll A, Kremer K, Briegel A, Izquierdo Martinez A, Jensen
GJ, Thanbichler M. 2017. LytM factors affect the recruitment of autolysins to the
cell division site in Caulobacter crescentus. Mol Microbiol 106:419-438.

20. Sham LT, Jensen KR, Bruce KE, Winkler ME. 2013. Involvement of FtsE ATPase
and FtsX extracellular loops 1 and 2 in FtsEX-PcsB complex function in cell
division of Streptococcus pneumoniae D39. MBio 4.

21. Sham LT, Barendt SM, Kopecky KE, Winkler ME. 2011. Essential PcsB putative
peptidoglycan hydrolase interacts with the essential FtsXSpn cell division protein
in Streptococcus pneumoniae D39. Proc Natl Acad Sci U S A 108:E1061-9.

22. Bartual SG, Straume D, Stamsas GA, Munoz IG, Alfonso C, Martinez-Ripoll M,
Havarstein LS, Hermoso JA. 2014. Structural basis of PcsB-mediated cell
separation in Streptococcus pneumoniae. Nat Commun 5:3842.

23. Meisner J, Montero Llopis P, Sham LT, Garner E, Bernhardt TG, Rudner DZ.
2013. FtsEX is required for CwlO peptidoglycan hydrolase activity during cell wall
elongation in Bacillus subtilis. Mol Microbiol 89:1069-83.

24. Yang DC, Peters NT, Parzych KR, Uehara T, Markovski M, Bernhardt TG. 2011.
An ATP-binding cassette transporter-like complex governs cell-wall hydrolysis at
the bacterial cytokinetic ring. Proc Natl Acad Sci U S A 108:E1052-60.

25. Yang DC, Tan K, Joachimiak A, Bernhardt TG. 2012. A conformational switch

controls cell wall-remodelling enzymes required for bacterial cell division. Mol

Microbiol 85:768-81.

26. Du S, Pichoff S, Lutkenhaus J. 2016. FtsEX acts on FtsA to regulate divisome

assembly and activity. Proc Natl Acad Sci U S A 113:E5052-61.

27. Schmidt KL, Peterson ND, Kustusch RJ, Wissel MC, Graham B, Phillips GJ,

Weiss DS. 2004. A predicted ABC transporter, FtsEX, is needed for cell division

in Escherichia coli. J Bacteriol 186:785-93.

28. Reddy M. 2007. Role of FtsEX in cell division of Escherichia coli: viability of ftsEX

mutants is dependent on functional SufI or high osmotic strength. J Bacteriol

189:98-108.

29. Fu Y, Bruce KE, Rued B, Winkler ME, Giedroc DP. 2016. 1H, 13C, 15N

resonance assignments of the extracellular loop 1 domain (ECL1) of

Streptococcus pneumoniae D39 FtsX, an essential cell division protein. Biomol

NMR Assign 10:89-92.

30. Zweckstetter M. 2008. NMR: prediction of molecular alignment from structure

using the PALES software. Nat Protoc 3:679-90.

31. Ishima R. 2014. CPMG relaxation dispersion. Methods Mol Biol 1084:29-49.

32. Bajaj R, Bruce KE, Davidson AL, Rued BE, Stauffacher CV, Winkler ME. 2016.

Biochemical characterization of essential cell division proteins FtsX and FtsE that

mediate peptidoglycan hydrolysis by PcsB in Streptococcus pneumoniae.

Microbiologyopen 5:738-752.

33.   Sung CK, Li H, Claverys JP, Morrison DA. 2001. An rpsL cassette, janus, for
592   gene replacement through negative selection in Streptococcus pneumoniae. Appl
593   Environ Microbiol 67:5190-6.

594   34.   Jacobsen FE, Kazmierczak KM, Lisher JP, Winkler ME, Giedroc DP. 2011.
595   Interplay between manganese and zinc homeostasis in the human pathogen
596   Streptococcus pneumoniae. Metallomics 3:38-41.

597   35.   Martin JE, Lisher JP, Winkler ME, Giedroc DP. 2017. Perturbation of manganese
598   metabolism disrupts cell division in Streptococcus pneumoniae. Mol Microbiol
599   104:334-348.

600   36.   Locher KP. 2016. Mechanistic diversity in ATP-binding cassette (ABC)
601   transporters. Nat Struct Mol Biol 23:487-93.

602   37.   Rees DC, Johnson E, Lewinson O. 2009. ABC transporters: the power to
603   change. Nat Rev Mol Cell Biol 10:218-27.

604   38.   Crow A, Greene NP, Kaplan E, Koronakis V. 2017. Structure and
605   mechanotransmission mechanism of the MacB ABC transporter superfamily.
606   Proc Natl Acad Sci U S A 114:12572-12577.

607   39.   Ottiger M, Delaglio F, Bax A. 1998. Measurement of J and dipolar couplings from
608   simplified two-dimensional NMR spectra. J Magn Reson 131:373-8.

609   40.   Yamazaki T, Yoshida M, Nagayama K. 1993. Complete assignments of magnetic
610   resonances of ribonuclease H from Escherichia coli by double- and triple-
611   resonance 2D and 3D NMR spectroscopies. Biochemistry 32:5656-69.

612   41.   Lee W, Tonelli M, Markley JL. 2015. NMRFAM-SPARKY: enhanced software for
613   biomolecular NMR spectroscopy. Bioinformatics 31:1325-7.

42.   Vranken WF, Boucher W, Stevens TJ, Fogh RH, Pajon A, Llinas M, Ulrich EL,

Markley JL, Ionides J, Laue ED. 2005. The CCPN data model for NMR

spectroscopy: development of a software pipeline. Proteins 59:687-96.

43.   Maciejewski MW, Schuyler AD, Gryk MR, Moraru, II, Romero PR, Ulrich EL,

Eghbalnia HR, Livny M, Delaglio F, Hoch JC. 2017. NMRbox: A Resource for

Biomolecular NMR Computation. Biophys J 112:1529-1534.

44.   Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeifer J, Bax A. 1995. NMRPipe: a

multidimensional spectral processing system based on UNIX pipes. J Biomol

NMR 6:277-93.

45.   Kabsch W. 2010. Xds. Acta Crystallogr D Biol Crystallogr 66:125-32.

46.   Evans PR, Murshudov GN. 2013. How good are my data and what is the

resolution? Acta Crystallogr D Biol Crystallogr 69:1204-14.

47.   Matthews BW. 1968. Some crystal forms of bovine chymotrypsinogen B and

chymotrypsinogen A. J Mol Biol 33:499-501.

48.   Adams PD, Afonine PV, Bunkoczi G, Chen VB, Davis IW, Echols N, Headd JJ,

Hung LW, Kapral GJ, Grosse-Kunstleve RW, McCoy AJ, Moriarty NW, Oeffner

R, Read RJ, Richardson DC, Richardson JS, Terwilliger TC, Zwart PH. 2010.

PHENIX: a comprehensive Python-based system for macromolecular structure

solution. Acta Crystallogr D Biol Crystallogr 66:213-21.

49.   Emsley P, Lohkamp B, Scott WG, Cowtan K. 2010. Features and development of

Coot. Acta Crystallogr D Biol Crystallogr 66:486-501.

50.   Chen VB, Arendall WB, 3rd, Headd JJ, Keedy DA, Immormino RM, Kapral GJ,

Murray LW, Richardson JS, Richardson DC. 2010. MolProbity: all-atom structure

validation for macromolecular crystallography. Acta Crystallogr D Biol Crystallogr

66:12-21.

51.   Lanie JA, Ng WL, Kazmierczak KM, Andrzejewski TM, Davidsen TM, Wayne KJ,

Tettelin H, Glass JI, Winkler ME. 2007. Genome sequence of Avery's virulent

serotype 2 strain D39 of Streptococcus pneumoniae and comparison with that of

unencapsulated laboratory strain R6. J Bacteriol 189:38-51.

52.   Land AD, Tsui HC, Kocaoglu O, Vella SA, Shaw SL, Keen SK, Sham LT, Carlson

EE, Winkler ME. 2013. Requirement of essential Pbp2x and GpsB for septal ring

closure in Streptococcus pneumoniae D39. Mol Microbiol 90:939-55.

53.   Tsui HC, Zheng JJ, Magallon AN, Ryan JD, Yunck R, Rued BE, Bernhardt TG,

Winkler ME. 2016. Suppression of a deletion mutation in the gene encoding

essential PBP2b reveals a new lytic transglycosylase involved in peripheral

peptidoglycan synthesis in Streptococcus pneumoniae D39. Mol Microbiol

100:1039-65.

TABLE 1. Structural statistics for NMR solution structure of
$FtsX_{ECL1}$ (From the ensemble of 20 best NMR structures).

NMR structure (PDB code)	6MK7
(BMRB code)	30523

NMR distance and angle restraints	
Total NOE-based distance constraints	1711
intraresidue (i=j)	232
sequential ($|i\text{-}j| = 1$)	499
medium-range ($1 < |i\text{-}j| < 5$)	365
long-Range ($|i\text{-}j| \geq 4$)	615
Maximum distance violation	0.49 Å
Dihedral angle restraints	110
Maximum dihedral angle violation	7°
Total number of RDCs used (measured)	82 (112)
Q factor	0.11
	(0.39)
Correlation (experimental to calculated)	0.99
	(0.89)

RMS deviations from idealized covalent geometry	
Covalent Bond Lengths (Å)	0.011 Å
Covalent Angle Values (°)	1.3 °
Ramachandran analysis [a,b]	
Most favored (%)	86.5
Additionally allowed (%)	8.4
Generously allowed (%)	1.8
Disallowed (%)	3.3
RMSD Values	
Backbone atoms (all)	1.5 Å
All heavy atoms (all)	1.9 Å
Backbone atoms (core[a])	0.4 Å
All heavy atoms (core[a])	0.9 Å

655   ^aCalculated over the core of the structure, excluding the helical

656   lobe and β-hairpin, residues 59-65, 89-108, 134-165. ^bAs

657   computed by Procheck.

660   **TABLE 2.** Crystallographic data collection and refinement statistics.[a]

	FtsX$_{ECL}$:D1	FtsX$_{ECL}$:D2	FtsX$_{ECL}$:D3
Data collection			
Wavelength (Å)	0.97934	0.97934	0.97934
Space group	P 4_3 2_1 2	P 4_3 2_1 2	P 4_3 2_1 2
Unit cell *a, b, c* (Å)	75.87, 75.87, 97.93	84.61, 84.61, 106.19	75.49, 75.49, 95.31
Unit cell α,β,γ (°)	90, 90, 90	90, 90, 90	90, 90, 90
T (K)	100	100	100
X-ray source	Synchrotron	Synchrotron	Synchrotron
Resolution range (Å)	47.05–(2.05–2.0)	44.97(2.38–2.3)	46.58–(2.23–2.16)
Unique reflections	19992 (1436)	17781 (1700)	15360 (1496)
Completeness (%)	100 (100)	100 (100)	99.85 (99.93)
Multiplicity	24.4 (25.9)	22.1 (22.5)	16.6 (21.8)
$R_{pim}{}^{d}$	0.013 (0.535)	0.007 (0.123)	0.010 (0.241)
$<I/\sigma(I)>$	21.9 (1.5)	41.7 (4.6)	24.6 (2.55)
CC1/2	1.00 (0.67)	1.00 (0.98)	1.00 (0.89)
Refinement			
Resolution range (Å)	41.14–2.0	42.31–2.3	46.58–2.16
$R_{work}/R_{free}{}^{c}$	0.24/ 0.28	0.26/ 0.30	0.25/ 0.31
No. Atoms			
Protein	1888	1732	1856
Water	25	21	23
Ligand	16	47	33
R.m.s. deviations			
Bond length (Å)	0.008	0.009	0.010
Bond angles (°)	1.27	1.22	1.29

Ramachandran			
Favored/outliers (%)	95.65/0.43	93.20/2.43	92.41/0.00
Monomers per AU	2	2	2
PDB code	6HE6	6HEE	6HFX

[a]Values between parentheses correspond to the highest resolution shells

[b]$R_{pim} = \Sigma_{hkl}[1/(N-1)]$ 1/2 $\Sigma_i \mid I_i(hkl) - [I(hkl)] \mid / \Sigma_{hkl}\Sigma_i I_i(hkl)$, where $\Sigma_i I_i(hkl)$ is the i-th

measurement of reflection hkl, [I(hkl)] is the weighted mean of all measurements

and N is the redundancy for the hkl reflection. [c]$R_{work}/R_{free} = \Sigma_{hkl} \mid F_o - F_c \mid / \Sigma_{hkl} \mid F_o \mid$,

where F_c is the calculated and F_o is the observed structure factor amplitude of

reflection hkl for the working / free (5%) set, respectively.

TABLE 3. Thermodynamic parameters of wild-type and mutant $FtsX_{ECL1}$ from direct analysis of isothermal titration calorimetry (ITC).[a]

Experiment	Protein	K_A (x 10^4 M^{-1})	K_D (µM)	ΔH (kcal mol^{-1})	ΔS (kcal mol^{-1} K^{-1})
ITC	$FtsX_{ECL1}$	3.0 ± 0.5	34 ± 6	-2.1 ± 0.2	13.4
ITC	L115A, M119A $FtsX_{ECL1}$	N/A[b]	N/A[b]	N/A[c]	N/A[c]

[a] Solution conditions: 50 mM sodium phosphate, 50 mM NaCl, 0.5 mM EDTA, pH 7.0 at 25 °C. The chemical model used to fit the data is indicated in *Supplemental Materials and Methods.* [b]No observed heat.

675 **TABLE 4.** Amino acid changes made *in vivo* to disrupt the FtsX$_{ECL1}$-PcsB interaction.[a]

	Categories of Amino Acid Changes Made		
	Amino Acid Change	Defect in Shape and Growth	Location in FtsX$_{ECL1}$
Class I	E109A	Shape Only	α2 helix
	E109Q	No	α2 helix
	Q111A	No[b]	α2 helix
	L115A	Shape Only	α2 helix
	M119A	Shape Only	α2 helix
	W123A	Shape Only	Loop[c]
	F126A	Shape Only	Loop[c]
	N131A	Shape Only	Loop[c]
	N131D	Shape Only	Loop[c]
Class II	E109A, N131A	Shape Only	α2 helix, Loop[c]
	Q111A, L115C	Shape Only	α2 helix
	L115A, M119A	Yes	α2 helix
	E109A, Q111A, L115A, M119A	Yes	α2 helix
	F126A, W123A, N131A	Yes	Loop[c]
Class III (Gly$_3$Ser)$_2$[d]	Residue 51	Yes	post-TM1
	Residue 78	Shape Only	β-hairpin
	Residue 173	N/A[e]	pre-TM2

[a]See Table S1 for corresponding strain number and strain construction for each

amino acid change or insertion. [b]In the absence of zinc, this strain had no

morphology defects. Morphology defects were observed at 3 h when the strain was

expressing both the wild-type copy of FtsX and FtsX(Q111A). [c]Loop refers to the

unstructured loop after $\alpha 2$ helix in the helical lobe of the $FtsX_{ECL1}$ structure. [d]These

strains an insertion of the amino acid sequence GGGSGGGS after the indicated

residue. [e]N/A: This strain did not express FtsX with $(Gly_3Ser)_2$ insertion. See Fig. 4E

for Western blot data.

Figure 1: The structure of FtsX_ECL1 from *Streptococcus pneumoniae*. A) Stereo view of the 20 conformers of the FtsX_ECL1 NMR structure as backbone traces, with helices shown in blue and β-strands shown in yellow. N- and C- termini of the domain are indicated by the residue numbers 57 and 166, respectively. B) Cartoon representation of the FtsX_ECL1 structure obtained by X-ray crystallography (chain B in FtsX_ECD-D1) in which the different secondary structure elements have been numbered and labeled. N- and C- termini of the domain are indicated and correspond to residue numbers 49 and 166, respectively. C) Stereo view showing changes in helical lobe upon interaction with detergent 1. Apo-form (chain B) is colored in pale brown and holo-form (chain A) is colored in green. Relevant residues affected by the presence of the detergent are depicted in capped sticks. Polar interaction represented by dashed line. D) Surface representation of the FtsX_ECL1 crystal structure in which the three different detergent molecules are superimposed and shown in sticks (detergent 1, 2 and 3 are colored in green orange and purple, respectively). E) An overlay of the backbone traces for the six FtsX_ECL1 crystal structures (in black) with the backbone traces of the 20 FtsX_ECL1 NMR conformers (in green) F) Structural superposition of the crystal structure of FtsX_ECL1 from *S. pneumoniae* obtained by crystallography (colored in cyan) and the crystal structure of FtsX_ECD from *M. tuberculosis* (PDB 4N8N, colored in orange). Secondary structure elements observed in the *M. tuberculosis* structure are labeled. Some residues in both structures are shown as sticks and numbered (see main text). Cys residues involved in disulfide bond formation in *M. tuberculosis* FtsX_ECD are colored in yellow. See main text for details. G) Sequence alignment of ECL1 domains from *S. pneumoniae* (Spn ECL1) and *M. tuberculosis* (Mtb ECD). Secondary structure elements are indicated and numbered.

Figure 2: FtsX$_{ECL1}$ binds PcsB-CC. A) Significant chemical shifts and peak height changes upon ^{1}H-^{15}N HSQC titration of 50 µM ^{15}N FtsX$_{ECL1}$ with 100 µM unlabeled PcsB-CC map to the lower lobe of FtsX$_{ECL1}$. Chemical shift (Δδ, ppm) and peak height changes are mapped as color gradients on the FtsX$_{ECL1}$ structure, orange to red and light grey to blue, respectively. L115 and M119 α carbons are shown as spheres on the upper image. Peaks that overlap in the ^{1}H-^{15}N HSQC spectra are colored white. Chemical shift and peak height change upon addition of 2 molar equivalents of PcsB-CC to FtsX$_{ECL1}$ are mapped to the structure. B) Representative titration of PcsB-CC with wild-type (WT) FtsX$_{ECL1}$ or L115A/M119A FtsX$_{ECL1}$ as monitored by ITC. Conditions: 50 mM potassium phosphate, 50 mM NaCl, 0.5 mM EDTA, pH 7.0 at 25.0 °C. Top panel, corrected ITC data; bottom panel, kcal/mole of injection vs. molar ratio. The black line overlapping the WT data indicates the best fit to a one-site binding model. Fitting parameters are summarized in Table 3. A red line drawn through the L115A/M119A FtsX$_{ECL1}$ data is for reference.

Figure 3: Multiple amino acid changes in the lower lobe of FtsX_ECL1 cause morphological and growth defects. A) Amino acid changes made mapped to the structure of FtsX_ECL1. The α carbon of each residue is shown as a colored sphere. The orange to red coloring on the FtsX_ECL1 structure represents the peak height changes from ^{1}H-^{15}N HSQC spectra upon addition of 2 molar equivalents of PcsB-CC to FtsX_ECL1 B) Representative growth curve of strains with amino acid changes in the lower lobe of FtsX_ECL1, compared to the growth of a FtsX depletion strain. These strains were grown in the presence or absence of 0.45 mM ZnCl_2 supplemented with 0.045 mM MnSO_4 (indicated as +Zn). Strains shown are as follows: black circle, D39 *rpsL1* Δ*cps* wild-type parent (1, IU1824); grey circle, IU1824 +Zn; dark blue square, D39 *rpsL1* Δ*cps* Δ*ftsX*::P-*aad9*//*bgaA*::*tet*-P_Zn-*ftsX*+ (2, IU12376); light blue square, IU12376 +Zn; red triangle, D39 *rpsL1* Δ*cps* *ftsX*(E109A, Q111A, L115A, M119A)//*bgaA*::*tet*-P_Zn-*ftsX*+ (3, IU12861); pink triangle, IU12861 +Zn; dark green inverted triangle, D39 *rpsL1* Δ*cps* *ftsX*(F126A, W123A, N131A)//*bgaA*::*tet*-P_Zn-*ftsX*+ (4, IU12864); light green inverted triangle, IU12864 +Zn. This growth curve was repeated three times with similar results. C) Representative images of strains at 6 hours growth. The genotype of the strain shownis indicated above each panel. No Zn or +Zn indicates if Zn/Mn mixture was added. % indicates the percentage of cells in the population that are morphologically similar to the images shown. Greater than 50 cells per strain, condition, and experimental repeat were analyzed. These experiments were performed three times independently with similar results. Scale bar shown is equal to 1 μM. D) Length, width, aspect ratio, and relative cell volume of strains at 3 hours and 6 hours growth. Strains are indicated according to numbering in panel B. Greater than 50 cells were measured per strain and condition over two experimental replicates. For statistical analysis, a Kruskal-Wallis test (one-way ANOVA) with Dunn's multiple comparison post-test was used to determine if length, width, aspect ratio, and relative cell volume were significantly different between strains and conditions. ns=non significant, * = p<0.05, ** = p<0.005, *** = p<0.0005.

Figure 4: FtsX with amino acid changes are expressed at near wild-type levels. Representative blots of α-FtsX and α-MreC (Western blotting control for loading) are shown, with the genotype indicated under each lane. Expected molecular weight (MW) for FtsX is 34.2 kDa, expected MW for MreC is 29.7 kDa. Samples were grown without Zn and harvested at 6 hours growth. Westerns were imaged as described in *Supplemental Materials and Methods*. A) FtsX(E109A, Q111A, L115A, M119A) and FtsX(W123A, F126A, N131A) are expressed at or above wild-type levels without zinc. Lane 1, D39 *rpsL1* Δ*cps* (IU1824); lane 2, D39 *rpsL1* Δ*cps* ftsX(E109A, Q111A, L115A M119A)//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU12861); lane 3, D39 *rpsL1* Δ*cps ftsX*(W123A, F126A, N131A)//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU12864); lane 4, D39 *rpsL1* Δ*cps* Δ*ftsX*//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU13461). B) FtsX(Q111A, L115C) and FtsX(L115A, M119A) are expressed at near wild-type levels without zinc. Lane 1, D39 *rpsL1* Δ*cps* (IU1824); lane 2, D39 *rpsL1* Δcps *ftsX*(Q111A, L115C)//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU13064); lane 3, D39 *rpsL1* Δ*cps ftsX*(L115A, M119A)//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU13066); lane 4, D39 *rpsL1* Δ*cps* Δ*ftsX*//*bgaA*::*tet*-P_Zn-*ftsX*+. C) FtsX (L115A) and FtsX (M119A) are expressed at near wild-type levels without zinc. Lane 1, D39 *rpsL1* Δ*cps* (IU1824); lane 2, D39 *rpsL1* Δ*cps ftsX*(L115A)//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU12521); lane 3, D39 *rpsL1* Δ*cps ftsX*(M119A)//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU12637); lane 4, D39 *rpsL1* Δ*cps* Δ*ftsX*//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU13461). D) FtsX (E109Q) and FtsX (N131D) are expressed at near wild-type levels without zinc. Lane 1, D39 *rpsL1* Δ*cps* (IU1824); lane 2, D39 *rpsL1* Δ*cps ftsX*(E109Q)//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU13088); lane 3, D39 *rpsL1* Δ*cps ftsX*(N131D)//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU13089); lane 4, D39 *rpsL1* Δ*cps* Δ*ftsX*//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU13461). E) FtsX with (Gly_4Ser)_2 after residue 51 is expressed, whereas FtsX with (Gly_4Ser)_2 after residue 173 is not expressed. These are referred to as FtsX N-term ECL1 (Gly_4Ser)_2 and FtsX C-terminal ECL1 (Gly_4Ser)_2, respectively. Lane 1, D39 *rpsL1* Δ*cps* (IU1824); lane 2, D39 *rpsL1* Δ*cps ftsX N-term ECL1 (Gly_4 Ser)_2*//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU12629); lane 3, D39 *rpsL1* Δ*cps ftsX C-term ECL1 (Gly_4 Ser)_2*//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU12869); lane 4, D39 *rpsL1* Δ*cps* Δ*ftsX*//*bgaA*::*tet*-P_Zn-*ftsX*+ (IU13461). These experiments were performed two to three times independently.

Figure 5: Model for the activation of PcsB by FtsX$_{ECL1}$. A) FtsEX dimerizes to form the active complex. PcsB is secreted into the extracellular milieu. Attraction of PcsB to the area of active FtsX complexes might be mediated by its propensity to interact with membranes (Bajaj et. al, 2017). The ECL1 and ECL2 loops are indicated on FtsX. FtsX$_{ECL1}$ is shown in green, with the β-hairpin and α-helical lobe shaded in yellow and blue, respectively. B) After formation of the active complex, ATP hydrolysis by FtsE causes a conformational change in FtsX. PcsB interacts with FtsX$_{ECL1}$ via its coiled coil domain, and this interaction causes activation of the peptidoglycan hydrolytic activity of PcsB. PcsB, along with other factors in the cell, allow for cell division to proceed normally. Functional FtsX, FtsE, and PcsB are all required for efficient cell division..