1   **Male and female bees show large differences in floral preference**
2

3   Michael Roswell mroswell.rutgers@gmail.com
4   Graduate program in ecology and evolution, Rutgers University
5   14 College Farm Road, New Brunswick, NJ 08904
6
7   Jonathan Dushoff
8   Department of biology, McMaster University
9   1280 Main St. West, Hamilton, Ontario ON L8S 4K1
10
11   Rachael Winfree
12   Department of ecology, evolution, and natural resources, Rutgers University
13   14 College Farm Road, New Brunswick, NJ 08904

Abstract

1. Intraspecific variation in foraging niche can drive food web dynamics and ecosystem processes. Field studies and theoretical analysis of plant-pollinator interaction networks typically focus on the partitioning of the floral community between pollinator species, with little attention paid to intraspecific variation among plants or foraging bees. In other systems, male and female animals exhibit different, cascading, impacts on interaction partners. Although the foraging ecology of male bees is little known, we expect foraging preferences to differ between male and female bees, which could strongly impact plant-pollinator interaction outcomes.

2. We designed an observational study to evaluate the strength and prevalence of sexually dimorphic foraging preferences in bees.

3. We observed bees visiting flowers in semi-natural meadows in New Jersey, USA. To detect differences in flower use against a shared background resource availability, we maximized the number of interactions observed within narrow spatio-temporal windows. To distinguish observed differences in bee use of flower species, which can reflect abundance patterns and sampling effects, from underlying differences in bee preferences, we analyzed our data with both a permutation-based null model and random effects models.

4. We found that the diets of male and female bees of the same species were often as dissimilar as the diets of bees of different species. Furthermore, we demonstrate differences in preference between male and female bees, and identify plant species that are particularly attractive to each sex. We show that intraspecific differences in preference can be robustly identified within interactions between hundreds of species, without precisely quantifying resource availability, and despite high phenological turnover.

5. Given the large differences in flower use and preference between male and female bees, ecological sex differences should be integrated into studies of bee demography, plant pollination, and coevolutionary relationships between flowers and insects.

Key Words
dimorphism, dissimilarity, Morisita-Horn, phenology, plant-pollinator interaction, pollination, pollinator habitat, preference

Introduction
Intraspecific variation in traits and behavior, including foraging niche, has important consequences for species interactions and conservation (Bolnick et al., 2011; Durell, 2000). Sexual dimorphism is a large source of individual niche variation, and an important factor in plant-animal interactions, such as seed dispersal (Zwolak, 2018). Sexual dimorphism underlies adaptation, speciation, and the way in which animals exploit their ecological niche (Butler, Sawyer, & Losos, 2007; Temeles, Miller, & Rifkin, 2010). Morphological, behavioral, and life-history dimorphisms can also drive the form and function of ecosystems, for example when predator sex ratio drives the community composition of lower

60 trophic levels, shown experimentally (Start & De Lisle, 2018) and in nature,
61 where responses extended to water chemistry as well (Fryxell, Arnett, Apgar,
62 Kinnison, & Palkovacs, 2015).
63
Though ecological dimorphisms were first studied in vertebrates (Selander,
1966), they are common across taxa, including insects (Shine, 1989).
Surprisingly, in bees (Hymenoptera, Apoidea) for which both foraging (P.
Willmer, 2011) and sexual dimorphism (Alcock et al., 1978) have been well
studied, sexually dimorphic foraging has rarely been documented. Intraspecific
variation in floral preference is well known in social (Heinrich, 1979) and to a
lesser extent, solitary bee species (Bruninga-Socolar, Crone, & Winfree, 2016;
Tur, Vigalondo, Trøjelsgaard, Olesen, & Traveset, 2014), yet most community-
level studies focus on species-level interaction networks, and furthermore, on the
foraging preferences of only female bees.
74
Male bees differ from their better-studied female counterparts in their life history
and ecology. Female bees construct, maintain, provision, and defend nests,
whereas male bees primarily seek mates (P. G. Willmer & Stone, 2004). Both
sexes drink floral nectar for their own caloric needs, but only females collect
pollen to provision young, and thus forage at greater rates. While the pollen from
each flower species (the term we use throughout for the flowers from a species
of plant) tends to be morphologically and nutritionally distinct, interspecific
variation in the chemical composition of nectar is comparatively subtle (Cane &
Sipes, 2006). Thus, we expect the foraging ecology, including floral preferences,
of male and female bees to differ as well.
85
Male bees prove to be important pollinators when studied, both in specialized oil-
or scent-collecting pollination systems (Eltz et al., 2007; Etl, Franschitz, Aguiar,
Schönenberger, & Dötterl, 2017; Janzen, 1971) and also when males are
foraging for nectar and pollen (Cane, 2002; Cane, Sampson, & Miller, 2011;
Ogilvie & Thomson, 2015). Male bees may also be particularly relevant for bee
conservation. Males may be limiting in declining populations, either because
genetic diversity is necessary for the development of female offspring as a result
of complementary sex determination, or because mate or sperm limitation results
from poor male condition (Elias, Dorn, & Mazzi, 2010; Straub et al., 2016). As the
dispersing sex in most bee species, males may be crucial for gene flow and
metapopulation persistence even when they are not locally limiting (López-Uribe,
Morreale, Santiago, & Danforth, 2015; Ulrich, Perrin, & Chapuisat, 2009).
98
Foraging niche is only partly described by resource use. Indeed, resource
preferences may be more important than use alone in many contexts, including
conservation. Preference—the use of a resource in excess of its relative
availability—is challenging to measure, because both resource use and
availability must be known. Floral resource availability for pollinators is
particularly hard to quantify outside an experimental context because the
appropriate scale and units of floral resource availability are unclear. The

composition, amount, and supply rate of pollen and nectar per flower, the number
of flowers per inflorescence, of inflorescences per individual, and the number and
distribution of individual plants over the square kilometers of a bee's foraging
range are all important components of availability (Hicks et al., 2016).
Furthermore, floral availability can change rapidly over time. However,
differences in flower use between bees foraging at the same place and time
indicate differences in preference, which may occur between species, or between
individuals of the same species.
114
In this study, we assess differences between floral preferences of male and
female bees in the wild. We collected bees foraging on flowers in meadows in
New Jersey, USA. In order to observe preference differences, we collected as
many individuals as possible during replicated, short (3-day) windows, during
which we assumed floral availability and bee abundance were constant at each
site. We compare the species composition of flowers visited by males and
females of the most common bee species across the entire study as a naïve
measure of differences in preference between the sexes. Then, using random
effects models, we assess when differential flower species *use* by male and
female bees likely arises from sex-specific floral *preference*, as opposed to
shifting overlap between foragers and floral resources (i.e. changes in *availability*
without differences in preference)*.* Specifically, we ask
127
128   1) How much do male and female bee diets overlap?
129   2) To what degree are particular flower species disproportionately visited by
130   bees of one sex?
131   3) To what extent are differences in floral use driven by preference, rather
132   than phenological differences between male and female bees?
133
Materials and Methods

Study design and data collection

Because absolute preference is nearly impossible to observe outside of an
experiment, we designed our study to reveal differences in preference between
groups of bees. In order to collect a large number of males and females from
many native bee species, we selected six meadows (sites) in New Jersey, USA
with a high abundance and diversity of flowers. These meadows were managed
for pollinator-attractive, summer-blooming forbs through seed addition, and a
combination of mowing, burning, and weed removal. Most flower species present
in the meadows are native to the eastern United States. We collected our data
during peak bloom and maximum day length (6 June to 20 August, 2016), and
during good weather (sunny enough for observers to see their own shadow, no
precipitation). We visited each site for three consecutive good weather days over
five evenly spaced sampling rounds in the 11-week period of our study. In all
analyses, we assume that bees and flowers detected at a site within one 3-day
sampling round co-occurred. In contrast, we assume that turnover of both plant

species in bloom and bee species activity can occur in the ~10 days between
sampling rounds.
154
During each 3-day sampling round, an observer walked parallel transects
through the meadow (which ranged in size from 0.8–2.2 ha; mean=1.4 ha),
observing every open flower within a moving 1-m semicircle, and net-collecting
any bee seen actively foraging, which we defined as contacting anthers or
collecting nectar from a flower (Fig. S1). We collected all bee species except
Apis mellifera L.*,* the domesticated western honey bee, because *Apis* males do
not forage. Observations began as soon as pollinator activity picked up in the
morning (7–9 am) and continued into the late afternoon or evening until pollinator
activity slowed substantially. Observers sampled nearly continuously, in 30-
minute timed collection bouts with short breaks in between. If inclement weather
precluded a minimum of six 30-minute sampling bouts in a day, we added an
additional day to the sampling round as soon as weather permitted.
167
Flower species were identified in the field by the data collector. Bee species were
identified using a dissecting microscope and published keys; Jason Gibbs
(University of Manitoba), Joel Gardner (University of Manitoba), and Sam Droege
(USGS) assisted with identification for bees in the genera *Andrena, Anthophora,*
Coelioxys, Halictus, Heriades, Hoplitis, Hylaeus, Lasioglossum, Megachile,
Melissodes, Nomada, Osmia, Pseudoanthidium, Ptilothrix, Sphecodes, Stelis,
and *Triepeolus,* and at least one of them confirmed voucher specimens for every
species. We determined every specimen to species except for the following four
complexes: Most bees in the genus *Nomada* with bidentate mandibles (*ruficornis*
group) were treated as one species. All specimens from the *Hylaeus* species
complex that includes *Hylaeus affinis*, *H. modestus*, and at least one additional
species, informally dubbed "species A*,*" were treated as a single species,
denoted *Hylaeus affinis-modestus*, because females cannot be reliably
distinguished. There is a cryptic species in the genus *Halictus* unlikely to occur in
our area, *Halictus poeyi*, which is not morphologically distinct from *H. ligatus*; we
treat all specimens in this complex as *Halictus ligatus*. We could not confidently
separate all specimens of the two closely related *Lasioglossum* species
Lasioglossum hitchensi and *L. weemsi*. Thus, we treat all specimens from either
species as one, denoted *Lasioglossum hitchensi-weemsi.* All bee specimens are
curated in the Winfree lab collection at Rutgers University, and the data used in
this paper are available from the Dryad Digital Repository
http://dx.doi.org/XXXXXXX (Roswell et al.)
190
191　**Analytical methods**
192
193　We performed all statistical analyses and simulations using R 3.5.1 (R Core
194　Team, 2018).
195
196　*1)　How much do male and female bee diets overlap?*
197

To compare the diets of male and female bees, we used the Morisita-Horn index
of resource overlap (Horn, 1966; Morisita, 1959). This dissimilarity index
compares the proportion of all female bees found on each flower species to the
proportion of all male bees found on each flower species. In other words, it
compares the contribution of each flower species to female diets (where this term
includes the food that females collect for themselves and also to feed to young)
to the contribution of the same flower species to male diets. The Morisita-Horn
index ranges from zero (completely similar) to one (maximally dissimilar), and
has several good properties for our purposes. First, it uses proportions, placing
visits from male and female bees on the same scale, even though most visits
come from females. Second, it is much more sensitive to large proportions than
to small ones, thereby down-weighting the contribution of flower species for
which we have little information. Third, the Morisita-Horn estimates are resilient
to undersampling and uneven sample size between groups (Barwell, Isaac, &
Kunin, 2015).
213
To determine whether the male-female differences we observed exceeded those
expected by chance, we compared the observed compositional dissimilarity
between flower visits from male and from female bees to dissimilarity measures
from a null model that randomly permuted the bee sex associated with each
flower-visit record. This permutation holds constant the total number of male and
of female visits, and the total number of visits to each flower species from both
sexes combined (Fig. S2). The range of dissimilarity values from this simulation
is the difference we would observe in our sample, if there were no true difference
in flower species use between males and females of the same bee species. We
evaluated the hypothesis that male and female diets overlap less than would be
expected by chance; thus we use a one-sided alpha of $p<0.05$. We iterated this
null simulation 9999 times, which was sufficient to stabilize p-values near our
chosen alpha (North, Curtis & Sham 2002). When the observed dissimilarity was
greater than 9500 of the 9999 simulated dissimilarities, we concluded that we
had detected a difference in the pattern of floral visitation between conspecific
male and female bees, given the observed diet breadth and abundance of each
sex.
231
To compare the diet overlap we observed between sexes to a meaningful
benchmark, interspecific diet overlap, we repeated the same null model analysis,
this time comparing females of the focal species to females of other species. We
performed one analysis for each bee species for which we collected at least 20
visitation records for each sex (19 species). This sample size threshold is
arbitrary, but null model variance shrinks with sample size, such that apparent
patterns for species with smaller sample sizes are rarely interpretable (Fig. S3).
Because we analyze 19 bee species, females of each species are compared to
18 others. We then compared the male-female difference (observed minus mean
null dissimilarity in flower communities visited) to the analogous species-species
difference (observed minus null dissimilarity).
243

244 For this analysis, which evaluates holistic differences between male and female
245 bees of the same species, we combined observations across the full season and
246 all sites. This allows us to observe foraging niche differences that are driven by
247 flower and/ or bee phenology, in addition to any sex-specific floral preference.
248
249 *2) To what degree are particular flower species disproportionately visited by bees*
250 *of one sex?*
251
This analysis uses our entire data set of 153 bee species to determine whether
particular flower species are disproportionately visited by male or female bees,
and whether the answer varies by bee species. We can infer a preference
difference between male and female bees for a flower species when predicted
odds of visitors to that flower species being male are especially high or low. To
do this, we use a random effects model in which bee sex is the response, and
flower species, bee species, site, and their interactions are random effects. We
statistically control for variation in the overall sex ratio across bee species
through a random intercept of bee species, and variation in sex ratios across
sites, through random intercepts for site, and the site-bee species and site-flower
species interactions. Because it is unlikely that, within bee species, sex ratios at
birth vary greatly across space, any variability attributed to site terms would likely
result from differential overlap of bee foraging activity and flower bloom across
space.
266
We call this model the "summed model" because we sum interactions observed
across the entire season (all five sampling rounds) at each site. In the summed
model, the relationship between phenological overlap and the odds of flower-
visiting bees being male would be incorporated into the species effects. This
perspective is helpful for considering flower species' contributions to the overall
diets of male versus female bees. We fit the model with the R package lme4
(Bates, Maechler, & Walker, 2016) with the following call:
274
275 **Summed model**
276 lme4::glmer(bee_sex ~ (1|site)+ (1|flower_species)+ (1|bee_species)+
277   (1|flower_species:bee_species)+ (1|site:bee_species)+
278   (1|site:flower_species), family="binomial", data=data)
279
We included bee species and site as random, rather than fixed, effects to directly
compare the variability in bee sex associated with each of these predictors to the
variability associated with flower species (preference). Comparing the overall
variability across these groups was more important to us than assessing
predictions on a per-site or per-bee-species basis. We fit flower species, the
primary covariate of interest, as a random effect to facilitate model fitting (fewer
degrees of freedom) as well as interpretation. In our summed model, we included
all two-way interactions, but omitted the three-way interaction, bee species by
flower species by site. Although the sort of context-dependent preference this
term could represent (e.g. males from bee species *1* prefer flower *A* at one site

(relative to females), but shun it at another) may exist in nature, it is unlikely we
would estimate it accurately in our model.
292
We confirmed model convergence by comparing several fitting methods using
the allFit function in lme4 (Bates et al., 2016), which all showed similar parameter
estimates (Table S1). We tested whether residuals from our model fit were
overdispersed using Bolker's function "overdisp" (Bolker, 2017), and visually
assessed our additivity assumptions with binned residual plots (Gelman & Hill,
2007) (Fig. S4).
299
300   **3)** *To what extent are differences in floral use driven by preference, rather than*
301   *phenological differences between male and female bees?*
302
Over the 11 weeks of our study, we observed turnover in bee species, in flower
bloom, and within-bee species changes in sex ratio. Therefore, phenological
overlap between male versus female bees and the bloom period of particular
flower species, rather than preference of those bees for those flowers, may
explain much of the variation in sex ratio we observed across visitors. In question
3, we are explicitly interested in distinguishing sex-specific diet *preferences* from
variable *use* resulting from seasonal resource availability and male vs. female
abundance. We do this in the "seasonal model" by incorporating sampling round
(our measure of phenology) as an additional random intercept effect, along with
random intercepts for the interactions between sampling round and the other
covariates. We chose to include sampling round as a random effect because this
enables direct comparison to all other terms in both models. We ignored the
three- and four-way interactions between bee species, flower species, and other
covariates. We fit this model with the following call in the R package lme4, with
new terms in bold:
318
Seasonal model
glmer(bee_sex ~ (1|site)+ (1|flower_species)+ (1|bee_species)+
(1|flower_species:bee_species)+ (1|site:bee_species)+
(1|site:flower_species)**+ (1|sampling_round)+ (1|site:sampling_round)+**
(1|flower_species:sampling_round)+
(1| bee_species:sampling_round)+
(1|site:bee_species:sampling_round)+
(1|site:flower_species:sampling_round), family="binomial", data=data)
327
Our index of preference for both the *summed model* and the *seasonal*
model is the change in odds that a bee is male when the flower species it visits is
given. To describe the importance of model terms, we calculated a bootstrapped
median odds ratio using code from Seth (Seth, 2017), which gives the expected
difference in odds that a flower-visiting bee is male between levels of a predictor
(Merlo et al., 2006). For example, a median odds ratio of five for the main effect
of sampling round would indicate that the odds of a flower-visiting bee being
male differ by about a factor of five between sampling rounds, while a median

odds ratio of one would indicate that the odds of a flower-visiting bee being male do not change across rounds. If the median odds ratio is large for flower species in both models, we could say that there are intrinsic (i.e. not simply phenological) properties of flower species identity that male or female bees prefer. If flower species is a strong predictor of bee sex in the summed model but not in the seasonal one, we would still conclude that flower species often contribute more strongly to the diet of one sex than the other, though these differences may not arise due to differing preferences. If the sampling round terms have large median odds ratios, then accounting for phenology is critical for identifying differences in preference in addition to differences in use.

Results

In total we collected 18,698 bee specimens belonging to 152 bee species (table S2) from a total of 109 flower species (table S3), which together comprised 1417 unique species-species interactions. Roughly 18% of specimens were male (n=3372). Thus, the overall ratio of male to female bees we collected was 0.22, although this ratio varied markedly between flower species (Fig. 1).

How much do male and female bee diets overlap?

We found that male and female bee diets overlap significantly less than would be expected at random (Fig. 2), and that the differences in diet composition between male and female bees of several species were of similar magnitude to the differences in diet between species of bee (Fig. 3).

To what degree are particular flower species disproportionately visited by bees of one sex?

The sex ratio of flower-visiting bees varied across species of flower (Fig. 2). After controlling for bee species identity (the strongest predictor of sex in our models, Fig. 4), and site, we still found that some flower species received a disproportionate number of male bee visitors (Figs 4–5). The median odds ratio for the main effect of flower species was 3.6 (bootstrapped CI 3.0–4.2) in our summed model, indicating that, typically, the visitor sex ratio differs between two flower species by more than a factor of 3. Furthermore, we observed sex-based differences in flower use specific to particular bee species: the median odds ratio for the flower species by bee species interaction in our summed model was nearly as large (median=3.1, bootstrapped CI 3.0–3.3) as the main effect of flower species. By contrast, sex ratios are not expected to differ between sites (median odds ratio for main effect of site=1).

To what extent are differences in floral use driven by preference, rather than phenological differences between male and female bees?

The flower species blooming in our system turned over throughout our 11-week

sampling period, with several highly visited species blooming for only one of the
three months during which we sampled. This turnover, along with potential sex-
specific bee flight seasons, means that differences in diet between male and
female bees could reflect seasonal availability and use, without also indicating
preference differences between the sexes. Indeed, phenology predicts bee sex
somewhat, with the odds of a flower-visiting bee being male expected to change
by a factor of 1.5 (bootstrapped CI 1.1–1.9) between sampling rounds (Fig. 4).
Phenological patterns of male vs. female flight seasons vary across bee species;
the median odds ratio for the bee species by sampling round interaction is 2.2
(bootstrapped CI 2.1–2.3) (Fig. 4). Even after accounting for these effects,
however, there remains a strong association between the species of flower a bee
visits and its sex (Figs 4–5). The relative effects of each flower species on the
sex of its visitors were changed very little by accounting for phenology; Pearson
and Spearman correlations between the random effect of flower species in the
seasonal model and the same random effect in the simpler summed model were
both 0.98. In addition to finding overall preference difference between male and
females, we found evidence for bee-species-specific difference in floral
preferences between the sexes (median odds ratios in both models for the bee
species by flower species interaction > 2.8).
401
402 **Discussion**
403
We found strong differences between the flower species preferences of male and
female bees. The difference in floral visits between male and female bees of the
same species was similar in magnitude to differences between females of
different species. The partitioning of the floral community among bee species is a
primary focus of pollination ecology and ecological network analysis (Bascompte
& Jordano, 2014), but male bees are typically disregarded or lumped together
with their female counterparts. Our study suggests this may represent an
important oversight. Further, our study provides strong confirmation of the few
studies that investigate the foraging behavior of male bees, which found that
males play a unique role in plant pollination (Cane, 2002; Cane et al., 2011;
Ogilvie & Thomson, 2015; Pascarella, 2010). Our result also implies that male
bees contribute substantially to the complexity of plant-pollinator networks in
nature, and that network analyses might benefit from separating males and
females into different nodes (Bolnick et al., 2011; Zwolak, 2018).
418
Phenology, a previously reported mechanism for distinct use of floral resources
by male and female bees (Ogilvie & Thomson, 2015; Robertson, 1925),
explained some variation in the sex ratio of flower-visiting bees, but was less
important than flower species identity over the period of our study. We expected
to find an effect of phenology because both the identity of the flower species
blooming within sites, and also the sex ratio of foragers within bee species, vary
across the season. Males emerge first in most solitary bees; for social species,
initial broods usually consist primarily of female workers, then males and
reproductive females emerge at the end of the colony cycle (P. G. Willmer &

Stone, 2004). Surprisingly, however, phenology only weakly predicted the sex of
flower-visiting bees. This is despite the fact that, as predicted by natural history,
the sampling round(s) in which males were relatively more prevalent depended
on bee species (the bee species by sampling round interaction was much bigger
than the sampling round main effect; Fig. 4). This indicates that our evidence for
floral preference differences between male and female bees was robust to
accounting for seasonal turnover in flower species bloom, bee species flight
seasons, and the sex ratios within bee species.
436
Whereas female bees collect both nectar and pollen, male bees forage primarily
for nectar to fuel flight. Thus, we predicted that male bees would avoid flowers
that produce no nectar. Indeed, in both our models, the predicted odds of a bee
visiting a nectar-less flower species being male were approximately half that of a
bee visiting a flower species that produces nectar (Fig. S5). A second biological
difference between male and female bees is that adult male bee activities orient
around mate seeking (Alcock et al., 1978). These behaviors, such as patrolling
routes (Barrows, 1976) or seeking flowers visited by conspecific females (Rossi,
Nonacs, & Pitts-Singer, 2010) could generate differences from females via
complementarity (males visiting flower species not visited by females), or
nestedness (one sex primarily visiting a subset of species visited by the other).
We found evidence for both (Fig. S6). Divergent floral preferences between
sexes may reflect nutritional needs or mating behavior, or simply biases resulting
from previous flower encounters, or visual or olfactory sensitivities that differ
between the sexes (Robert, Frasnelli, Collett, & de Ibarra, 2016; Somanathan,
Borges, Warrant, & Kelber, 2017; Streinzer, Kelber, Pfabigan, Kleineidam, &
Spaethe, 2013).
454
While natural and semi-natural habitats are critical landscape elements for many
bee species (Senapathi, Goddard, Kunin, & Baldock, 2016), what constitute
suitable and/ or limiting resources within these habitats remains less clear (De
Palma et al., 2015). Flowers, which provide food for adult and larval bees, are
likely among them (Roulston & Goodell, 2011). Whether floral diversity *per se*
tends to benefit individual pollinator taxa remains unclear (Spiesman, Bennett,
Isaacs, & Gratton, 2017; Sutter, Jeanneret, Bartual, Bocci, & Albrecht, 2017).
However, complementary flower species use between the sexes implies a
mechanism by which a bee species could benefit from a diversity of flower
choices. In addition, it is currently unknown how the distinct foraging niches of
male bees mediate either the robustness of pollinator communities to species
loss and environmental perturbations (Brosi & Briggs, 2013; Ramos-Jiliberto,
Valdovinos, Moisset de Espanés, & Flores, 2012; Tur et al., 2014), or the
effectiveness of different habitat ameliorations (Rundlöf, Persson, Smith, &
Bommarco, 2014; Rusterholtz & Erhardt, 2000; Williams & Lonsdorf, 2018). This
study suggests that both questions warrant further investigation.
471
Patterns in bee-flower interaction data can arise from the sampling process itself
(Blüthgen, 2010; Fründ, McCann, & Williams, 2016). Our analyses control for

these patterns. To evaluate diet overlap, we used a dissimilarity index that
downweights rare diet items, and implemented a null model that accounts for
differences that could arise from sampling and abundance effects. To evaluate
preference, we used random effects models that incorporated all (nearly 19,000)
observations, and shrank extreme values for rarely observed species-species
interactions towards the global mean for each effect. Thus, our estimates for sex-
specific preferences should be robust to the inevitable under-sampling of rarer
taxa. Establishing differences in preference between categories of bees such as
males and females, even when resource availability is seasonal and difficult to
quantify, is possible using methods such as these, though absolute preference
remains elusive.
485
Pollination ecology and pollinator conservation still face the question of how
important sexually dimorphic foraging is. Does it enhance or reduce the stability
of bee populations? Should pollinator restorations explicitly include "male bee"
flowers and "female bee" flowers? Are floral traits under selection to favor female
versus male visitors? While our study does not answer these questions, by
showing that the diets and preferences of male bees commonly differ from those
of their female conspecifics, we suggest they are worthy of future study.
493
494 **Author contributions**
495
496 MR and RW conceived the ideas and designed field methodology; MR, JD, and
497 RW designed statistical methodology. MR collected the data; MR, RW, and JD
498 analyzed the data; MR and RW led manuscript drafting. All authors contributed
499 critically to the drafts and gave final approval for publication.
500

501 **Acknowledgements**
502
We thank Daniel Cariveau and Neal Williams for study design suggestions, and
Jason Gibbs, Tina Harrison, and Dylan Simpson for comments on an earlier
draft. A USDA Conservation Innovation Grant to the Xerces Society for
Invertebrate Biology (lead PI) and RW (senior investigator) funded data
collection. MR was supported by an NSF Graduate Research Fellowship.
Cameron Kanterman and Riva Letchinger helped collect field data; Kurtis
Himmler, Tiffany Bennet, and professional taxonomists Jason Gibbs, Joel
Gardiner, and Sam Droege helped with bee species determinations. Mercer
County, the Institute for Advanced Study, Somerset County, and the Raritan
Headwaters Association provided site permission.
513

References

Alcock, J., Barrows, E. M., Gordh, G., Hubbard, L. J., Kirkendall, L., Pyle, D. W., … Zalom, F. G. (1978). The ecology and evolution of male reproductive behaviour in the bees and wasps. *Zoological Journal of the Linnean Society*, *64*(4), 293–326. doi:10.1111/j.1096-3642.1978.tb01075.x

Barrows, E. M. (1976). Mating behavior in halictine bees (Hymenoptera: Halictidae): II. microterritorial and patrolling behavior in males of Lasioglossum rohweri. *Zeitschrift Fur Tierpsychologie-Journal of Comparative Ethology*, *40*(4), 377–389.

Barwell, L. J., Isaac, N. J. B., & Kunin, W. E. (2015). Measuring beta-diversity with species abundance data. *Journal of Animal Ecology*, 1112–1122. doi:10.1111/1365-2656.12362

Bascompte, J., & Jordano, P. (2014). Mutualistic networks in time and space. In J. Bascompte & P. Jordano (Eds.), *Mutualistic networks* (pp. 87–106). Princeton University Press. Retrieved from http://www.jstor.org/stable/j.ctt5hhnpq.8

Bates, D., Maechler, M., & Walker, S. (2016). Package "lme4." *CRAN Repository*. doi:10.18637/jss.v067.i01

Blüthgen, N. (2010). Why network analysis is often disconnected from community ecology: A critique and an ecologist's guide. *Basic and Applied Ecology*, *11*(3), 185–195. doi:10.1016/j.baae.2010.01.001

Bolker, B. M. (2017). GLMM FAQ. Retrieved from http://bbolker.github.io/mixedmodels-misc/glmmFAQ.html

Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., … Vasseur, D. A. (2011). Why intraspecific trait variation matters in community ecology. *Trends in Ecology and Evolution*, *26*(4), 183–192. doi:10.1016/j.tree.2011.01.009

Brosi, B. J., & Briggs, H. M. (2013). Single pollinator species losses reduce floral fidelity and plant reproductive function. *Proceedings of the National Academy of Sciences*, *110*(32), 13044–8. doi:10.1073/pnas.1307438110

Bruninga-Socolar, B., Crone, E. E., & Winfree, R. (2016). The role of floral density in determining bee foraging behavior: a natural experiment. *Natural Areas Journal*, *36*(4), 392–399. doi:10.3375/043.036.0406

Butler, M. A., Sawyer, S. A., & Losos, J. B. (2007). Sexual dimorphism and adaptive radiation in Anolis lizards. *Nature*, *447*(7141), 202–5. doi:10.1038/nature05774

Cane, J. H. (2002). Pollinating bees (Hymenoptera: Apiformes) of U.S. alfalfa compared for rates of pod and seed set. *Journal of Economic Entomology*, *95*(1), 22–27. doi:10.1603/0022-0493-95.1.22

Cane, J. H., Sampson, B. J., & Miller, S. A. (2011). Pollination value of male bees: The specialist bee Peponapis pruinosa (Apidae) at Summer Squash (Cucurbita pepo). *Environmental Entomology*, *40*(3), 614–620. doi:10.1603/EN10084

Cane, J. H., & Sipes, S. S. (2006). Characterizing floral specialization by bees: Analytical methods and a revised lexicon for oligolecty. In *Plant–pollinator*

560   *interactions: From specialization to generalization*. doi:10.1007/BF00937854

561 De Palma, A., Kuhlmann, M., Roberts, S. P. M., Potts, S. G., Börger, L., Hudson,

562   L. N., … Purvis, A. (2015). Ecological traits affect the sensitivity of bees to

563   land-use pressures in European agricultural landscapes. *Journal of Applied*

564   *Ecology*, *52*(6), 1567–1577. doi:10.1111/1365-2664.12524

565 Durell, S. E. A. L. V. dit. (2000). Individual feeding specialization in shorebirds:

566   population consequences and conservation implications. *Biological Reviews*

567   *of the Cambridge Philosophical Society*, *75*(2000), 503–518.

568 Elias, J., Dorn, S., & Mazzi, D. (2010). No evidence for increased extinction

569   proneness with decreasing effective population size in a parasitoid with

570   complementary sex determination and fertile diploid males. *BMC*

571   *Evolutionary Biology*, *10*(1), 366. doi:10.1186/1471-2148-10-366

572 Eltz, T., Zimmermann, Y., Haftmann, J., Twele, R., Francke, W., Quezada-Euan,

573   J. J. G., & Lunau, K. (2007). Enfleurage, lipid recycling and the origin of

574   perfume collection in orchid bees. *Proceedings. Biological Sciences / The*

575   *Royal Society*, *274*(1627), 2843–8. doi:10.1098/rspb.2007.0727

576 Etl, F., Franschitz, A., Aguiar, A. J. C., Schönenberger, J., & Dötterl, S. (2017). A

577   perfume-collecting male oil bee? Evidences of a novel pollination system

578   involving Anthurium acutifolium (Araceae) and Paratetrapedia chocoensis

579   (Apidae, Tapinotaspidini). *Flora: Morphology, Distribution, Functional*

580   *Ecology of Plants*, *232*, 7–15. doi:10.1016/j.flora.2017.02.020

581 Fründ, J., McCann, K. S., & Williams, N. M. (2016). Sampling bias is a challenge

582   for quantifying specialization and network structure: Lessons from a

583   quantitative niche model. *Oikos*, *125*(4), 502–513. doi:10.1111/oik.02256

584 Fryxell, D. C., Arnett, H. A., Apgar, T. M., Kinnison, M. T., & Palkovacs, E. P.

585   (2015). Sex ratio variation shapes the ecological effects of a globally

586   introduced freshwater fish. *Proceedings of the Royal Society B*, *282*(1817),

587   20151970. doi:10.1098/rspb.2015.1970

588 Gelman, A., & Hill, J. (2007). *Data analysis using regression and multilevel/*

589   *hierarchical models*. New York: Cambridge University Press.

590 Heinrich, B. (1979). "Majoring" and "Minoring" by foraging bumblebees, Bombus

591   vagans: an experimental analysis. *Ecology*, *60*(2), 245–255.

592 Hicks, D. M., Ouvrard, P., Baldock, K. C. R., Baude, M., Goddard, M. A., Kunin,

593   W. E., … Stone, G. N. (2016). Food for pollinators: Quantifying the nectar

594   and pollen resources of urban flower meadows. *PLoS ONE*, *11*(6).

595   doi:10.1371/journal.pone.0158117

596 Horn, H. S. (1966). Measurement of "Overlap" in comparative ecological studies.

597   *The American Naturalist*, *100*(914), 419–424. Retrieved from

598   http://www.jstor.org/stable/2459242

599 Janzen, D. H. (1971). Euglossine bees as long-distance pollinators of tropical

600   plants. *Science*, *171*(3967), 203–205. doi:10.1126/science.171.3967.203

601 López-Uribe, M. M., Morreale, S. J., Santiago, C. K., & Danforth, B. N. (2015).

602   Nest suitability, fine-scale population structure and male-mediated dispersal

603   of a solitary ground nesting bee in an urban landscape. *PLoS ONE, 10*(5),

604   1–20. doi:10.1371/journal.pone.0125719

605 Merlo, J., Chaix, B., Ohlsson, H., Beckman, A., Johnell, K., Hjerpe, P., … Larsen,

K. (2006). A brief conceptual tutorial of multilevel analysis in social epidemiology: Using measures of clustering in multilevel logistic regression to investigate contextual phenomena. *Journal of Epidemiology and Community Health*, *60*(4), 290–297. doi:10.1136/jech.2004.029454

Morisita, M. (1959). Measuring of interspecific association and similarity between communities. *Memoirs of the Faculty of Science, Kyushu University, Series E (Biology)*. Retrieved from http://reference.morisita.gr.jp/paper_pdf/56.pdf

Ogilvie, J. E., & Thomson, J. D. (2015). Male bumble bees are important pollinators of a late-blooming plant. *Arthropod-Plant Interactions*, *9*(2), 205–213. doi:10.1007/s11829-015-9368-x

Pascarella, J. B. (2010). Pollination biology of *Gelsemium sempervirens* L. (Ait.) (Gelsemiaceae): do male and female *Habropoda laboriosa* F. (Hymenoptera, Apidae) differ in pollination efficiency? *Journal of Apicultural Research*, *49*(2), 170–176. doi:10.3896/IBRA.1.49.2.05

R Core Team. (2018). R: A Language and Environment for Statistical Computing. Vienna, Austria. Retrieved from https://www.r-project.org/

Ramos-Jiliberto, R., Valdovinos, F. S., Moisset de Espanés, P., & Flores, J. D. (2012). Topological plasticity increases robustness of mutualistic networks. *Journal of Animal Ecology*, *81*(4), 896–904. doi:10.1111/j.1365-2656.2012.01960.x

Robert, T., Frasnelli, E., Collett, T. S., & de Ibarra, N. H. (2016). Male bumblebees perform learning flights on leaving a flower but not when leaving their nest. *The Journal of Experimental Biology*, *220*, 930–937. doi:10.1242/jeb.151126

Robertson, C. (1925). Heterotropic Bees. *Ecology*, *6*(4), 412–436. Retrieved from http://www.jstor.org/stable/1941781

Rossi, B. H., Nonacs, P., & Pitts-Singer, T. L. (2010). Sexual harassment by males reduces female fecundity in the alfalfa leafcutting bee, Megachile rotundata. *Animal Behaviour*, *79*(1), 165–171. doi:10.1016/j.anbehav.2009.10.023

Roulston, T. H., & Goodell, K. (2011). The role of resources and risks in regulating wild bee populations. *Annual Review of Entomology*, *56*, 293–312. doi:10.1146/annurev-ento-120709-144802

Rundlöf, M., Persson, A. S., Smith, H. G., & Bommarco, R. (2014). Late-season mass-flowering red clover increases bumble bee queen and male densities. *Biological Conservation*, *172*, 138–145. doi:10.1016/j.biocon.2014.02.027

Rusterholtz, H.-P., & Erhardt, A. (2000). Can nectar properties explain sex-specific flower preferences in the Adonis Blue butterfly Lysandra bellargus? *Ecological Entomology*, *25*, 81–90.

Selander, R. K. (1966). Sexual Dimorphism and Differential Niche Utilization in Birds. *The Condor*, *68*(2), 113–151. doi:10.2307/1365712

Senapathi, D., Goddard, M. A., Kunin, W. E., & Baldock, K. C. R. (2016). Landscape impacts on pollinator communities in temperate systems: evidence and knowledge gaps. *Functional Ecology*. doi:10.1111/1365-2435.12809

Seth, M. (2017). How to implement credible 95% interval for median odds ratio

using JAGS? Retrieved May 9, 2018, from
https://stats.stackexchange.com/q/318912
Shine, R. (1989). Ecological Causes for the Evolution of Sexual Dimorphism: A
Review of the Evidence. *The Quarterly Review of Biology*, *64*(4), 419–461.
Somanathan, H., Borges, R. M., Warrant, E. J., & Kelber, A. (2017). Visual
adaptations for mate detection in the male carpenter bee Xylocopa
tenuiscapa. *PLoS ONE*, *12*(1). doi:10.1371/journal.pone.0168452
Spiesman, B. J., Bennett, A., Isaacs, R., & Gratton, C. (2017). Bumble bee
colony growth and reproduction depend on local flower dominance and
natural habitat area in the surrounding landscape. *Biological Conservation*,
206, 217–223. doi:10.1016/j.biocon.2016.12.008
Start, D., & De Lisle, S. (2018). Sexual dimorphism in a top predator
(Notophthalmus viridescens) drives aquatic prey community assembly.
Proceedings of the Royal Society B, *285*. doi:10.1098/rspb.2018.1717
Straub, L., Villamar-Bouza, L., Bruckner, S., Chantawannakul, P., Gauthier, L.,
Khongphinitbunjong, K., … Williams, G. R. (2016). Neonicotinoid
insecticides can serve as inadvertent insect contraceptives. *Proceedings of*
the Royal Society - B, *283*(20160506). doi:10.1098/rspb.2016.0506
Streinzer, M., Kelber, C., Pfabigan, S., Kleineidam, C. J., & Spaethe, J. (2013).
Sexual dimorphism in the olfactory system of a solitary and a eusocial bee
species. *Journal of Comparative Neurology*, *521*(12), 2742–2755.
doi:10.1002/cne.23312
Sutter, L., Jeanneret, P., Bartual, A. M., Bocci, G., & Albrecht, M. (2017).
Enhancing plant diversity in agricultural landscapes promotes both rare bees
and dominant crop-pollinating bees through complementary increase in key
floral resources. *Journal of Applied Ecology*, *54*(6), 1856–1864.
doi:10.1111/1365-2664.12907
Temeles, E. J., Miller, J. S., & Rifkin, J. L. (2010). Evolution of sexual dimorphism
in bill size and shape of hermit hummingbirds (Phaethornithinae): a role for
ecological causation. *Philosophical Transactions of the Royal Society B*,
365(1543), 1053–1063. doi:10.1098/rstb.2009.0284
Tur, C., Vigalondo, B., Trøjelsgaard, K., Olesen, J. M., & Traveset, A. (2014).
Downscaling pollen-transport networks to the level of individuals. *Journal of*
Animal Ecology, *83*(1), 306–317. doi:10.1111/1365-2656.12130
Ulrich, Y., Perrin, N., & Chapuisat, M. (2009). Flexible social organization and
high incidence of drifting in the sweat bee, Halictus scabiosae. *Molecular*
Ecology, *18*(8), 1791–1800. doi:10.1111/j.1365-294X.2009.04154.x
Williams, N. M., & Lonsdorf, E. V. (2018). Selecting cost-effective plant mixes to
support pollinators. *Biological Conservation*, *217*, 195–202.
doi:10.1016/j.biocon.2017.10.032
Willmer, P. (2011). *Pollination and floral ecology*. Princeton: Princeton University
Press.
Willmer, P. G., & Stone, G. N. (2004). Behavioral, ecological, and physiological
determinants of the activity patterns of bees. *Advances in the Study of*
Behavior, *34*, 347–466.
Zwolak, R. (2018). How intraspecific variation in seed-dispersing animals matters

for plants. *Biological Reviews, 93*, 897–913. doi:10.1111/brv.12377
699

Figure 1. The sex ratio (M:F) of flower-visiting bees varies across flower species.
Each red point represents a flower species, the first three letters of the Latin
genus and species names for the flower species label each point. The x-axis is
the number of bees collected from that species, the y-axis is the ratio of male to
female bees collected from the flower. Flower species that received >19 visits are
plotted (n=54). Blue lines are smooth fits to the 97.5th and 2.5th percentiles of the
binomial distribution given by the observed ratio of males to females in our
overall dataset (M/F=0.22; i.e. M/(M+F)=0.18). This distribution represents our
expectation for random variation in sex ratio across flower species, if the sex
ratio of flower-visiting bees is independent of flower species identity (male and
female bees exhibit the same floral preferences), and remains nearly constant
across time and space.
713

Figure 2: Flower visit patterns of male and female bees of the same species differed significantly. Red points are observed Morisita-Horn dissimilarities between flower communities visited by all male and all female bees of a particular species across all sites and sampling rounds. Black points are the mean dissimilarity (gray bars, 95% CI) from a permutation-based null model that randomly shuffles the sex associated with each visit record, maintaining the total number of males, females, and overall combined visits to each floral species.

Figure 3. The diets of male and female bees of the same species can be as dissimilar as the diets of females of two different bee species. Dissimilarities in this figure are the observed statistic minus, for each pairwise comparison, the mean dissimilarity in the null model. Each panel focuses on a bee species (panel name) and shows: above the label "sex", observed diet dissimilarity between male and female bees of the focal species, minus the average null dissimilarity resulting from randomly permuting the sex identity of each visit record; above the label "sp", observed diet dissimilarity between female bees of the focal species and each other bee species, minus the average null dissimilarity resulting from randomly permuting the species identity of each visit record.

Figure 4. Flower species, along with bee species, predicts the sex of visiting bees, which suggests floral preferences differ between male and female bees. Flower species is an important predictor of bee sex even after accounting for phenology (seasonal model). For each term ("bee"= bee species, "flower"=flower species, "round"=sampling round) in each model, the median odds ratio (+/- 95% bootstrapped credible interval) indicates the expected difference in odds that a flower-visiting bee is male between two levels. For example, a median odds ratio of 3.7 for the flower species term means the odds of a visitor being male are expected to differ by a factor of 3.7 between two randomly selected species of flower.

Figure 5. Male bee preferences for and against flower species vary across
flower species. Each point is the conditional mode of the random effects
prediction (the random-effects analog to an estimate), for a flower species that
received at least 20 visits, on the logit scale. Zero represents the odds of a visitor
being male on a random flower, and -2 or 2 indicates a ~7 fold decrease or
increase in those odds, given flower species identity. Error bars are the square
root of the conditional variances on the conditional mode * 1.96, and can be
interpreted as the expected range in which the random effect for a particular
flower truly lies, analogous to 1.96 times the standard error of the mean for a
fixed effect.
757

758
Figure S1: Sampling scheme. (a) The six study sites in central New Jersey,
USA. (b) Schematic sampling diagram (not to scale). One observer walked
parallel 2m transects covering the entire sampling area. Each 30-minute
sampling bout resumed where the previous one left off; observers typically
covered the entire meadow once over a 3-day sampling round. (c) The
southwestern-most site in peak bloom.

Figure S2. Schematic cartoon of our simulation for the dissimilarity values
associated with our null hypothesis that diets of male and female bees do not
differ. (a) Each collection record for each bee species associates the sex of an
individual bee to the flower species from which it was collected. (b) To compute
the dissimilarity between males and females, we compare all visits to each flower
species from males (purple vector) to all visits to each flower species from
females (green vector). (c) The Morisita-Horn index summarizes the differences
between the two vectors as a value between 0 (identical) and 1 (maximally
dissimilar). (d) For our null model, we shuffle the sex column from our
observation table. (e) This produces two null vectors. The row and column sums
for the matrices in (b) and (c) are identical, but the elements can differ. (f) For our
null model, we compute the dissimilarity between the null vectors. We repeated
steps d-f 9999 times to generate confidence intervals for the null hypothesis that
the sex of a visiting bee is unrelated to the flower species it is collected from.
When comparing the flower species visited by different species of bee, we
conducted an analysis identical except that rather than comparing two sexes of
the same species, we compared two species of the same sex (i.e. exchanging
"sex" and "species" throughout figure S1).

Figure S3. Effect size for diet dissimilarity is independent of sample size, while
standardized effect is strongly driven by the number of individuals of the sex with
the fewest records. a) Observed Morisita-Horn dissimilarity in flower communities
visited by male and female bees of a single species, minus average null
dissimilarity vs. the number of records for the less frequently observed sex. b)
Observed minus null dissimilarity in composition of flowers visited by male and
female bees of a single species, scaled by the variation in the null model, versus
the number of records for the less frequently observed sex.

summed model

seasonal model

795
Figure S4. Binned residual plots for each model show minor violation of the
additivity assumption. Residuals and predicted values on the probability scale.

Figure S5. Seasonal model predictions are consistent with the hypothesis that
male bees avoid flower species that do not produce nectar, relative to females.
Each point is the random effect prediction (change in odds that a bee visiting that
flower is male) for a flower species.
803

807 **Figure S6**. Dissimilarity in flower communities visited by male and female bees
808 arise due to complementarity in addition to nestedness patterns. For each bee
809 species, the proportion of male (blue) and female (red) visits to each flower
810 species that received >10% of at least one sex's visits are pictured. Due to
811 omitted flower species, bars may sum to <1.
812

  Table S1. Model convergence confirmed based on similar parameter estimates across fitting routines. For each model,
  the estimate for each term is given for each of 6 fitting algorithms in the R package lme4. Subsequent analyses used
  parameter estimates in yellow, in both cases tied for the highest estimated likelihood with other very similar fits.

term	model	bobyqa	Nelder_Mead	nlminbw	optimx.L-BFGS-B	nloptwrap.NLOPT_LN_NELDERMEAD	nloptwrap.NLOPT_LN_BOBYQA
intercept	summed	-2.43	-2.43	-2.43	-2.43	-2.43	-2.43
bee species	summed	2.04	2.04	2.04	2.04	2.04	2.04
flower species	summed	1.40	1.40	1.40	1.40	1.40	1.40
site	summed	0.00	0.00	0.00	0.00	0.00	0.00
bee species:flower species	summed	1.21	1.21	1.21	1.21	1.21	1.21
site:bee species	summed	0.62	0.62	0.62	0.62	0.62	0.62
site:flower species	summed	0.61	0.61	0.61	0.61	0.61	0.61
intercept	seasonal	-2.38	-2.45	-2.38	-2.38	-2.45	-2.45
bee species	seasonal	2.09	2.14	2.09	2.09	2.13	2.13
flower species	seasonal	1.25	1.27	1.25	1.25	1.27	1.27
site	seasonal	0.00	0.00	0.00	0.00	0.00	0.00
bee species:flower species	seasonal	1.09	1.10	1.09	1.09	1.10	1.10
site:bee species	seasonal	0.46	0.47	0.46	0.46	0.47	0.47
site:flower species	seasonal	0.35	0.35	0.35	0.35	0.35	0.35
sampling round	seasonal	0.38	0.36	0.38	0.38	0.36	0.36
sampling round:bee species	seasonal	0.83	0.84	0.83	0.83	0.84	0.84
sampling round:flower species	seasonal	0.00	0.00	0.00	0.00	0.00	0.00
sampling round:site	seasonal	0.29	0.29	0.29	0.29	0.29	0.29
sampling round:site:bee species	seasonal	0.60	0.60	0.60	0.60	0.60	0.60
sampling round:site:flower species	seasonal	0.28	0.29	0.28	0.28	0.28	0.28

817
818
819
820 **Table S2.** Bee species with number of female and male specimens collected.
821 *This table will be removed from the final submission when data are deposited on*
822 *Dryad.*

family	genus	species	females	males
Andrenidae	*Andrena*	*brevipalpis*	1	0
Andrenidae	*Andrena*	*carlini*	3	0
Andrenidae	*Andrena*	*commoda*	3	0
Andrenidae	*Andrena*	*cressonii*	16	0
Andrenidae	*Andrena*	*fragilis*	2	0
Andrenidae	*Andrena*	*hippotes*	4	0
Andrenidae	*Andrena*	*imitatrix*	6	0
Andrenidae	*Andrena*	*krigiana*	14	0
Andrenidae	*Andrena*	*nasonii*	13	0
Andrenidae	*Andrena*	*nuda*	2	0
Andrenidae	*Andrena*	*pruni*	6	0
Andrenidae	*Andrena*	*robertsonii*	8	0
Andrenidae	*Andrena*	*rudbeckiae*	8	11
Andrenidae	*Andrena*	*rugosa*	1	0
Andrenidae	*Andrena*	*spiraeana*	1	0
Andrenidae	*Andrena*	*vicina*	6	0
Andrenidae	*Andrena*	*wilkella*	277	59
Andrenidae	*Andrena*	*wilmattae*	2	0
Andrenidae	*Calliopsis*	*andreniformis*	4	1
Apidae	*Anthophora*	*abrupta*	4	0
Apidae	*Anthophora*	*terminalis*	3	2
Apidae	*Bombus*	*auricomus*	1	0
Apidae	*Bombus*	*bimaculatus*	577	175
Apidae	*Bombus*	*citrinus*	0	5
Apidae	*Bombus*	*fervidus*	18	0
Apidae	*Bombus*	*griseocollis*	681	815
Apidae	*Bombus*	*impatiens*	2358	105
Apidae	*Bombus*	*perplexus*	22	36
Apidae	*Bombus*	*vagans*	14	2
Apidae	*Ceratina*	*calcarata*	1417	133
Apidae	*Ceratina*	*dupla*	151	19
Apidae	*Ceratina*	*mikmaqi*	130	5
Apidae	*Ceratina*	*strenua*	285	13

Apidae	*Melissodes*	*agilis*	0	7
Apidae	*Melissodes*	*bimaculatus*	9	1
Apidae	*Melissodes*	*denticulatus*	7	73
Apidae	*Melissodes*	*desponsus*	1	7
Apidae	*Melissodes*	*subillatus*	31	6
Apidae	*Melissodes*	*trinodis*	1	7
Apidae	*Nomada*	*articulata*	4	0
Apidae	*Nomada*	*bidentate_gr*	7	0
Apidae	*Nomada*	*erigeronis*	1	0
Apidae	*Nomada*	*lehighensis*	1	0
Apidae	*Nomada*	*maculata*	2	0
Apidae	*Nomada*	*pygmaea*	15	0
Apidae	*Ptilothrix*	*bombiformis*	0	1
Apidae	*Triepeolus*	*cressonii*	0	1
Apidae	*Triepeolus*	*eliseae*	1	0
Apidae	*Triepeolus*	*remigatus*	1	0
Apidae	*Xylocopa*	*virginica*	137	13
Colletidae	*Hylaeus*	*affinis_modestus*	1376	363
Colletidae	*Hylaeus*	*fedorica*	1	0
Colletidae	*Hylaeus*	*leptocephalus*	1	3
Colletidae	*Hylaeus*	*mesillae*	575	173
Halictidae	*Agapostemon*	*sericeus*	5	5
Halictidae	*Agapostemon*	*virescens*	203	76
Halictidae	*Augochlora*	*pura*	1036	377
Halictidae	*Augochlorella*	*aurata*	397	39
Halictidae	*Augochlorella*	*persimilis*	434	116
Halictidae	*Augochloropsis*	*metallica*	121	40
Halictidae	*Dufourea*	*novaeangliae*	0	1
Halictidae	*Halictus*	*confusus*	174	35
Halictidae	*Halictus*	*ligatus*	2432	160
Halictidae	*Halictus*	*parallelus*	6	18
Halictidae	*Halictus*	*rubicundus*	31	19
Halictidae	*Lasioglossum*	*abanci*	6	0
Halictidae	*Lasioglossum*	*admirandum*	15	0
Halictidae	*Lasioglossum*	*anomalum*	17	0
Halictidae	*Lasioglossum*	*atwoodi*	7	1
Halictidae	*Lasioglossum*	*birkmanni*	1	0
Halictidae	*Lasioglossum*	*bruneri*	6	4
Halictidae	*Lasioglossum*	*callidum*	54	0

Halictidae	Lasioglossum	cattellae	14	4
Halictidae	Lasioglossum	coeruleum	2	0
Halictidae	Lasioglossum	coreopsis	1	0
Halictidae	Lasioglossum	coriaceum	14	0
Halictidae	Lasioglossum	cressonii	16	5
Halictidae	Lasioglossum	ellisiae	0	3
Halictidae	Lasioglossum	ephialtum	1	0
Halictidae	Lasioglossum	foxii	2	2
Halictidae	Lasioglossum	fuscipenne	9	0
Halictidae	Lasioglossum	gotham	74	2
Halictidae	Lasioglossum	hitchensi_weemsi	152	27
Halictidae	Lasioglossum	illinoense	70	7
Halictidae	Lasioglossum	imitatum	462	15
Halictidae	Lasioglossum	leucocomum	2	0
Halictidae	Lasioglossum	leucozonium	2	0
Halictidae	Lasioglossum	nigroviride	2	0
Halictidae	Lasioglossum	oblongum	4	2
Halictidae	Lasioglossum	obscurum	7	1
Halictidae	Lasioglossum	oceanicum	104	23
Halictidae	Lasioglossum	oenotherae	1	0
Halictidae	Lasioglossum	paradmirandum	50	0
Halictidae	Lasioglossum	pectorale	3	0
Halictidae	Lasioglossum	pilosum	2	0
Halictidae	Lasioglossum	platyparium	2	3
Halictidae	Lasioglossum	rozeni	15	11
Halictidae	Lasioglossum	smilacinae	4	0
Halictidae	Lasioglossum	subviridatum	5	1
Halictidae	Lasioglossum	tegulare	31	2
Halictidae	Lasioglossum	trigeminum	44	0
Halictidae	Lasioglossum	truncatum	2	0
Halictidae	Lasioglossum	versatum	681	93
Halictidae	Lasioglossum	viridatum	11	2
Halictidae	Lasioglossum	zephyrum	12	1
Halictidae	Sphecodes	atlantis	0	1
Halictidae	Sphecodes	dichrous	3	5
Halictidae	Sphecodes	heraclei	10	5
Megachilidae	Anthidiellum	notatum	4	1
Megachilidae	Anthidium	manicatum	7	8
Megachilidae	Anthidium	oblongatum	18	19

Megachilidae	*Coelioxys*	*alternatus*	1	2
Megachilidae	Coelioxys	banksi	1	0
Megachilidae	Coelioxys	germanus	0	1
Megachilidae	Coelioxys	hunteri	0	1
Megachilidae	Coelioxys	modestus	0	1
Megachilidae	Coelioxys	obtusiventris	1	0
Megachilidae	Coelioxys	octodentatus	1	1
Megachilidae	Coelioxys	porterae	0	1
Megachilidae	Coelioxys	sayi	2	6
Megachilidae	Heriades	carinatus	31	2
Megachilidae	Heriades	leavitti	1	6
Megachilidae	Heriades	variolosus	10	0
Megachilidae	Hoplitis	pilosifrons	46	1
Megachilidae	Hoplitis	producta	8	0
Megachilidae	Hoplitis	spoliata	2	1
Megachilidae	Lithurgus	chrysurus	0	6
Megachilidae	Megachile	brevis	25	3
Megachilidae	Megachile	campanulae	6	18
Megachilidae	Megachile	exilis	11	29
Megachilidae	Megachile	frugalis	26	6
Megachilidae	Megachile	gemula	4	2
Megachilidae	Megachile	georgica	1	0
Megachilidae	Megachile	inimica	4	0
Megachilidae	Megachile	integra	1	0
Megachilidae	Megachile	melanophaea	0	1
Megachilidae	Megachile	mendica	21	56
Megachilidae	Megachile	montivaga	15	9
Megachilidae	Megachile	petulans	0	2
Megachilidae	Megachile	pugnata	2	3
Megachilidae	Megachile	rotundata	11	8
Megachilidae	Megachile	sculpturalis	17	32
Megachilidae	Megachile	xylocopoides	2	1
Megachilidae	Osmia	albiventris	3	0
Megachilidae	Osmia	atriventris	9	0
Megachilidae	Osmia	bucephala	21	0
Megachilidae	Osmia	distincta	7	0
Megachilidae	Osmia	georgica	5	0
Megachilidae	Osmia	pumila	30	0
Megachilidae	Pseudoanthidium	nanum	0	1

Megachilidae	*Stelis*	*lateralis*	1	0
Megachilidae	*Stelis*	*louisae*	1	2

824 **Table S3.** Number of male and female visitors to each plant species, and bias
825 towards attracting male bee visitors. This bias is the random effect prediction
826 from the seasonal model, which indicates the change in log(odds) that a visiting
827 bee is male when the species of flower it visits is given; greater values indicate
828 male bias.

family	genus	species	female visits	male visits	random effect
Alliaceae	*Allium*	*vineale*	3	0	-0.224
Apiaceae	*Daucus*	*carota*	1778	350	0.526
Apiaceae	*Eryngium*	*yuccifolium*	2	0	-0.225
Apiaceae	*Sanicula*	*canadensis*	1	0	-0.049
Apocynaceae	*Apocynum*	*cannabinum*	283	92	0.754
Asclepidaceae	*Asclepias*	*incarnata*	7	0	-0.736
Asclepidaceae	*Asclepias*	*syriaca*	28	89	0.929
Asclepidaceae	*Asclepias*	*tuberosa*	114	26	0.423
Asteraceae	*Achillea*	*millefolium*	472	36	-0.004
Asteraceae	*Bidens*	*trichosperma*	1	0	-0.015
Asteraceae	*Carduus*	*nutans*	1	0	-0.366
Asteraceae	*Centuarea*	*stoebe*	321	50	0.265
Asteraceae	*Cichorium*	*intybus*	104	10	0.516
Asteraceae	*Cirsium*	*arvense*	351	96	0.858
Asteraceae	*Cirsium*	*discolor*	5	2	0.349
Asteraceae	*Cirsium*	*vulgare*	112	16	0.121
Asteraceae	*Conyza*	*canadensis*	32	10	1.097
Asteraceae	*Coreopsis*	*lanceolata*	21	0	-0.857
Asteraceae	*Coreopsis*	*tinctoria*	1	0	-0.069
Asteraceae	*Crepis*	*capillaris*	6	0	-0.236
Asteraceae	*Doellingeria*	*umbellata*	1	0	-0.052
Asteraceae	*Echinacea*	*purpurea*	86	45	0.309
Asteraceae	*Erechtites*	*hieraciifolius*	130	203	1.878
Asteraceae	*Erigeron*	*annuus*	26	3	0.266
Asteraceae	*Erigeron*	*strigosus*	712	119	1.037
Asteraceae	*Euthamia*	*graminifolia*	50	18	1.467
Asteraceae	*Eutrochium*	*maculatum*	461	166	1.368
Asteraceae	*Gaillardia*	*aristata*	4	0	-0.274
Asteraceae	*Helianthus*	*angustifolius*	11	0	-0.411

Asteraceae	*Helianthus*	*strumosus*	1	1	0.517
Asteraceae	*Heliopsis*	helianthoides	186	49	0.763
Asteraceae	*Hieracium*	pilosella	2	0	-0.148
Asteraceae	*Krigia*	*biflora*	19	0	-0.542
Asteraceae	*Lactuca*	serriola	4	0	-0.257
Asteraceae	*Leucanthemum*	vulgare	405	20	-0.349
Asteraceae	*Liatris*	spicata	186	127	1.137
Asteraceae	*Ratibida*	pinnata	539	121	0.005
Asteraceae	*Rudbeckia*	hirta	1174	189	0.645
Asteraceae	*Solidago*	altissima	8	0	-0.586
Asteraceae	*Solidago*	canadensis	8	2	0.816
Asteraceae	*Solidago*	gigantea	107	9	0.613
Asteraceae	*Solidago*	juncea	636	77	1.106
Asteraceae	*Solidago*	rugosa	1	0	-0.015
Asteraceae	*Symphyotrichum*	novae-angliae	25	6	0.451
Asteraceae	*Vernonia*	noveboracensis	52	37	-0.045
Brassicaceae	*Barbarea*	vulgaris	3	0	-0.114
Campanulaceae	*Lobelia*	inflata	12	3	1.147
Campanulaceae	*Lobelia*	siphilitica	2	0	-0.034
Caryophyllaceae	*Dianthus*	armeria	3	0	-0.280
Commelinaceae	*Tradescantia*	ohiensis	34	1	-0.805
Convulvulaceae	*Calystegia*	silvatica	5	0	-0.538
Cornaceae	*Swida*	amomum	4	0	-0.250
Cornaceae	*Swida*	racemosa	12	1	0.546
Fabaceae	*Baptisia*	tinctoria	19	5	-0.566
Fabaceae	*Chamaecrista*	fasciculata	246	2	-1.514
Fabaceae	*Desmodium*	paniculatum	6	1	-0.138
Fabaceae	*Lotus*	corniculatus	142	33	-0.231
Fabaceae	*Melilotus*	albus	20	9	0.571
Fabaceae	*Melilotus*	officinalis	41	21	1.309
Fabaceae	*Securigera*	varia	38	1	-0.654
Fabaceae	*Senna*	hebecarpa	5	5	1.678
Fabaceae	*Trifolium*	aureum	1	0	-0.033
Fabaceae	*Trifolium*	campestre	365	39	0.106
Fabaceae	*Trifolium*	hybridum	11	7	0.739
Fabaceae	*Trifolium*	pratense	192	20	-0.869
Fabaceae	*Trifolium*	repens	130	6	-0.847
Fabaceae	*Vicia*	tetrasperma	1	0	-0.048
Gentianaceae	*Sabatia*	angularis	1	0	-0.034

Hypericaceae	*Hypericum*	*perforatum*	223	10	-1.095
Hypericaceae	*Hypericum*	*punctatum*	1	0	-0.176
Iridaceae	*Sisyrinchium*	*angustifolium*	9	0	-0.351
Lamiaceae	*Clinopodium*	*vulgare*	64	3	-0.854
Lamiaceae	*Monarda*	*fistulosa*	1397	400	-0.131
Lamiaceae	*Monarda*	*punctata*	0	1	1.242
Lamiaceae	*Nepeta*	*cataria*	99	81	1.270
Lamiaceae	*Prunella*	*vulgaris*	17	5	0.296
Lamiaceae	*Pycnanthemum*	*muticum*	398	59	0.811
Lamiaceae	*Pycnanthemum*	*tenuifolium*	1113	421	0.609
Lamiaceae	*Pycnanthemum*	*verticillatum*	5	8	0.886
Lamiaceae	*Teucrium*	*canadense*	3	0	-0.104
Loniceraceae	*Lonicera*	*japonica*	1	0	-0.131
Lythraceae	*Lythrum*	*salicaria*	364	38	-0.016
Onagraceae	*Oenothera*	*biennis*	2	0	-0.143
Onagraceae	*Oenothera*	*fruticosa*	2	1	0.221
Oxalidaceae	*Oxalis*	*stricta*	8	0	-0.249
Phytolaccaceae	*Phytolacca*	*americana*	108	74	1.580
Plantaginaceae	*Plantago*	*lanceolata*	147	0	-1.997
Poaceae	*Glyceria*	*grandis*	1	0	-0.187
Poaceae	*Sorghastrum*	*nutans*	1	0	-0.092
Polygonaceae	*Fallopia*	*convolvulus*	3	4	1.044
Polygonaceae	*Persicaria*	*pensylvanica*	4	0	-0.391
Polygonaceae	*Persicaria*	*setacea*	3	1	0.158
Ranunculaceae	*Ranunculus*	*hispidus*	1	0	-0.057
Rosaceae	*Drymocallis*	*arguta*	22	8	0.604
Rosaceae	*Potentilla*	*recta*	56	1	-0.628
Rosaceae	*Rosa*	*carolina*	71	1	-1.412
Rosaceae	*Rosa*	*multiflora*	5	0	-0.264
Rosaceae	*Rubus*	*flagellaris*	1	0	-0.027
Rosaceae	*Rubus*	*pensilvanicus*	7	0	-0.241
Rubiaceae	*Galium*	*mollugo*	4	0	-0.369
Scrophulariaceae	*Linaria*	*vulgaris*	275	4	-1.633
Scrophulariaceae	*Penstemon*	*digitalis*	862	48	-0.847
Scrophulariaceae	*Penstemon*	*hirsutus*	36	0	-0.822
Scrophulariaceae	*Verbascum*	*blattaria*	15	0	-0.721
Scrophulariaceae	*Verbascum*	*thapsus*	129	1	-1.085
Solanaceae	*Solanum*	*carolinense*	14	0	-0.518
Verbenaceae	*Verbena*	*hastata*	8	3	0.927

| Verbenaceae | *Verbena* | *simplex* | 11 | 3 | 0.482 |
| Verbenaceae | *Verbena* | *urticifolia* | 58 | 71 | 2.118 |