1   **Male and female bees show large differences in floral preference**
2

3   Michael Roswell mroswell.rutgers@gmail.com
4   Graduate program in ecology and evolution, Rutgers University
5   14 College Farm Road, New Brunswick, NJ 08904
6
7   Jonathan Dushoff
8   Department of biology, McMaster University
9   1280 Main St. West, Hamilton, Ontario ON L8S 4K1
10
11   Rachael Winfree
12   Department of ecology, evolution, and natural resources, Rutgers University
13   14 College Farm Road, New Brunswick, NJ 08904

Abstract

1.   Efforts to conserve and support wild bees, which are important pollinators of wild plants and food crops, often rely on planting pollinator-attractive flowering plants to augment forage resources. Bee preferences for particular plant taxa should guide plant inclusion in these plantings. Thus far, studies of bee preference focused on identifying the plant species preferred by different bee species. However, strong natural history differences between male and female bees may create differences in foraging preference even within species. The foraging ecology of male bees is little known.

2.   We observed bees visiting flowers in semi-natural meadows in New Jersey, USA, and ask whether male and female bees prefer different species of flowers. To detect differences in flower use against a shared background resource availability, we maximized the number of interactions observed within narrow spatio-temporal windows. To distinguish observed differences in bees' floral use, which can reflect neutral, abundance-based patterns and sampling effects, from underlying differences in bee preferences, we analyzed our data with both a permutation-based null model and random effects models.

3.   We found that the diets of male and female bees of the same species were often as dissimilar as the diets of bees of different species. We identify plant species that are particularly attractive to male bees.

Synthesis and applications: Given that male and female bees use and prefer different species of flower, seed producers and land managers can choose how to balance prioritization of male- or female-preferred flower species in bee habitat enhancements, according to restoration targets.

Key Words

Introduction

Most species of land plants rely on animal pollinators for sexual reproduction (Ollerton, Winfree, & Tarrant, 2011), and human food systems also require animal pollinators (Potts et al., 2016). This dependence on pollinators motivates significant efforts to conserve and augment bee populations. While natural and semi-natural habitats are critical landscape elements for many bee species (Senapathi, Goddard, Kunin, & Baldock, 2016), what constitute suitable and/ or limiting resources within these habitats remains less clear (De Palma et al., 2015). Flowers, which provide food for adult and larval bees, are likely among them (Roulston & Goodell, 2011). Currently, in the United States and Europe, pollinator conservation focuses on planting bee-attractive flowering plants (Tonietto & Larkin, 2018).

Determining the appropriate plant taxa for inclusion in pollinator habitat plantings is a central challenge in pollinator conservation (Williams & Lonsdorf, 2018). The

links between bee demography and flowers in the landscape remain poorly
known (Dicks et al., 2015; Spiesman, Bennett, Isaacs, & Gratton, 2017).
Nonetheless, assuming that pollinators tend to prefer the plants that increase
their fitness, pollinator preference can guide which taxa to plant. Preference—the
use of a resource in excess of its relative availability—is challenging to measure,
because both resource use and availability must be known. Floral resource
availability for pollinators is particularly hard to quantify outside an experimental
context because the appropriate scale and units of floral resource availability are
unclear. The composition, amount, and supply rate of pollen and nectar per
flower, the number of flowers per inflorescence, of inflorescences per individual,
and the number and distribution of individual plants over the square kilometers of
a bee's foraging range are important, but distinct, components of availability
(Hicks et al., 2016). Furthermore, floral availability can change rapidly over time.
However, differences in flower use between bees foraging at the same place and
time indicate differences in preference, which may occur between species, or
between individuals of the same species.
76
Intraspecific variation in traits and behavior, including foraging niche, has
important consequences for species interactions (Thomson & Chittka, 2001), and
conservation (Bolnick et al., 2011; Durell, 2000). A widespread source of
intraspecific variation, sexual dimorphism, serves as a platform for studying
adaptation, speciation, and the way in which animals exploit their ecological
niche (Butler, Sawyer, & Losos, 2007; Temeles, Miller, & Rifkin, 2010).
Morphological, behavioral, and life-history dimorphisms can also drive the form
and function of ecosystems (Fryxell, Arnett, Apgar, Kinnison, & Palkovacs, 2015;
Ogilvie & Thomson, 2015). Although intraspecific variation in floral preference is
well known in social (Heinrich, 1979) and to a lesser extent, solitary bee species
(Bruninga-Socolar, Crone, & Winfree, 2016; Tur, Vigalondo, Trøjelsgaard,
Olesen, & Traveset, 2014), previous studies focused on female bees.
89
Male bees differ from their better-studied female counterparts in their life history
and ecology. Female bees construct, maintain, provision, and defend nests,
whereas male bees primarily seek mates (Willmer & Stone, 2004). Most
importantly, while both sexes drink floral nectar for their own caloric needs, only
females collect pollen to provision young. While the pollen from each flower
species (the term we use throughout for the flowers from a species of plant)
tends to be morphologically and nutritionally distinct, interspecific variation in the
chemical composition of nectar is comparatively subtle (Cane & Sipes, 2006).
Thus, we expect the foraging ecology, including floral preferences, of male and
female bees to differ as well.
100
Current guidelines for restoring pollinator habitat are based on studies involving
only or primarily females, and thus overlook the needs of male bees. This is
because most bee studies quantify foragers, which, because females need to
forage for both themselves and offspring, tend to be female. The foraging and
pollination ecology of male bees is little known, yet male bees prove to be

important pollinators when studied (Ogilvie & Thomson, 2015). Male bees are
also relevant to bee conservation efforts. Males may be limiting in declining
populations (Elias, Dorn, & Mazzi, 2010; Straub et al., 2016), and, as the
dispersing sex in most species, males may be crucial for gene flow and
metapopulation persistence even when they are not locally limiting (López-Uribe,
Morreale, Santiago, & Danforth, 2015; Ulrich, Perrin, & Chapuisat, 2009).
112
In this study, we assess differences between floral preferences of male and
female bees in the wild. We collected bees foraging on flowers in meadows in
New Jersey, USA. In order to observe preference differences, we collected as
many individuals as possible during replicated short (3-day) windows, during
which we assumed floral availability and bee abundance were constant at each
site. We compare the species composition of flowers visited by males and
females of the most common bee species across the entire study as a naïve
measure of differences in preference between the sexes. Then, using random
effects models, we assess when differential flower species *use* by male and
female bees likely to arises from sex-specific floral *preference*, as opposed to
shifting overlap between foragers and floral resources (i.e. changes in *availability*
without differences in *preference*). Specifically, we ask

1) How much do male and female bee diets overlap?
2) To what degree are particular flower species disproportionately visited by
bees of one sex?
3) To what extent are apparent differences in preference driven by
phenological differences between male and female bees?

Materials and Methods

Study design and data collection

Because absolute preference is nearly impossible to observe outside of an
experiment, we designed our study to reveal differences in preference between
groups of bees. In order to collect a large number of males and females from
many native bee species, we selected six meadows (sites) in New Jersey, USA
with a high abundance and diversity of flowers. These meadows were managed
for pollinator-attractive, summer-blooming forbs through seed addition, and a
combination of mowing, burning, and weed removal. Most flower species present
in the meadows are native to the eastern United States. We collected our data
during peak bloom and maximum day length (6 June to 20 August, 2016), and
during good weather (sunny enough for observers to see their own shadow, no
precipitation). We visited each site for three consecutive good weather days over
five evenly spaced sampling rounds in the 11-week period of our study. In all
analyses, we assume that bees and flowers detected at a site within one 3-day
sampling round co-occurred. In contrast, we assume that turnover of both plant
species in bloom and bee species activity can occur in the ~10 days between
sampling rounds.

152
During each 3-day sampling round, an observer walked parallel transects
through the meadow (which ranged in size from 0.8–2.2 ha; mean=1.4 ha),
observing every open flower within a moving 1-m semicircle, and net-collecting
any bee seen actively foraging, which we defined as contacting anthers or
collecting nectar from a flower (Fig. S1). We collected all bee species except
Apis mellifera L.*,* the domesticated western honey bee, because *Apis* males do
not forage. Observations began as soon as pollinator activity picked up in the
morning (7–9 am) and continued into the late afternoon or evening until pollinator
activity slowed substantially. Observers sampled nearly continuously, in 30-
minute timed collection bouts with short breaks in between. If inclement weather
precluded a minimum of six 30-minute sampling bouts in a day, we added an
additional day to the sampling round as soon as weather permitted.
165
Flower species were identified in the field by the data collector. Bee species were
identified using a dissecting microscope and published keys; Jason Gibbs
(University of Manitoba), Joel Gardner (University of Manitoba), and Sam Droege
(USGS) assisted with identification for bees in the genera *Andrena, Anthophora,*
Coelioxys, Halictus, Heriades, Hoplitis, Hylaeus, Lasioglossum, Megachile,
Melissodes, Nomada, Osmia, Pseudoanthidium, Ptilothrix, Sphecodes, Stelis,
and *Triepeolus,* and at least one of them confirmed voucher specimens for every
species. We determined every specimen to species except for the following four
complexes: Most bees in the genus *Nomada* with bidentate mandibles (*ruficornis*
group) were treated as one species. All specimens from the *Hylaeus* species
complex that includes *Hylaeus affinis*, *H. modestus*, and at least one additional
species, informally dubbed "species A*,*" were treated as a single species,
denoted *Hylaeus affinis-modestus*, because females cannot be reliably
distinguished. There is a cryptic species in the genus *Halictus* unlikely to occur in
our area, *Halictus poeyi*, which is not morphologically distinct from *H. ligatus*; we
treat all specimens in this complex as *Halictus ligatus*. We could not confidently
separate all specimens of the two closely related *Lasioglossum* species
Lasioglossum hitchensi and *L. weemsi*. Thus, we treat all specimens from either
species as one, denoted *Lasioglossum hitchensi-weemsi.* All bee specimens are
curated in the Winfree lab collection at Rutgers University, and the data used in
this paper are available from the Dryad Digital Repository
http://dx.doi.org/XXXXXXX (Roswell et al.)
188
189 **Analytical methods**
190
191 We performed all statistical analyses and simulations using R 3.5.1 (R Core
192 Team, 2018).
193
194   1) How much do male and female bee diets overlap?
195
To compare the diets of male and female bees, we used the Morisita-Horn index
of resource overlap (Horn, 1966; Morisita, 1959). This dissimilarity index

compares the proportion of all female bees found on each flower species to the
proportion of all male bees found on each flower species. In other words, it
compares the contribution of each flower species to female diets (where this term
includes the food that females collect for themselves and also to feed to young)
to the contribution of the same flower species to male diets. The Morisita-Horn
index ranges from zero (completely similar) to one (maximally dissimilar), and
has several good properties for our purposes. First, it uses proportions, placing
visits from male and female bees on the same scale, even though most visits
come from females. Second, it is much more sensitive to large proportions than
to small ones, thereby down-weighting the contribution of flower species for
which we have little information. Third, the Morisita-Horn estimates are resilient
to undersampling and uneven sample size between groups (Barwell, Isaac, &
Kunin, 2015).
211
To determine whether the male-female differences we observed exceeded those
expected by chance, we compared the observed compositional dissimilarity
between flower visits from male and from female bees to dissimilarity measures
from a null model that randomly permuted the bee sex associated with each
flower-visit record. This permutation holds constant the total number of male and
of female visits, and the total number of visits to each flower species from both
sexes combined (Fig. S2). The range of dissimilarity values from this simulation
is the difference we would observe in our sample, if there were no true difference
in flower species use between males and females of the same bee species. We
evaluated the hypothesis that male and female diets overlap less than would be
expected by chance; thus we use a one-sided alpha of $p<0.05$. We iterated this
null simulation 9999 times, which was sufficient to stabilize p-values near our
chosen alpha (North, Curtis & Sham 2002). When the observed dissimilarity was
greater than 9500 of the 9999 simulated dissimilarities, we concluded that we
had detected a difference in the pattern of floral visitation between conspecific
male and female bees, given the observed diet breadth and abundance of each
sex.
229
To compare the diet overlap we observed between sexes to a meaningful
benchmark, interspecific diet overlap, we repeated the same null model analysis,
this time comparing females of the focal species to females of other species. We
performed one analysis for each bee species for which we collected at least 20
visitation records for each sex (19 species). This sample size threshold is
arbitrary, but null model variance shrinks with sample size, such that apparent
patterns for species with smaller sample sizes are rarely interpretable (Fig. S3).
Because we analyze 19 bee species, females of each species are compared to
18 others. We then compared the male-female difference (observed minus mean
null dissimilarity in flower communities visited) to the analogous species-species
difference (observed minus null dissimilarity).
241
242   For this analysis, which evaluates holistic differences between male and female
243   bees of the same species, we combined observations across the full season and

244 all sites. This allows us to observe foraging niche differences that are driven by
245 flower and/ or bee phenology, in addition to any sex-specific floral preference.
246
247 **2)** *To what degree are particular flower species disproportionately visited by bees*
248 *of one sex?*
249
This analysis uses our entire data set of 153 bee species to determine whether
particular flower species are disproportionately visited by male or female bees,
and whether the answer varies by bee species. We can infer a preference
difference between male and female bees for a flower species when predicted
odds of visitors to that flower species being male are especially high or low. To
do this, we use a random effects model in which bee sex is the response, and
flower species, bee species, site, and their interactions are random effects. We
statistically control for variation in the overall sex ratio across bee species
through the term (1|bee_species), and variation in sex ratios across sites,
through the terms (1|site), (1|site:bee_species), and (1|site:flower_species).
Because it is unlikely that, within bee species, sex ratios at birth vary greatly
across space, any variability attributed to site terms would likely result from
differential overlap of bee foraging activity and flower bloom across space.
263
We call this model the "summed model" because we sum interactions observed
across the entire season (all five sampling rounds) at each site. In the summed
model, the relationship between phenological overlap and the odds of flower-
visiting bees being male would be incorporated into the species effects. This
perspective is helpful for considering flower species' contributions to the overall
diets of male versus female bees. We fit the model with the R package lme4
(Bates, Maechler, & Walker, 2016) with the following call:
271
272 ***Summed model***
273 lme4::glmer(bee_sex ~ (1|site)+ (1|flower_species)+ (1|bee_species)+
274   (1|flower_species:bee_species)+ (1|site:bee_species)+
275   (1|site:flower_species), family="binomial", data=data)
276
We included bee species and site as random, rather than fixed, effects to directly
compare the variability in bee sex associated with each of these predictors to the
variability associated with flower species (preference). Comparing the overall
variability across these groups was more important to us than assessing
predictions on a per-site or per-bee-species basis. We fit flower species, the
primary covariate of interest, as a random effect to facilitate model fitting (fewer
degrees of freedom) as well as interpretation. In our summed model, we included
all two-way interactions, but omitted the three-way interaction, bee species by
flower species by site. Although the sort of context-dependent preference this
term could represent (e.g. males from bee species *1* prefer flower *A* at one site
(relative to females), but shun it at another) may exist in nature, it is unlikely we
would estimate it accurately in our model.
289

We confirmed model convergence by comparing several fitting methods using
the allFit function in lme4 (Bates et al., 2016), which all showed similar parameter
estimates (Table S1). We tested whether residuals from our model fit were
overdispersed using Bolker's function "overdisp" (Bolker, 2017), and visually
assessed our additivity assumptions with binned residual plots (Gelman & Hill,
2007) (Fig. S4).
296
297 *3) To what extent are apparent differences in preference driven by phenological*
298 *differences between male and female bees?*
299
Over the 11 weeks of our study, we observed turnover in bee species, in flower
bloom, and within-bee species changes in sex ratio. Therefore, phenological
overlap between male versus female bees and the bloom period of particular
flower species, rather than preference of those bees for those flowers, may
explain much of the variation in sex ratio we observed across visitors. In question
3, we are explicitly interested in distinguishing sex-specific diet *preferences* from
variable *use* resulting from seasonal resource availability and male vs. female
abundance. We do this in the "seasonal model" by incorporating sampling round
(our measure of phenology) as an additional random intercept effect, along with
random intercepts for the interactions between sampling round and the other
covariates. We chose to include sampling round as a random effect because this
enables direct comparison to all other terms in both models. We ignored the
three- and four-way interactions between bee species, flower species, and other
covariates. We fit this model with the following call in the R package lme4, with
new terms in bold:
315
Seasonal model
glmer(bee_sex ~ (1|site)+ (1|flower_species)+ (1|bee_species)+
(1|flower_species:bee_species)+ (1|site:bee_species)+
(1|site:flower_species)**+ (1|sampling_round)+ (1|site:sampling_round)+**
(1|flower_species:sampling_round)+
(1| bee_species:sampling_round)+
(1|site:bee_species:sampling_round)+
(1|site:flower_species:sampling_round), family="binomial", data=data)
324
Our index of preference for both the *summed model* and the *seasonal*
model is the change in odds that a bee is male when the flower species it visits is
given. To describe the importance of model terms, we calculated a bootstrapped
median odds ratio using code from Seth (Seth, 2017), which gives the expected
difference in odds that a flower-visiting bee is male between levels of a predictor
(Merlo et al., 2006). For example, a median odds ratio of five for the main effect
of sampling round would indicate that the odds of a flower-visiting bee being
male differ by about a factor of five between sampling rounds, while a median
odds ratio of one would indicate that the odds of a flower-visiting bee being male
do not change across rounds. If the median odds ratio is large for flower species
in both models, we could say that there are intrinsic (i.e. not simply phenological)

properties of flower species identity that male or female bees prefer. If flower species is a strong predictor of bee sex in the summed model but not in the seasonal one, we would still conclude that flower species often contribute more strongly to the diet of one sex than the other, though these differences may not arise due to differing preferences. If the sampling round terms have large median odds ratios, then accounting for phenology is critical for identifying differences in preference in addition to differences in use.

Results

In total we collected 18,687 bee specimens belonging to 152 bee species (table S2) from a total of 109 flower species (table S3), which together comprised 1417 unique species-species interactions. Roughly 18% of specimens were male (n=3369). Thus, the overall ratio of male to female bees we collected was 0.22, although this ratio varied markedly between flower species (Fig. 1).

How much do male and female bee diets overlap?

We found that male and female bee diets overlap significantly less than would be expected at random (Fig. 2), and that the differences in diet composition between male and female bees of several species were of similar magnitude to the differences in diet between species of bee (Fig. 3).

To what degree are particular flower species disproportionately visited by bees of one sex?

The sex ratio of flower-visiting bees varied across species of flower (Fig. 2). After controlling for bee species identity (the strongest predictor of sex in our models, Fig. 4), and site, we still found that some flower species received a disproportionate number of male bee visitors (Figs 4–5). The median odds ratio for the main effect of flower species was 3.6 (bootstrapped CI 3.0–4.2) in our summed model, indicating that typically, the visitor sex ratio differs between two flower species by more than a factor of 3. Furthermore, we observed sex-based differences in flower use specific to particular bee species: the median odds ratio for the flower species by bee species interaction in our summed model was nearly as large (median=3.1, bootstrapped CI 3.0–3.3) as the main effect of flower species. By contrast, sex ratios are not expected to differ between sites (median odds ratio for main effect of site=0).

3) To what extent are apparent differences in preference driven by phenological differences between male and female bees?

The flower species blooming in our system turned over throughout our 11-week sampling period, with several highly visited species blooming for only one of the three months during which we sampled. This turnover, along with potential sex-specific bee flight seasons, means that differences in diet between male and

female bees could reflect seasonal availability and use, without also indicating preference differences between the sexes. Indeed, phenology predicts bee sex somewhat, with the odds of a flower-visiting bee being male expected to change by a factor of 1.5 (bootstrapped CI 1.1–1.9) between sampling rounds (Fig. 4). Phenological patterns of male vs. female flight seasons vary across bee species; the median odds ratio for the bee species by sampling round interaction is 2.2 (bootstrapped CI 2.1–2.3) (Fig. 4). Even after accounting for these effects, however, there remains a strong association between the species of flower a bee visits and its sex (Figs 4–5). The relative effects of each flower species on the sex of its visitors were changed very little by accounting for phenology; Pearson and Spearman correlations between the random effect of flower species in the seasonal model and the same random effect in the simpler summed model were both 0.98. In addition to finding overall preference difference between male and females, we found evidence for bee-species-specific difference in floral preferences between the sexes (median odds ratios in both models for the bee species by flower species interaction > 2.8).

Discussion

We found strong differences between the flower species preferences of male and female bees. The difference in floral visits between male and female bees was similar in magnitude to visitation differences between females of different species. The partitioning of the floral community among bee species is a primary focus of pollination ecology and ecological network analysis (Bascompte & Jordano, 2014), but male bees are typically disregarded or lumped together with their female counterparts. Our study suggests this may represent an important oversight. Further, our study provides strong confirmation of the few studies that investigate the foraging behavior of male bees, which found that males play a unique role in plant pollination (Cane, Sampson, & Miller, 2011; Ogilvie & Thomson, 2015). Our result also implies that male bees contribute substantially to the complexity of plant-pollinator networks in nature, and that network analyses might benefit from separating males and females into different nodes (Bolnick et al., 2011).

Designing plant mixes to support and augment pollinator populations may also benefit from considering the foraging needs of males and females separately (Rundlöf, Persson, Smith, & Bommarco, 2014). Whether floral diversity *per se* benefits individual pollinator taxa remains murky (Spiesman et al., 2017; Sutter, Jeanneret, Bartual, Bocci, & Albrecht, 2017). However, complementary flower-species use by the sexes provides a new potential link between landscape floral diversity and bee populations. In addition, it is currently unknown how the distinct foraging niches of male bees mediate either the robustness of pollinator communities to species loss and environmental perturbations (Brosi & Briggs, 2013; Ramos-Jiliberto, Valdovinos, Moisset de Espanés, & Flores, 2012; Tur et al., 2014), or the effectiveness of different habitat ameliorations (Rundlöf et al., 2014; Rusterholtz & Erhardt, 2000). This study suggests that both questions

428   warrant further investigation.
429
Phenology, a previously reported mechanism for distinct use of floral resources
by male and female bees (Ogilvie & Thomson, 2015; Robertson, 1925),
explained some variation in the sex ratio of flower-visiting bees, but was less
important than flower species identity over the period of our study. We expected
to find an effect of phenology because both the identity of the flower species
blooming within sites, and also the sex ratio of foragers within bee species, vary
across the season. Males emerge first in most solitary bees; for social species,
initial broods usually consist primarily of female workers, then males and
reproductive females emerge at the end of the colony cycle (Willmer & Stone,
2004). Surprisingly, however, phenology only weakly predicted the sex of flower-
visiting bees. This is despite the fact that, as predicted by natural history, the
sampling round(s) in which males were relatively more prevalent depended on
bee species (the bee species by sampling round interaction was much bigger
than the sampling round main effect; Fig. 4). This indicates that our evidence for
floral preference differences between male and female bees was robust to
accounting for seasonal turnover in flower species bloom, bee species flight
seasons, and the sex ratios within bee species.
447
Whereas female bees collect both nectar and pollen, male bees forage primarily
for nectar to fuel flight. Thus, we predicted that male bees would avoid flowers
that produce no nectar. Indeed, in both our models, the predicted odds of a bee
visiting a nectar-less flower species being male were approximately half that of a
bee visiting a flower species that produces nectar (Fig. S5). A second biological
difference between male and female bees is that adult male bee activities orient
around mate seeking (Alcock et al., 1978). These behaviors, such as patrolling
routes (Barrows, 1976) or seeking flowers visited by conspecific females (Rossi,
Nonacs, & Pitts-Singer, 2010) could generate differences from females via
complementarity (males visiting flower species not visited by females), or
nestedness (one sex primarily visiting a subset of species visited by the other).
We found evidence for both (Fig. S6). Divergent floral preferences between
sexes may reflect nutritional needs or mating behavior, or simply biases resulting
from previous flower encounters, or visual or olfactory sensitivities that differ
between the sexes (Robert, Frasnelli, Collett, & de Ibarra, 2016; Somanathan,
Borges, Warrant, & Kelber, 2017; Streinzer, Kelber, Pfabigan, Kleineidam, &
Spaethe, 2013).
465
Patterns in bee-flower interaction data can arise from the sampling process itself
(Blüthgen, 2010; Fründ, McCann, & Williams, 2016). Our analyses control for
these patterns. To evaluate diet overlap, we used a dissimilarity index that
downweights rare diet items, and implemented a null model that accounts for
differences that could arise from sampling and abundance effects. To evaluate
preference, we used random effects models that incorporated all (nearly 19,000)
observations, and shrank extreme values for rarely observed species-species
interactions towards the global mean for each effect. Thus, our estimates for sex-

specific preferences should be robust to the inevitable undersampling of rarer
taxa. Establishing differences in preference between categories of bees such as
males and females, even when resource availability is seasonal and difficult to
quantify, is possible using methods such as these, though absolute preference
remains elusive.
479
Pollination ecology and pollinator conservation still face the question of how
important sexually dimorphic foraging is. Does it enhance or reduce the stability
of bee populations? Should pollinator restorations explicitly include "male bee"
flowers and "female bee" flowers? Are floral traits under selection to favor female
versus male visitors? While our study does not answer these questions, in
showing that the diets and preferences of male bees commonly differ from those
of their female conspecifics, we suggest they are worthy of future study.
487

488 **Author contributions**
489
490 MR and RW conceived the ideas and designed field methodology; MR, JD, and
491 RW designed statistical methodology. MR collected the data; MR, RW, and JD
492 analyzed the data; MR and RW led manuscript drafting. All authors contributed
493 critically to the drafts and gave final approval for publication.
494

Acknowledgements
We thank Daniel Cariveau and Neal Williams for study design suggestions, and
Jason Gibbs and Tina Harrison for comments on an earlier draft. A USDA
Conservation Innovation Grant to the Xerces Society for Invertebrate Biology
(lead PI) and RW (senior investigator) funded data collection. MR was supported
by an NSF Graduate Research Fellowship. Cameron Kanterman and Riva
Letchinger helped collect field data; Kurtis Himmler, Tiffany Bennet, and
professional taxonomists Jason Gibbs, Joel Gardiner, and Sam Droege helped
with bee species determinations. Mercer County, the Institute for Advanced
Study, Somerset County, and the Raritan Headwaters Association provided site
permission.

References

Alcock, J., Barrows, E. M., Gordh, G., Hubbard, L. J., Kirkendall, L., Pyle, D. W.,
 … Zalom, F. G. (1978). The ecology and evolution of male reproductive
 behaviour in the bees and wasps. *Zoological Journal of the Linnean Society*,
 64(4), 293–326. doi:10.1111/j.1096-3642.1978.tb01075.x

Barrows, E. M. (1976). Mating behavior in halictine bees (Hymenoptera:
 Halictidae): II. microterritorial and patrolling behavior in males of
 Lasioglossum rohweri. *Zeitschrift Fur Tierpsychologie-Journal of
 Comparative Ethology*, *40*(4), 377–389.

Barwell, L. J., Isaac, N. J. B., & Kunin, W. E. (2015). Measuring beta-diversity
 with species abundance data. *Journal of Animal Ecology*, 1112–1122.
 doi:10.1111/1365-2656.12362

Bascompte, J., & Jordano, P. (2014). Mutualistic networks in time and space. In
 J. Bascompte & P. Jordano (Eds.), *Mutualistic networks* (pp. 87–106).
 Princeton University Press. Retrieved from
 http://www.jstor.org/stable/j.ctt5hhnpq.8

Bates, D., Maechler, M., & Walker, S. (2016). Package "lme4." *CRAN
 Repository*. doi:10.18637/jss.v067.i01

Blüthgen, N. (2010). Why network analysis is often disconnected from community
 ecology: A critique and an ecologist's guide. *Basic and Applied Ecology*,
 11(3), 185–195. doi:10.1016/j.baae.2010.01.001

Bolker, B. M. (2017). GLMM FAQ. Retrieved from
 http://bbolker.github.io/mixedmodels-misc/glmmFAQ.html

Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak,
 M., … Vasseur, D. A. (2011). Why intraspecific trait variation matters in
 community ecology. *Trends in Ecology and Evolution*, *26*(4), 183–192.
 doi:10.1016/j.tree.2011.01.009

Brosi, B. J., & Briggs, H. M. (2013). Single pollinator species losses reduce floral
 fidelity and plant reproductive function. *Proceedings of the National
 Academy of Sciences*, *110*(32), 13044–8. doi:10.1073/pnas.1307438110

Bruninga-Socolar, B., Crone, E. E., & Winfree, R. (2016). The role of floral
 density in determining bee foraging behavior: a natural experiment. *Natural
 Areas Journal*, *36*(4), 392–399. doi:10.3375/043.036.0406

Butler, M. A., Sawyer, S. A., & Losos, J. B. (2007). Sexual dimorphism and
 adaptive radiation in Anolis lizards. *Nature*, *447*(7141), 202–5.
 doi:10.1038/nature05774

Cane, J. H., Sampson, B. J., & Miller, S. A. (2011). Pollination value of male
 bees: The specialist bee Peponapis pruinosa (Apidae) at Summer Squash
 (Cucurbita pepo). *Environmental Entomology*, *40*(3), 614–620.
 doi:10.1603/EN10084

Cane, J. H., & Sipes, S. S. (2006). Characterizing floral specialization by bees:
 Analytical methods and a revised lexicon for oligolecty. In *Plant–pollinator
 interactions: From specialization to generalization*. doi:10.1007/BF00937854

De Palma, A., Kuhlmann, M., Roberts, S. P. M., Potts, S. G., Börger, L., Hudson,
 L. N., … Purvis, A. (2015). Ecological traits affect the sensitivity of bees to
 land-use pressures in European agricultural landscapes. *Journal of Applied*

Ecology, *52*(6), 1567–1577. doi:10.1111/1365-2664.12524
Dicks, L. V., Baude, M., Roberts, S. P. M., Phillips, J., Green, M., & Carvell, C.
(2015). How much flower-rich habitat is enough for wild pollinators?
Answering a key policy question with incomplete knowledge. *Ecological*
Entomology, *40*(S1), 22–35. doi:10.1111/een.12226
Durell, S. E. A. L. V. dit. (2000). Individual feeding specialization in shorebirds:
population consequences and conservation implications. *Biological Reviews*
of the Cambridge Philosophical Society, *75*(2000), 503–518.
Elias, J., Dorn, S., & Mazzi, D. (2010). No evidence for increased extinction
proneness with decreasing effective population size in a parasitoid with
complementary sex determination and fertile diploid males. *BMC*
Evolutionary Biology, *10*(1), 366. doi:10.1186/1471-2148-10-366
Fründ, J., McCann, K. S., & Williams, N. M. (2016). Sampling bias is a challenge
for quantifying specialization and network structure: Lessons from a
quantitative niche model. *Oikos*, *125*(4), 502–513. doi:10.1111/oik.02256
Fryxell, D. C., Arnett, H. A., Apgar, T. M., Kinnison, M. T., & Palkovacs, E. P.
(2015). Sex ratio variation shapes the ecological effects of a globally
introduced freshwater fish. *Proceedings of the Royal Society B*, *282*(1817),
20151970. doi:10.1098/rspb.2015.1970
Gelman, A., & Hill, J. (2007). *Data analysis using regression and multilevel/*
hierarchical models. Cambridge. doi:10.2277/0521867061
Heinrich, B. (1979). "Majoring" and "Minoring" by foraging bumblebees, Bombus
vagans: an experimental analysis. *Ecology*, *60*(2), 245–255.
Hicks, D. M., Ouvrard, P., Baldock, K. C. R., Baude, M., Goddard, M. A., Kunin,
W. E., … Stone, G. N. (2016). Food for pollinators: Quantifying the nectar
and pollen resources of urban flower meadows. *PLoS ONE, 11*(6).
doi:10.1371/journal.pone.0158117
Horn, H. S. (1966). Measurement of "Overlap" in comparative ecological studies.
The American Naturalist, *100*(914), 419–424. Retrieved from
http://www.jstor.org/stable/2459242
López-Uribe, M. M., Morreale, S. J., Santiago, C. K., & Danforth, B. N. (2015).
Nest suitability, fine-scale population structure and male-mediated dispersal
of a solitary ground nesting bee in an urban landscape. *PLoS ONE, 10*(5),
1–20. doi:10.1371/journal.pone.0125719
Merlo, J., Chaix, B., Ohlsson, H., Beckman, A., Johnell, K., Hjerpe, P., … Larsen,
K. (2006). A brief conceptual tutorial of multilevel analysis in social
epidemiology: Using measures of clustering in multilevel logistic regression
to investigate contextual phenomena. *Journal of Epidemiology and*
Community Health, *60*(4), 290–297. doi:10.1136/jech.2004.029454
Morisita, M. (1959). Measuring of interspecific association and similarity between
communities. *Memoirs of the Faculty of Science, Kyushu University, Series*
E (Biology). Retrieved from http://reference.morisita.gr.jp/paper_pdf/56.pdf
Ogilvie, J. E., & Thomson, J. D. (2015). Male bumble bees are important
pollinators of a late-blooming plant. *Arthropod-Plant Interactions*, *9*(2), 205–
213. doi:10.1007/s11829-015-9368-x
Ollerton, J., Winfree, R., & Tarrant, S. (2011). How many flowering plants are

pollinated by animals? *Oikos*, *120*(3), 321–326. doi:10.1111/j.1600-
0706.2010.18644.x
Potts, S. G., Imperatriz-Fonseca, V. L., Ngo, H. T., Biesmeijer, J. C., Breeze, T.
D., Dicks, L. V, … Viana, B. F. (2016). *The assessment report on pollinators,*
pollination and food production. Summary for policymakers.
R Core Team. (2018). R: A Language and Environment for Statistical Computing.
Vienna, Austria. Retrieved from https://www.r-project.org/
Ramos-Jiliberto, R., Valdovinos, F. S., Moisset de Espanés, P., & Flores, J. D.
(2012). Topological plasticity increases robustness of mutualistic networks.
Journal of Animal Ecology, *81*(4), 896–904. doi:10.1111/j.1365-
2656.2012.01960.x
Robert, T., Frasnelli, E., Collett, T. S., & de Ibarra, N. H. (2016). Male
bumblebees perform learning flights on leaving a flower but not when leaving
their nest. *The Journal of Experimental Biology*, *220*, 930–937.
doi:10.1242/jeb.151126
Robertson, C. (1925). Heterotropic Bees. *Ecology*, *6*(4), 412–436. Retrieved from
http://www.jstor.org/stable/1941781
Rossi, B. H., Nonacs, P., & Pitts-Singer, T. L. (2010). Sexual harassment by
males reduces female fecundity in the alfalfa leafcutting bee, Megachile
rotundata. *Animal Behaviour*, *79*(1), 165–171.
doi:10.1016/j.anbehav.2009.10.023
Roulston, T. H., & Goodell, K. (2011). The role of resources and risks in
regulating wild bee populations. *Annual Review of Entomology*, *56*, 293–
312. doi:10.1146/annurev-ento-120709-144802
Rundlöf, M., Persson, A. S., Smith, H. G., & Bommarco, R. (2014). Late-season
mass-flowering red clover increases bumble bee queen and male densities.
Biological Conservation, *172*, 138–145. doi:10.1016/j.biocon.2014.02.027
Rusterholtz, H.-P., & Erhardt, A. (2000). Can nectar properties explain sex-
specific flower preferences in the Adonis Blue butterly Lysandra bellargus?
Ecological Entomology, *25*, 81–90.
Senapathi, D., Goddard, M. A., Kunin, W. E., & Baldock, K. C. R. (2016).
Landscape impacts on pollinator communities in temperate systems:
evidence and knowledge gaps. *Functional Ecology*. doi:10.1111/1365-
2435.12809
Seth, M. (2017). How to implement credible 95% interval for median odds ratio
using JAGS? Retrieved May 9, 2018, from
https://stats.stackexchange.com/q/318912
Somanathan, H., Borges, R. M., Warrant, E. J., & Kelber, A. (2017). Visual
adaptations for mate detection in the male carpenter bee Xylocopa
tenuiscapa. *PLoS ONE*, *12*(1). doi:10.1371/journal.pone.0168452
Spiesman, B. J., Bennett, A., Isaacs, R., & Gratton, C. (2017). Bumble bee
colony growth and reproduction depend on local flower dominance and
natural habitat area in the surrounding landscape. *Biological Conservation*,
206, 217–223. doi:10.1016/j.biocon.2016.12.008
Straub, L., Villamar-Bouza, L., Bruckner, S., Chantawannakul, P., Gauthier, L.,
Khongphinitbunjong, K., … Williams, G. R. (2016). Neonicotinoid

insecticides can serve as inadvertent insect contraceptives. *Proceedings of*
the Royal Society - B, 283(20160506). doi:10.1098/rspb.2016.0506

Streinzer, M., Kelber, C., Pfabigan, S., Kleineidam, C. J., & Spaethe, J. (2013).
Sexual dimorphism in the olfactory system of a solitary and a eusocial bee
species. *Journal of Comparative Neurology, 521*(12), 2742–2755.
doi:10.1002/cne.23312

Sutter, L., Jeanneret, P., Bartual, A. M., Bocci, G., & Albrecht, M. (2017).
Enhancing plant diversity in agricultural landscapes promotes both rare bees
and dominant crop-pollinating bees through complementary increase in key
floral resources. *Journal of Applied Ecology, 54*(6), 1856–1864.
doi:10.1111/1365-2664.12907

Temeles, E. J., Miller, J. S., & Rifkin, J. L. (2010). Evolution of sexual dimorphism
in bill size and shape of hermit hummingbirds (Phaethornithinae): a role for
ecological causation. *Philosophical Transactions of the Royal Society B,*
365(1543), 1053–1063. doi:10.1098/rstb.2009.0284

Thomson, J. D., & Chittka, L. (2001). Pollinator individuality: when does it matter?
In L. Chittka & J. D. Thomson (Eds.), *Cognitive Ecology of Pollination:*
Animal Behaviour and Floral Evolution (pp. 191–213). New York: Cambridge
University Press. doi:10.1017/CBO9780511542268.011

Tonietto, R. K., & Larkin, D. J. (2018). Habitat restoration benefits wild bees: A
meta-analysis. *Journal of Applied Ecology, 55*(2), 582–590.
doi:10.1111/1365-2664.13012

Tur, C., Vigalondo, B., Trøjelsgaard, K., Olesen, J. M., & Traveset, A. (2014).
Downscaling pollen-transport networks to the level of individuals. *Journal of*
Animal Ecology, 83(1), 306–317. doi:10.1111/1365-2656.12130

Ulrich, Y., Perrin, N., & Chapuisat, M. (2009). Flexible social organization and
high incidence of drifting in the sweat bee, Halictus scabiosae. *Molecular*
Ecology, 18(8), 1791–1800. doi:10.1111/j.1365-294X.2009.04154.x

Williams, N. M., & Lonsdorf, E. V. (2018). Selecting cost-effective plant mixes to
support pollinators. *Biological Conservation, 217*(April 2017), 195–202.
doi:10.1016/j.biocon.2017.10.032

Willmer, P. G., & Stone, G. N. (2004). Behavioral, ecoclogical, and physiological
determinants of the activity patterns of bees. *Advances in the Study of*
Behavior, 34(Section II), 47–466.

Figure 1. The sex ratio (M:F) of flower-visiting bees varies across flower species.
Each red point represents a flower species, the first three letters of the Latin
genus and species names for the flower species label each point. The x-axis is
the number of bees collected from that species, the y-axis is the ratio of male to
female bees collected from the flower. Flower species that received >19 visits are
plotted (n=54). Blue lines are smooth fits to the 97.5th and 2.5th percentiles of the
binomial distribution given by the observed ratio of males to females in our
overall dataset (M/F=0.22; i.e. M/(M+F)=0.18). This distribution represents our
expectation for random variation in sex ratio across flower species, if the sex
ratio of flower-visiting bees is independent of flower species identity (male and
female bees exhibit the same floral preferences), and remains nearly constant
across time and space.
693
694

Figure 2: Flower visit patterns of male and female bees of the same species differed significantly. Red points are observed Morisita-Horn dissimilarities between flower communities visited by all male and all female bees of a particular species across all sites and sampling rounds. Black points are the mean dissimilarity (gray bars, 95% CI) from a permutation-based null model that randomly shuffles the sex associated with each visit record, maintaining the total number of males, females, and overall combined visits to each floral species.

Figure 3. The diets of male and female bees of the same species can be as dissimilar as the diets of females of two different bee species. Dissimilarities in this figure are the observed statistic minus, for each pairwise comparison, the mean dissimilarity in the null model. Each panel focuses on a bee species (panel name) and shows: above the label "sex", observed diet dissimilarity between male and female bees of the focal species, minus the average null dissimilarity resulting from randomly permuting the sex identity of each visit record; above the label "sp", observed diet dissimilarity between female bees of the focal species and each other bee species, minus the average null dissimilarity resulting from randomly permuting the species identity of each visit record.

718
Figure 4. Flower species, along with bee species, predicts the sex of visiting
bees, which suggests floral preferences differ between male and female bees.
Flower species is an important predictor of bee sex even after accounting for
phenology (seasonal model). For each term ("bee"= bee species, "flower"=flower
species, "round"=sampling round) in each model, the median odds ratio (+/- 95%
bootstrapped credible interval) indicates the expected difference in odds that a
flower-visiting bee is male between two levels. For example, a median odds ratio
of 3.7 for the flower species term means the odds of a visitor being male are
expected to differ by a factor of 3.7 between two randomly selected species of
flower.

Figure 5. Male bee preferences for and against flower species vary across flower species. Each point is the conditional mode of the random effects prediction (the random-effects analog to an estimate), for a flower species that received at least 20 visits, on the logit scale. Zero represents the odds of a visitor being male on a random flower, and -2 or 2 indicates a ~7 fold decrease or increase in those odds, given flower species identity. Error bars are the square root of the conditional variances on the conditional mode * 1.96, and can be interpreted as the expected range in which the random effect for a particular flower truly lies, analogous to 1.96 times the standard error of the mean for a fixed effect.

742
Figure S1: Sampling scheme. (a) The six study sites in central New Jersey,
USA. (b) Schematic sampling diagram (not to scale). One observer walked
parallel 2m transects covering the entire sampling area. Each 30-minute
sampling bout resumed where the previous one left off; observers typically
covered the entire meadow once over a 3-day sampling round. (c) The
southwestern-most site in peak bloom.

749
750
751

Figure S2. Schematic cartoon of our simulation for the dissimilarity values
associated with our null hypothesis that diets of male and female bees do not
differ. (a) Each collection record for each bee species associates the sex of an
individual bee to the flower species from which it was collected. (b) To compute
the dissimilarity between males and females, we compare all visits to each flower
species from males (purple vector) to all visits to each flower species from
females (green vector). (c) The Morisita-Horn index summarizes the differences
between the two vectors as a value between 0 (identical) and 1 (maximally
dissimilar). (d) For our null model, we shuffle the sex column from our
observation table. (e) This produces two null vectors. The row and column sums
for the matrices in (b) and (c) are identical, but the elements can differ. (f) For our
null model, we compute the dissimilarity between the null vectors. We repeated
steps d-f 9999 times to generate confidence intervals for the null hypothesis that
the sex of a visiting bee is unrelated to the flower species it is collected from.
When comparing the flower species visited by different species of bee, we
conducted an analysis identical except that rather than comparing two sexes of
the same species, we compared two species of the same sex (i.e. exchanging
"sex" and "species" throughout figure S1).

Figure S3. Effect size for diet dissimilarity is independent of sample size, while standardized effect is strongly driven by the number of individuals of the sex with the fewest records. a) Observed Morisita-Horn dissimilarity in flower communities visited by male and female bees of a single species, minus average null dissimilarity vs. the number of records for the less frequently observed sex. b) Observed minus null dissimilarity in composition of flowers visited by male and female bees of a single species, scaled by the variation in the null model, versus the number of records for the less frequently observed sex.

summed model

seasonal model

 Figure S4. Binned residual plots for each model show minor violation of the additivity assumption. Residuals and predicted values on the probability scale.

Figure S5. Seasonal model predictions are consistent with the hypothesis that male bees avoid flower species that do not produce nectar, relative to females. Each point is the random effect prediction (change in odds that a bee visiting that flower is male) for a flower species.

Figure S6. Dissimilarity in flower communities visited by male and female bees arise due to complementarity in addition to nestedness patterns. For each bee species, the proportion of male (blue) and female (red) visits to each flower species that received >10% of at least one sex's visits are pictured. Due to omitted flower species, bars may sum to <1.

798 **Table S1.** Model convergence confirmed based on similar parameter estimates across fitting routines. For each model,
799 the estimate for each term is given for each of 6 fitting algorithms in the R package lme4. Subsequent analyses used
800 parameter estimates in yellow, in both cases tied for the highest estimated likelihood with other very similar fits.

term	model	bobyqa	Nelder_Mead	nlminbw	optimx.L-BFGS-B	nloptwrap.NLOPT_LN_NELDERMEAD	nloptwrap.NLOPT_LN_BOBYQA
intercept	summed	-2.435	-2.435	-2.435	-2.435	-1.753	-1.753
bee species	summed	2.043	2.042	2.043	2.042	2.112	2.112
flower species	summed	1.403	1.402	1.403	1.402	1.397	1.397
site	summed	0.000	0.000	0.000	0.000	0.000	0.000
bee species:flower species	summed	1.209	1.209	1.209	1.209	1.193	1.193
site:bee species	summed	0.624	0.624	0.624	0.625	0.616	0.616
site:flower species	summed	0.606	0.606	0.606	0.605	0.619	0.619
intercept	seasonal	-2.379	-2.470	-2.379	-2.379	-2.449	-2.449
bee species	seasonal	2.091	2.216	2.090	2.091	2.132	2.132
flower species	seasonal	1.251	1.291	1.251	1.251	1.267	1.267
site	seasonal	0.000	0.020	0.000	0.000	0.003	0.003
bee species:flower species	seasonal	1.095	1.120	1.095	1.095	1.101	1.101
site:bee species	seasonal	0.465	0.473	0.465	0.465	0.466	0.466
site:flower species	seasonal	0.353	0.349	0.353	0.353	0.354	0.354
sampling round	seasonal	0.375	0.400	0.375	0.375	0.365	0.365
sampling round:bee species	seasonal	0.831	0.840	0.831	0.831	0.836	0.836
sampling round:flower species	seasonal	0.000	0.006	0.000	0.000	0.006	0.006
sampling round:site	seasonal	0.288	0.298	0.288	0.288	0.289	0.289
sampling round:site:bee species	seasonal	0.596	0.611	0.596	0.596	0.598	0.598
sampling round:site:flower species	seasonal	0.281	0.294	0.281	0.281	0.282	0.282

805 **Table S2.** Bee species with number of female and male specimens collected.
806 *This table will be removed from the final submission when data are deposited on*
807 *Dryad.*

family	genus	species	females	males
Andrenidae	*Andrena*	*brevipalpis*	1	0
Andrenidae	*Andrena*	*carlini*	3	0
Andrenidae	*Andrena*	*commoda*	3	0
Andrenidae	*Andrena*	*cressonii*	16	0
Andrenidae	*Andrena*	*fragilis*	2	0
Andrenidae	*Andrena*	*hippotes*	4	0
Andrenidae	*Andrena*	*imitatrix*	6	0
Andrenidae	*Andrena*	*krigiana*	14	0
Andrenidae	*Andrena*	*nasonii*	13	0
Andrenidae	*Andrena*	*nuda*	2	0
Andrenidae	*Andrena*	*pruni*	6	0
Andrenidae	*Andrena*	*robertsonii*	8	0
Andrenidae	*Andrena*	*rudbeckiae*	8	11
Andrenidae	*Andrena*	*rugosa*	1	0
Andrenidae	*Andrena*	*spiraeana*	1	0
Andrenidae	*Andrena*	*vicina*	6	0
Andrenidae	*Andrena*	*wilkella*	277	59
Andrenidae	*Andrena*	*wilmattae*	2	0
Andrenidae	*Calliopsis*	*andreniformis*	4	1
Apidae	*Anthophora*	*abrupta*	4	0
Apidae	*Anthophora*	*terminalis*	3	2
Apidae	*Bombus*	*auricomus*	1	0
Apidae	*Bombus*	*bimaculatus*	576	175
Apidae	*Bombus*	*citrinus*	0	5
Apidae	*Bombus*	*fervidus*	18	0
Apidae	*Bombus*	*griseocollis*	681	814
Apidae	*Bombus*	*impatiens*	2358	105
Apidae	*Bombus*	*perplexus*	22	36
Apidae	*Bombus*	*vagans*	14	2
Apidae	*Ceratina*	*calcarata*	1417	133
Apidae	*Ceratina*	*dupla*	151	19
Apidae	*Ceratina*	*mikmaqi*	130	5
Apidae	*Ceratina*	*strenua*	285	13

Apidae	*Melissodes*	*agilis*	0	7
Apidae	*Melissodes*	*bimaculatus*	9	1
Apidae	*Melissodes*	*denticulatus*	7	73
Apidae	*Melissodes*	*desponsus*	1	7
Apidae	*Melissodes*	*subillatus*	31	6
Apidae	*Melissodes*	*trinodis*	1	7
Apidae	*Nomada*	*articulata*	4	0
Apidae	*Nomada*	*bidentate_gr*	7	0
Apidae	*Nomada*	*erigeronis*	1	0
Apidae	*Nomada*	*lehighensis*	1	0
Apidae	*Nomada*	*maculata*	2	0
Apidae	*Nomada*	*pygmaea*	15	0
Apidae	*Ptilothrix*	*bombiformis*	0	1
Apidae	*Triepeolus*	*cressonii*	0	1
Apidae	*Triepeolus*	*eliseae*	1	0
Apidae	*Triepeolus*	*remigatus*	1	0
Apidae	*Xylocopa*	*virginica*	137	13
Colletidae	*Hylaeus*	*affinis_modestus*	1372	363
Colletidae	*Hylaeus*	*fedorica*	1	0
Colletidae	*Hylaeus*	*leptocephalus*	1	3
Colletidae	*Hylaeus*	*mesillae*	575	173
Halictidae	*Agapostemon*	*sericeus*	5	5
Halictidae	*Agapostemon*	*virescens*	203	75
Halictidae	*Augochlora*	*pura*	1035	376
Halictidae	*Augochlorella*	*aurata*	397	39
Halictidae	*Augochlorella*	*persimilis*	434	116
Halictidae	*Augochloropsis*	*metallica*	121	40
Halictidae	*Dufourea*	*novaeangliae*	0	1
Halictidae	*Halictus*	*confusus*	174	35
Halictidae	*Halictus*	*ligatus*	2432	160
Halictidae	*Halictus*	*parallelus*	6	18
Halictidae	*Halictus*	*rubicundus*	31	19
Halictidae	*Lasioglossum*	*abanci*	6	0
Halictidae	*Lasioglossum*	*admirandum*	15	0
Halictidae	*Lasioglossum*	*anomalum*	17	0
Halictidae	*Lasioglossum*	*atwoodi*	7	1
Halictidae	*Lasioglossum*	*birkmanni*	1	0
Halictidae	*Lasioglossum*	*bruneri*	6	4
Halictidae	*Lasioglossum*	*callidum*	54	0

Halictidae	Lasioglossum	cattellae	14	4
Halictidae	Lasioglossum	coeruleum	2	0
Halictidae	Lasioglossum	coreopsis	1	0
Halictidae	Lasioglossum	coriaceum	14	0
Halictidae	Lasioglossum	cressonii	16	5
Halictidae	Lasioglossum	ellisiae	0	3
Halictidae	Lasioglossum	ephialtum	1	0
Halictidae	Lasioglossum	foxii	2	2
Halictidae	Lasioglossum	fuscipenne	9	0
Halictidae	Lasioglossum	gotham	74	2
Halictidae	Lasioglossum	hitchensi_weemsi	152	27
Halictidae	Lasioglossum	illinoense	70	7
Halictidae	Lasioglossum	imitatum	462	15
Halictidae	Lasioglossum	leucocomum	2	0
Halictidae	Lasioglossum	leucozonium	2	0
Halictidae	Lasioglossum	nigroviride	2	0
Halictidae	Lasioglossum	oblongum	4	2
Halictidae	Lasioglossum	obscurum	7	1
Halictidae	Lasioglossum	oceanicum	104	23
Halictidae	Lasioglossum	oenotherae	1	0
Halictidae	Lasioglossum	paradmirandum	50	0
Halictidae	Lasioglossum	pectorale	3	0
Halictidae	Lasioglossum	pilosum	2	0
Halictidae	Lasioglossum	platyparium	2	3
Halictidae	Lasioglossum	rozeni	15	11
Halictidae	Lasioglossum	smilacinae	4	0
Halictidae	Lasioglossum	subviridatum	5	1
Halictidae	Lasioglossum	tegulare	31	2
Halictidae	Lasioglossum	trigeminum	44	0
Halictidae	Lasioglossum	truncatum	2	0
Halictidae	Lasioglossum	versatum	681	93
Halictidae	Lasioglossum	viridatum	11	2
Halictidae	Lasioglossum	zephyrum	12	1
Halictidae	Sphecodes	atlantis	0	1
Halictidae	Sphecodes	dichrous	3	5
Halictidae	Sphecodes	heraclei	10	5
Megachilidae	Anthidiellum	notatum	4	1
Megachilidae	Anthidium	manicatum	7	8
Megachilidae	Anthidium	oblongatum	18	19

Megachilidae	*Coelioxys*	*alternatus*	1	2
Megachilidae	*Coelioxys*	*banksi*	1	0
Megachilidae	*Coelioxys*	*germanus*	0	1
Megachilidae	*Coelioxys*	*hunteri*	0	1
Megachilidae	*Coelioxys*	*modestus*	0	1
Megachilidae	*Coelioxys*	*obtusiventris*	1	0
Megachilidae	*Coelioxys*	*octodentatus*	1	1
Megachilidae	*Coelioxys*	*porterae*	0	1
Megachilidae	*Coelioxys*	*sayi*	2	6
Megachilidae	*Heriades*	*carinatus*	31	2
Megachilidae	*Heriades*	*leavitti*	1	6
Megachilidae	*Heriades*	*variolosus*	10	0
Megachilidae	*Hoplitis*	*pilosifrons*	46	1
Megachilidae	*Hoplitis*	*producta*	8	0
Megachilidae	*Hoplitis*	*spoliata*	2	1
Megachilidae	*Lithurgus*	*chrysurus*	0	6
Megachilidae	*Megachile*	*brevis*	25	3
Megachilidae	*Megachile*	*campanulae*	6	18
Megachilidae	*Megachile*	*exilis*	11	29
Megachilidae	*Megachile*	*frugalis*	26	6
Megachilidae	*Megachile*	*gemula*	4	2
Megachilidae	*Megachile*	*georgica*	1	0
Megachilidae	*Megachile*	*inimica*	4	0
Megachilidae	*Megachile*	*integra*	1	0
Megachilidae	*Megachile*	*melanophaea*	0	1
Megachilidae	*Megachile*	*mendica*	21	56
Megachilidae	*Megachile*	*montivaga*	15	9
Megachilidae	*Megachile*	*petulans*	0	2
Megachilidae	*Megachile*	*pugnata*	2	3
Megachilidae	*Megachile*	*rotundata*	11	8
Megachilidae	*Megachile*	*sculpturalis*	17	32
Megachilidae	*Megachile*	*xylocopoides*	2	1
Megachilidae	*Osmia*	*albiventris*	3	0
Megachilidae	*Osmia*	*atriventris*	9	0
Megachilidae	*Osmia*	*bucephala*	21	0
Megachilidae	*Osmia*	*distincta*	7	0
Megachilidae	*Osmia*	*georgica*	5	0
Megachilidae	*Osmia*	*pumila*	30	0
Megachilidae	*Pseudoanthidium*	*nanum*	0	1

family	genus	species		
Megachilidae	*Stelis*	*lateralis*	1	0
Megachilidae	*Stelis*	*louisae*	1	2

809 **Table S3.** Number of male and female visitors to each plant species, and bias
810 towards attracting male bee visitors. This bias is the random effect prediction
811 from the seasonal model, which indicates the change in log(odds) that a visiting
812 bee is male when the species of flower it visits is given; greater values indicate
813 male bias.

family	genus	species	female visitors	male visitors	random effect
Verbenaceae	*Verbena*	*urticifolia*	58	71	2.12
Asteraceae	*Erechtites*	*hieraciifolius*	130	203	1.88
Fabaceae	*Senna*	*hebecarpa*	5	5	1.68
Phytolaccaceae	*Phytolacca*	*americana*	108	74	1.58
Asteraceae	*Euthamia*	*graminifolia*	50	18	1.47
Asteraceae	*Eutrochium*	*maculatum*	461	166	1.37
Fabaceae	*Melilotus*	*officinalis*	41	21	1.31
Lamiaceae	*Nepeta*	*cataria*	99	81	1.27
Lamiaceae	*Monarda*	*punctata*	0	1	1.24
Campanulaceae	*Lobelia*	*inflata*	12	3	1.15
Asteraceae	*Liatris*	*spicata*	186	127	1.14
Asteraceae	*Solidago*	*juncea*	636	77	1.11
Asteraceae	*Conyza*	*canadensis*	32	10	1.10
Polygonaceae	*Fallopia*	*convolvulus*	3	4	1.04
Asteraceae	*Erigeron*	*strigosus*	712	119	1.04
Asclepidaceae	*Asclepias*	*syriaca*	28	89	0.93
Verbenaceae	*Verbena*	*hastata*	8	3	0.93
Lamiaceae	*Pycnanthemum*	*verticillatum*	5	8	0.89
Asteraceae	*Cirsium*	*arvense*	351	96	0.86
Asteraceae	*Solidago*	*canadensis*	8	2	0.82
Lamiaceae	*Pycnanthemum*	*muticum*	398	59	0.81
Asteraceae	*Heliopsis*	*helianthoides*	186	49	0.76
Apocynaceae	*Apocynum*	*cannabinum*	283	92	0.75
Fabaceae	*Trifolium*	*hybridum*	11	7	0.74
Asteraceae	*Rudbeckia*	*hirta*	1174	189	0.65
Asteraceae	*Solidago*	*gigantea*	107	9	0.61
Lamiaceae	*Pycnanthemum*	*tenuifolium*	1113	421	0.61
Rosaceae	*Drymocallis*	*arguta*	22	8	0.60
Fabaceae	*Melilotus*	*albus*	20	9	0.57

Cornaceae	*Swida*	*racemosa*	12	1	0.55
Apiaceae	*Daucus*	*carota*	1778	350	0.53
Asteraceae	*Helianthus*	*strumosus*	1	1	0.52
Asteraceae	*Cichorium*	*intybus*	104	10	0.52
Verbenaceae	*Verbena*	*simplex*	11	3	0.48
Asteraceae	*Symphyotrichum*	*novae-angliae*	25	6	0.45
Asclepidaceae	*Asclepias*	*tuberosa*	114	26	0.42
Asteraceae	*Cirsium*	*discolor*	5	2	0.35
Asteraceae	*Echinacea*	*purpurea*	86	45	0.31
Lamiaceae	*Prunella*	*vulgaris*	17	5	0.30
Asteraceae	*Erigeron*	*annuus*	26	3	0.27
Asteraceae	*Centuarea*	*stoebe*	321	50	0.26
Onagraceae	*Oenothera*	*fruticosa*	2	1	0.22
Polygonaceae	*Persicaria*	*setacea*	3	1	0.16
Asteraceae	*Cirsium*	*vulgare*	112	16	0.12
Fabaceae	*Trifolium*	*campestre*	365	39	0.11
Asteraceae	*Ratibida*	*pinnata*	539	121	0.01
Asteraceae	*Achillea*	*millefolium*	472	36	0.00
Asteraceae	*Bidens*	*trichosperma*	1	0	-0.01
Asteraceae	*Solidago*	*rugosa*	1	0	-0.01
Lythraceae	*Lythrum*	*salicaria*	364	38	-0.02
Rosaceae	*Rubus*	*flagellaris*	1	0	-0.03
Fabaceae	*Trifolium*	*aureum*	1	0	-0.03
Gentianaceae	*Sabatia*	*angularis*	1	0	-0.03
Campanulaceae	*Lobelia*	*siphilitica*	2	0	-0.03
Asteraceae	*Vernonia*	*noveboracensis*	52	37	-0.05
Fabaceae	*Vicia*	*tetrasperma*	1	0	-0.05
Apiaceae	*Sanicula*	*canadensis*	1	0	-0.05
Asteraceae	*Doellingeria*	*umbellata*	1	0	-0.05
Ranunculaceae	*Ranunculus*	*hispidus*	1	0	-0.06
Asteraceae	*Coreopsis*	*tinctoria*	1	0	-0.07
Poaceae	*Sorghastrum*	*nutans*	1	0	-0.09
Lamiaceae	*Teucrium*	*canadense*	3	0	-0.10
Brassicaceae	*Barbarea*	*vulgaris*	3	0	-0.11
Lamiaceae	*Monarda*	*fistulosa*	1397	400	-0.13
Loniceraceae	*Lonicera*	*japonica*	1	0	-0.13
Fabaceae	*Desmodium*	*paniculatum*	6	1	-0.14
Onagraceae	*Oenothera*	*biennis*	2	0	-0.14
Asteraceae	*Hieracium*	*pilosella*	2	0	-0.15

Hypericaceae	*Hypericum*	*punctatum*	1	0	-0.18
Poaceae	*Glyceria*	*grandis*	1	0	-0.19
Alliaceae	*Allium*	*vineale*	3	0	-0.22
Apiaceae	*Eryngium*	*yuccifolium*	2	0	-0.23
Fabaceae	*Lotus*	*corniculatus*	142	33	-0.23
Asteraceae	*Crepis*	*capillaris*	6	0	-0.24
Rosaceae	*Rubus*	*pensilvanicus*	7	0	-0.24
Oxalidaceae	*Oxalis*	*stricta*	8	0	-0.25
Cornaceae	*Swida*	*amomum*	4	0	-0.25
Asteraceae	*Lactuca*	*serriola*	4	0	-0.26
Rosaceae	*Rosa*	*multiflora*	5	0	-0.26
Asteraceae	*Gaillardia*	*aristata*	4	0	-0.27
Caryophyllaceae	*Dianthus*	*armeria*	3	0	-0.28
Asteraceae	*Leucanthemum*	*vulgare*	405	20	-0.35
Iridaceae	*Sisyrinchium*	*angustifolium*	9	0	-0.35
Asteraceae	*Carduus*	*nutans*	1	0	-0.37
Rubiaceae	*Galium*	*mollugo*	4	0	-0.37
Polygonaceae	*Persicaria*	*pensylvanica*	4	0	-0.39
Asteraceae	*Helianthus*	*angustifolius*	11	0	-0.41
Solanaceae	*Solanum*	*carolinense*	14	0	-0.52
Convulvulaceae	*Calystegia*	*silvatica*	5	0	-0.54
Asteraceae	*Krigia*	*biflora*	19	0	-0.54
Fabaceae	*Baptisia*	*tinctoria*	19	5	-0.57
Asteraceae	*Solidago*	*altissima*	8	0	-0.59
Rosaceae	*Potentilla*	*recta*	56	1	-0.63
Fabaceae	*Securigera*	*varia*	38	1	-0.65
Scrophulariaceae	*Verbascum*	*blattaria*	15	0	-0.72
Asclepidaceae	*Asclepias*	*incarnata*	7	0	-0.74
Commelinaceae	*Tradescantia*	*ohiensis*	34	1	-0.81
Scrophulariaceae	*Penstemon*	*hirsutus*	36	0	-0.82
Fabaceae	*Trifolium*	*repens*	130	6	-0.85
Scrophulariaceae	*Penstemon*	*digitalis*	862	48	-0.85
Lamiaceae	*Clinopodium*	*vulgare*	64	3	-0.85
Asteraceae	*Coreopsis*	*lanceolata*	21	0	-0.86
Fabaceae	*Trifolium*	*pratense*	192	20	-0.87
Scrophulariaceae	*Verbascum*	*thapsus*	129	1	-1.08
Hypericaceae	*Hypericum*	*perforatum*	223	10	-1.09
Rosaceae	*Rosa*	*carolina*	71	1	-1.41
Fabaceae	*Chamaecrista*	*fasciculata*	246	2	-1.51

| Scrophulariaceae | *Linaria* | *vulgaris* | 275 | 4 | -1.63 |
| Plantaginaceae | *Plantago* | *lanceolata* | 147 | 0 | -2.00 |