Pacific geoduck (*Panopea generosa)* resilience to natural pH variation

Laura H. Spencer[a], Micah Horwith[b], Alexander T. Lowe[c], Yaamini R. Venkataraman[a], Emma Timmins-Schiffman[d], Brook L. Nunn[d], Steven B. Roberts[a]

[a]University of Washington, School of Aquatic and Fishery Sciences, 1122 NE Boat St, Seattle, WA 98105, United States
[b]Washington State Department of Natural Resources, 1111 Washington St SE, MS 47027, Olympia, WA 98504, United States
[c]University of Washington, Biological Sciences, 24 Kincaid Hall, Seattle, WA 98105, United States
[d]University of Washington, Genome Sciences, William H. Foege Hall, 3720 15th Ave NE, Seattle, WA 98195, United States

Corresponding Author: Steven B. Roberts, sr320@uw.edu

Abstract

Pacific geoduck aquaculture is a growing industry, however, little is known about how geoduck respond to varying environmental conditions, or how the industry will fare under projected climate conditions. To understand how geoduck production may be impacted by low pH associated with ocean acidification, multi-faceted environmental heterogeneity needs to be included to understand species and community responses. In this study, eelgrass habitats and environmental heterogeneity across four estuarine bays were leveraged to examine low pH effects on geoduck under different natural regimes, using targeted proteomics to assess physiology. Juvenile geoduck were deployed in eelgrass and adjacent unvegetated habitats for 30 days while pH, temperature, dissolved oxygen, and salinity were monitored. Across the four bays, pH was lower in unvegetated habitats compared to eelgrass habitats. However this did not impact geoduck growth, survival, or proteomic abundance patterns in gill tissue. Temperature and dissolved oxygen differences across all locations corresponded to differences in growth and targeted protein abundance patterns. Specifically, three protein abundance levels (trifunctional-enzyme β-subunit, puromycin-sensitive aminopeptidase, and heat shock protein 90-α) and shell growth positively correlated with dissolved oxygen variability and inversely correlated with mean temperature. These results demonstrate that geoduck may be resilient to low pH in a natural setting, but other abiotic factors (i.e. temperature, dissolved oxygen variability) may have a greater influence on geoduck physiology. In addition this study contributes to the understanding of how eelgrass patches influences water chemistry.

Key words

Aquaculture, comparative physiology, ocean acidification, Panopea generosa, proteomics

Introduction

The Pacific geoduck, *Panopea generosa,* is native to the North American Pacific Coast and is a burgeoning aquaculture species with strong overseas demand as a luxury commodity (Coan et al. 2000; Shamshak and King 2015; Vadopalas et al. 2010). As the largest burrowing clam in the world, cultured geoduck reach upwards of 180mm and are harvested after growing approximately 6-7 years in sub- or intertidal sediment (Vadopalas et al. 2015; Washington DNR website 2017; Washington Sea Grant 2013). The long grow-out period and high per-animal value highlights the importance of site selection for farmers to maximize investment; however, there remains a paucity of data on the optimal environmental conditions for geoduck aquaculture.

As marine calcifiers, geoduck may be vulnerable to ocean acidification due to their reliance on calcite and aragonite (forms of calcium carbonate) for shell secretion (Orr et al. 2005; Weiss et al. 2002), both of which become less biologically available as seawater pH declines with pCO_2 enrichment (Feely et al. 2008). While there are no ocean acidification studies on *Panopea* clams to date, a growing body of research on marine calcifiers generally indicates that projected low pH will shift organisms' physiology to the detriment of species-wide

abundances and distributions (Pörtner 2008; Pörtner and Farrell 2008). However, broad
generalizations of how ocean acidification affects calcifiers are few due to varying pH sensitivity
between taxa (Gazeau et al. 2007; Ries et al. 2009) and life stage (Kurihara 2008; Kroeker et al.
2010). For example, in the deeply studied oyster genus *Crassostrea*, Miller et al. (2009) found
that larvae of two species varied in their response to elevated pCO_2, as calcification rates were
significantly reduced in the Eastern oyster (*C. virginica*), but the Suminoe oyster (*C. ariakesnsis*)
showed no negative response. Thus, lessons learned from other bivalve species cannot directly
be applied to geoduck.
The effect of low pH on cultured geoduck needs to be explored to help the aquaculture
industry make informed site selection, selective breeding, and investment decisions. For
practical application, geoduck ocean acidification studies should best replicate the natural
environment in which they are grown. Ninety percent of global production occurs in the Puget
Sound estuary of Washington State, where environmental drivers vary between subbasin,
season, and diurnal cycle (Moore et al. 2008; Shamshak and King 2015). This habitat
heterogeneity exposes geoduck to a variety of secondary stressors when outplanted. Similarly,
there is substantial evidence that low pH is not occurring in isolation, but rather in conjunction
with changes in other environmental drivers such as temperature (meta-analyses: Byrne and
Przeslawski 2013; Harvey et al. 2013; Kroeker et al. 2013), dissolved oxygen (Gobler et al.
2014), and salinity (Przeslawski et al. 2015). Thus, single-stressor studies are limited in their
predictive capacity of response to broad scale environmental change. For example, an additive,
negative effect of elevated pCO_2 and temperature was observed in juvenile giant fluted clam
survival (*Tridacna squamosa*) (Watson et al. 2012). Another consideration is the incorporation of
naturally-occurring diurnal pH variability into ocean acidification studies, as variable pH can
have differing effects on marine calcifiers compared to persistent low pH (Review, Boyd et al.
2016). Porcelain crabs, for example, exposed to diurnally variable pH and temperature
conditions demonstrated significantly slower metabolism than when crabs were exposed to less
variability, or to temperature or pH variability alone (Paganini et al. 2014).
To best predict the effect of ocean acidification on geoduck aquaculture, this project
deployed geoduck in variable environmental conditions and leveraged the natural pH
differences between eelgrass and unvegetated habitats in Washington State estuaries. Ocean
acidification studies are increasingly exploiting naturally low pH systems to monitor the
environmental heterogeneity alongside test organisms, in hydrothermal vents (Tunnicliffe et al.
2009; Kerrison et al. 2011), shallow CO_2 seeps (Duquette et al. 2017), coastal upwelling regions
and eutrophic estuaries (Howarth et al. 2011; Thomsen et al. 2013). Compared to controlled
laboratory studies, these deployment studies can uniquely incorporate natural ranges and daily
cycles in air exposure, temperature, pH, dissolved oxygen, salinity, and food availability. For
instance, Ringwood and Keppler (2002) deployed hard clams (*Mercenaria mercenaria*) in the
Charleston Harbor estuary in South Carolina while collecting physical-chemical parameters.
They observed that while salinity was the primary determinant of growth, pH was also important
particularly when salinity was low, and when pH dropped below 7.5, a nuanced finding that is
more likely to be captured in a natural environment.
Estuaries along the United States Pacific Coast are ideal, natural mesocosms for
examining the effect of ocean acidification on commercially vital calcifiers, as they contain
dense macroalgae beds (Bulthuis 1995), environmental conditions that vary considerably

between subbasins (Banas et al. 2004; Moore et al. 2008), and have rich communities of native
and cultured shellfish (Dethier et al. 2006; Miller et al. 2009; Washington Sea Grant 2015).
Furthermore, coastal estuaries have already shifted towards lower pH and warmer temperature
averages, and are projected to continue along this trend (Abatzoglou et al. 2013; Busch et al.
2013; Doney et al. 2007; Feely et al. 2012, 2010, 2008; Mote and Salathé 2010). The buffering
capacity of macroalgae (seagrass meadows, kelp forests) allows for block-designed
experiments to examine the effect of pH, while controlling for varying background environments
and maintaining diurnal fluctuations (Middelboe and Hansen 2007; Palacios and Zimmerman
2007; Wahl et al. 2018).
In order to better inform geoduck aquaculture practices, we set out to examine how low
pH and other natural variation in environmental conditions influence geoduck growth and
physiology, using native eelgrass (*Zostera marina*) as a primary determinant of water chemistry.
Physiology was evaluated with a unique two-phase proteomics approach using Selected
Reaction Monitoring, with targets identified using Data Independent Acquisition, and selected
based on prior environmental stress response studies.
Ocean acidification contributes to an elevation of reactive oxygen species in marine
invertebrates (Tomanek 2015). Reactive oxygen species (ROS), or free-radicals, result in
oxidative stress and in addition to low pH, higher ROS levels are associated with other
environmental stressors including temperature, oxygen variability, salinity, and heavy metals
(Review, Lushchak, 2011). Upregulation of anti-oxidants such as catalase, peroxiredoxins, and
superoxide dismutase (among others) have consistently been observed in bivalves under low
pH and heat stress (Tomanek et al. 2011; Matozzo et al. 2013; Hu et al. 2015), and under heavy
metal exposure (Giarratano et al. 2014). In addition to ROS response, ocean acidification is
thought to elicit a broader and generic molecular stress response in marine bivalves. Notably,
the inducible heat shock proteins are associated with response to hypercapnia, in addition to
acute heat, inflammation, and heavy metals, as they act as chaperones to recognize and bind to
unfolded or improperly folded proteins (Bozaykut et al. 2014). Induction of HSP90, for example,
has been universally observed thus far in bivalve species (Fabbri et al. 2008). Metabolic
function is also altered under low pH, hypoxia, and salinity stress, generally shifting to anaerobic
metabolism to minimize the mitochondrial ROS production associated with aerobic metabolism
(Tomanek 2014).
In addition to antioxidant, metabolic, and general stress-response proteins, this study
targeted proteins involved in mitotic growth, detoxification, acid-base balance, and ion regulation
(Table 2), all quantified simultaneously to characterize the physiological response in the Pacific
geoduck under variable pH environments. This novel application demonstrates the advances in
proteomic research and the potential it has to improve aquaculture production.

Methods

Experimental Design

Panopea generosa juveniles (14.0 ± 0.85 mm) were used in this experiment. Animals
were of the same cohort, hatched from broodstock harvested from Puget Sound in Washington
State, and reared in a commercial facility in Dabob Bay, WA in controlled conditions (18°C,
salinity of 30ppt and pH 8.2). Geoduck were out-planted in four bays throughout Western
Washington State from June 21 to July 21, 2016: Fidalgo Bay (FB), Port Gamble Bay (PG), and
Case Inlet (CI) in Puget Sound, and Willapa Bay (WB) located off the southwest Pacific Coast of
Washington (Table 1, Figure 1). All locations were selected based on the criteria that both *Z.*
marina eelgrass beds ("eelgrass"), and unvegetated sediment ("unvegetated") habitats were
present. Clams were placed in 10 cm diameter polymerizing vinyl chloride (PVC) pipes buried in
sediment with 5 cm exposed; this method replicates aquaculture techniques. Five clams were
placed in each of the 3 tubes in both the eelgrass and unvegetated habitat, with a total of 30
clams across 6 tubes per bay. Pipes were covered with a protective mesh exclosure to limit
predation. The replicate structures surrounded and were equidistant to a suite of water quality
sensors capturing pH (Honeywell Durafet II Electrode, in custom-built housing), salinity (via
conductivity, Dataflow Systems Ltd. Odyssey Conductivity and Temperature Logger), dissolved
oxygen (Precision Measurement Engineering MiniDOT Logger), and temperature (via dissolved
oxygen probes). Sensors were modified for submersible, autonomous data collection, and
logged at 10-minute intervals for the duration of the 30-day outplant.

Figure 1: Geoduck juveniles were deployed for 30 days in 2 habitats (eelgrass beds, unvegetated) within 4 bays in Western Washington State

Bay	Eelgrass Habitat	Unvegetated Habitat
Fidalgo Bay	48° 28' 52.8312" N 122° 35' 0.7044" W	48° 28' 54.0876" N 122° 35' 0.708" W
Port Gamble Bay	47° 50' 52.7388" N 122° 34' 58.5084" W	47° 50' 33.6336" N 122° 35' 1.7952" W
Case Inlet	47° 21' 30.3808" N 122° 47' 47.2182" W	47° 21' 28.5721" N 122° 47' 44.7457" W
Willapa Bay	46° 29' 42.288" N 124° 1' 35.472" W	46° 29' 40.124" N 124° 1' 34.0882" W

Table 1: Coordinates for geoduck placement.

Collection and Sampling

Animals were collected during low tide and transferred on wet ice to shore where mortality and size were recorded. Live animals were dissected, and ctenidia tissue was isolated and flash-frozen in an ethanol-dry ice bath. Ctenidia was selected for proteomic analysis due to its direct interaction with the environment, importance in gas and ion regulation, and its implication in environmental stress response (Timmins-Schiffman et al., 2014, Matozzo et al. 2013, Zhang et al. 2015, Thompson et al. 2015). During sampling all instruments were sterilized between samples with bleach then ethanol, and rinsed with nanopure water. Samples were held on dry ice while transported back to the lab and stored at -80ºC.

Environmental and Growth Data

Temperature, pH, salinity and dissolved oxygen data were compared between outplant locations. Outliers for all environmental parameters were removed, as determined using Tukey Interquartile Range (IQR) method (Tukey 1977), excluding data outside the inner fence (1.5*IQR). Tidal charts from WWW Tide/Current Predictor and salinity data (<20ppt) were also used to remove DO and pH data corresponding to periods of tidal exposure. Four probe failures occurred during deployment (Supplemental Table 1, *failed*) and these data were not included in the analysis (pH at Port Gamble-eelgrass, salinity at Port Gamble-unvegetated & Case Inlet-eelgrass, DO in Fidalgo Bay-eelgrass). Salinity data from two additional locations was not reliably measured (Willapa Bay-unvegetated, Fidalgo Bay-unvegetated), so habitat comparisons were not performed for salinity data. For each parameter at each location, daily mean and daily standard deviation time series were calculated. Relative growth for each animal was determined as (L_f-L_i) , where L_f= final geoduck shell length, L_i = average initial geoduck shell length within each exclosure (n=5). Differences in growth and environmental parameters between habitat

187 were compared using 2-way analysis of variance (ANOVA) applied to regression models (value
188 ~ habitat*bay). Bays and ad-hoc regions (north vs. south bays) were tested using 1-way
189 ANOVA. Pairwise comparisons were tested with the t-statistic. Significance for all tests was
190 defined as $P < 0.05$, corrected for multiple comparisons using the Bonferroni correction.
191

Protein Analysis

Protein Preparation

Relative protein abundance was ultimately assessed in a two-phase proteomics approach using
Selected Reaction Monitoring (SRM), with targets identified using Data Independent Acquisition
(DIA). Tissues were prepared separately for DIA and SRM, both following the protocol in
Timmins-Schiffman et al. (2014) with a few exceptions. For DIA, 8 ctenidia tissue samples were
analyzed, one sample from each location and habitat: FB-eelgrass (G048), FB-unvegetated
(G058), PGB-eelgrass (G077), PGB-unvegetated (G068), CI-eelgrass (G010), CI-unvegetated
(G018), WB-eelgrass (G131), WB-unvegetated (G119). For SRM, new ctenidia samples were
examined, 12 individuals per bay (Fidalgo Bay, Port Gamble Bay, Case Inlet, Willapa Bay), with
6 from each habitat (eelgrass, unvegetated) for 48 samples total. Tissue was homogenized with
sterile plastic pestle in 100 µl lysis buffer (50 mM NH_4HCO_3, 6M urea solution) and sonicated
with Sonic Dismembrator (Fisher Scientific, Model 120) at 50% amplitude for ten seconds, three
times. Protein concentration was quantified via Pierce™ BCA Protein Assay Kit (ThermoFisher
Scientific, Waltham, MA USA).
207

Mini-Trypsin Digestion

Aliquots of protein (30 µg for DIA, 100 µg for SRM) were suspended in Lysis Buffer (50 mM
NH_4HCO_3 + 6 M urea solution) to a total volume of 100 µl followed by: 1) a 1 hour incubation at
37°C with 200 mM Tris(2-carboxyethyl)phosphine (2.5µl) and 1.5 M Tris at pH 8.8 (6.6 µl); 2) 1
hour at room temperature in dark with 200 mM iodoacetamide (20 µl); 3) 1 hour at room
temperature with 200 mM diothiothreitol (20 µl); 4) 1 hour at room temperature with 2 ug/µl Lysyl
Endopeptidase (Lys-C, Wako Chemicals) (3.3 µg); 5) overnight at room temperature in 25 mM
NH_4HCO_3 (800 µl) + high pressure liquid chromatography grade methanol (200 µl) + Pierce
Trypsin Protease, MS Grade (1 ug/µl, Thermo Scientific) at 1:30 enzyme:protein ratio (3.3 µg).
Samples were evaporated to near dryness at 4°C using a CentriVap Benchtop Vacuum
Concentrator.
219

Desalting

221 Samples were desalted to isolate peptides using MacroSpin Columns (Nest Group, 50-450 µl,
222 Peptide Protein C18). Peptides were reconstituted in 5% acetonitrile + 0.1% trifluoroacetic acid
223 (TFA)
224

(100 µl), then 10% formic acid (70 µl) was added to achieve pH ≤2. Columns were washed with
60% acetonitrile + 0.1% TFA (Solvent A, 200 µl) four times, then equilibrated with 5%
acetonitrile + 0.1% TFA (Solvent B, 200 µl) three times. Peptides were bound to the column by
running the digest through the column twice, followed by peptide elution with two additions each
of Solvent A (100 ul). Columns were spun for 3 minutes at 3000 rpm on VWR Galaxy 16DH
digital microcentrifuge at each stage. Samples were evaporated to near dryness at 4°C, then
reconstituted in the Final Solvent (3% acetonitrile + 0.1% formic acid) (60 µl for 0.5 µg/µl final
concentration of protein, and 50 µl for 2 µg/µl final concentration for DIA & SRM, respectively).
233
Peptide sample preparation and internal standard
Final mixtures for mass spectrometry included 3.13 fmol/µl Peptide Retention Time Calibration
mixture (PRTC), 0.33 µg/µl and 0.5 µg/µl peptides for DIA and SRM, respectively, in Final
Solvent for 15 µl total volume. To confirm that peptides were quantified correctly in SRM, 10 µg
from 5 randomly selected geoduck peptide samples were pooled, and 8 dilutions were prepared
by combining with oyster peptides at known percentages of total protein content (10%, 13.3%,
20%, 40%, 60%, 80%, 87.7%, 90%) and analyzed alongside other samples.

Data Independent Acquisition

Data acquisition
Data Independent Acquisition (DIA) was performed to assess global protein abundance patterns
and to identify consistently detectable peptides for SRM targets. Eight samples, one per
deployment location, were analyzed in technical duplicates via liquid chromatography tandem
mass spectrometry (LC-MS/MS) with the Thermo Scientific™ Orbitrap Fusion Lumos™
Tribrid™. Prior to sample analysis, the 30 cm analytical column and 3 cm trap were packed in-
house with with C18 beads (Dr. Maisch HPLC, Germany, 0.3 µm). For each sample, 3 µl of
geoduck peptides (1.0 µg) + PRTC peptides was injected and analyzed in MS1 over 400-900
m/z range, in 5 *m/z* overlapping isolation windows from 450-850 *m/z* with 15K resolution in MS2.
Final Solvent blanks were run between each geoduck peptide injection to ensure against
peptide carry-over. Lumos MS/MS method and sequence files are available in the project
repository (Spencer et al. 2019), and data are available via ProteomeXchange with identifier
PXD012266.
255
Protein identification and analysis
Proteins were inferred using an assembled, translated, and annotated *P. generosa* gonad
transcriptome (combined male and female) (Timmins-Schiffman et al. 2017; DOI:
10.17605/OSF.IO/3XF6M). Transcriptome peptides were queried against those detected by the
Lumos MS/MS using PEptide-Centric Analysis (PECAN) (Ting 2017) to create a peptide
spectral library (.blib type file). DIA raw files were first demultiplexed using MSConvert
(ProteoWizard version 3.0.10738, 2017-04-12) (Chambers et al. 2012) with filters set to vendor
centroiding for msLevels [2,3] (--"peakPicking true 1-2"), and optimizing overlapping spectra
("demultiplex optimization-overlap only"). The transcriptome fasta file was tryptic digested *in*
silico in Protein Digestion Simulator (version 2.2.6471.25262), set to Fully Tryptic from 400-6000
fragment mass range, 5 minimum residues allowed, 3 maximum missed cleavages and peak

matching thresholds set to 5 ppm mass tolerance, and 0.05 ppm NET tolerance. Skyline version
3.7.0.11317 (MacLean et al. 2010) automatically selected transition peaks and quantified
peptide abundances using peak area integration. All PRTC peptide peak selections were
manually verified and corrected. Skyline peak selection error rate was calculated by manually
checking chromatograms from 100 proteins across all DIA samples. Auto-selected peaks were
assigned correct or incorrect selection based on transition retention time alignment across
replicates, using PRTC peptides as a reference. Transition peak area, which is assumed to
correlate to peptide transition abundance, was exported from Skyline for analysis in R version
2.4-3 (R Core Team 2016). Abundance was normalized by the total ion current (TIC) for each
injection. Technical replicate, bay and habitat differences were assessed to inform SRM
analysis via non-metric multidimensional scaling (NMDS) analysis using `metaNMDS` in the
vegan package (Oksanen et al. 2016) on log(x+1) transformed abundances using a Bray-Curtis
dissimilarity matrix. Technical replicate spectral abundances clustered together on NMDS plots,
thus were averaged across each sample. Bay and habitat differences in global abundance were
visually but not quantitatively analyzed (Supplemental Figure 4).

Selected Reaction Monitoring

Target selection

Thirteen proteins were selected for SRM targets (Table 2). First, candidate proteins (~200) were
selected based on biological function listed in the Universal Protein Knowledgebase (Apweiler et
al. 2004) and evidence of stress response in bivalves from the scientific literature. Candidate
proteins were screened for detectability using DIA results. Selected proteins were required to
have ≥3 high quality peptides, each with ≥3 transitions, present in all DIA biological and
technical replicate data. Quality peptides had uniform peak morphology and retention time in
Skyline across replicates. A total of 49 peptides were selected for SRM: 39 to quantify 13 target
proteins (116 transitions), and 10 for internal standard (30 transitions). A full list of transition
targets are published on https://panoramaweb.org/e0TsuK.url and available in the project
repository (Spencer et al. 2019).

SRM Protein Targets	Top Blast Hit Uniprot SpID (E-value)	Peptide sequences
heat shock protein 90-alpha	P30946 (0)	GVVDSEDLPLNISR EVVQSSAFVER DSSTMGYMAAK
heat shock protein 70	Q91233 (0)	TTPSYVAFNDTER NAVVTVPAYFNDAQR IINEPTAAALAYGLDK
superoxide dismutase	P28757 (1e-57)	THGAPTDEER ISLTGPHSIIGR TIVVHADVDDLGK
catalase	P00432 (0)	AGELGGSDPDYAMR LYSYSDTHR LTANIAGHLIGAQEFIQK
peroxiredoxin-1	Q6B4U9 (2e-95)	ALFIIDDK QITMNDLPVGR

		LVQAFQFTDK
puromycin-sensitive aminopeptidase	Q11011 (0)	LNSGSVGVYR SLTENFVTEEQAK SIQQSVENIR
protein disulfide-isomerase	P07237 (0)	NNKPSDYQGGR DNVVVIGFFK MDSTANEVEDVK
ras-related protein rab-11B	O35509 (1e-85)	VVLVGDSGVGK STIGVEFATR AQLWDTAGQER
sodium/potassium-transporting ATPase subunit alpha	Q13733 (9e-62)	TVIEPMAGDGLR MVTGDNVNTAR LLDQVWPDLR
glycogen phosphorylase (muscle form)	Q9WUB3 (0)	APNSFNLR VLYPNDNFFEGK TSFDAFPDK
trifunctional enzyme β-Subunit (mitochondrial)	O46629 (0)	AAQDNGLLTDVLAYK ALELGLKPK FNLWGGSLSLGHPFGATGV R
cytochrome P450	P00185 (7e-39)	IITRPFNVNGLLAYDSR WLDESGVFLPEEHPSR QSLLPFGATGPR
arachidonate 5-lipoxygenase	P09917 (2e-94)	APGLPAQIK MDVEGTLPEDLK GLGLGGVPGQNGK

295   **Table 2:** SRM proteins targets selected based on biological function and detectability across DIA samples
296

Data acquisition
SRM samples were analyzed on a Vantage Triple-Stage Quadrupole Mass Spectrometer
(Thermo Scientific, San Jose, CA, USA), and injected by a nanoACQUITY UPLC® system
(Waters, Milford, MA, USA) at random in two technical replicates. For each sample, 2 µl of
peptides + PRTC solution containing 1.0 µg of geoduck peptides was injected, trapped on a 3
cm pre-column and separated on a 30 cm analytical column using a chromatography gradient of
2-60% acetonitrile over 60 minutes. Columns were prepared as in DIA (above). Samples were
injected in randomized groups of 5, followed by a Peptide Retention Time Calibration (PRTC)
plus bovine serum albumin peptides (BSA) standard, then Final Solvent blank. Vantage MS
sequence and method files are available in the project repository (Spencer et al. 2019).
307

Protein identification and quality assurance
Peptides were identified and quantified via Skyline-daily version 3.7.1.11357 (MacLean et al.
2010). Raw SRM files were imported into a Skyline-daily project along with the target protein
transitions, and the spectral library (.blib file) created previously in the DIA Protein Identification
step. SRM peptides were verified by regressing PRTC peptide retention time (RT) in SRM
against retention time in DIA. A fitted model from PRTC peptides predicted RT of protein target
peptides. Where necessary, peak selection and boundaries were manually adjusted for SRM
peptide chromatograms, and actual RT were regressed against predict RT to confirm correct
selection ($F_{(1,38)}$: 5768, p-value: < 2.2e-16, Adjusted R-squared: 0.9933) (Supplemental Figure

5). Transition peak area, defined henceforth as abundance, was exported from Skyline for
further analysis in R (R Core Team 2016). Abundance results from the separate serial dilution
samples were used to remove peptides that did not adhere to the dilution curve. Briefly, dilution
abundances (exported from Skyline) for each transition were normalized by the most dilute
sample abundance, then regressed against predicted ratios. Peptides with slope coefficient
$0.2<x<1.5$ and adjusted $R^2 >0.7$ were included in analysis. Ten of the 39 peptides were
discarded from the dataset based on dilution standards results (Supplemental Figure 6). To
determine and remove disparate technical replicates, NMDS analysis was performed as
described above. Technical replicates with ordination distance >0.2 were removed, and only
samples with two technical replicates were preserved for analysis (Supplemental Figure 7).
Thirteen technical replicates from different samples and all replicates from three sample were
discarded, for 84% technical replicate and 94% biological replicate retention. Within samples,
transitions with coefficients of variation (CV) > 40% between technical replicates were also
discarded (2% of all transitions across 21 samples). In final dataset for differential analysis, 10
proteins, 26 peptides, and 77 transitions were retained. Mean transition abundance was
calculated for replicates, with zero in the place of n/a values, which Skyline generates for
replicates without peaks. Transition abundances within each peptide were summed for a total
peptide abundance before analyzing for differential abundance.
335
Differential protein analysis
After data quality screening, peptide abundance was analyzed for differences between locations
and habitats. NMDS plots visualized patterns in peptide abundances by bay and habitat as
described above. Global peptide abundance was compared between bay and habitats using
two-way ANOVA on log-transformed abundances. For protein-specific comparisons, peptide
abundances were grouped by protein, box-cox transformed (Box and Cox 1964) and normality
confirmed via qqplot (Wickham 2017). Two-way ANOVA tested abundances for each protein
between eelgrass and unvegetated habitats within and between bays. Pairwise comparisons for
differentially abundant proteins were tested with the t-statistic. Peptides within proteins were
regressed against each other to confirm stable abundance patterns. For all statistical analyses,
significance was defined as alpha ≤ 0.05, corrected for multiple comparisons using the
Bonferroni correction.
348
Correlative analysis
To understand how environmental and biometric parameters covaried, Pearson's product-
moment correlation and scatter plots were assessed between protein abundances, growth, and
environmental summary statistics (mean and variance). Each protein was assessed
independently. Due to salinity probe malfunction, salinity data were not included in correlation
tests.
355
356 All analyses were performed in RStudio version 1.1.383 (R Core Team 2016). R scripts and
357 notebooks (Spencer et al. 2019), raw data (ProteomeXchange PXD012266), and Skyline project
358 files (https://panoramaweb.org/e0TsuK.url) are publicly available.

Results

Environmental & Growth Data

Mean pH differed significantly between habitats across all bays ($F_{(1,206)}=180.0$, $p=1.1e-28$) (Figure 2). During the deployment, pH was recorded from 6.71 to 8.34, with mean pH 7.86±0.15 in eelgrass, and 7.51±0.25 in unvegetated habitats (means are for all locations). Variability in pH was significantly different among bays ($F_{(3,206}=43.8$, $p=1.0e-20$). Variability did not differ between habitats across all bays, but differences were detected between habitats within Case Inlet and within Willapa Bay (less variable in eelgrass, Supplemental Table 1). The locations with the highest and lowest daily mean pH were Fidalgo Bay-eelgrass (7.90±0.19) and Port Gamble Bay-unvegetated, respectively (7.32±0.25). On average across all locations, pH fluctuated daily by 0.46±0.23 pH units. Considerable heterogeneity among bays was observed in the other environmental parameters. Mean temperature was significantly different among all bays ($F_{(3,236)}=129.4$, $p=2.2e-48$), and temperature decreased with latitude (coldest in northernmost Fidalgo Bay, warmest in southernmost Willapa bay). Temperature did not differ between habitats within bays (Supplemental Figure 1). Dissolved oxygen (DO) varied among bays in both daily standard deviation ($F_{(3,210)}=132.8$, $p=4.6e-47$) and mean ($F_{(3,210)}=56.7$, $p=1.1e-25$). DO variability was substantially higher in the two northern bays (SD was 5.6 and 3.9 mg/L in FB, PGB), as compared to the southern bays (2.5 and 1.4mg/L in CI, WB). Across all bays, DO variability did not differ between habitats, but did differ within Case Inlet and Fidalgo Bay (Supplemental Table 1 & Figure 2). Mean salinity differed by bay ($F_{(3,136)}=254.3$, $p=2.3e-54$), with the largest differences between Fidalgo Bay (mean 29.9 ppt) and the other three bays (mean 23.4-27.0 ppt) (Supplemental Figure 3). Growth significantly differed between northern and southern bays ($F_{(1,97)}=54.8$, $P=4.9-11$), but not between habitats either within or across all bays. Geoduck in Fidalgo Bay and Port Gamble Bay grew larger compared to Willapa Bay, and Case Inlet (Figure 3). Survival did not differ among locations (Supplemental Table 1).

Figure 2: Daily mean pH in eelgrass (dashed lines) and unvegetated (solid lines) across bays during geoduck deployment. Gray ribbons denote daily standard deviation. Data from Port Gamble bay eelgrass are not included due to probe failure.

Figure 3: Geoduck shell growth after 30 days across Willapa Bay (WB), Case Inlet (CI), Port Gamble Bay (PG), and Fidalgo Bay (FB), where -U and -E represent unvegetated (empty boxes) and eelgrass habitats (filled boxes), respectively. Growth is relative to the mean initial shell length within deployment groups (n=5 per group, 3 groups per location). Boxes contain all biological replicates lying within the interquartile range (IQR), with median growth indicated by line in middle of boxes. Whiskers extend to the largest value no greater than 1.5*IQR, and dots indicate outliers beyond 1.5*IQR. Geoduck that did not survive deployment are not included. Growth differed significantly between southern bays (WB, CI) and northern bays (PG, FB) (p=4.9e-11) but not between habitats within bays.

Protein Detection and Variance

In DIA, a total of 298,345 peptide transitions were detected from 30,659 distinct peptides across
the 8 samples (one sample per habitat from each bay). These peptides were associated with
8,077 proteins, and more than half of the proteins (4,252) were annotated using Universal
Protein Resource database (UniProt). Automated peak selection (Skyline) success rate was
71%.
In SRM, the final dataset after screening included 10 proteins, 26 peptides, and 77
transitions. The 3 proteins fully removed from the dataset were heat shock protein 70,
peroxiredoxin-1, and ras-related rab. The SRM mean coefficients of variation (CV) of technical
replicate abundances across all transitions decreased from 18.2% to 9.6% after screening.
Transition abundance CV within bays ranged from 24.9% to 83.2% with mean 50.1%, and within
deployment locations CV ranged from 11.6% to 93.0% with mean 48.9% (Supplemental Table
3). Within proteins, regression analysis indicated that peptide abundances from the same
protein differed slightly, however the relative abundances across samples was consistent. This
indicates a small degree of background variability in peptide detection, digestion, or stability
within proteins that applied to all samples (Pep1xPep2: R^2_A=0.985, coefficient=0.682;

Pep1xPep3: R^2_A=0.990, coefficient=0.954; Pep2xPep3: R^2_A=0.990, coefficient=0.954).

Protein Abundance Differences

None of the 10 targeted proteins were differentially abundant between habitats within or across
bays (Figure 4, Supplemental Table 2). NMDS plots of all transitions in DIA and those targeted
in SRM revealed clustering of overall proteomic response by bay (Supplemental Figures 4 & 8).
In SRM, Fidalgo Bay and Port Gamble Bay samples clustered together (henceforth "northern
bays"), and some overlap between Case Inlet and Willapa Bay ("southern bays") indicated
similar protein abundances within these ad-hoc regions (Supplemental Figure 8). This was
verified from the ANOVA results, which detected significant abundance differences between
northern and southern bays for three proteins: HSP90-α (HSP90) ($F_{(1,133)}$=20.5, p-adj=1.8e-
4), trifunctional-enzyme subunit β-subunit (TEβ) ($F_{(1,88)}$=11.1, p-adj=0.018), and puromycin-
sensitive aminopeptidase (PSA) ($F_{(1,130)}$=9.11, p-adj=0.043). HSP90 and TEβ abundances
were also significantly different between bays (respectively: $F_{(3,131)}$= 7.80, p-adj=0.0011;
$F_{(3,345)}$= 5.19, p-adj=0.034), but these differences were driven by regional differences, as post-
hoc tests detected no differences between Case Inlet and Willapa Bay (southern), or beteen
Fidalgo Bay and Port Gamble Bay (northern). For the three differentially abundant proteins,
abundances were lowest in Case Inlet (southernmost in Puget Sound) followed by WIllapa Bay
(southernmost overall), then Port Gamble Bay, and highest in Fidalgo Bay (northernmost).

Figure 4: Boxplots of protein mean spectral abundances (mean of 2 or 3 peptides targeted for each protein) for
Fidalgo Bay (FB), Port Gamble Bay (PG), Case Inlet (CI), and Willapa Bay (WB), where -U and -E represent
unvegetated (white boxes) and eelgrass (filled boxes) habitats, respectively. For each location, n=5 or 6 geoduck.
Boxes contain all biological replicates lying within the interquartile range (IQR), with median abundances indicated by
line in middle of boxes. Whiskers extend to the largest value no greater than 1.5*IQR, and dots indicate outliers
beyond 1.5*IQR. Protein abundance ranges (y-axes) vary between proteins. Differentially abundant proteins between
region and bay are indicated by (**), and region only by (*). No protein abundances were significantly different
between habitats.
428
Correlation between Environment, Abundance, and Growth
Growth positively correlated with peptide abundance in all but 2 of the 10 targeted proteins (no
correlation with catalase and superoxide dismutase), including the three proteins that were
differentially abundant between bays (Table 3). Growth also correlated with most environmental
parameters (excluding salinity SD). Heat Shock Protein 90 correlated positively with dissolved
oxygen SD. Mean and SD pH did not correlate with any peptide abundance patterns or growth.
None of the environmental parameters, nor growth, correlated significantly with peptide
abundances pooled across all proteins.
437

	Growth	heat shock protein 90	puromycin sensitive aminopeptidase	trifunctional enzyme β-subunit
Growth	---	**r=0.53** **p=5.54e-11**	**r=0.46** **p=1.42e-7**	**r=0.43** **p=9.70e-7**
T_{mean}	**r= -0.39** **p=8.11e-21**	r= -0.36 p=0.060	r= -0.24 p=0.48	r= -0.21 p=0.70
T_{sd}	**r= 0.39** **p=2.10e-20**	r= 0.24 p=0.42	r= 0.15 p=1	r= 0.11 p= 1
DO_{mean}	**r= 0.45** **p=2.12e-28**	r= 0.26 p=0.34	r=0.18 p=0.91	r=0.21 p=0.66
DO_{sd}	**r= 0.48** **p=2.64e-32**	**r=0.41** **p=0.021**	r=0.31 p=0.16	r=0.31 p=0.15
pH_{mean}	**r= -0.32** **p=1.08e-12**	r= -0.09 p=1	r=-0.02 p=1	r=-0.03 p=1
pH_{sd}	**r= 0.30** **p=8.72e-11**	r= 0.22 p=0.73	r=0.06 p=1	r=0.04 p=1

Table 3: Correlation analysis results between growth, environmental
parameters, and peptide abundance (z-transformed). Correlation coefficient
r shown with p-values adjusted via Bonferroni correction (number of
comparisons). Correlations deemed significant are in bold.

Discussion

This study tested the effects of varying pH on geoduck, a valuable aquaculture species
in a natural setting, and confirmed that *Zostera marina* eelgrass can effectively alter local pH
during warm summer months (June and July). We have shown that ocean acidification research
on cultured shellfish can augment findings from controlled laboratory studies with field
deployments to incorporate natural variability and relevant environmental drivers associated
with an organism's habitat. Targeted proteomics was assessed alongside growth and
environmental data for an integrated view of how geoduck respond to varying environmental
conditions. Proteomics is a powerful approach suitable for comparative physiological studies of
non-model, marine organisms (Tomanek, 2014). Using a two step method, this study detected
substantially more proteins (8,077) compared to the previous geoduck study using Data
Dependent Acquisition (3,651) (Timmins-Schiffman et al. 2017). This produced a valuable
protein catalogue for future projects, as researchers can now skip directly to the targeted SRM
phase to greatly reduce the cost and time associated with a discovery analysis.
Geoduck exhibited no phenotypic differences between pH conditions, counter to our
predictions. We predicted that pH would be higher within eelgrass habitats, creating a refuge
against the less alkaline surrounding waters and reducing oxidative stress. Concordantly,
proteins involved in the oxidative stress response would be less abundant inside eelgrass
habitats (such as superoxide dismutase, peroxiredoxin-1, catalase, and HSP), possibly
translating to differential growth as less energy would be used to counter the pH stress. While
pH in eelgrass habitats was found to be consistently higher in this study, no differences in
abundance of selected peptides, growth or survival were found between habitats across all four
bays. This suggests that juvenile geoduck may tolerate a wide pH range in the context of the
natural environment in which they are cultured.
Earlier studies on other clam species point to some degree of pH tolerance, but also
describe complex responses to low pH that vary between metrics, species, and when secondary
stressors are applied (Ries et al. 2009; Ringwood and Keppler 2002; G. G. Waldbusser et al.
2010). For example, juvenile carpet shell clams (*Ruditapes decussatus*) under ambient (pH 8.2)
and reduced pH (7.8, 7.5) for 75 days displayed no difference in size, weight, or calcification
rate (Range et al. 2011), but other physiological parameters (clearance, ingestion, respiration,
ammonia excretion) differed at day 87 (Fernández-Reiriz et al. 2011). In the hard clam
Mercenaria mercenaria, protein oxidation, biomineralization, and standard metabolic rate (SMR,
measured as resting oxygen consumption) in adults were largely unaffected by hypercapnia
alone, but when combined with elevated temperature SMR increased and shell strength
decreased (Ivanina et al. 2013; Matoo et al. 2013). Interestingly, the baltic clam (*Macoma
balthica*) grew significantly larger in low pH (7.35 vs. 7.85 for 29 days), and were largest when
combined with low dissolved oxygen (3.0 mg/L vs. 8.5 mg/L) (Jansson et al. 2015). Geoduck
metrics examined in this study were not affected by varying pH, but other physiological
parameters (metabolic rate, biomineralization, reproductive development, cytoskeleton), and
other tissues such as mantle or hepatopancreas, may be affected and should be examined in
future studies.
The complex, mixed responses exhibited in clam species may, in part, be a function of
local adaptation to varying environmental drivers. Pacific geoduck are native to the Puget

Sound, a region that experiences regular episodes of low pH in certain areas and has significant
diel and monthly pH variability (Busch et al. 2013; Feely et al. 2008, 2010). Thus, the species
may have evolved under selective pressure to withstand periods of low pH. The native
Northeast Pacific Coast oyster, *Ostrea lurida,* also shows signs of pH tolerance as veliger larvae
compared to the non-native Pacific oyster (*Crassostrea gigas*) (Waldbusser et al. 2016), a stage
primarily found to be vulnerable in other calcifying species (for reviews see Byrne and
Przeslawski 2013; Kurihara 2008). Geoduck are also infaunal organisms, extending their long
siphons into the water column for feeding and retreating to deep burrows during low tide or
when disturbed (Goodwin and Pease 1987). Sediment and burrow chemistry, while influenced
by the overlying water column, can have lower pH due to aerobic microbial activity, another
potential source of selective pressure shaping this giant clam's pH tolerance (Gattuso and
Hansson 2011; Widdicombe and Spicer 2008). An important future step is to assess the relative
influence of sediment pH and overlying water column pH on burrowing calcifiers. This is
particularly applicable when comparing habitats that likely have varying bacterial communities
and activity.
While pH was not a universal predictor of geoduck phenotype in this study, mean
temperature and dissolved oxygen variability correlated significantly with biometric parameters
and separated into two groups: northern bays (Fidalgo and Port Gamble Bays), and southern
bays (Case Inlet, and Willapa Bay). Geoduck grew less (or not at all, in Case Inlet) and had
lower levels of targeted proteins in the southern bays, which were warmer with less variable
dissolved oxygen content.
Temperatures in the southern bays (16-18°C) during the deployment dates may have
exceeded optimal conditions for juvenile geoduck, resulting in elevated metabolism and less
energy available for growth (Newell and Branch 1980). Similar temperature-dependent growth
was observed in *M. mercenaria*, where shell calcification rate was highest between 12.8-15.2°C,
above which growth negatively correlated with temperature (except for a secondary peak at
23.9°C) (Storr et al. 1982). In *P. generosa*, Goodwin (1973) reported that temperature for
normal larval development is between 6-16°C. In adults, the optimal hatchery temperature for
reproduction is relatively low (appr. 11°C), and at the highest experimental temperature (19°C)
gonad did not regenerate after an initial spawning event (Marshall et al. 2012). Arney et al.
(2015) found that in early juveniles (<3.5mm), growth increased with temperature within 7-19°C
when fed ad libitum. However, organic weight accumulation (total body ash-free dry weight) was
highest between 11-15°C, indicating that the optimal juvenile temperature may be approximately
15°C. In the present study, geoduck grew fastest in cooler, northern bays (15°C), but stress
protein abundances (e.g. HSP90) did not suggest an acute thermal stress in the warmer,
southern bays (abundances were inversely related to temperature). Southern bays may have
exceeded the geoduck upper pejus temperature but remained below acute-stress, which could
explain the reduced growth in those locations without a proteomic signal. Conversely, as tissues
were collected at day 30, a heat stress signal could have been captured with earlier or more
frequent samples. A thermal performance curve for *P. generosa* under natural feeding levels
would be valuable for aquaculture siting, but these data suggest that cooler summer
temperatures are more suitable for culturing geoduck.
Dissolved oxygen (DO) variability may be an indirect indicator of geoduck performance
as it is often correlated to phytoplankton biomass (Khangaonkar et al. 2012). Less DO

fluctuation in the southern bays could be an indicator of less phytoplankton biomass, translating
to lower food availability (Anderson and Taylor 2001; Bergondo et al. 2005; Winter et al. 1975).
While we were unable to monitor chlorophyll during the outplant, both southern bays, Willapa
Bay and Case Inlet, may have phytoplankton populations that are controlled by shellfish grazers
due to long residence times and aquaculture activity (Banas et al. 2007; Washington Sea Grant
2015). It is possible that food availability was different between northern and southern locations
during the outplant period (June-July), and could be the underlying cause of higher growth and
abundances of selected proteins in the northern locations (Carmichael, Shriver, and Valiela
2004; Liu et al. 2016; Loosanoff and Davis 1963), although this warrants additional data
collection.

Conclusion

This is the first study to investigate geoduck performance alongside varying pH conditions, and
contributes a geoduck ctenidia peptide database useful for quantifying multiple proteins
simultaneously. The primary finding is that geoduck aquaculture may be less impacted by ocean
acidification compared to other environmental stressors, for example ocean warming. Geoduck
ocean acidification research is in its infancy, and these results are a snapshot into geoduck
physiology at one developmental stage, using one tissue type (ctenidia), with individuals from
one genetic pool, and with present-day pH levels in Washington State. To best inform current
and future geoduck aquaculture, further foundational studies are needed to elucidate the
variability in the species' pH limits in conjunction with more acute environmental stressors, and
expanded to include other key tissues and functions (e.g. mantle for shell secretion), and whole-
animal physiological studies (e.g. metabolic rate, reproductive development).
This study also demonstrates applied use of systems such as eelgrass beds in estuaries
to test pH effects in a natural system. There is growing interest in using macroalgae as an
ocean acidification bioremediation tool, also known as phytoremediation (Greiner et al. 2013;
Hendriks et al. 2014; Washington State Blue Ribbon Panel on Ocean Acidification 2012; Groner
et al. 2018). Incorporating seagrass into shellfish-pH interaction studies can help evaluate the
potential for merging mariculture with shellfish aquaculture to improve growing conditions for
vulnerable cultured species.

Acknowledgements

Our gratitude to the following people who assisted with this project: Grace Crandall, Kaitlyn
Mitchell and Jose Guzman assisted with protein extraction. Jarrett Egertson contributed to DIA
design. Austin Keller adapted MSConvert to correctly demultiplex and convert DIA files. Han-Yin
Yang, Brian Searle and Sean Bennett assisted with running PECAN. Brittany Taylor and Taylor
Shellfish Farms provided geoducks. Thank you to anonymous reviewers for the helpful
feedback.
562

Reference List

Abatzoglou, John T., David E. Rupp, and Philip W. Mote. 2013. "Seasonal Climate Variability and Change in the Pacific Northwest of the United States." *Journal of Climate* 27 (5): 2125–42.

Anderson, Timothy H., and Gordon T. Taylor. 2001. "Nutrient Pulses, Plankton Blooms, and Seasonal Hypoxia in Western Long Island Sound." *Estuaries* 24 (2): 228–43.

Apweiler, Rolf, Amos Bairoch, Cathy H. Wu, Winona C. Barker, Brigitte Boeckmann, Serenella Ferro, Elisabeth Gasteiger, et al. 2004. "UniProt: The Universal Protein Knowledgebase." *Nucleic Acids Research* 32 (Database issue): D115–19.

Arney, Bianca, Wenshan Liu, Ian Forster, R. Scott Mckinley, and Christopher M. Pearce. 2015. "Temperature and Food-Ration Optimization in the Hatchery Culture of Juveniles of the Pacific Geoduck Panopea Generosa." *Journal of Shellfish Research* 34 (1): 39–54.

Banas, N. S., B. M. Hickey, P. MacCready, and J. A. Newton. 2004. "Dynamics of Willapa Bay, Washington: A Highly Unsteady, Partially Mixed Estuary." *Journal of Physical Oceanography* 34 (11): 2413–27.

Banas, N. S., B. M. Hickey, J. A. Newton, and J. L. Ruesink. 2007. "Tidal Exchange, Bivalve Grazing, and Patterns of Primary Production in Willapa Bay, Washington, USA." *Marine Ecology Progress Series* 341: 123–39.

Bergondo, Deanna L., Dana R. Kester, Heather E. Stoffel, and Wendy L. Woods. 2005. "Time-Series Observations during the Low Sub-Surface Oxygen Events in Narragansett Bay during Summer 2001." *Marine Chemistry* 97 (1): 90–103.

Box, G. E. P., and D. R. Cox. 1964. "An Analysis of Transformations." *Journal of the Royal Statistical Society. Series B, Statistical Methodology* 26 (2): 211–52.

Boyd, Philip W., Christopher E. Cornwall, Andrew Davison, Scott C. Doney, Marion Fourquez, Catriona L. Hurd, Ivan D. Lima, and Andrew McMinn. 2016. "Biological Responses to Environmental Heterogeneity under Future Ocean Conditions." Global Change Biology 22 (8): 2633–50.

Bozaykut, Perinur, Nesrin Kartal Ozer, and Betul Karademir. 2014. "Regulation of Protein Turnover by Heat Shock Proteins." *Free Radical Biology & Medicine* 77 (December): 195–209.

Bulthuis, Douglas A. 1995. "Distribution of Seagrasses in a North Puget Sound Estuary: Padilla Bay, Washington, USA." *Aquatic Botany* 50 (1): 99–105.

Busch, D. Shallin, Chris J. Harvey, and Paul McElhany. 2013. "Potential Impacts of Ocean Acidification on the Puget Sound Food Web." *ICES Journal of Marine Science: Journal Du Conseil* 70 (4): 823–33.

Byrne, Maria, and Rachel Przeslawski. 2013. "Multistressor Impacts of Warming and Acidification of the Ocean on Marine Invertebrates' Life Histories." *Integrative and Comparative Biology* 53 (4): 582–96.

Carmichael, R. H., Andrea C. Shriver, and I. Valiela. 2004. "Changes in Shell and Soft Tissue Growth, Tissue Composition, and Survival of Quahogs, Mercenaria Mercenaria, and Softshell Clams, Mya Arenaria, in Response to Eutrophic-Driven Changes in Food Supply and Habitat." *Journal of Experimental Marine Biology and Ecology* 313 (1): 75–104.

Chambers, Matthew C., Brendan Maclean, Robert Burke, Dario Amodei, Daniel L. Ruderman, Steffen Neumann, Laurent Gatto, et al. 2012. "A Cross-Platform Toolkit for Mass

Spectrometry and Proteomics." *Nature Biotechnology* 30 (10): 918–20.

Coan, Eugene V., Paul Valentich Scott, and Frank R. Bernard. 2000. *Bivalve Seashells of*
Western North America: Marine Bivalve Mollusks from Arctic Alaska to Baja California.
Santa Barbara Museum of Natural History.

Dethier, Megan N., Tom Mumford, Tom Leschine, Kurt Presh, Si Simenstad, Hugh Shipman,
Doug Myers, et al. 2006. "Native Shellfish in Nearshore Ecosystems of Puget Sound."
WASHINGTON UNIV SEATTLE. http://www.dtic.mil/docs/citations/ADA477852.

Doney, Scott C., Natalie Mahowald, Ivan Lima, Richard A. Feely, Fred T. Mackenzie, Jean-
Francois Lamarque, and Phil J. Rasch. 2007. "Impact of Anthropogenic Atmospheric
Nitrogen and Sulfur Deposition on Ocean Acidification and the Inorganic Carbon System."
Proceedings of the National Academy of Sciences of the United States of America 104
(37): 14580–85.

Duquette, Ashley, James B. McClintock, Charles D. Amsler, Alberto Pérez-Huerta, Marco
Milazzo, and Jason M. Hall-Spencer. 2017. "Effects of Ocean Acidification on the Shells of
Four Mediterranean Gastropod Species near a CO2 Seep." Marine Pollution Bulletin 124
(2): 917–28.

Fabbri, E., P. Valbonesi, and S. Franzellitti. 2008. "HSP Expression in Bivalves." *Invertebrate*
Survival Journal: ISJ 5 (135): e161.

Feely, Richard A., Simone R. Alin, Jan Newton, Christopher L. Sabine, Mark Warner, Allan
Devol, Christopher Krembs, and Carol Maloy. 2010. "The Combined Effects of Ocean
Acidification, Mixing, and Respiration on pH and Carbonate Saturation in an Urbanized
Estuary." *Estuarine, Coastal and Shelf Science* 88 (4): 442–49.

Feely, Richard A., Terrie Klinger, Jan A. Newton, and Meg Chadsey. 2012. *Scientific Summary*
of Ocean Acidification in Washington State Marine Waters. US Department of Commerce,
National Oceanic and Atmospheric Administration, Office of Oceanic and Atmospheric
Research.

Feely, Richard A., Christopher L. Sabine, J. Martin Hernandez-Ayon, Debby Ianson, and Burke
Hales. 2008. "Evidence for Upwelling of Corrosive' Acidified' Water onto the Continental
Shelf." *Science* 320 (5882): 1490–92.

Fernández-Reiriz, Ma José, Pedro Range, Xosé Antón Álvarez-Salgado, and Uxio Labarta.
2011. "Physiological Energetics of Juvenile Clams Ruditapes Decussatus in a High CO_2
Coastal Ocean." *Marine Ecology Progress Series* 433: 97–105.

Gattuso, Jean-Pierre, and Lina Hansson. 2011. *Ocean Acidification*. OUP Oxford.

Gazeau, Frédéric, Christophe Quiblier, Jeroen M. Jansen, Jean-Pierre Gattuso, Jack J.
Middelburg, and Carlo H. R. Heip. 2007. "Impact of Elevated CO2 on Shellfish
Calcification." *Geophysical Research Letters* 34 (7): L07603.

Giarratano, Erica, Mónica N. Gil, and Gabriela Malanga. 2014. "Biomarkers of Environmental
Stress in Gills of Ribbed Mussel Aulacomya Atra Atra (Nuevo Gulf, Northern Patagonia)."
Ecotoxicology and Environmental Safety 107 (September): 111–19.

Gobler, Christopher J., Elizabeth L. DePasquale, Andrew W. Griffith, and Hannes Baumann.
2014. "Hypoxia and Acidification Have Additive and Synergistic Negative Effects on the
Growth, Survival, and Metamorphosis of Early Life Stage Bivalves." PloS One 9 (1):
e83648.

Goodwin, C. L. 1973. "Effects of Salinity and Temperature on Embryos of the Geoduck Clam
(Panope generosa Gould)." *In Proceedings of the National Shellfisheries Association*,
63:93–95.

Goodwin, C. Lynn, and Bruce Pease. 1987. *The Distribution of Geoduck (Panope Abrupta)*
Size, Density, and Quality in Relation to Habitat Characteristics such as Geographic Area,
Water Depth, Sediment Type, and Associated Flora and Fauna in Puget Sound,
Washington. State of Washington, Department of Fisheries, Shellfish Division.

Greiner, Jill T., Karen J. McGlathery, John Gunnell, and Brent A. McKee. 2013. "Seagrass
      Restoration Enhances 'Blue Carbon' Sequestration in Coastal Waters." *PloS One* 8 (8):
      e72469.
Groner, Maya L., Colleen A. Burge, Ruth Cox, Natalie Rivlin, Mo Turner, Kathryn L. Van
      Alstyne, Sandy Wyllie-Echeverria, John Bucci, Philip Staudigel, and Carolyn S. Friedman.
      2018. "Oysters and Eelgrass: Potential Partners in a High pCO2 Ocean." *Ecology*, May.
      https://doi.org/10.1002/ecy.2393.
Harvey, Ben P., Dylan Gwynn-Jones, and Pippa J. Moore. 2013. "Meta-Analysis Reveals
      Complex Marine Biological Responses to the Interactive Effects of Ocean Acidification and
      Warming." Ecology and Evolution 3 (4): 1016–30.
Hendriks, Iris E., Y. S. Olsen, L. Ramajo, L. Basso, A. Steckbauer, T. S. Moore, J. Howard, and
      C. M. Duarte. 2014. "Photosynthetic Activity Buffers Ocean Acidification in Seagrass
      Meadows." *Biogeosciences* 11 (2): 333.
Howarth, Robert, Francis Chan, Daniel J. Conley, Josette Garnier, Scott C. Doney, Roxanne
      Marino, and Gilles Billen. 2011. "Coupled Biogeochemical Cycles: Eutrophication and
      Hypoxia in Temperate Estuaries and Coastal Marine Ecosystems." Frontiers in Ecology and
      the Environment 9 (1): 18–26.
Hu, Menghong, Lisha Li, Yanming Sui, Jiale Li, Youji Wang, Weiqun Lu, and Sam Dupont.
      2015. "Effect of pH and Temperature on Antioxidant Responses of the Thick Shell Mussel
      Mytilus Coruscus." Fish & Shellfish Immunology 46 (2): 573–83.
Ivanina, Anna V., Gary H. Dickinson, Omera B. Matoo, Rita Bagwe, Ashley Dickinson, Elia
      Beniash, and Inna M. Sokolova. 2013. "Interactive Effects of Elevated Temperature and
      CO2 Levels on Energy Metabolism and Biomineralization of Marine Bivalves Crassostrea
      Virginica and Mercenaria Mercenaria." *Comparative Biochemistry and Physiology. Part A,
      Molecular & Integrative Physiology* 166 (1): 101–11.
Jansson, Anna, Joanna Norkko, Sam Dupont, and Alf Norkko. 2015. "Growth and Survival in a
      Changing Environment: Combined Effects of Moderate Hypoxia and Low pH on Juvenile
      Bivalve Macoma Balthica." *Journal of Sea Research* 102 (August): 41–47.
Kerrison, Philip, Jason M. Hall-Spencer, David J. Suggett, Leanne J. Hepburn, and Michael
      Steinke. 2011. "Assessment of pH Variability at a Coastal CO2 Vent for Ocean Acidification
      Studies." Estuarine, Coastal and Shelf Science 94 (2): 129–37.
Khangaonkar, Tarang, Brandon Sackmann, Wen Long, Teizeen Mohamedali, and Mindy
      Roberts. 2012. "Simulation of Annual Biogeochemical Cycles of Nutrient Balance,
      Phytoplankton Bloom(s), and DO in Puget Sound Using an Unstructured Grid Model."
      Ocean Dynamics 62 (9): 1353–79.
Kroeker, Kristy J., Rebecca L. Kordas, Ryan N. Crim, and Gerald G. Singh. 2010. "Meta-
      Analysis Reveals Negative yet Variable Effects of Ocean Acidification on Marine
      Organisms." *Ecology Letters* 13 (11): 1419–34.
Kurihara, H. 2008. "Effects of CO2-Driven Ocean Acidification on the Early Developmental
      Stages of Invertebrates." *Marine Ecology Progress Series* 373 (December): 275–84.
Liu, W., C. M. Pearce, R. S. McKinley, and I. P. Forster. 2016. "Nutritional Value of Selected
      Species of Microalgae for Larvae and Early Post-Set Juveniles of the Pacific Geoduck
      Clam, Panopea Generosa." *Aquaculture* 452 (Supplement C): 326–41.
Loosanoff, Victor L., and Harry C. Davis. 1963. "Rearing of Bivalve Mollusks." In *Advances in
      Marine Biology*, edited by F. S. Russell, 1:1–136. Academic Press.
Lushchak, Volodymyr I. 2011. "Environmentally Induced Oxidative Stress in Aquatic Animals."
      Aquatic Toxicology 101 (1): 13–30.
MacLean, Brendan, Daniela M. Tomazela, Nicholas Shulman, Matthew Chambers, Gregory L.
      Finney, Barbara Frewen, Randall Kern, David L. Tabb, Daniel C. Liebler, and Michael J.
      MacCoss. 2010. "Skyline: An Open Source Document Editor for Creating and Analyzing

711   Targeted Proteomics Experiments." *Bioinformatics* 26 (7): 966–68.

712 Marshall, Robert, R. Scott McKinley, and Christopher M. Pearce. 2012. "Effect of Temperature

713   on Gonad Development of the Pacific Geoduck Clam (Panopea Generosa Gould, 1850)."

714   *Aquaculture* 338-341 (March): 264–73.

715 Matoo, Omera B., Anna V. Ivanina, Claus Ullstad, Elia Beniash, and Inna M. Sokolova. 2013.

716   "Interactive Effects of Elevated Temperature and CO2 Levels on Metabolism and Oxidative

717   Stress in Two Common Marine Bivalves (Crassostrea Virginica and Mercenaria

718   Mercenaria)." *Comparative Biochemistry and Physiology. Part A, Molecular & Integrative*

719   *Physiology* 164 (4): 545–53.

720  Matozzo, Valerio, Andrea Chinellato, Marco Munari, Monica Bressan, and Maria Gabriella

721   Marin. 2013. "Can the Combination of Decreased pH and Increased Temperature Values

722   Induce Oxidative Stress in the Clam Chamelea Gallina and the Mussel Mytilus

723   Galloprovincialis?" Marine Pollution Bulletin 72 (1): 34–40.

724 Middelboe, Anne Lise, and Per Juel Hansen. 2007. "High pH in Shallow-Water Macroalgal

725   Habitats." *Marine Ecology Progress Series* 338: 107–17.

726 Miller, A. Whitman, Amanda C. Reynolds, Cristina Sobrino, and Gerhardt F. Riedel. 2009.

727   "Shellfish Face Uncertain Future in High CO2 World: Influence of Acidification on Oyster

728   Larvae Calcification and Growth in Estuaries." *PloS One* 4 (5): e5661.

729 Moore, Stephanie K., Nathan J. Mantua, Jan A. Newton, Mitsuhiro Kawase, Mark J. Warner,

730   and Jonathan P. Kellogg. 2008. "A Descriptive Analysis of Temporal and Spatial Patterns of

731   Variability in Puget Sound Oceanographic Properties." *Estuarine, Coastal and Shelf*

732   *Science* 80 (4): 545–54.

733 Mote, Philip W., and Eric P. Salathé. 2010. "Future Climate in the Pacific Northwest." *Climatic*

734   *Change* 102 (1-2): 29–50.

735 Newell, R. C., and G. M. Branch. 1980. "The Influence of Temperature on the Maintenance of

736   Metabolic Energy Balance in Marine Invertebrates." In *Advances in Marine Biology*, edited

737   by J. H. S. Blaxter, Frederick S. Russell, and Maurice Yonge, 17:329–96. Academic Press.

738 Oksanen, J., F. Guillaume Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R.

739   Minchin, et al. 2016. *Vegan: Community Ecology Package* (version R package version 2.4-

740   1, 2016). https:// CRAN.R-project.org/package=vegan.

741 Orr, James C., Victoria J. Fabry, Olivier Aumont, Laurent Bopp, Scott C. Doney, Richard A.

742   Feely, Anand Gnanadesikan, et al. 2005. "Anthropogenic Ocean Acidification over the

743   Twenty-First Century and Its Impact on Calcifying Organisms." *Nature* 437 (September):

744   681.

745 Paganini, Adam W., Nathan A. Miller, and Jonathon H. Stillman. 2014. "Temperature and

746   Acidification Variability Reduce Physiological Performance in the Intertidal Zone Porcelain

747   Crab Petrolisthes Cinctipes." *The Journal of Experimental Biology* 217 (22): 3974–80.

748 Palacios, Sherry L., and Richard C. Zimmerman. 2007. "Response of Eelgrass Zostera Marina

749   to CO^2 Enrichment: Possible Impacts of Climate Change and Potential for Remediation of

750   Coastal Habitats." *Marine Ecology Progress Series* 344: 1–13.

751 Pörtner, Hans-O. 2008. "Ecosystem Effects of Ocean Acidification in Times of Ocean Warming:

752   A Physiologist's View." *Marine Ecology Progress Series* 373: 203–18.

753 Pörtner, Hans O., and Anthony P. Farrell. 2008. "Physiology and Climate Change." *Science* 322

754   (5902): 690–92.

755 Przeslawski, Rachel, Maria Byrne, and Camille Mellin. 2015. "A Review and Meta-Analysis of

756   the Effects of Multiple Abiotic Stressors on Marine Embryos and Larvae." Global Change

757   Biology 21 (6): 2122–40.

758 Range, P., M. A. Chícharo, R. Ben-Hamadou, D. Piló, D. Matias, S. Joaquim, A. P. Oliveira, and

759   L. Chícharo. 2011. "Calcification, Growth and Mortality of Juvenile Clams Ruditapes

760   Decussatus under Increased pCO2 and Reduced pH: Variable Responses to Ocean

Acidification at Local Scales?" *Journal of Experimental Marine Biology and Ecology* 396 (2): 177–84.

R Core Team. 2016. *R: A Language and Environment for Statistical Computing* (version 1.1.383). https://www.R-project.org/.

Ries, Justin B., Anne L. Cohen, and Daniel C. McCorkle. 2009. "Marine Calcifiers Exhibit Mixed Responses to CO2-Induced Ocean Acidification." *Geology* 37 (12): 1131–34.

Ringwood, Amy H., and Charles J. Keppler. 2002. "Water Quality Variation and Clam Growth: Is pH Really a Non-Issue in Estuaries?" *Estuaries* 25 (5): 901–7.

Shamshak, Gina Louise, and Jonathan R. King. 2015. "From Cannery to Culinary Luxury: The Evolution of the Global Geoduck Market." *Marine Policy* 55 (May): 81–89.

Spencer, Laura H., Yaamini Venkataraman, Emma Timmins-Schiffman, Brook L. Nunn, Steven B. Roberts, Micah Horwith, and Alexander T. Lowe. 2019. "Github Repo for "pacific Geoduck (panopea Generosa) Resilience to Natural Ph Variation"". figshare. doi:10.6084/m9.figshare.7562354.v1.

Storr, John F., Alexander L. Costa, and David A. Prawel. 1982. "Effects of Temperature on Calcium Deposition in the Hard-Shell Clam, Mercenaria Mercenaria." *Journal of Thermal Biology* 7 (1): 57–61.

Thompson, E. L., W. O'Connor, L. Parker, P. Ross, and D. A. Raftos. 2015. "Differential Proteomic Responses of Selectively Bred and Wild-Type Sydney Rock Oyster Populations Exposed to Elevated CO2." Molecular Ecology 24 (6): 1248–62.

Thomsen, Jörn, Isabel Casties, Christian Pansch, Arne Körtzinger, and Frank Melzner. 2013. "Food Availability Outweighs Ocean Acidification Effects in Juvenile Mytilus Edulis: Laboratory and Field Experiments." Global Change Biology 19 (4): 1017–27.

Timmins-Schiffman, Emma, William D. Coffey, Wilber Hua, Brook L. Nunn, Gary H. Dickinson, and Steven B. Roberts. 2014. "Shotgun Proteomics Reveals Physiological Response to Ocean Acidification in Crassostrea Gigas." BMC Genomics 15 (November): 951.

Timmins-Schiffman, Emma B., Grace A. Crandall, Brent Vadopalas, Michael E. Riffle, Brook L. Nunn, and Steven B. Roberts. 2017. "Integrating Discovery-Driven Proteomics and Selected Reaction Monitoring To Develop a Noninvasive Assay for Geoduck Reproductive Maturation." *Journal of Proteome Research* 16 (9): 3298–3309.

Ting, Ying Sonia. 2017. "Shifting the Paradigm: Peptide-Centric Analysis of Systematically Sampled Mass Spectrometry Data." https://digital.lib.washington.edu/researchworks/bitstream/handle/1773/38147/Ting_washington_0250E_16676.pdf?sequence=1&isAllowed=y.

Tomanek, Lars. 2011. "Environmental Proteomics: Changes in the Proteome of Marine Organisms in Response to Environmental Stress, Pollutants, Infection, Symbiosis, and Development." Annual Review of Marine Science 3: 373–99.

Tomanek, Lars, Marcus J. Zuzow, Anna V. Ivanina, Elia Beniash, and Inna M. Sokolova. 2011. "Proteomic Response to Elevated PCO2 Level in Eastern Oysters, Crassostrea Virginica: Evidence for Oxidative Stress." The Journal of Experimental Biology 214 (Pt 11): 1836–44.

Tomanek, Lars. 2014. "Proteomics to Study Adaptations in Marine Organisms to Environmental Stress." Journal of Proteomics 105 (Supplement C): 92–106.

Tomanek, Lars. 2015. "Proteomic Responses to Environmentally Induced Oxidative Stress." *The Journal of Experimental Biology* 218 (Pt 12): 1867–79.

Tukey, John W. 1977. "Exploratory Data Analysis." https://pdfs.semanticscholar.org/2f40/fd06bc9fd00a27437e14ed171e96e4fd9326.pdf.

Tunnicliffe, Verena, Kimberley T. A. Davies, David A. Butterfield, Robert W. Embley, Jonathan M. Rose, and William W. Chadwick Jr. 2009. "Survival of Mussels in Extremely Acidic Waters on a Submarine Volcano." Nature Geoscience 2 (April): 344.

Vadopalas, B., T. W. Pietsch, C. S. Friedman - Malacologia, and 2010. 2010. "… Proper Name
for the Geoduck: Resurrection of Panopea Generosa Gould, 1850, from the Synonymy of
Panopea Abrupta (Conrad, 1849)(Bivalvia: Myoida: Hiatellidae)." *BioOne*.
http://www.bioone.org/doi/abs/10.4002/040.052.0111.
Vadopalas, Brent, Jonathan P. Davis, and Carolyn S. Friedman. 2015. "Maturation, Spawning,
and Fecundity of the Farmed Pacific Geoduck Panopea Generosa in Puget Sound,
Washington." *Journal of Shellfish Research* 34 (1): 31–37.
Wahl, M., S. Schneider Covachã, V. Saderne, C. Hiebenthal, J. D. Müller, C. Pansch, and Y.
Sawall. 2018. "Macroalgae May Mitigate Ocean Acidification Effects on Mussel Calcification
by Increasing pH and Its Fluctuations: Biogenic Fluctuations Mitigate OA Effects."
Limnology and Oceanography 63 (1): 3–21.
Waldbusser, George G., Matthew W. Gray, Burke Hales, Chris J. Langdon, Brian A. Haley, Iria
Gimenez, Stephanie R. Smith, Elizabeth L. Brunner, and Greg Hutchinson. 2016. "Slow
Shell Building, a Possible Trait for Resistance to the Effects of Acute Ocean Acidification."
Limnology and Oceanography 61 (6): 1969–83.
Waldbusser, G. G., H. Bergschneider, and M. A. Green. 2010. "Size-Dependent pH Effect on
Calcification in Post-Larval Hard Clam Mercenaria Spp." *Marine Ecology Progress Series*
417 (November): 171–82.
Washington Sea Grant. 2013. "Final Report: Geoduck Aquaculture Research Program." *Report
to the Washington State Legislature. Washington Sea Grant Technical Report WSG-TR*,
13–03.
Washington Sea Grant. 2015. "Shellfish Aquaculture in Washington State." *Final Report to the
Washington State Legislature* 84.
https://pdfs.semanticscholar.org/b833/e0fcb8a0459697f94fd86b9848ee0e59c0a2.pdf.
Washington State Blue Ribbon Panel on Ocean Acidification. 2012. "Ocean Acidification: From
Knowledge to Action, Washington State's Strategic Response." 12-01-015 . Washington
Department of Ecology, Olympia, Washington.
https://fortress.wa.gov/ecy/publications/documents/1201015.pdf.
"Washington's Wild Geoduck Fishery | WA - DNR." n.d. Accessed November 20, 2017.
https://www.dnr.wa.gov/programs-and-services/aquatics/shellfish/washingtons-wild-
geoduck-fishery.
Watson, Sue-Ann, Southgate, Paul C., Miller, Gabrielle M., Moorhead, Jonathan A., and
Knauer, Jens. 2012. "Ocean acidification and warming reduce juvenile survival of the fluted
giant clam, Tridacna squamosa." *Molluscan Research*, 32 (3). pp. 177-180.
Weiss, Ingrid Maria, Noreen Tuross, Lia Addadi, and Steve Weiner. 2002. "Mollusc Larval Shell
Formation: Amorphous Calcium Carbonate Is a Precursor Phase for Aragonite." *The
Journal of Experimental Zoology* 293 (5): 478–91.
Wickham, Hadley. 2017. "ggplot2 - Elegant Graphics for Data Analysis (2nd Edition)." *Journal of
Statistical Software, Book Reviews* 77 (2): 1–3.
Widdicombe, Stephen, and John I. Spicer. 2008. "Predicting the Impact of Ocean Acidification
on Benthic Biodiversity: What Can Animal Physiology Tell Us?" *Journal of Experimental
Marine Biology and Ecology* 366 (1): 187–97.
Winter, D. F., K. Banse, and G. C. Anderson. 1975. "The Dynamics of Phytoplankton Blooms in
Puget Sound a Fjord in the Northwestern United States." *Marine Biology* 29 (2): 139–76.
"WWW Tide/Current Predictor - Site Selection." n.d. Accessed December 7, 2017.
http://tbone.biol.sc.edu/tide/.
Zhang, Yang, Jin Sun, Huawei Mu, Jun Li, Yuehuan Zhang, Fengjiao Xu, Zhiming Xiang,
Pei-Yuan Qian, Jian-Wen Qiu, and Ziniu Yu. 2015. "Proteomic Basis of Stress Responses in
the Gills of the Pacific Oyster Crassostrea Gigas." Journal of Proteome Research 14 (1):
304–17.

Supplemental Material

Supplemental Table 1. Outplant location metrics; FB=Fidalgo Bay, PG=Port Gamble, SR=Skokomish River Delta, CI=Case Inlet, WB=Willapa Bay; -E=Eelgrass, -U=Unvegetated. Δ Habitats, Δ Bays, Δ Regions: p-adjusted from 2-way ANOVA testing differences between habitats, bays, and ad-hoc regions (south=CI & WB, north=FB & PG). **Growth did not differ within ad-hoc regions. *Salinity probes failed fully or intermittently at 4 of the 8 locations; habitat salinities were not compared.

Deployment location	FB-E	FB-U	PG-E	PG-U	CI-E	CI-U	WB-E	WB-U	Δ Habitat	Δ Bays	Δ Regions
% Survival	87	80	80	93	67	60	93	100	p= 0.88	p=0.25	p=0.66
Mean Relative % Growth (± SD)	16.2 (±7.3)	23.3 (±12.5)	22.2 (±12.5)	21.1 (±13.3)	1.2 (±8.3)	-2.8 (±5.3)	10 (±16.2)	4.5 (±8.3)	p=1	**p= 2.4e-10	p= 4.9e-11
Mean pH (TS)	7.90	7.54	failed	7.32	7.87	7.63	7.81	7.55	p= 1.1e-28	p= 1.1e-27	p= 3.9e-3
pH Standard Deviation (TS)	0.19	0.23	failed	0.25	0.16	0.20	0.06	0.18	p=1	p= 1.0e-20	p= 1.6e-15
Mean DO (mg/L)	failed	10.9	10.8	11.9	10.4	8.3	8.4	8.8	p=1	p= 1.1e-25	p= 3.2e-19
DO Standard Deviation (mg/L)	failed	5.7	3.9	3.9	2.6	1.9	1.5	1.3	p=0.31	p= 4.6e-47	p= 2.8e-39
Mean Temp (°C)	15.1	15.1	15.3	15.2	16.6	16.4	18.2	18.2	p=1	p= 2.2e-48	p= 1.2e-36
Temp Standard Deviation (°C)	1.6	1.6	2.1	2.4	1.7	1.6	1.2	1.2	p=0.90	p=0.05	p=1
Mean Salinity (ppt)	30.2	*29.1	23.4	failed	failed	24.7	27.3	*24.4	*Not tested	*p= 2.3e-54	p= 0.30
Salinity Standard Deviation (ppt)	0.4	*0.34	1.3	failed	failed	1.2	0.70	*1.1	*Not tested	*p= 2.6e-3	p=0.12

Supplemental Figure 1: Daily mean temperature in eelgrass (dashed lines) and unvegetated (solid lines) across bays during geoduck deployment. Gray ribbons denote standard deviations per day.

Supplemental Figure 2: Daily mean DO in eelgrass (dashed lines) and unvegetated (solid lines) across bays during geoduck deployment. Gray ribbons denote standard deviations per day. Fidalgo Bay eelgrass probe failed towards the beginning of the outplant period.

Supplemental Figure 3: Daily mean salinity in eelgrass (dashed lines) and unvegetated (solid lines) across bays during geoduck deployment. Gray ribbons denote standard deviations per day. Salinity probes failed at several locations; no habitat comparisons were made.

Supplemental Table 2: SRM protein ANOVA results by Region (FB/PGB vs. CI/WB), Bay, and Habitat for all proteins combined, then each protein individually, with Pr(>F)-adjusted calculated via the Bonferroni Correction. Bold = significantly different abundance. Habitat was tested using 2-way ANOVA with abundance ~ Bay*Habitat.

	Comparison	Df	Sum Sq	Mean Sq	F value	Pr(>F)	P.adj
all peptides combined	**North vs. South**	**1**	**18.2**	**18.2**	**10.9**	**0.00010**	**0.0014**
	Bay	**3**	**34.2**	**11.4**	**7.0**	**0.00012**	**0.0017**
	Habitat	1	1.2	1.2	0.75	5.4	14
arachidonate	North vs. South	1,133	7.4	7.4	1.3	0.26	3.6
	Bay	3,131	60	20	3.7	0.014	0.20
	Habitat	1,130	2.2	2.2	0.40	0.53	7.4
catalase	North vs. South	1,128	8.6E-05	8.6E-05	0.46	0.50	7.0
	Bay	3,126	0.00019	6.4E-05	0.34	0.79	11
	Habitat	1,125	1.4E-08	1.4E-08	7.4E-05	0.99	14
cytochrome P450	North vs. South	1,133	0.0065	0.0065	0.96	0.33	4.6
	Bay	3,131	0.015	0.0051	0.76	0.52	7.3
	Habitat	1,130	0.0085	0.0085	1.3	0.26	3.7
glycogen phosphorylase	North vs. South	1,133	94	94	1.05	0.31	4.3
	Bay	3,131	459	153	1.7	0.16	2.3
	Habitat	1,130	114	114	1.3	0.26	3.6
HSP90-alpha	**North vs. South**	**1,133**	**421,443**	**421,443**	**21**	**1.3E-05**	**0.00018**
	Bay	**3,131**	**477,740**	**159,247**	**7.8**	**7.83E-05**	**0.0011**
	Habitat	1,130	567	567	0.028	0.87	12
protein disulfide isomerase	North vs. South	1,84	3.8	3.8	5.55	0.021	0.29
	Bay	3,82	4.6	1.5	2.2	0.10	1.4
	Habitat	1,81	0.01	0.006	0.0078	0.93	13

puromycin-sensitive	**North vs. South**	**1,130**	**0.94**	**0.94**	**9.1**	**0.0031**	**0.043**
	Bay	3,128	1.2	0.40	3.87	0.011	0.15
	Habitat	1,127	0.09	0.089	0.87	0.35	4.9
sodium/potassium transporting ATPase	North vs. South	1,88	0.65	0.65	0.42	0.52	7.2
	Bay	3,86	3.2	1.06	0.68	0.56	7.9
	Habitat	1,85	0.85	0.85	0.55	0.46	6.5
superoxide dismutase	North vs. South	1,66	0.0012	0.0012	1.8	0.18	2.5
	Bay	3,64	0.0020	0.00067	1.02	0.39	5.5
	Habitat	1,63	5.4E-05	5.4E-05	0.081	0.78	11
trifunctional enzyme β-subunit	**North vs. South**	**1,88**	**252**	**252**	**11.1**	**0.0013**	**0.018**
	Bay	**3,86**	**345**	**115**	**5.19**	**0.0024**	**0.034**
	Habitat	1,85	22	22	1.0	0.32	4.5

Supplemental Table 3: Mean coefficients of variation of SRM transition abundance for each protein, location

Protein	CI-B	CI-E	FB-B	FB-E	PG-B	PG-E	WB-B	WB-E
arachidonate 5-lipoxygenase	0.84	0.57	0.44	0.33	0.46	0.64	0.48	0.58
catalase	0.61	0.52	0.47	0.42	0.50	0.66	0.50	0.43
cytochrome P450	0.74	0.57	0.38	0.43	0.26	0.58	0.56	0.46
glycogen phosphorylase (muscle form)	0.55	0.49	0.35	0.35	0.37	0.61	0.51	0.47
heat shock protein 70	0.59	0.61	0.20	0.33	0.37	0.44	0.45	0.38
heat shock protein 90-alpha	0.70	0.62	0.32	0.40	0.45	0.52	0.50	0.49
protein disulfide isomerase	0.63	0.50	0.38	0.31	0.38	0.70	0.41	0.43
peroxiredoxin-1	0.56	0.46	0.31	0.33	0.42	0.46	0.40	0.36
puromycin-sensitive aminopeptidase	0.69	0.52	0.39	0.40	0.42	0.51	0.44	0.42
ras-related protein rab-11B	0.61	0.43	0.30	0.33	0.42	0.63	0.57	0.54
Na/K transporting ATPase	0.70	0.58	0.30	0.36	0.40	0.62	0.51	0.44
Superoxide dismutase	0.49	0.53	0.49	0.57	0.30	0.71	0.46	0.44
Trifunctional Enzyme β-subunit	0.61	0.48	0.41	0.38	0.46	0.63	0.52	0.53

Supplemental Figure 4: Non-metric Multi-Dimensional Scaling plot (NMDS) showing patterns of similarity among DIA peptide abundances between technical replicates (same symbol/color), deployment bay (same color), and north/south region, where each point represents one geoduck. Relative proximity of points represents overall degree of peptide abundance similarity.

Supplemental Figure 5: Linear regression model fit of PRTC internal standard peptides in SRM against DIA, used to confirm identity of targeted peptides in SRm by calculating adjustment in retention time between SRM and DIA.

Supplemental Figure 6 (Interactive, online): Dilution curve peptide abundance ratios regressed against predicted ratios from serial sample dilutions. Peptides with slope coefficient 0.2<x<1.5 and adjusted R2 >0.7 were included in analysis. Link to interactive figure: http://owl.fish.washington.edu/generosa/Generosa_DNR/Dilution-Curve-Transitions.html

Supplemental Figure 7 (Interactive, online): Non-metric Multi-Dimensional Scaling plot (NMDS) of SRM technical replicates. Link to interactive figure: http://owl.fish.washington.edu/generosa/Generosa_DNR/NMDS-technical-replicate.html

Supplemental Figure 8: Non-metric Multi-Dimensional Scaling plot (NMDS) showing patterns of similarity among targeted SRM peptide abundances between deployment locations, where each point represents one geoduck. Relative proximity of points represents overall degree of peptide abundance similarity.