Mapping human pressures across the planet uncovers anthropogenic threat complexes

D.E. Bowler[1-4*], A.D. Bjorkman[5], M. Dornelas[6], I.H. Myers-Smith[7], L. M. Navarro[2,8], A. Niamir[1], S.R. Supp[9], C. Waldock[10], M. Vellend[11], S. A. Blowes[2,12], K. Böhning-Gaese[1], H. Bruelheide[8,2], R. Elahi[13], L.H. Antão[6,14], J. Hines[2,15], F. Isbell[16], H.P. Jones[17], A.E. Magurran[6], J. S. Cabral[2,18], M. Winter[2,19], & A.E. Bates[10,20]

[1]Senckenberg Biodiversity and Climate Research Centre, Senckenberganlage 25, D-60325 Frankfurt am Main, Germany.

[2]German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany

[3]Institute of Biodiversity, Friedrich Schiller University Jena, Dornburger Straße 159, 07743 Jena, Germany

[4]UFZ – Helmholtz Centre for Environmental Research, Department of Ecosystem Services, Permoserstr. 15, 04318 Leipzig, Germany

[5]Section for Ecoinformatics & Biodiversity, Department of Bioscience, Aarhus University, DK-8000 Aarhus, Denmark.

[6]Centre for Biological Diversity, University of St Andrews, Sir Harold Mitchell Building, St Andrews, KY16 9TH Scotland, UK.

[7]School of GeoSciences, University of Edinburgh, Edinburgh EH9 3FF, UK

[8]Institute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle–Wittenberg, Am Kirchtor 1, 06108 Halle, Germany.

[9]Denison University, Data Analytics Program, Granville, OH 43023 USA

[10]Ocean and Earth Science, National Oceanography Centre Southampton, University of

Southampton, Waterfront Campus, European Way, Southampton, SO14 3ZH, United Kingdom

[11]Université de Sherbrooke, Sherbrooke, QC Canada J1K 2R1

[12]Department of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel

Aviv 69978, Israel

[13]Hopkins Marine Station, Stanford University, Pacific Grove, CA 93950, USA

[14]Department of Biology and CESAM, Universidade de Aveiro, Campus Universitário de

Santiago, 3810-193, Aveiro, Portugal

[15]Institute of Biology, Leipzig University, Deutscher Platz 5e, 04103 Leipzig, Germany

[16]Department of Ecology, Evolution, and Behavior, University of Minnesota Twin Cities, Saint

Paul, Minnesota 55108, USA

[17]Department of Biological Sciences and Institute for the Study of the Environment,

Sustainability, and Energy. Northern Illinois University, DeKalb, IL 60115, USA.

[18]Ecosystem Modeling, Centre for Computational and Theoretical Biology, University of

Würzburg, Würzburg, Germany

[19]Leipzig University, Ritterstraße, 26, 04109 Leipzig, Germany

[20]Department of Ocean Sciences, Memorial University of Newfoundland, St. John's A1C 5S7,

Canada

*Correspondence to: diana.e.bowler@gmail.com

Abstract

1. Climate change and other anthropogenic drivers of biodiversity change are unequally distributed across the world. Overlap in the distributions of different drivers have important implications for biodiversity change attribution and the potential for interactive effects. However, the spatial relationships among different drivers, and whether they differ between the terrestrial and marine realm has yet to be examined.

2. We compiled global gridded datasets on climate change, land-use, resource exploitation, pollution, alien species potential, and human population density. We used multivariate statistics to examine the spatial relationships amongst the drivers and to characterize the typical combinations of drivers experienced by different regions of the world.

3. We found stronger positive correlations among drivers in the terrestrial than in the marine realm, leading to areas with high intensities of multiple drivers on land. Climate change tended to be negatively correlated with other drivers in the terrestrial realm (e.g., in the tundra and boreal forest with high climate change but low human use and pollution), whereas the opposite was true in the marine realm (e.g., in the Indo-Pacific with high climate change and high fishing).

4. We show that different regions of the world can be defined by anthropogenic threat complexes (ATCs), distinguished by different sets of drivers with varying intensities. The ATCs can be used to test hypotheses about patterns of biodiversity change, especially in response to the joint effects of multiple drivers. More generally, our global analysis highlights the broad conservation priorities needed to mitigate the impacts of

64   anthropogenic change, with different priorities emerging on land and in the ocean, and in

65   different parts of the world.

Introduction

Human activities are reshaping biological communities and impacting ecosystem functioning across the Earth (Barnosky *et al.* 2011; Dornelas *et al.* 2014; Isbell *et al.* 2017). Meeting the global challenge of the conservation and sustainable use of nature requires not only quantifying biodiversity change, but also identifying the underlying causes of change (Tittensor *et al.* 2014; Isbell *et al.* 2017). Climate change, habitat change, exploitation, pollution and invasive alien species have been recognized as the most important and widespread direct causes of biodiversity change (Pereira, Navarro & Martins 2012; IPCC 2013; IPBES 2019). These five main drivers have been linked with changes in multiple dimensions of biodiversity, including genetic diversity, species' population sizes and the functional diversity of communities (Pereira, Navarro & Martins 2012). The impacts of these drivers on a biological community in a given region critically hinge on the extent of exposure to each driver, which refers to its local magnitude or change (such as the strength of climate change or intensity of land-use). An important, but so far underexplored, step towards understanding the global patterns of biodiversity change is characterizing the exposure patterns of biological communities to different types of environmental change.

Global maps of pressures such as the terrestrial human footprint (Sanderson *et al.* 2002; Venter *et al.* 2016), marine pressures (Halpern *et al.* 2008; Halpern *et al.* 2015a) and river threats (Vorosmarty *et al.* 2010) characterize the geographic hotspots of anthropogenic threats to biodiversity. These maps have estimated that at least 75% of terrestrial land has been exposed to some sort of land-use change (Venter *et al.* 2016), while the marine realm has been exposed to multiple pressures (Halpern *et al.* 2015a). The intensity of the terrestrial human footprint has been linked with spatial variation in ecological processes; for instance, reduced animal

movement was found in areas with a higher human footprint across different species (Tucker *et*

al. 2018). However, these global maps show the summed pressure of different drivers related to

human activities and ignore any relationships among them. Areas identified as having high

human pressure could be underlaid by different combinations of drivers with varying intensity,

each of which may have contrasting impacts on biodiversity. Hence, while useful, these

cumulative threat maps do not help understand the relative contribution of each driver to global

biodiversity change, nor the potential for interactive impacts among drivers.

The relative importance of different drivers for biodiversity change and ecosystem

services is a key component of both policy-oriented assessments such as IPBES framework

(Diaz *et al.* 2015) and conservation targets such as CBD Aichi Biodiversity Targets (Tittensor *et*

al. 2014). Hence, unpacking the spatial patterns of exposure of different drivers, and assessing

the extent to which they jointly act on communities, is an important line of research. Many

studies have examined the relationships between different anthropogenic drivers and biodiversity

change, such as change in biological communities (richness, composition and population

abundances), but they usually focus on only one or two drivers, such as climate change and land

use change (Sirami *et al.* 2017). Multiple drivers may be associated with each other, either

coincidentally (due to shared causes) or causally (when one driver affects the intensity of

another) (Geary *et al.* 2019). Moreover, when drivers co-occur, their impacts on communities

may be additive, or interact synergistically (total impact stronger when together) or

antagonistically (total impact weaker when together) (Cote, Darling & Brown 2016; Sirami *et al.*

2017). Nonetheless, few studies have examined the effects of multiple drivers on biodiversity

(Sirami *et al.* 2017; Mazor *et al.* 2018).

Here, we analyzed the spatial relationships among variables related to the main drivers of

biodiversity change, and show how they overlap in different biogeographic regions across the

entire surface of the world. The motivation for our study was primarily to advance the study of

the attribution of biodiversity change, especially regarding potential drivers of long-term changes

in species' populations and communities (Dornelas *et al.* 2014; Bowler *et al.* 2017; Dornelas *et*

al. 2019). Hence, our questions were focused towards the relevance of the spatial patterns of

drivers for biodiversity change rather than on explaining the spatial patterns in drivers

themselves. We determined the extent of driver overlap to assess the potential for different

drivers to either act alone on communities, rendering their specific fingerprints easier to detect,

or for drivers to act in combination, with the potential for interactive effects. For many drivers, it

can be hypothesized that exposure patterns may be inter-linked due to related local or regional

human activities, driven by local human population density (Ellis *et al.* 2010; Geary *et al.* 2019).

In contrast, climate change is expected to be distributed differently to other variables because it

is an outcome of processes at regional and global scales (IPCC 2013).

Towards these aims, we compiled global spatial gridded datasets on variables that

characterize different anthropogenic drivers (Tables 1 and S1–S2). Although the specific

variables differ among realms, we aligned each variable to the dominant drivers that are common

across both realms (Table 1). We quantified the strengths of the spatial relationships among the

intensities of the different variables related to climate change, habitat conversion and

exploitation (grouped together as 'human use'), pollution and potential for alien species

immigration. Based on these relationships, we defined 'anthropogenic threat complexes' that

typify the combinations of drivers impacting different regions of the world. Studies mapping

drivers of biodiversity change have so far considered the terrestrial and marine realms separately.

By employing a standardized analysis for both realms, our study also highlights similarities and

differences in anthropogenic environmental changes across the world, including across realms.

Methods

Approach to data selection

By drivers, we refer to anthropogenic drivers only and not biological processes, such as dispersal

and demographic rates, that more proximately lead to changes in species' abundances. Our

analysis required datasets that were (1) global; (2) spatially gridded at high-resolution and (3)

publicly available. We selected datasets on variables related to the main five drivers that were

included in previous studies of global environmental change in the context of biodiversity

change (Table 1) (Sanderson *et al.* 2002; Halpern *et al.* 2008). We further searched for data on

other relevant variables following the IUCN threats categories (Table S1) (Salafsky *et al.* 2008).

As we focused on a land versus ocean comparison, we did not specifically consider freshwater

threats (Vorosmarty *et al.* 2010). Variables representing drivers of biodiversity change were in

most cases based on data between 1990 and 2010 (except climate change, and forest loss; see

below). In total, we had datasets for 16 driver-related variables (Table 1). The terrestrial datasets

came from various sources (Table S2). Most of the marine datasets came from the landmark

study of Halpern et al. (Halpern *et al.* 2008). For interpretation and presentation purposes,

variables were grouped into five broad categories: climate change, habitat conversion,

exploitation, pollution and alien species potential. Because habitat conversion and exploitation

were difficult to classify separately across terrestrial and marine ecosystems, we combined both

into a single "human use" category.

Climate change

Climate change has multiple components (IPCC 2013), hence we characterized climate change

by several variables using global spatiotemporal gridded temperature data for the terrestrial

(Harris *et al.* 2014) and marine realm (Rayner *et al.* 2003). To calculate temperature trends, we

used data between 1950, proposed as the start of the Anthropocene (Waters *et al.* 2016), and

2010. Temperature trends were estimated by fitting simple linear regression models to annual

temperature means of each grid cell and extracting the coefficient for the effect of year.

Temperature divergence was inferred from the t-statistic of this linear regression, i.e., the trend

divided by its standard error, representing the significance of the trend. Velocity of climate

change (Loarie *et al.* 2009) was calculated as the ratio between the temporal temperature trend

and the local spatial gradient in temperature. Trends of extreme temperatures were calculated by

whichever was largest of the temporal trends in mean temperature of the warmest or coolest

month. We further included aridity for the terrestrial realm and ocean acidification for the marine

realm. The aridity trend was estimated by taking monthly and annual datasets on potential

evapotranspiration and precipitation, and calculating the monthly:annual ratio (Zorner *et al.*

2008), followed by the temporal trend of the annual monthly average of this ratio. Ocean

acidification, from the Halpern layers, was based on the change in aragonite saturation state

between 1870 and 2000–2009 (Halpern *et al.* 2008).

Human use

In the terrestrial realm, we collated human use variables related to different types of land

conversion or use: cropland, pasture land, cattle density, urban area and forest loss. Data on crop

land, pasture land and urban/built land-cover were taken from different databases, primarily

based on satellite imagery – crop land (Fritz *et al.* 2015), pasture land (Ramankutty *et al.* 2008)

and urban land (Friedl *et al.* 2010). We additionally included information on forest loss since

deforestation itself is a recognized threat (Barlow *et al.* 2016). Forest loss, based on FAO (Food

and Agriculture Association of the United Nations) wood harvest statistics, was calculated as the

loss of primary forest for the same time frame as our climate change statistics, i.e., between 1950

and 2010 (Hurtt *et al.* 2011). We also included data on cattle density, which was based on sub-

national livestock data that were statistically downscaled using multiple predictors (Robinson *et*

al. 2014). In the marine realm, human use variables were based on different commercial fishing

activities separated by gear types (e.g., dredging or cast nets), which determine their selectivity

and impact on the surrounding seascape (Halpern *et al.* 2008). These fishing types were pelagic

low-bycatch, pelagic high-bycatch, demersal habitat-modifying, demersal non-habitat-modifying

low-bycatch, and demersal non-habitat-modifying high bycatch. These data were based on FAO

and other commercial catch data sources and downscaled based on an ocean productivity model

(Halpern *et al.* 2008).

Pollution

Nitrogen from both fossil fuel combustion and agriculture is one is the biggest pollutants

impacting biodiversity (De Schrijver *et al.* 2011). We included data on nitrogen pollution for the

terrestrial realm in the form of atmospheric nitrogen (Dentener 2006) and fertilizer use (Potter *et*

al. 2010), and for the marine realm as fertilizer use (Halpern *et al.* 2008). We also included data

on pesticide use in both realms (Halpern *et al.* 2008; Vorosmarty *et al.* 2010), another important

component of agricultural intensification that negatively affects biodiversity (Geiger *et al.* 2010).

Country-specific estimates of fertilizer use and pesticide were downscaled to a raster grid by the

data providers according to crop land-cover maps; thus, these datasets were not fully independent

of the cropland data, but they still represent the best current knowledge of the spatial distribution

of these variables. We also included a layer estimating the extent of ocean pollution, based on the

distribution of shipping lanes (Halpern *et al.* 2008). Finally, we included night-time light

pollution detected by satellite imagery (Halpern *et al.* 2015a), which was also included in

previous terrestrial and marine threat maps (Halpern *et al.* 2015a; Venter *et al.* 2016).

Alien species potential

The spread of alien species is among the greatest threats to biodiversity and ecosystem services

(Blackburn, Bellard & Ricciardi 2019). Alien species are defined as species that are introduced

into areas beyond their historical range, usually through human transport, accidentally or

incidentally (Hulme 2009; Seebens *et al.* 2015). We used information on transport infrastructure

related to human movement and trade that depict possible species transportation pathways and

vectors (Hulme 2009; Davidson *et al.* 2018). Specifically, we used spatial datasets of transport

connectivity (including data on road and rail networks and navigable rivers) in the terrestrial

realm and cargo volume at ports in the marine realm (Table S1). While these do not represent

the only invasion pathways, they are commonly accepted proxies for human-mediated propagule

pressure, which is known to be among the most important determinant of alien species

establishment (Hulme 2009; Seebens *et al.* 2015).

We note two limitations of our alien species data. First, our proxy was for alien species in

general and not specifically for invasive and harmful alien species. Only a proportion of alien

species become invasive, but this proportion differs among taxa (Jeschke & Pyšek 2018).

Although the most harmful alien species have been identified for some regions (Katsanevakis *et*

al. 2014), this is an on-going process at the global level (Pagad *et al.* 2018). Hence, we refer to

this variable as alien species potential rather than specifically as invasive alien species. Second,

we only used a proxy for alien species propagule pressure and not direct measures of their

distribution. Our analysis required high-resolution global gridded maps, but these are not

available for alien species distributions for neither realm. However, information on alien species

distribution was available at a regional, sub-national and national levels for some taxonomic

groups, including birds (Dyer *et al.* 2017) and plants (van Kleunen *et al.* 2019). To assess the

validity of our connectivity proxy, we used these datasets, representing taxa with low and high

mobility, to test the correlation between alien species richness and mean connectivity at the

spatial scale of the distribution data. We found a significant rank correlation for both datasets

(birds, $\rho=0.42$; plants, $\rho=0.46$, see Fig. S1 for more details), suggesting our proxy represents a

reasonable approximation of alien species potential.

Human population

We also included "human population density" as a separate driver (CIESIN 2017) accounting for

the effects of human activities not falling into the other categories (Salafsky *et al.* 2008).

Including human population density also allowed quantitative tests of the relationship between

human population density and the other drivers.

Justification for layer exclusion

We did not use data for some variables that were previously included in the terrestrial human

footprint or the Halpern layers. The human footprint includes data on roads, railways and

navigable waterways (Venter *et al.* 2016). Although we did not separately include these data,

these data has been taken into account in the terrestrial connectivity variable (for invasions). In

the marine realm, we excluded a shipping lane variable since the ocean pollution variable was

already based on the distribution of shipping lanes (Halpern *et al.* 2008). Additional available

marine layers that we did not use were: UV radiation, oils rigs (based on night lights, already

included), inorganic pollution (highly correlated with other land-based coastal pollutants that

were already included) and artisanal fishing (data poor and mostly modelled) (Halpern *et al.*

2008).

Geographic region data

To understand how the driver distributions compared with classic biome distributions that are

already used in biodiversity research, we also extracted geographic region data on biomes. Data

on the spatial distribution of terrestrial biomes (including deserts, temperate broadleaf forest and

boreal forest among other) were taken from WWF (Olson *et al.* 2001). Marine regions were

defined by combining coastal/shelf region polygon data – MEOW (Spalding *et al.* 2007) and

ocean polygon data (naturalearthdata.com). We did not use marine ecosystem data as used by

others (Halpern *et al.* 2015a) because the ecosystems spatially overlapped in our 2-D global

raster grid, when, in reality, these different ecosystems occur at different depths in the water

column.

Data processing

We harmonized each dataset to a standard global grid. The resolutions of the original datasets
were approximately at a 100 km square grid (or 1°) or finer resolution; hence, we aggregated all
datasets to a standard grid of 100 km square grid cells by taking the mean value of the grid cells.
Atmospheric nitrogen deposition was only available at a coarser resolution (see Table S1);
however, we disaggregated this also to 100 km. Datasets were bound between latitudes of -58
and 78 (due to poorer data at extreme latitudes) and re-projected onto a common equal-area map
projection (Eckert IV; ESPG = 54012). Missing values in some of the human activity datasets
(primarily in pesticide use) occurred in remote regions (e.g., very high latitudes and deserts) with
likely absent or low variable values (since they were in areas with no crop cover) and were
imputed as zero. Greenland was excluded due to missing data in several of the datasets.

Because each dataset comprised data in different units (e.g., temperature data in °C and
fertilizer data in kg/ha), it was not possible to directly compare their absolute values. Instead, we
ranked scaled (between 0 and 1) values of each dataset for ease of interpretation (Fig. S2 shows
the distributions of the original values of each variable and Fig. S3 shows global maps of the
ranked and scaled data). This processing also reduced the large skew in the absolute values of
many datasets. For most datasets, larger values reflected a greater magnitude, and thus higher
potential exposure of biodiversity to that variable. Transformations were needed in only one case
– we inverted terrestrial accessibility (i.e., $values^{-1}$). For climate change metrics, we only used
positive values to focus on the drivers of warming and drying. By using ranks, we avoided
making any complex assumptions about the relationships between the absolute values of each

driver variable and its impact on organisms. We rather assumed that all variables were similarly

important and that higher variable values would have a stronger impact on biodiversity.

Data analysis

To examine the relationships among the intensities of the 16 different variables, we calculated

Spearman's rank correlation coefficients (ρ) for each pairwise combination of variables across all

grid cells in each realm. We chose this statistic because it is equivalent to the commonly used

Pearson's correlation on ranked data, which was consistent with our aforementioned data

processing. We used Dutilleul's modified t-test to account for spatial autocorrelation before

testing the significance of the correlations (Dutilleul, Pelletier & Alpargu 2008). We also used

Moran's I and correlograms to assess the extent of spatial autocorrelation within each variable.

For the marine realm, correlations were also examined separately for grid cells whose centroid

overlapped with oceanic or coastal regions. To assess the importance of the drivers in different

parts of the world, we compared the average intensity of each of the five main drivers (climate

change, human use, human population, pollution and alien species potential) for different

geographic regions. We first calculated the mean driver value for each grid cell, i.e., averaging

across associated driver-related variables (Table 1). We then plotted the distribution of these

mean values across all grid cells within each terrestrial biome and marine region.

We used k-medoid clustering, with the partitioning around the medoids algorithm with

Euclidean distances (Maechler *et al.* 2018), for clustering grid cells according to their extent of

exposure of all the variables. To focus the clustering on the main axes of variation, we first used

PCA to reduce the number of variables in driver groups with multiple variables, i.e., climate

change, human use and pollution. We performed the clustering with two or three PCA

components (whichever captured at least 75% of the variation) for each of these drivers along

with the variables for human population density and potential alien species. We selected the

number of clusters by comparing the changes in cluster silhouette width with increasing cluster

number; however, we limited the cluster number to <10. To slightly smooth the maps, we used a

moving window to assign each cell the mode of its 3 x 3 cell neighborhood. Although, driver

combinations vary in a continuous manner, we chose a clustering method that produces discrete

grouping to provide the simplest description of the main groupings in the data. Finally, to repack

the datasets into cumulative driver maps, we summed the number of variables for which each

grid cell was in the upper 10% of values (based on values greater than zero). Analyses were run

in R v. 3.4.1 (R Core Team 2018), using the packages raster (Hijmans 2017), SpatialPack

(Vallejos, Osorio & Bevilacqua 2018) and cluster (Maechler *et al.* 2018).

Sensitivity analyses

To examine the effect of the grain size, we repeated the data processing steps except harmonized

the datasets to a global grid of 500 km resolution and repeated the correlation analysis (similar

results were obtained – see Fig. S4). To check the effects of ranking the data values, we repeated

the data processing steps by logging the values (to the base 10) rather than ranking them, after

bounding extreme values to the upper and lower 2.5% quantiles. This alternative data

transformation does not affect the correlation coefficients because Spearman's correlations only

uses the ranks of the data. We repeated our remaining analysis with this alternative

transformation, calculating the average variable intensities for different terrestrial and marine

regions, and the clustering analysis (generally similar results were obtained – see Fig. S5 and

S6). Since the distributions are still skewed after logging instead of ranking, the patterns are

strongly affected by extreme values with this approach, especially in the marine realm.

Results

We found that drivers of biodiversity change were more spatially coupled in the terrestrial than

in the marine realm (Fig. 1, Fig. S7). On land, 40% of the possible pairwise relationships

between variables (excluding climate change-related variables) showed positive correlation

strengths of at least 0.7. Thus, terrestrial areas with high intensities of one variable also tended to

have high intensities of other variables. Moreover, correlations were found between different

types of drivers; for instance, high crop land-cover was associated with high pollution, high alien

potential and high human population density. Conversely, in the marine realm, we found fewer

correlations – only 15% of the possible pairwise relationships (excluding climate change-related

variables) showed a strong positive correlation (> 0.7) – and these relationships were mostly

within, rather than between, different driver types; for instance, among different types of human

use (e.g., different types of demersal fishing; Fig. 1). Across all variables, oceanic regions

showed fewer correlations compared to coastal regions (Fig. S8). Spatial autocorrelation was

present in all variables and tended to reach greater distances in the marine human-uses and

climate-change variables (Figs S9 and S10), and shorter distances in the coastal-based marine

variables. However, the correlations among drivers were statistically significant (all $P<0.05$ for

links shown in Fig. 1) after accounting for autocorrelation. In neither realm were there strong

negative correlations among variables (Fig. S11 shows the full correlation matrix).

Strong correlations between climate change and non-climatic drivers (human use,

pollution and alien species) were not observed in either realm (Fig. 1, Fig. S11); however, there

were still significant weak correlations, with the direction of these correlations differing

systematically between realms (Fig. 2). Temperature change, and its velocity, was negatively

associated with the average intensity of other variables in the terrestrial realm (Fig. 2;

temperature change: $\rho = -0.26$, $P<0.01$; velocity: $\rho = -0.22$, $P<0.01$), but positively associated

with the average intensity of other variables in the marine realm (Fig. 2; temperature change: $\rho =$

0.20, $P=0.06$; velocity: $\rho = 0.24$, $P<0.05$). Pollution, cattle density and human population density

were especially negatively correlated with temperature change in the terrestrial realm while

human use (demersal and pelagic fishing) was especially positively correlated with temperature

change in the marine realm (Fig. 2).

Consistent with the negative association between climate change and human use in the

terrestrial realm, terrestrial biomes exposed to some of the strongest climate change, such as the

tundra, boreal forest and desert, have experienced the lowest intensities of human use (Fig. 3,

Fig. S12). By contrast, other terrestrial biomes, such as tropical dry broadleaf forest and tropical

coniferous forest, have had high intensities of human use, pollution and invasions but lower

intensities of climate change. Temperate broadleaf and mixed forest is distinct in the terrestrial

realm by experiencing higher than average intensities of all drivers (Fig. 3). In the marine realm,

central and western Indo-Pacific were most exposed to multiple drivers, including rapid climate

change and multiple human uses, especially fishing (Fig. 3). The temperate Northern Atlantic,

which includes the North Sea, has been also strongly exposed to multiple drivers. By contrast,

the Pacific Ocean been exposed to the lowest intensities of multiple drivers.

The cluster analysis defined five terrestrial and six marine regions according to their

similarity of exposure to the different driver variables (Fig. 4). These exposure patterns can be

regarded as 'anthropogenic threat complexes' (ATC) that characterize the typical combinations

of environmental change emerging at large scales. ATCs I and VI (dark blue-grey regions in Fig.

4) represent terrestrial and marine areas with high exposure to multiple drivers, especially non-

climatic drivers such as human use, pollution and alien potential. ATCs II and VII are regions

(medium blue-grey) exposed to medium intensities of many variables, including climate change.

ATCs IV and IX (red regions) were similarly exposed to medium intensities of many variables,

but higher intensities of climate change. ATCs III and VIII (light blue) were exposed to

relatively weaker climate change than to non-climatic drivers; by contrast, ATCs V and X

(orange regions) were exposed to relatively greater climate change than to non-climatic drivers.

ATC XI represents areas (light orange) that were exposed to low intensities of most variables

and was only found in the marine realm. The largest terrestrial ATC was V (34% of terrestrial

grid cells), covering regions exposed to medium-high climate change and lower intensities of

other drivers. The largest marine ATC was IX (23% of marine grid cells), including regions

exposed to both medium human use, especially fishing, and medium-high climate change.

The global maps (Fig. 5) highlight the areas exposed to high intensities of multiple

drivers and connect the ATCs to previous cumulative human impact maps produced separately

for the terrestrial (Sanderson *et al.* 2002; Venter *et al.* 2016) and marine realms (Halpern *et al.*

2008; Halpern *et al.* 2015a). Terrestrial regions with the highest cumulative intensities across

multiple drivers tended to be within ATC I, covering parts of Europe, Northeastern America,

India as well as Northern and Eastern China (Fig. 5). In the marine realm, the highest cumulative

intensities were found within ATCs VI and IX, especially in the North Sea of the Northern

Atlantic, East China Sea and central Indo-Pacific. Terrestrial regions with the lowest cumulative

intensities of multiple drivers were commonly within ATCs III and V, covering parts of South

America, Africa, Russia and Australia. In the marine realm, the lowest cumulative intensities

were within ATCs X and XI, in the Indian Ocean, the southern Atlantic Ocean and Pacific

Ocean.

Discussion

Our findings suggest that many drivers of biodiversity change are unlikely to act alone, but rather

to jointly impact biological communities. In the terrestrial realm, drivers that tend to be

simultaneously present in a region were associated with agricultural activities (including land

conversion to cropland, use of land for cattle rearing, use of pesticides/fertilizer), human

population density (urban cover and light pollution) and potential for alien species. These drivers

have especially high spatial overlap in the temperate broadleaf and mixed forests of eastern

North America, Europe and China. In the marine realm, drivers that tend to spatially overlap

include different types of pollutants and the potential for alien species, with the coastal regions of

the Indo-Pacific and Northern Atlantic (especially the North Sea) showing the highest overlap.

Taken together with recent studies that highlight the lack of research on interactive effects of

drivers (Sirami *et al.* 2017; Mazor *et al.* 2018), our findings emphasize the widespread relevance

of further research on understanding how multiple drivers combine to impact biological

communities.

Correlations among drivers were more frequent in the terrestrial realm than in the marine

realm. Spatial relationships among drivers in the terrestrial realm were likely based on different

activities supporting proximal human populations, leading to simultaneous pressures from land-

use change and pollution (Ellis *et al.* 2010). Indeed, Venter et al. (2016) already linked spatial

variation in the human footprint to land suitable for agriculture. By contrast, in the marine realm,

different types of human uses were based on different fishing activities that largely occur in

different areas, leading to lower spatial overlap among them. For instance, demersal fisheries,

such as bottom-trawling that disturbs the seabed, mostly occurs over the continental shelf; while,

pelagic fisheries, such as longline fishing for species such as tuna, can be either continental or

oceanic. Coastal regions were intermediate in patterns between terrestrial and oceanic regions,

suggesting that local human populations contribute to the differences between the two realms

(Halpern *et al.* 2015b). Different histories of human activities in the terrestrial and marine realm

likely also plays a role. Human exploitation in the terrestrial realm is more advanced, spanning

millennia, while technological advances have only recently allowed greater exploitation of the

world's oceans (Knapp *et al.* 2017).

Climate change emerged from our analysis as a spatially distinct driver of biodiversity

change. In both realms, climate change was only weakly associated with other drivers (human

use, pollution and alien species), as expected based on the broad spatial scale at which carbon

emissions affect climate. This finding suggests climate change impacts should be easier to isolate

in statistical models from the impacts of other drivers, and that fingerprints of climate change

impacts on species abundances, range limits and community compositions (Parmesan & Yohe

2003; Poloczanska *et al.* 2013) may be easier to detect than those of other drivers. In the

terrestrial realm, climate change tended to be greater in areas with lower human use. Indeed,

high-latitude terrestrial regions that are experiencing pronounced climate change (IPCC 2013;

Pithan & Mauritsen 2014) have historically undergone less human settlement and agriculture. In

areas where non-climatic drivers are weak, climate change has the potential to be the dominant

driver of change, for instance in deserts, tundra and boreal forests of the terrestrial realm and in

the south Atlantic Ocean and Indian Ocean of the marine realm.

Our classification of Anthropogenic Threat Complexes helps regard anthropogenic

environmental change as a series of at least 11 large-scale natural experiments across the globe.

The ATCs highlight where future research on the joint impact of multiple drivers, and their

potentially interactive effects, could be most usefully directed to have the widest relevance for

the study of biodiversity change. They suggest, for instance, that interactive effects between

climate change and fishing should be further studied in the Indo-Pacific and North Sea, where

both rapid temperature change and intense fishing have been occurring (Ramirez *et al.* 2017). In

the terrestrial realm, biological communities in Europe may be of special interest to study the

joint impacts of climate change, land use and pollution on different ecosystems because of the

strong intensities of multiple drivers.

Attribution of observed biodiversity change to the underlying anthropogenic drivers may

be most successful if focused on complexes of environmental change, rather than on each

variable individually. Because of the correlations among drivers, relationships between a driver

and a metric of biodiversity change may in fact be caused by a correlated driver. Although

spatial heterogeneity at smaller spatial scales (e.g., neighboring sites that differ in only one

driver, such as land-use change) can estimate the local effect of drivers such as habitat

conversion (Newbold *et al.* 2015), correlated large-scale drivers can affect regional species pools

and hence still influence local community dynamics (Harrison & Cornell 2008). Analysis of the

relationship between ATCs and species' declines may help identify the most harmful

combinations of drivers. The ATCs could be resolved to finer spatial scales and also used to

inform the design of biodiversity observatories that treat driver exposure as a natural experiment.

Observatories could be selected along a given driver gradient, keeping all but this driver

constant. Alternatively, study regions that are most suitable to study a specific driver could be

selected from within geographic clusters dominated by the driver of interest, to reduce the

confounding effects of other drivers in the landscape.

Large-scale information on drivers is most immediately relevant for global conservation

policy (Tittensor *et al.* 2014; IPBES 2019). For biodiversity conservation at specific locations,

local data on the magnitudes of different types of anthropogenic pressures are essential to decide

on the most suitable management plan. However, even in this latter context, there are a number

of advantages of having knowledge on the large-scale patterns of drivers. First, these large-scale

patterns allow local management to be modified according to the wider anthropogenic land- or

seascape context (Harrison & Cornell 2008). Whilst landscape ecology and habitat connectivity

already inform conservation planning, knowledge of all drivers present in the region could

influence local decision-making by enabling prediction of possible future threats and biodiversity

change at local-scales, e.g., large-scale land-use change affecting regional species pools, with

consequences for local community changes through dispersal (Hansen & DeFries 2007).

Secondly, by characterizing regions of the world in terms of the nature of anthropogenic

environmental change, the ATCs suggest how information and data might be pooled and

synthesized across regions, and even across realms. Regions exposed to the same ATC,

regardless of location, would benefit from exchanging knowledge about prioritization strategies

and management of the multiple drivers, as well as implementing cross-border and inter-regional

strategies to minimize their impact (Bonebrake *et al.* 2019).

Our findings are clearly dependent on the included layers – we selected layers based on

their past inclusion in global threat maps and their linkage to IUCN threat categories. However,

many of the recognized IUCN threats to biodiversity (Salafsky *et al.* 2008; Joppa *et al.* 2016)

were not available as high-resolution spatial datasets at a global scale, which was necessary for

our analysis. These threats include energy production and mining, hunting, and other forms of

human disturbance (Salafsky *et al.* 2008). Data availability was especially limited for the marine

realm and the threat from invasive alien species globally. Rather than use proxy variables of

species transport and thus of propagule pressure, spatially-explicit maps of the number of

invasive alien species would have improved our analysis. The Global Register of Introduced and

Invasive Species (www.griis.org) is working towards better knowledge of the distribution of

invasive species (Pagad *et al.* 2018). Our study only focused on spatial patterns and did not

explicitly consider the consequences of changes in human activities over time. Ongoing projects

such as the Copernicus project (http://www.copernicus.eu/) will greatly increase the availability

of high-resolution spatiotemporal datasets on different variables in the coming years, enabling

better attribution of biodiversity change to the underlying drivers.

Quantifying exposure to environmental change is the first step towards determining

which species, in which places, are most impacted by human activities. However, the realized

outcome of different drivers on biodiversity will ultimately depend on a combination of both the

magnitude of exposure to drivers and species' sensitivities to environmental change (Foden *et al.*

2013). While exposure characterizes the amount of environmental change (e.g., temperature

change in °C), sensitivity refers to the tolerance of the biological community to that driver (e.g.,

abundance (or individual fitness) change per °C temperature change). We intentionally focused

on exposure patterns so that our results were not species-specific and therefore potentially

relevant for any taxa or ecosystem. Unlike exposure, sensitivities vary among taxa according to

characteristics such as their life history, traits and niche breadth (Sunday *et al.* 2015) and

therefore need to be examined separately for different taxa. Hence, despite similar exposure

patterns, we can expect a diversity of biodiversity responses within each ATC due to variation in

species' sensitivities.

Our macroecological approach to mapping the drivers of biodiversity change contributes

to the development of broad conservation policy targeted toward the mitigation of specific driver

complexes. Development of conservation strategies that simultaneously attempt to tackle

multiple drivers are likely to be more efficient in the long-term (Bonebrake *et al.* 2019). The

ATC framework emphasizes the fact that multiple drivers usually affect local and regional

biological communities and discourages the prevalent simplistic focus on one or two drivers.

Much more research should focus on understanding the joint impacts of multiple drivers and how

different drivers interact on biological communities. Our findings help direct some of these

future studies by identifying which drivers most commonly overlap, and in which regions of the

world. Finally, by taking a cross-realm approach, we hope to encourage information exchange

across regions of the world that are exposed to similar suites of drivers, regardless of

environmental realm, and the development of joined-up conservation policies across national

borders and the terrestrial-marine interface.

Data availability

Table S2 shows the sources of each dataset and links to where each dataset can be downloaded.

Datasets produced during our analysis (raster layers shown in Figures 4 and 5) are available as

georeferenced TIFF files in the SOM.

Code availability
539

540   R script to harmonize the raster to a standard grid is found here:

541   https://github.com/bowlerbear/harmonizeRasters

542   R script for the subsequent analysis is found here:

543   https://github.com/bowlerbear/geographyDrivers

Acknowledgments:

This paper arose from discussions at the sChange Workshop (www.idiv.de/schange), which was supported by sDiv, the synthesis centre of iDiv, the German Centre for Integrative Biodiversity Research Halle-Jena-Leipzig. We thank all other participants of this workshop for the stimulating discussion and compilation of data that directly fed into this project. We thank the Associate Editor and two anonymous reviewers for comments on the manuscript. We also thank Suzanne Fritz, Mary O'Connor, Bob O'Hara and Gergana Dasklova for comments on a previous version of the manuscript. DB, JSC, JH, SAB & MW were funded by the German Research Foundation (DB: Grant no BO 1221/23-1; JSC, JH, LMN, SAB & MW: via iDiv: FZT 118). SRS was funded by the National Science Foundation, USA (NSF 1400911). LHA was supported by Fundação para a Ciência e Tecnologia, Portugal (POPH/FSE SFRH/BD/90469/2012). ADB was supported by The Danish Council for Independent Research - Natural Sciences (DFF 4181-00565). MD was funded by the Scottish Funding Council (MASTS, grant reference HR09011) and MD & AEM by the ERC project BioTIME (250189) and BioCHANGE (727440). CW was supported by the Natural Environmental Research Council (grant number NE/L002531/).

Author contributions

DB performed the analyses and wrote the first outline of the paper with AEB. All authors designed the study and helped draft the manuscript.

Fig. 1 Strong and positive relationships among anthropogenic drivers of biodiversity change. Each grey link between two variables represents a significant positive correlation with strength >0.7 of the variable intensities across 100 square km grids covering each realm. We find a higher number of correlations between drivers in the terrestrial versus the marine realm.

Fig. 2 Relationships between climate change and non-climatic drivers (human use, pollution and alien species). The length of each bar represents the correlation coefficient between each variable and temperature change (upper bar, air or sea surface temperature – SST) or VOCC (lower bar, velocity of climate change). We find negative correlations in the terrestrial realm and positive correlations in the marine realm. * denotes statistical significance after accounting for spatial autocorrelation.

Fig. 3 Regions of the terrestrial and marine realms are exposed to distinct combinations of drivers. The violin plots show the distribution of values for each driver in each region. Violins with a median greater than the global median of each driver (the dashed zero line; median of biome values) are colored in a darker shade. Regions are presented in declining order of total driver exposure: regions at the top are exposed to the greater intensities of drivers and those at the bottom are exposed to the least. Names of the terrestrial regions were shortened for presentation purposes. Figure S12 shows the full distributions for each individual driver variable in each region as well as gives the full names of the terrestrial regions.

Fig. 4 Geography of the Anthropocene. Different geographic regions of the world are exposed to different Anthropogenic Threat Complexes (labelled I to XI). In the world map, each colour in each realm denotes a region exposed to the same ATC. The bar charts below are a legend for the colours of the map that describe the magnitudes of climate change and non-climatic drivers in each ATC. Similar ATCs across realms are shaded with the same colour. The bars represent the lower and upper quartiles of the intensities of variables within each driver group (climate change or non-climatic drivers that included human use, pollution and alien species potential). Fig. S13 provides the legend showing the full suite of variable intensities in each ATC.

Fig. 5 Regions of the world exposed to high intensities of multiple drivers. The main map

shows the number of the 16 driver variables for which each grid cell was in the highest 10% of

values within each realm. Regions in the darkest orange are exposed to high intensities of

multiple variables, while those in off-white are exposed to lower intensities (i.e., within the 90%

quantile of values) of all variables. The smaller plots below show the same for each of the

separate drivers. Larger versions of the driver plots are presented in Fig. S14.

Table 1 Anthropogenic drivers of biodiversity change and their respective variables based on available global spatial datasets (Tables S1 and S2). Variables in the same line do not necessarily represent the equivalent variable in each realm.

Anthropogenic driver of biodiversity change	Associated variables	
	Terrestrial	**Marine**
Climate Change	Temperature trend Temperature divergence Change in climate extremes Velocity of climate change Aridity trend	Temperature trend Temperature divergence Change in climate extremes Velocity of climate change Ocean acidification
Human use (land/sea use or change, resource extraction, exploitation)	Crop cover Pasture cover Urban cover Forest loss Livestock density	Destructive demersal fishing Low by-catch demersal fishing High by-catch demersal fishing Low by-catch pelagic fishing High by-catch pelagic fishing
Human population density	Population density	Coastal population density
Pollution	Atmospheric nitrogen deposition Nitrogen fertilizer application Pesticide application Light pollution	Ocean pollution Fertilizer coastal pollution Pesticide coastal pollution Light pollution
Alien potential	Connectivity (transport infrastructure)	Port cargo volume

Barlow, J., Lennox, G.D., Ferreira, J., Berenguer, E., Lees, A.C., Mac Nally, R., Thomson, J.R., Ferraz, S.F.D., Louzada, J., Oliveira, V.H.F., Parry, L., Solar, R.R.D., Vieira, I.C.G., Aragao, L., Begotti, R.A., Braga, R.F., Cardoso, T.M., de Oliveira, R.C., Souza, C.M., Moura, N.G., Nunes, S.S., Siqueira, J.V., Pardini, R., Silveira, J.M., Vaz-de-Mello, F.Z., Veiga, R.C.S., Venturieri, A. & Gardner, T.A. (2016) Anthropogenic disturbance in tropical forests can double biodiversity loss from deforestation. *Nature,* **535,** 144-+.

Barnosky, A.D., Matzke, N., Tomiya, S., Wogan, G.O.U., Swartz, B., Quental, T.B., Marshall, C., McGuire, J.L., Lindsey, E.L., Maguire, K.C., Mersey, B. & Ferrer, E.A. (2011) Has the Earth's sixth mass extinction already arrived? *Nature,* **471,** 51-57.

Blackburn, T.M., Bellard, C. & Ricciardi, A. (2019) Alien versus native species as drivers of recent extinctions. *Frontiers in Ecology and the Environment,* **17,** 203-207.

Bonebrake, T.C., Guo, F., Dingle, C., Baker, D.M., Kitching, R.L. & Ashton, L.A. (2019) Integrating Proximal and Horizon Threats to Biodiversity for Conservation. *Trends in Ecology & Evolution*.

Bowler, D.E., Hof, C., Haase, P., Kroncke, I., Schweiger, O., Adrian, R., Baert, L., Bauer, H.G., Blick, T., Brooker, R.W., Dekoninck, W., Domisch, S., Eckmann, R., Hendrickx, F., Hickler, T., Klotz, S., Kraberg, A., Kuhn, I., Matesanz, S., Meschede, A., Neumann, H., O'Hara, R., Russell, D.J., Sell, A.F., Sonnewald, M., Stoll, S., Sundermann, A., Tackenberg, O., Turkay, M., Valladares, F., van Herk, K., van Klink, R., Vermeulen, R., Voigtlander, K., Wagner, R., Welk, E., Wiemers, M., Wiltshire, K.H. & Bohning-Gaese, K. (2017) Cross-realm assessment of climate change impacts on species' abundance trends. *Nature Ecology & Evolution,* **1,** 7.

CIESIN (2017) Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 10. Palisades, NY: NASA Socioeconomic Data and Applications Center (SEDAC). (ed. C.f.I.E.S.I.N.-C.-C. University).

Cote, I.M., Darling, E.S. & Brown, C.J. (2016) Interactions among ecosystem stressors and their importance in conservation. *Proceedings of the Royal Society B-Biological Sciences,* **283**.

Davidson, I.C., Scianni, C., Minton, M.S. & Ruiz, G.M. (2018) A history of ship specialization and consequences for marine invasions, management and policy. *Journal of Applied Ecology,* **55,** 1799-1811.

De Schrijver, A., De Frenne, P., Ampoorter, E., Van Nevel, L., Demey, A., Wuyts, K. & Verheyen, K. (2011) Cumulative nitrogen input drives species loss in terrestrial ecosystems. *Global Ecology and Biogeography,* **20,** 803-816.

Dentener, F.J. (2006) Global Maps of Atmospheric Nitrogen Deposition, 1860, 1993, and 2050. ORNL DAAC. Oak Ridge, Tennessee, USA.

Diaz, S., Demissew, S., Carabias, J., Joly, C., Lonsdale, M., Ash, N., Larigauderie, A., Adhikari, J.R., Arico, S., Baldi, A., Bartuska, A., Baste, I.A., Bilgin, A., Brondizio, E., Chan, K.M.A., Figueroa, V.E., Duraiappah, A., Fischer, M., Hill, R., Koetz, T., Leadley, P., Lyver, P., Mace, G.M., Martin-Lopez, B., Okumura, M., Pacheco, D., Pascual, U., Perez, E.S., Reyers, B., Roth, E., Saito, O., Scholes, R.J., Sharma, N., Tallis, H., Thaman, R., Watson, R., Yahara, T., Hamid, Z.A., Akosim, C., Al-Hafedh, Y., Allahverdiyev, R., Amankwah, E., Asah, S.T., Asfaw, Z., Bartus, G., Brooks, L.A., Caillaux, J., Dalle, G., Darnaedi, D., Driver, A., Erpul, G., Escobar-Eyzaguirre, P., Failler, P., Fouda, A.M.M., Fu, B., Gundimeda, H., Hashimoto, S., Homer, F., Lavorel, S., Lichtenstein, G., Mala, W.A., Mandivenyi, W., Matczak, P., Mbizvo, C., Mehrdadi, M., Metzger, J.P., Mikissa, J.B., Moller, H., Mooney, H.A., Mumby, P., Nagendra, H., Nesshover, C., Oteng-Yeboah, A.A., Pataki, G., Roue, M., Rubis, J., Schultz, M., Smith, P., Sumaila, R., Takeuchi, K., Thomas, S., Verma, M., Yeo-Chang, Y. & Zlatanova, D. (2015) The IPBES Conceptual Framework - connecting nature and people. *Current Opinion in Environmental Sustainability,* **14,** 1-16.

Dornelas, M., Gotelli, N.J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C. & Magurran, A.E. (2014) Assemblage Time Series Reveal Biodiversity Change but Not Systematic Loss. *Science,* **344,** 296-299.

Dornelas, M., Gotelli, N.J., Shimadzu, H., Moyes, F., Magurran, A.E. & McGill, B.J. (2019) A balance of winners and losers in the Anthropocene. *Ecology Letters,* **22,** 847-854.

Dutilleul, P., Pelletier, B. & Alpargu, G. (2008) Modified F tests for assessing the multiple correlation between one spatial process and several others. *Journal of Statistical Planning and Inference,* **138,** 1402-1415.

Dyer, E.E., Cassey, P., Redding, D.W., Collen, B., Franks, V., Gaston, K.J., Jones, K.E., Kark, S., Orme, C.D.L. & Blackburn, T.M. (2017) The Global Distribution and Drivers of Alien Bird Species Richness. *Plos Biology,* **15**.

Ellis, E.C., Goldewijk, K.K., Siebert, S., Lightman, D. & Ramankutty, N. (2010) Anthropogenic transformation of the biomes, 1700 to 2000. *Global Ecology and Biogeography,* **19,** 589-606.

Foden, W.B., Butchart, S.H.M., Stuart, S.N., Vie, J.C., Akcakaya, H.R., Angulo, A., DeVantier, L.M., Gutsche, A., Turak, E., Cao, L., Donner, S.D., Katariya, V., Bernard, R., Holland, R.A., Hughes, A.F., O'Hanlon, S.E., Garnett, S.T., Sekercioglu, C.H. & Mace, G.M. (2013) Identifying the World's Most Climate Change Vulnerable Species: A Systematic Trait-Based Assessment of all Birds, Amphibians and Corals. *Plos One,* **8**.

Friedl, M.A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A. & Huang, X. (2010) MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets, 2001-2012, Collection 5.1 IGBP Land Cover. Boston University, Boston, MA, USA.

Fritz, S., See, L., McCallum, I., You, L.Z., Bun, A., Moltchanova, E., Duerauer, M., Albrecht, F., Schill, C., Perger, C., Havlik, P., Mosnier, A., Thornton, P., Wood-Sichra, U., Herrero, M., Becker-Reshef, I., Justice, C., Hansen, M., Gong, P., Aziz, S.A., Cipriani, A., Cumani, R., Cecchi, G., Conchedda, G., Ferreira, S., Gomez, A., Haffani, M., Kayitakire, F., Malanding, J., Mueller, R., Newby, T., Nonguierma, A., Olusegun, A., Ortner, S., Rajak, D.R., Rocha, J., Schepaschenko, D., Schepaschenko, M., Terekhov, A., Tiangwa, A., Vancutsem, C., Vintrou, E., Wu, W.B., van der Velde, M., Dunwoody, A., Kraxner, F. & Obersteiner, M. (2015) Mapping global cropland and field size. *Global Change Biology,* **21,** 1980-1992.

Geary, W.L., Nimmo, D.G., Doherty, T.S., Ritchie, E.G. & Tulloch, A.I.T. (2019) Threat webs: reframing the co-occurrence and interactions of threats to biodiversity. *Journal of Applied Ecology*.

Geiger, F., Bengtsson, J., Berendse, F., Weisser, W.W., Emmerson, M., Morales, M.B., Ceryngier, P., Liira, J., Tscharntke, T., Winqvist, C., Eggers, S., Bommarco, R., Part, T., Bretagnolle, V., Plantegenest, M., Clement, L.W., Dennis, C., Palmer, C., Onate, J.J., Guerrero, I., Hawro, V., Aavik, T., Thies, C., Flohre, A., Hanke, S., Fischer, C., Goedhart, P.W. & Inchausti, P. (2010) Persistent negative effects of pesticides on biodiversity and biological control potential on European farmland. *Basic and Applied Ecology,* **11,** 97-105.

Halpern, B.S., Frazier, M., Potapenko, J., Casey, K.S., Koenig, K., Longo, C., Lowndes, J.S., Rockwood, R.C., Selig, E.R., Selkoe, K.A. & Walbridge, S. (2015a) Spatial and temporal changes in cumulative human impacts on the world's ocean. *Nature Communications,* **6**.

Halpern, B.S., Longo, C., Lowndes, J.S.S., Best, B.D., Frazier, M., Katona, S.K., Kleisner, K.M., Rosenberg, A.A., Scarborough, C. & Selig, E.R. (2015b) Patterns and Emerging Trends in Global Ocean Health. *Plos One,* **10**.

Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C., Bruno, J.F., Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin, E.M.P., Perry, M.T., Selig, E.R., Spalding, M., Steneck, R. & Watson, R. (2008) A global map of human impact on marine ecosystems. *Science,* **319,** 948-952.

Hansen, A.J. & DeFries, R. (2007) Ecological mechanisms linking protected areas to surrounding lands. *Ecological Applications,* **17,** 974-988.

Harris, I., Jones, P.D., Osborn, T.J. & Lister, D.H. (2014) Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset. *International Journal of Climatology,* **34,** 623-642.

Harrison, S. & Cornell, H. (2008) Toward a better understanding of the regional causes of local community richness. *Ecology Letters,* **11,** 969-979.

Hijmans, R.J. (2017) raster: Geographic Data Analysis and Modeling. R package version
2.6-7. https://CRAN.R-project.org/package=raster.

Hulme, P.E. (2009) Trade, transport and trouble: managing invasive species pathways in an era of globalization. *Journal of Applied Ecology,* **46,** 10-18.

Hurtt, G.C., Chini, L.P., Frolking, S., Betts, R.A., Feddema, J., Fischer, G., Fisk, J.P., Hibbard, K., Houghton, R.A., Janetos, A., Jones, C.D., Kindermann, G., Kinoshita, T., Goldewijk, K.K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D.P. & Wang, Y.P. (2011) Harmonization of land-use scenarios for the period 1500-2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Climatic Change,* **109,** 117-161.

IPBES (2019) Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. IPBES secretariat, Bonn, Germany.

IPCC (2013) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Isbell, F., Gonzalez, A., Loreau, M., Cowles, J., Diaz, S., Hector, A., Mace, G.M., Wardle, D.A., O'Connor, M.I., Duffy, J.E., Turnbull, L.A., Thompson, P.L. & Larigauderie, A. (2017) Linking the influence and dependence of people on biodiversity across scales. *Nature,* **546,** 65-72.

Jeschke, J. & Pyšek, P. (2018) *Ten rules*. CAB International, Wallingford.

Joppa, L.N., O'Connor, B., Visconti, P., Smith, C., Geldmann, J., Mann, M.H., Watson, J.E.M., Butchart, S.H.M., Virah-Sawmy, M., Halpern, B.S., Ahmed, S.E., Balmford, A., Sutherland, W.J., Harfoot, M., Hilton-Taylor, C., Foden, W., Di Minin, E., Pagad, S., Genovesi, P., Hutton, J. & Burgess, N.D. (2016) Filling in biodiversity threat gaps. *Science,* **352,** 416-418.

Katsanevakis, S., Wallentinus, I., Zenetos, A., Leppakoski, E., Cinar, M.E., Ozturk, B., Grabowski, M., Golani, D. & Cardoso, A.C. (2014) Impacts of invasive alien marine species on ecosystem services and biodiversity: a pan-European review. *Aquatic Invasions,* **9,** 391-423.

Knapp, S., Schweiger, O., Kraberg, A., Asmus, H., Asmus, R., Brey, T., Frickenhaus, S., Gutt, J., Kuhn, I., Liess, M., Musche, M., Portner, H.O., Seppelt, R., Klotz, S. & Krause, G. (2017) Do drivers of biodiversity change differ in importance across marine and terrestrial systems - Or is it just different research communities' perspectives? *Science of the Total Environment,* **574,** 191-203.

Loarie, S.R., Duffy, P.B., Hamilton, H., Asner, G.P., Field, C.B. & Ackerly, D.D. (2009) The velocity of climate change. *Nature,* **462,** 1052-U1111.

Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M. & Hornik, K. (2018) cluster: Cluster Analysis Basics and Extensions. R package version 2.0.7-1.

Mazor, T., Doropoulos, C., Schwarzmueller, F., Gladish, D.W., Kumaran, N., Merkel, K., Di Marco, M. & Gagic, V. (2018) Global mismatch of policy and research on drivers of biodiversity loss. *Nature Ecology & Evolution,* **2,** 1071-+.

Newbold, T., Hudson, L.N., Hill, S.L.L., Contu, S., Lysenko, I., Senior, R.A., Borger, L., Bennett, D.J., Choimes, A., Collen, B., Day, J., De Palma, A., Diaz, S., Echeverria-Londono, S., Edgar, M.J., Feldman, A., Garon, M., Harrison, M.L.K., Alhusseini, T., Ingram, D.J., Itescu, Y., Kattge, J., Kemp, V., Kirkpatrick, L., Kleyer, M., Correia, D.L.P., Martin, C.D., Meiri, S., Novosolov, M., Pan, Y., Phillips, H.R.P., Purves, D.W., Robinson, A., Simpson, J., Tuck, S.L., Weiher, E., White, H.J., Ewers, R.M., Mace, G.M., Scharlemann, J.P.W. & Purvis, A. (2015) Global effects of land use on local terrestrial biodiversity. *Nature,* **520,** 45-+.

Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V.N., Underwood, E.C., D'Amico, J.A., Itoua, I., Strand, H.E., Morrison, J.C., Loucks, C.J., Allnutt, T.F., Ricketts, T.H., Kura, Y., Lamoreux, J.F., Wettengel, W.W., Hedao, P. & Kassem, K.R. (2001) Terrestrial ecoregions of the worlds: A new map of life on Earth. *Bioscience,* **51,** 933-938.

Pagad, S., Genovesi, P., Carnevali, L., Schigel, D. & McGeoch, M.A. (2018) Data Descriptor: Introducing the Global Register of Introduced and Invasive Species. *Scientific Data,* **5**.

Parmesan, C. & Yohe, G. (2003) A globally coherent fingerprint of climate change impacts across natural systems. *Nature,* **421,** 37-42.

Pereira, H.M., Navarro, L.M. & Martins, I.S. (2012) Global Biodiversity Change: The Bad, the Good, and the Unknown. *Annual Review of Environment and Resources, Vol 37* (eds A. Gadgil & D.M. Liverman), pp. 25-+.

Pithan, F. & Mauritsen, T. (2014) Arctic amplification dominated by temperature feedbacks in contemporary climate models. *Nature Geoscience,* **7,** 181-184.

Poloczanska, E.S., Brown, C.J., Sydeman, W.J., Kiessling, W., Schoeman, D.S., Moore, P.J., Brander, K., Bruno, J.F., Buckley, L.B., Burrows, M.T., Duarte, C.M., Halpern, B.S., Holding, J., Kappel, C.V., O'Connor, M.I., Pandolfi, J.M., Parmesan, C., Schwing, F., Thompson, S.A. & Richardson, A.J. (2013) Global imprint of climate change on marine life. *Nature Climate Change,* **3,** 919-925.

Potter, P., Ramankutty, N., Bennett, E.M. & Donner, S.D. (2010) Characterizing the Spatial Patterns of Global Fertilizer Application and Manure Production. *Earth Interactions,* **14,** 1-22.

R Core Team (2018) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Ramankutty, N., Evan, A.T., Monfreda, C. & Foley, J.A. (2008) Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles,* **22**.

Ramirez, F., Afan, I., Davis, L.S. & Chiaradia, A. (2017) Climate impacts on global hot spots of marine biodiversity. *Science Advances,* **3**.

Rayner, N.A., Parker, D.E., Horton, E.B., Folland, C.K., Alexander, L.V., Rowell, D.P., Kent, E.C. & Kaplan, A. (2003) Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *Journal of Geophysical Research-Atmospheres,* **108**.

Robinson, T.P., Wint, G.R.W., Conchedda, G., Van Boeckel, T.P., Ercoli, V., Palamara, E., Cinardi, G., D'Aietti, L., Hay, S.I. & Gilbert, M. (2014) Mapping the Global Distribution of Livestock. *Plos One,* **9**.

Salafsky, N., Salzer, D., Stattersfield, A.J., Hilton-Taylor, C., Neugarten, R., Butchart, S.H.M., Collen, B., Cox, N., Master, L.L., O'Connor, S. & Wilkie, D. (2008) A standard lexicon for biodiversity conservation: Unified classifications of threats and actions. *Conservation Biology,* **22**, 897-911.

Sanderson, E.W., Jaiteh, M., Levy, M.A., Redford, K.H., Wannebo, A.V. & Woolmer, G. (2002) The human footprint and the last of the wild. *Bioscience,* **52**, 891-904.

Seebens, H., Essl, F., Dawson, W., Fuentes, N., Moser, D., Pergl, J., Pyšek, P., van Kleunen, M., Weber, E., Winter, M. & Blasius, B. (2015) Global trade will accelerate plant invasions in emerging economies under climate change. *Global Change Biology,* **21**, 4128-4140.

Sirami, C., Caplat, P., Popy, S., Clamens, A., Arlettaz, R., Jiguet, F., Brotons, L. & Martin, J.L. (2017) Impacts of global change on species distributions: obstacles and solutions to integrate climate and land use. *Global Ecology and Biogeography,* **26**, 385-394.

Spalding, M.D., Fox, H.E., Halpern, B.S., McManus, M.A., Molnar, J., Allen, G.R., Davidson, N., Jorge, Z.A., Lombana, A.L., Lourie, S.A., Martin, K.D., McManus, E., Molnar, J., Recchia, C.A. & Robertson, J. (2007) Marine ecoregions of the world: A bioregionalization of coastal and shelf areas. *Bioscience,* **57**, 573-583.

Sunday, J.M., Pecl, G.T., Frusher, S., Hobday, A.J., Hill, N., Holbrook, N.J., Edgar, G.J., Stuart-Smith, R., Barrett, N., Wernberg, T., Watson, R.A., Smale, D.A., Fulton, E.A., Slawinski, D., Feng, M., Radford, B.T., Thompson, P.A. & Bates, A.E. (2015) Species traits and climate velocity explain geographic range shifts in an ocean-warming hotspot. *Ecology Letters,* **18**, 944-953.

Tittensor, D.P., Walpole, M., Hill, S.L.L., Boyce, D.G., Britten, G.L., Burgess, N.D., Butchart, S.H.M., Leadley, P.W., Regan, E.C., Alkemade, R., Baumung, R., Bellard, C., Bouwman, L., Bowles-Newark, N.J., Chenery, A.M., Cheung, W.W.L., Christensen, V., Cooper, H.D., Crowther, A.R., Dixon, M.J.R., Galli, A., Gaveau, V., Gregory, R.D., Gutierrez, N.L., Hirsch, T.L., Hoft, R., Januchowski-Hartley, S.R., Karmann, M., Krug, C.B., Leverington, F.J., Loh, J., Lojenga, R.K., Malsch, K., Marques, A., Morgan, D.H.W., Mumby, P.J., Newbold, T., Noonan-Mooney, K., Pagad, S.N., Parks, B.C., Pereira, H.M., Robertson, T., Rondinini, C., Santini, L., Scharlemann, J.P.W., Schindler, S., Sumaila, U.R., Teh, L.S.L., van Kolck, J., Visconti, P. & Ye, Y.M. (2014) A mid-term analysis of progress toward international biodiversity targets. *Science,* **346**, 241-244.

Tucker, M.A., Bohning-Gaese, K., Fagan, W.F., Fryxell, J.M., Van Moorter, B., Alberts, S.C., Ali, A.H., Allen, A.M., Attias, N., Avgar, T., Bartlam-Brooks, H., Bayarbaatar, B., Belant, J.L., Bertassoni, A., Beyer, D., Bidner, L., van Beest, F.M., Blake, S., Blaum, N., Bracis, C., Brown, D., de Bruyn, P.J.N., Cagnacci, F., Calabrese, J.M., Camilo-Alves, C., Chamaille-Jammes, S., Chiaradia, A., Davidson, S.C., Dennis, T., DeStefano, S., Diefenbach, D., Douglas-Hamilton, I., Fennessy, J., Fichtel, C., Fiedler, W., Fischer, C., Fischhoff, I., Fleming, C.H., Ford, A.T., Fritz, S.A., Gehr, B., Goheen, J.R., Gurarie, E., Hebblewhite, M., Heurich, M., Hewison, A.J.M., Hof, C., Hurme, E., Isbell, L.A., Janssen, R., Jeltsch, F., Kaczensky, P., Kane, A., Kappeler, P.M., Kauffman, M., Kays, R., Kimuyu, D., Koch, F., Kranstauber, B., LaPoint, S., Leimgruber, P., Linnell, J.D.C., Lopez-Lopez, P., Markham, A.C., Mattisson, J., Medici, E.P., Mellone, U., Merrill, E., Mourao, G.D., Morato, R.G., Morellet, N., Morrison, T.A., Diaz-Munoz, S.L., Mysterud, A., Nandintsetseg, D., Nathan, R., Niamir, A., Odden, J., O'Hara, R.B., Oliveira-Santos, L.G.R., Olson, K.A., Patterson, B.D., de Paula, R.C., Pedrotti, L., Reineking, B., Rimmler, M., Rogers, T.L., Rolandsen, C.M., Rosenberry, C.S., Rubenstein, D.I., Safi, K., Said, S., Sapir, N., Sawyer, H., Schmidt, N.M., Selva, N., Sergiel, A., Shiilegdamba, E., Silva, J.P., Singh, N., Solberg, E.J., Spiegel, O., Strand, O., Sundaresan, S., Ullmann, W., Voigt, U., Wall, J., Wattles, D., Wikelski, M., Wilmers, C.C., Wilson, J.W., Wittemyer, G., Zieba, F., Zwijacz-Kozica, T. & Mueller, T. (2018) Moving in the Anthropocene: Global reductions in terrestrial mammalian movements. *Science,* **359**, 466-469.

Vallejos, R., Osorio, F. & Bevilacqua, M. (2018) *Spatial Relationships Between Two Georeferenced Variables: with Applications in R*. Springer, New York.

van Kleunen, M., Pyšek, P., Dawson, W., Essl, F., Kreft, H., Pergl, J., Weigelt, P., Stein, A., Dullinger, S., Konig, C., Lenzner, B., Maurel, N., Moser, D., Seebens, H., Kartesz, J., Nishino, M., Aleksanyan, A., Ansong, M., Antonova, L.A., Barcelona, J.F., Breckle, S.W., Brundu, G., Cabezas, F.J., Cardenas, D., Cardenas-Toro, J., Castano, N., Chacon, E., Chatelain, C., Conn, B., de Sa Dechoum, M., Dufour-Dror, J.M., Ebel, A.L.,

Figueiredo, E., Fragman-Sapir, O., Fuentes, N., Groom, Q.J., Henderson, L., Inderjit, Jogan, N., Krestov, P., Kupriyanov, A., Masciadri, S., Meerman, J., Morozova, O., Nickrent, D., Nowak, A., Patzelt, A., Pelser, P.B., Shu, W.S., Thomas, J., Uludag, A., Velayos, M., Verkhosina, A., Villasenor, J.L., Weber, E., Wieringa, J.J., Yazlik, A., Zeddam, A., Zykova, E. & Winter, M. (2019) The Global Naturalized Alien Flora (GloNAF) database. *Ecology,* **100**.

Venter, O., Sanderson, E.W., Magrach, A., Allan, J.R., Beher, J., Jones, K.R., Possingham, H.P., Laurance, W.F., Wood, P., Fekete, B.M., Levy, M.A. & Watson, J.E.M. (2016) Sixteen years of change in the global terrestrial human footprint and implications for biodiversity conservation. *Nature Communications,* **7**.

Vorosmarty, C.J., McIntyre, P.B., Gessner, M.O., Dudgeon, D., Prusevich, A., Green, P., Glidden, S., Bunn, S.E., Sullivan, C.A., Liermann, C.R. & Davies, P.M. (2010) Global threats to human water security and river biodiversity. *Nature,* **467,** 555-561.

Waters, C.N., Zalasiewicz, J., Summerhayes, C., Barnosky, A.D., Poirier, C., Galuszka, A., Cearreta, A., Edgeworth, M., Ellis, E.C., Ellis, M., Jeandel, C., Leinfelder, R., McNeill, J.R., Richter, D.D., Steffen, W., Syvitski, J., Vidas, D., Wagreich, M., Williams, M., An, Z.S., Grinevald, J., Odada, E., Oreskes, N. & Wolfe, A.P. (2016) The Anthropocene is functionally and stratigraphically distinct from the Holocene. *Science,* **351,** 137-+.

Zorner, R.J., Trabucco, A., Bossio, D.A. & Verchot, L.V. (2008) Climate change mitigation: A spatial analysis of global land suitability for clean development mechanism afforestation and reforestation. *Agriculture Ecosystems & Environment,* **126,** 67-80.