The geography of the Anthropocene differs between the land and the sea

D.E. Bowler[1*], A.D. Bjorkman[2], M. Dornelas[3], I. Myers-Smith[4], L. M. Navarro[5,6], A. Niamir[1], S.R. Supp[7], C. Waldock[8], M. Vellend[9], S. A. Blowes[5,10], K. Böhning-Gaese[1], H. Bruelheide[6,5], R. Elahi[11], L.H. Antão[3,12], J. Hines[5,13], F. Isbell[14], H.P. Jones[15], A.E. Magurran[3], J. S. Cabral[5,16], M. Winter[5,17], & A.E. Bates[8,18]

[1]Senckenberg Biodiversity and Climate Research Centre, Senckenberganlage 25, D-60325 Frankfurt am Main, Germany.

[2]Section for Ecoinformatics & Biodiversity, Department of Bioscience, Aarhus University, DK-8000 Aarhus, Denmark.

[3]Centre for Biological Diversity, University of St Andrews, Sir Harold Mitchell Building, St Andrews, KY16 9TH Scotland, UK.

[4]School of GeoSciences, University of Edinburgh, Edinburgh EH9 3FF, UK

[5]German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany

[6]Institute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle–Wittenberg, Am Kirchtor 1, 06108 Halle, Germany.

[7]Denison University, Data Analytics Program, Granville, OH 43023 USA

[8]Ocean and Earth Science, National Oceanography Centre Southampton, University of Southampton, Waterfront Campus, European Way, Southampton, SO14 3ZH, United Kingdom

[9]Université de Sherbrooke, Sherbrooke, QC Canada J1K 2R1

[10]Department of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv 69978, Israel

[11]Hopkins Marine Station, Stanford University, Pacific Grove, CA 93950, USA

[12]Department of Biology and CESAM, Universidade de Aveiro, Campus Universitário de Santiago, 3810-193, Aveiro, Portugal

[13]Institute of Biology, Leipzig University, Deutscher Platz 5e, 04103 Leipzig, Germany

[14]Department of Ecology, Evolution, and Behavior, University of Minnesota Twin Cities, Saint Paul, Minnesota 55108, USA

[15]Department of Biological Sciences and Institute for the Study of the Environment, Sustainability, and Energy. Northern Illinois University, DeKalb, IL 60115, USA.

[16]Ecosystem Modeling, Centre for Computational and Theoretical Biology, University of Würzburg, Würzburg, Germany

[17]Leipzig University, Ritterstraße, 26, 04109 Leipzig, Germany

[18]Department of Ocean Sciences, Memorial University of Newfoundland, St. John's A1C 5S7, Canada

*Correspondence to: diana.e.bowler@gmail.com

Present address: Norsk institutt for naturforskning – NINA, Postadresse: Postboks 5685 Sluppen, 7485 Trondheim, Norway

Abstract

Climate change and other anthropogenic drivers of biodiversity change are unequally distributed across the world. Despite the implications for biodiversity change, it is unknown if the geographic patterns of drivers differ between the terrestrial and marine realm. Using global datasets on human population density, land and resource exploitation, pollution, species invasions, and climate change, we found stronger positive correlations among drivers in the terrestrial than in the marine realm, leading to areas of especially intense human impact on land. Climate change tended to be negatively correlated with other drivers in the terrestrial realm whereas the opposite was true in the marine realm. We show that different regions of the world are exposed to distinct 'anthropogenic threat complexes', comprising suites of drivers of varying intensities. Our global analysis highlights the broad conservation priorities needed to mitigate the drivers shaping biodiversity responses to anthropogenic change.

Introduction

Human activities are reshaping biological communities and impacting ecosystem functioning across the Earth[1-4]. Meeting the global challenge of the conservation and sustainable use of nature requires not only quantifying biodiversity change, but also identifying the underlying causes of change[5]. An essential first step towards determining these causes is characterizing the exposure patterns of biological communities to environmental change. Considerable effort has been devoted to mapping the magnitude of environmental changes that are affecting biodiversity. Global maps produced by this research, such as the Human Footprint[6-10], have played an important role in highlighting the geographic hotspots of biodiversity threats. However, these maps show the summed pressure of different drivers and ignore any relationships among them. Understanding the relationships among drivers is essential for disentangling the relative importance of climate change and other human drivers for biodiversity change and ecosystem services, which is a key component of both policy-oriented assessments, such as IPBES[11], and conservation targets, such as Aichi Biodiversity Targets[5]. Moreover, studies mapping drivers of biodiversity change[6-10] have so far considered the terrestrial and marine realms separately. Identifying similarities and differences in anthropogenic environmental change across the world, including across realms, would contribute towards a more general understanding of the global pattern of biodiversity change as well as help identify regions over which knowledge and information could be shared and synthesized to mitigate impacts. For the first time, we examined the relationships between drivers of biodiversity change across the entire surface of the world. Based on these relationships, we defined 'anthropogenic threat complexes' that typify the combinations of drivers impacting different regions of the world.

We quantified the strengths of the relationships among the intensities of different variables

related to the dominant direct anthropogenic drivers of biodiversity change – climate change,

habitat conversion and exploitation, pollution and species invasions[12-14]. We conducted our

analysis at the global scale to identify the most general patterns emerging across ecosystems. By

employing a standardized analysis for both the terrestrial and marine realms, we could compare

the patterns in each. Multiple drivers may act in the same areas due to related local or regional

human activities, especially in the terrestrial realm. In contrast, climate change is expected to be

distributed differently than other variables because it is an outcome of processes at regional and

global scales[12]. Based on this, we expected the correlations among drivers to differ, with

implications for the typical combinations of drivers emerging in different regions of the world.

We tested two main hypotheses on these correlations: 1) the intensities of many drivers are more

strongly and positively correlated in the terrestrial compared with marine realm because of closer

proximate human influences and 2) climate change-related variables are spatially decoupled

from variables related to human populations due to the different scale at which the underlying

processes act.

We selected global spatial gridded datasets on variables that characterize dimensions of the

different anthropogenic drivers at some time point between 1950 and 2010 (Tables 1 and S1–

S2). We focused on variables that had previously been deemed of sufficient global importance to

be included in maps of environmental change for each realm[6-9]. Although the specific variables

differ among realms, we aligned each variable to one of the dominant drivers that are common

across both realms (Table 1). We also included several climate change metrics that were not

included in some of the previous global maps[6,7]. For climate change-related variables, we used

time-series data to calculate temporal trends. In most other cases, only single time-slice data

were available, but trends could also be calculated for land cover variables. For simplicity,

habitat conversion and exploitation were grouped together as 'human use'. We ranked and scaled

the values of each variable between 0 and 1 to enable comparison. To investigate the

relationships among these variables, we calculated Spearman's rank correlation coefficients (ρ)

for each pair-wise combination of variables within each realm. We used a modified t-test[15] to

account for spatial autocorrelation. To better understand the causes and consequences of these

correlations, we examined the magnitude of the different drivers in different terrestrial biomes

and marine regions and we used a clustering algorithm to classify regions of the world according

to their pattern of exposure to the different drivers. Finally, we determined which regions of the

world have been exposed to the highest intensities of multiple drivers.

Results

Consistent with our first hypothesis, we found that drivers of biodiversity change were more

spatially coupled in the terrestrial than in the marine realm (Fig. 1). On land, 33% of the possible

pair-wise relationships between variables (excluding climate change-related variables) showed

positive correlation strengths of at least 0.7. Thus, terrestrial areas with high intensities of one

variable also tended to have high intensities of other variables. Moreover, correlations were

found between different types of drivers. High cropland cover was associated with high

pollution, high accessibility, high human population density and rapid increases in urban land

cover. Conversely, in the marine realm, we found fewer correlations – only 15% of the possible

pair-wise relationships (excluding climate change-related variables) showed a strong positive

correlation (> 0.7) – and they were mostly within, rather than between, different driver types; for

instance, among different types of human use (e.g., different fishing types; Fig. 1). Oceanic

regions showed fewer correlations compared to coastal regions (Fig. S3). Spatial autocorrelation

was present in all variables and tended to reach greater distances in the marine human-uses and

climate-change variables (Figs S4 and S5), and shorter distances in the coastal-based marine

variables, but the correlations remained statistical significant (all P<0.05 for links in Fig. 1) after

accounting for autocorrelation.

Climate change emerged from our analysis as a spatially distinct driver of biodiversity change,

with contrasting relationships to other drivers in the terrestrial and marine realms. In neither

realm were there strong correlations between climate change and other drivers (Fig. 1),

supporting our second hypothesis based on the broader spatial scale at which carbon emissions

affect climate[12]. However, we did find some weak correlations, with the direction of these

correlations differing between realms (Fig. 2). Temperature change was negatively associated

with the average intensity of other variables in the terrestrial realm (ρ = -0.25, P<0.01, Fig. 2),

but positively associated with the average intensity of other variables in the marine realm (ρ =

0.20, P<0.05; Fig. 2). Terrestrial biomes exposed to strong climate change, such as the tundra,

boreal forest and deserts, have experienced relatively low human use; while terrestrial biomes,

such as tropical dry broadleaf forest, with high intensities of human use, pollution and invasions

have had lower intensities of climate change (Fig. 3). In contrast, marine areas exposed to strong

climate change have been also strongly exposed to other drivers (Fig. 3). The central and western

Indo-Pacific emerged as regions particularly at risk by being exposed to rapid climate change as

well as multiple human uses.

Using cluster analysis, we defined five terrestrial and six marine regions according to their

similarity of exposure to the different driver variables (Fig. 4). These exposure patterns can be

regarded as 'anthropogenic threat complexes' (ATC) that characterize the typical combinations

of environmental change. ATCs I and VI represent terrestrial and marine areas ranked with

higher exposure to climate change than to other drivers (red regions in Fig. 4), while the reverse

is true for ATCs IV and X (grey regions). ATCs II and III (terrestrial) and VII and VIII (marine)

are regions exposed to relatively high intensities of many variables (orange and blue-grey

regions) while ATCs V and XI (light blue regions) are areas generally exposed to lower

intensities of most variables.

We further used our analysis to produce a fully global map showing areas exposed to high

intensities of multiple drivers (Fig. 5). This helps understand the connection between the ATCs

and previous cumulative human impact maps produced separately for the terrestrial[6,7] and marine

realms[8,9]. Regions with the highest cumulative intensities across all variables tended to be within

ATCs III and IV (terrestrial), areas with especially high pollution, and VIII (marine), coastal

areas with high intensities of almost all drivers. By contrast, regions with the lowest cumulative

intensities include ATCs I and V (terrestrial) and VI and XI (marine), which have lower human

uses, pollution and invasions, but still could have high exposure to climate change.

Discussion

Correlations among drivers have important implications because they indicate that regional

biological communities are often jointly impacted by different pressures. Strong correlations are

also likely to hinder attempts to disentangle the contributions of different drivers to biodiversity

change. Although spatial heterogeneity at smaller spatial scales (e.g., neighboring sites with

different land cover) can be used to estimate the local effect of drivers such as habitat

conversion[16], correlated large-scale drivers affecting regional species pools may still influence

local community dynamics[17]. Spatial relationships between different land use changes in the

terrestrial realm were expected based on the land requirements to support proximal human

populations[18]. In the marine realm, different human uses (i.e., fisheries) largely occur in different

areas (for instance, demersal fisheries mostly occur over the continental shelf, whilst pelagic

fisheries can be either continental or oceanic), explaining the weaker correlations. Coastal

regions were intermediate in patterns between terrestrial and oceanic regions, suggesting that the

prevalence of human presence may contribute to the differences between the two realms[19]. As

we found fewer strong correlations among different driver variables in the marine realm,

separating the effects of different drivers may be more feasible in marine, especially in open

ocean, ecosystems.

As climate change is only weakly associated with other drivers, there is considerable opportunity

to disentangle climate change impacts from those of other drivers. The weak association means

that climate change affects biological communities exposed to both strong and weak intensities

of other drivers. In the case when other drivers are weak, climate change has the potential to be

the dominant driver of change. Even though other drivers are also spatially variable, their

variability tends to be spatially correlated. Consequently, climate change impacts on species

abundances, range limits and community compositions[20,21] may be easier to isolate than those of

other drivers. Indeed, high-latitude regions, such as the tundra and boreal forests, with low

human-use but pronounced climate change[12,22] have historically undergone less human

settlement and agriculture. Warming of ocean temperatures is affected by additional factors

compared with air temperatures, especially ocean currents[12,23], which likely contributes to the

rapid temperature change in the Indo-Pacific, also an area of intense fishing activity[24]. Hence,

locations in which climate change is the main driver of change in a community are likely to be

more common in terrestrial communities.

Given the strong spatial correlations among many drivers of biodiversity change, attributing

biodiversity change to drivers is likely to be most successful if focused on complexes of

environmental change, rather than on each variable individually. Our classification of ATCs

helps regard anthropogenic environmental change as a series of at least 11 'natural experiments'

across the globe. In particular, the ATCs highlight which environmental changes have the most

opportunity to jointly influence communities. When multiple drivers simultaneously act on a

community, they could have additive, synergistic or antagonistic effects[25-27]. The differential

associations of drivers, summarized by the proposed ATCs, provide an informed baseline for

further studies aiming at understanding the effects of multiple drivers on biodiversity and

ecosystem services. Because drivers related to human-use, pollution and species invasions have

great potential to co-occur, understanding these interactive effects has widespread importance.

As climate change is occurring globally[12], interactive effects of climate change and other drivers

also have the potential to be widespread. However, we find that there is greater spatial overlap

between high intensities of climate change and other drivers in the marine realm. The ATCs

could be further used in macroecological studies of driver impacts. For example, examination of

the relationships between the ATCs and the distributions of threatened species or local/regional

estimates of biodiversity change may help to identify the most harmful combination of drivers.

Estimating the specific effects of individual drivers may also be aided by considering our ATCs

(or driver clusters identified by a similar approach). Study regions that are most suitable to

isolate the effects of a specific driver could be selected from within geographic clusters

dominated by the driver of interest, to reduce the confounding effects of other drivers in the

landscape. Moreover, our approach could be used to design the spatial sampling of quasi-

experimental observatories in future monitoring programs. Observatories could be selected along

different driver gradients (keeping all but one driver constant) or within different driver

combinations. Long-term data from such observatories could greatly advance our understanding

of the underlying causes of biodiversity change.

Quantifying exposure to environmental change is the first step towards understanding how life

on Earth is being reshaped in the Anthropocene and more specifically for determining which

species, in which places, are or will be most exposed to human activities. The impacts of

different drivers on biodiversity will depend on a combination of the magnitude of exposure to

drivers and species' sensitivities to environmental change[28]. We intentionally focused on

exposure patterns so that our results are not species-specific and are therefore potentially relevant

for any taxa or ecosystem. Unlike exposure, sensitivities vary among taxa according to

characteristics such as their life history, traits and niche breadth among others[29] and therefore

should be examined separately for different taxa. We also avoided making any complex

assumptions about the relationships between the absolute levels of each driver variable and its

impact on organisms, rather we assumed that all variables were similarly important.

Management at specific locations is clearly aided by assessing the local magnitudes of different

drivers. However, there are a number of advantages of knowledge on the general patterns in how

different drivers combine at larger-scales. First, these large-scale patterns allow local

management to be modified according to the wider anthropogenic land- or seascape context,

which affects the regional species pool and hence potentially biodiversity changes at smaller-

scales[17]. Second, managers may only have access to partial data at local scales – the typical

combinations of drivers that we identify can help managers predict the extent to which other

drivers should be of concern. Finally, by characterizing regions of the world in terms of the

nature of environmental change, our ATCs suggest how information and data might be pooled

and synthesized across regions, and even across realms[30]. Regions exposed to the same ATC,

regardless of location, would benefit from exchanging knowledge about prioritization strategies

and management of the multiple drivers, as well as implementing cross-border strategies to

minimize their impact.

Global impact assessments and mitigation policy can be better informed by explicitly

incorporating the coupled and decoupled drivers of current and future biodiversity change. Our

macroecological approach to mapping the drivers of biodiversity change contributes to the

development of broad conservation policy targeted toward the mitigation of specific driver

complexes. The main drivers affecting nature and nature's contribution to people are one of the

overarching components of the IPBES framework and assessments[11,31] while attribution of

climate change impacts is a core chapter of the IPCC report[32]. Our findings are especially

relevant for the IPBES Global Assessment that seeks knowledge on global-level linkages.

Monitoring progress towards the Aichi Biodiversity Targets and the Sustainable Development

Goals also requires a clear understanding of the different drivers and their inter-causal

relationships[5]. The empirical spatial relationships among drivers indicate the strength to which

they are affected by common processes and hence are likely to have inter-linked impacts. A

better understanding of these associations, and their integration within biodiversity models could

improve the quality of the projections made with scenarios of future global change, by

considering how the full suite of drivers might change over time[33].

Data on global drivers of biodiversity change are still limited[34]. Further global datasets on driver

variables in the ocean, such as plastic pollution, would be especially valuable, and allow deeper

examination of the relationships among drivers in the marine realm. Spatially-explicit maps of

the number of invasive species would also have improved our analysis. These data are now often

available at national or sometimes regional scales[35] but have not been downscaled or modelled

explicitly across the world. We followed the approach of Halpern et al.[8,9] and inferred the

pressure from invasive species through transport connectivity for both the terrestrial and marine

realm; however, our overall findings are not affected by the inclusion of this variable. For

estimation of historical temporal trends in land cover between 1950-2010, we used the land use

harmonization (LUH2) dataset of annual predictions of land-use states, which is based on diverse

empirical data along with interpolation assumptions[36], and hence are not exact data. However,

we regarded all datasets as the best current estimates of each driver-related variable. For most

drivers, it is currently only possible to examine spatially-explicit trends over recent, short time-

scales[7,8]. Ongoing projects, such as the Copernicus project (http://www.copernicus.eu/), will

greatly increase the availability of spatiotemporally-explicit, high resolution datasets on different

variables[37] in the coming years for further study of the relationships between drivers and for

attribution of biodiversity change to the underlying drivers.

Anthropogenic impacts are now pervasive across the globe, even reaching ecosystems that have

so far avoided major exposure to the effects of direct human activity. However, their effects are

not spatially homogeneous. Terrestrial and marine communities differ in their exposure to

anthropogenic drivers, with greater spatial coupling of threats in the terrestrial realm, likely

driven by proximate human populations. The relationships that we document here have

implications for the spatial patterning of biodiversity change across the world. Especially on

land, climate change can act as a lone driver of biodiversity change across more communities

than other drivers. This means that its unique fingerprints may be easier to detect than those of

other drivers that more often act in combination, regardless of its relative importance. A central

focus of modern ecology is to understand global patterns of biodiversity change. Yet, all too

often, scientists and managers are reading, citing, and focusing on system and realm-specific

influences of global change drivers[30]. By a cross-realm approach, we hope to encourage

information exchange across regions of the world that are exposed to similar suites of drivers,

regardless of environmental realm, and the development of joined-up conservation policies

across the terrestrial-marine interface.

Methods

Data selection

Our analysis is only made possible by the great efforts of researchers to develop and make

publicly available gridded global datasets on environmental change variables[10,36,38-46], especially

Halpern et al. for the marine realm[8,9]. We selected variables that were regarded to be sufficiently

global important to be included in other studies on global drivers of change[6-9], even if these

impacts did not cover the whole area within each realm. The terrestrial datasets came from

various sources (see Tables 1 and S1 for all datasets). Most of the marine datasets came from the

landmark study of Halpern et al.[9]. Even though a more recent set of these marine layers is

available[8], we used the layers from their first article[9] because the time-period of the data was

closer to that of the other datasets (i.e., before 2010). For the terrestrial realm, we could calculate

both a trend (i.e., average change in area per year) for the period 1950-2010[36] as well as a current

value for some variables (e.g., urban and crop land cover in 2000/2001)[40,41]. We decided to

include both in the analysis since the underlying processes generating the current area of cover

may precede 1950 and therefore may have a different distribution than the recent trend in cover

since 1950 (justified by the fact that they were not strongly correlated). In fact, only urban land

cover trend displayed any strong correlations with the other variables in our analysis. For forest

cover, only forest cover trend (i.e., loss) based on FAO wood harvest statistics[36] was included as

an anthropogenic pressure.

Further spatial datasets that were potentially relevant but subsequently found to be highly

correlated (>0.9) with another dataset, and were present over slightly smaller areas than their

correlated partners, were excluded on the basis of redundancy: phosphorus fertilizer application

(correlated with N fertilizer application) in the terrestrial realm, and pesticide (correlated with

fertilizer) and shipping (correlated with ocean pollution) in the marine realm (see Table S1).

Although not correlated, we obtained similar results (in terms of relationships with other drivers)

for pasture cover and pasture cover trend, so we used only the latter in our analysis to ensure an

even number in the variables tested in the terrestrial and marine realms. Some datasets were not

entirely independent. In both the terrestrial and marine realms, country-specific estimates of land

pollution (pesticides and nitrogen fertilizer use) were downscaled by the data providers

according to the distribution of cropland[10,45]. Also in the marine realm, national estimates of

artisanal fishing had been downscaled based on assumptions regarding the distance from coastal

human populations[9]. We regarded each dataset as the best current estimate of the spatial patterns

for each variable. The Land-Use Harmonization dataset (used for trends in forest, crop, pasture

and urban land-use – see Table S1)[36] also used human population data for downscaling but we

only used this dataset to calculate trends in land-use.

Data on the spatial distribution of terrestrial biomes were taken from WWF[47] and marine regions

were obtained by combining coastal region polygon data – MEOW[48] and ocean polygon data

(naturalearthdata.com).

Data organization

For interpretation and presentation purposes, variables were grouped by which global driver of

change they were most directly related to, i.e., climate change, habitat conversion, exploitation,

pollution or species invasions. Because habitat conversion and exploitation were difficult to class

separately across the terrestrial and marine realms, we combined both into a single "human use"

category. There are no high-resolution spatial maps of invasive species richness; however, we

used maps of human transport connectivity, based on the assumption that they are a proxy of

human-mediated propagule pressure (e.g., related to human movement and trade) of alien

species, which is known to be an important determinant of invasion success[49,50] and is an

approach following others[8,9]. We used spatial datasets of accessibility based on transport

infrastructure in the terrestrial realm and cargo volume at ports in the marine realm (Table

S1)[9,46]. We also included "human population density"[42] as a separate driver accounting for the

effects of human activities not falling into the other categories (e.g., tourism/recreation

activities), as well as to determine the relationship of human population density with other

drivers.

Data processing

For most datasets, the data were collected from one or a few years (at some point during 1990–

2010) and were available as pooled data into a single time point (see Table S1 for more details).

For data available as a time series (see those marked by * in Table 1), we calculated temporal

trends for each raster grid as the regression coefficient of a year effect in a linear regression for

1950–2010, reflecting change during the recently defined Anthropocene that is estimated to have

started in 1950[51]. Aridity trend was estimated by taking monthly and annual datasets on potential

evapotranspiration and precipitation, and calculating their ratio[52], and finally the temporal trend

of the annual monthly average of this ratio. Velocity of climate change was calculated following

ref.[53]. We also calculated two additional climate change metrics: temperature divergence –

following ideas by ref.[54], which was inferred from the t-static of the linear regression (i.e.,

temperature trend in °C divided by its standard error), and also trends of extreme temperatures

(whichever was largest of the temporal trends in temperature of the warmest or coolest month).

Missing values in some of the human activity datasets were in remote regions (e.g., very high

latitudes) with likely absent or low variable values and were imputed as zero. However, datasets

were also bounded to an extent of -179, 179, -58, 78 (xmin, xmax, ymin, ymax) to avoid map

edge effects. Greenland was also excluded due to missing data in several of the datasets.

Next, we harmonized each dataset to a standard global grid. The resolutions of the original

datasets were approximately at a 100 km square grid (or 1°) or finer resolution, with the

exception of atmospheric nitrogen deposition[43]. Thus, we chose to aggregate all datasets to a

standard grid of 100 km square grid cells. This aggregation was done by summing the values of

grid cells or by taking the median of values (in the case of trend variables, accessibility and

ocean acidification). Datasets were then re-projected onto a common equal-area map projection

(Eckert IV; ESPG = 54012). Because each dataset comprised data in different units, it was not

possible to directly compare the absolute values among all the datasets. Instead, the values of

each dataset were ranked because of the highly-skewed distributions and scaled between 0 and 1

for ease of interpretation (Fig. S9 show the distributions of the original values of each variable

and Fig. S10 shows global maps of the ranked and scaled data). For all datasets, larger values

reflected a greater potential exposure of that variable on biodiversity. Transformations were

needed in two cases to achieve this – we inverted terrestrial accessibility (i.e., $values^{-1}$) and

changed the sign of the forest trend values (i.e., $values \times -1$).

Data analysis

Spearman's rank correlation coefficients (ρ) were calculated for each pair-wise combination of

variables in each realm at the global-level. This statistic is robust to data processing decisions

because it only uses rankings of the data values and is equivalent to the commonly used

Pearson's correlation on ranked data. We used Dutilleul's modified t-test to account for spatial

autocorrelation in each dataset before testing the significance of the correlations[15]. For the

marine realm, these correlations were also examined separately for grid cells whose centroid

overlapped with oceanic or coastal regions. We examined Moran's I and correlograms to

determine the spatial extent of autocorrelation within each variable and its statistical

significance[55].

To compare the relative importance of different drivers for different regions of each realm, we

calculated the difference between each region's average (weighting each grid cell by the

coverage of each region) and the average across all regions in each realm. We did not use marine

ecosystem data as used by others[9] because the ecosystems spatially overlapped in our coarse 2-D

global raster grid, when, in reality, they are at different depths in the water column. We used k-

medoid clustering, using the partitioning around the medoids algorithm with Manhattan

distances[56], for clustering grid cells according to their extent of exposure of all variables. To

make the number of variables per driver more comparable, we first used principal components

analysis on drivers with multiple associated variables (climate change, human use and pollution)

to produce a reduced number of variables (two axes for climate change and pollution, three axes

for human use) that explained most (>75%) of the variation in each. We then applied the cluster

analysis to a dataset of these PCA variables with the human population density and species

invasion variables. We selected the number of clusters by comparing the changes in dissimilarly

and cluster silhouette width with increasing cluster number. However, we limited the cluster

number so that it was less than 10 and so that each cluster was linked to a different dominant

driver variable. To slightly smooth the maps, we used a moving window to assign each cell the

mode of its 3 x 3 cell neighborhood. Although, driver combinations vary in a continuous manner,

we chose a clustering method that produces discrete grouping to provide the simplest description

of the main groupings in the data. Finally, to identify which regions of the world were exposed to

high intensities of multiple drivers, we identified and summed the number of variables for which

a grid cell was in the upper 10% of values (based on all values greater than zero) of each

variable.

Sensitivity analyses

To examine the effect of the grain size of our global grid, we repeated the data processing steps

except harmonizing the datasets to global grids of different resolutions (800, 400, 200, 100 and

50 [terrestrial only] square km grid) and repeated the analysis of correlations (similar results

were obtained – see Fig. S11). To check the effects of ranking the data values because of the

skewed data distributions, we repeated the data processing steps by logging the values (to the

base 10) rather than ranking them, after bounding values above the upper and lower 2.5% of

quantiles to the values of the upper and lower 2.5% quantiles. This alternative data

transformation does not affect the correlation coefficients because in any case Spearman's

correlations only uses the ranks of the data. We repeated our remaining analysis with this

alternative transformation, calculating the average variable intensities for different terrestrial and

marine regions, and the clustering analysis (similar results were obtained – see Fig. S12 and

S13).

Data availability

Table S1 shows the sources of each dataset and links to where each dataset can be downloaded.

Datasets produced during our analysis (raster layers shown in Figures 4 and 5) are available as

georeferenced TIFF files in the SOM.

Code availability
408

409   R script to harmonize the raster to a standard grid is found here:

410   https://github.com/bowlerbear/harmonizeRasters

411   R script for the subsequent analysis is found here:

412   https://github.com/bowlerbear/geographyDrivers

References

Pereira, H. M. *et al.* Scenarios for Global Biodiversity in the 21st Century. *Science* **330**, 1496-1501, doi:10.1126/science.1196624 (2010).

Barnosky, A. D. *et al.* Has the Earth's sixth mass extinction already arrived? *Nature* **471**, 51-57, doi:10.1038/nature09678 (2011).

Dornelas, M. *et al.* Assemblage Time Series Reveal Biodiversity Change but Not Systematic Loss. *Science* **344**, 296-299, doi:10.1126/science.1248484 (2014).

Isbell, F. *et al.* Linking the influence and dependence of people on biodiversity across scales. *Nature* **546**, 65-72, doi:10.1038/nature22899 (2017).

Tittensor, D. P. *et al.* A mid-term analysis of progress toward international biodiversity targets. *Science* **346**, 241-244, doi:10.1126/science.1257484 (2014).

Sanderson, E. W. *et al.* The human footprint and the last of the wild. *Bioscience* **52**, 891-904, doi:10.1641/0006-3568(2002)052[0891:thfatl]2.0.co;2 (2002).

Venter, O. *et al.* Sixteen years of change in the global terrestrial human footprint and implications for biodiversity conservation. *Nature Communications* **7**, doi:10.1038/ncomms12558 (2016).

Halpern, B. S. *et al.* Spatial and temporal changes in cumulative human impacts on the world's ocean. *Nature Communications* **6**, doi:10.1038/ncomms8615 (2015).

Halpern, B. S. *et al.* A global map of human impact on marine ecosystems. *Science* **319**, 948-952, doi:10.1126/science.1149345 (2008).

Vorosmarty, C. J. *et al.* Global threats to human water security and river biodiversity. *Nature* **467**, 555-561, doi:10.1038/nature09440 (2010).

Diaz, S. *et al.* The IPBES Conceptual Framework - connecting nature and people. *Current Opinion in Environmental Sustainability* **14**, 1-16, doi:10.1016/j.cosust.2014.11.002 (2015).

IPCC. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge University Press, 2013).

Butchart, S. H. M. *et al.* Global Biodiversity: Indicators of Recent Declines. *Science* **328**, 1164-1168, doi:10.1126/science.1187512 (2010).

Pereira, H. M., Navarro, L. M. & Martins, I. S. in *Annual Review of Environment and Resources, Vol 37* Vol. 37 *Annual Review of Environment and Resources* (eds A. Gadgil & D. M. Liverman) 25-+ (2012).

Dutilleul, P., Pelletier, B. & Alpargu, G. Modified F tests for assessing the multiple correlation between one spatial process and several others. *Journal of Statistical Planning and Inference* **138**, 1402-1415, doi:10.1016/j.jspi.2007.06.022 (2008).

Newbold, T. *et al.* Global effects of land use on local terrestrial biodiversity. *Nature* **520**, 45-+, doi:10.1038/nature14324 (2015).

Harrison, S. & Cornell, H. Toward a better understanding of the regional causes of local community richness. *Ecology Letters* **11**, 969-979, doi:10.1111/j.1461-0248.2008.01210.x (2008).

Ellis, E. C., Goldewijk, K. K., Siebert, S., Lightman, D. & Ramankutty, N. Anthropogenic transformation of the biomes, 1700 to 2000. *Global Ecology and Biogeography* **19**, 589-606, doi:10.1111/j.1466-8238.2010.00540.x (2010).

Halpern, B. S. *et al.* Patterns and Emerging Trends in Global Ocean Health. *Plos One* **10**, doi:10.1371/journal.pone.0117863 (2015).

Parmesan, C. & Yohe, G. A globally coherent fingerprint of climate change impacts across natural systems. *Nature* **421**, 37-42, doi:10.1038/nature01286 (2003).

Poloczanska, E. S. *et al.* Global imprint of climate change on marine life. *Nature Climate Change* **3**, 919-925, doi:10.1038/nclimate1958 (2013).

Pithan, F. & Mauritsen, T. Arctic amplification dominated by temperature feedbacks in contemporary climate models. *Nature Geoscience* **7**, 181-184, doi:10.1038/ngeo2071 (2014).

Burrows, M. T. *et al.* The Pace of Shifting Climate in Marine and Terrestrial Ecosystems. *Science* **334**, 652-655, doi:10.1126/science.1210288 (2011).

Ramirez, F., Afan, I., Davis, L. S. & Chiaradia, A. Climate impacts on global hot spots of marine biodiversity. *Science Advances* **3**, doi:10.1126/sciadv.1601198 (2017).

Hof, C., Araujo, M. B., Jetz, W. & Rahbek, C. Additive threats from pathogens, climate and land-use change for global amphibian diversity. *Nature* **480**, 516-U137, doi:10.1038/nature10650 (2011).

26 Garcia-Valdes, R., Svenning, J. C., Zavala, M. A., Purves, D. W. & Araujo, M. B. Evaluating the combined effects of climate and land-use change on tree species distributions. *Journal of Applied Ecology* **52**, 902-912, doi:10.1111/1365-2664.12453 (2015).

27 Travis, J. M. J. Climate change and habitat destruction: a deadly anthropogenic cocktail. *Proceedings of the Royal Society B-Biological Sciences* **270**, 467-473, doi:10.1098/rspb.2002.2246 (2003).

28 Foden, W. B. *et al.* Identifying the World's Most Climate Change Vulnerable Species: A Systematic Trait-Based Assessment of all Birds, Amphibians and Corals. *Plos One* **8**, doi:10.1371/journal.pone.0065427 (2013).

29 Sunday, J. M. *et al.* Species traits and climate velocity explain geographic range shifts in an ocean-warming hotspot. *Ecology Letters* **18**, 944-953, doi:10.1111/ele.12474 (2015).

30 Webb, T. J. Marine and terrestrial ecology: unifying concepts, revealing differences. *Trends in Ecology & Evolution* **27**, 535-541, doi:10.1016/j.tree.2012.06.002 (2012).

31 Kok, M. T. J. *et al.* Biodiversity and ecosystem services require IPBES to take novel approach to scenarios. *Sustainability Science* **12**, 177-181, doi:10.1007/s11625-016-0354-8 (2017).

32 Cramer, W. *et al.* in *Climate Climate Change 2014 : Impacts, Adaptation, and Vulnerability*   (Cambridge University Press, 2014).

33 Rosa, I. M. D. *et al.* Multiscale scenarios for nature futures. *Nature Ecology & Evolution* **1**, 1416-1419, doi:10.1038/s41559-017-0273-9 (2017).

34 Joppa, L. N. *et al.* Filling in biodiversity threat gaps. *Science* **352**, 416-418, doi:10.1126/science.aaf3565 (2016).

35 Dawson, W. *et al.* Global hotspots and correlates of alien species richness across taxonomic groups. *Nature Ecology & Evolution* **1**, doi:10.1038/s41559-017-0186 (2017).

36 Hurtt, G. C. *et al.* Harmonization of land-use scenarios for the period 1500-2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Climatic Change* **109**, 117-161, doi:10.1007/s10584-011-0153-2 (2011).

37 Skidmore, A. K. *et al.* Agree on biodiversity metrics to track from space. *Nature* **523**, 403-405, doi:10.1038/523403a (2015).

38 Harris, I., Jones, P. D., Osborn, T. J. & Lister, D. H. Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset. *International Journal of Climatology* **34**, 623-642, doi:10.1002/joc.3711 (2014).

39 Rayner, N. A. *et al.* Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *Journal of Geophysical Research-Atmospheres* **108**, doi:10.1029/2002jd002670 (2003).

40 Ramankutty, N., Evan, A. T., Monfreda, C. & Foley, J. A. Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles* **22**, doi:10.1029/2007gb002952 (2008).

41 Friedl, M. A. *et al.* MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets, 2001-2012, Collection 5.1 IGBP Land Cover. (Boston University, Boston, MA, USA, 2010).

42 Center for International Earth Science Information Network - CIESIN - Columbia University.   (NASA Socioeconomic Data and Applications Center (SEDAC), Palisades, NY, 2017).

43 Dentener, F. J.   (Oak Ridge, Tennessee, USA, 2006).

44 Hurtt, G., Chini, L., Sahajpal, R. & Frolking, S. Harmonization of global land-use change and management for the period 850-2100". Geoscientific Model Development. ((in prep)).

45 Potter, P., Ramankutty, N., Bennett, E. M. & Donner, S. D. Characterizing the Spatial Patterns of Global Fertilizer Application and Manure Production. *Earth Interactions* **14**, 1-22 (2010).

46 Nelson, A.   (2008).

47 Olson, D. M. *et al.* Terrestrial ecoregions of the worlds: A new map of life on Earth. *Bioscience* **51**, 933-938, doi:10.1641/0006-3568(2001)051[0933:teotwa]2.0.co;2 (2001).

48 Spalding, M. D. *et al.* Marine ecoregions of the world: A bioregionalization of coastal and shelf areas. *Bioscience* **57**, 573-583, doi:10.1641/b570707 (2007).

49 Seebens, H. *et al.* Global trade will accelerate plant invasions in emerging economies under climate change. *Global Change Biology* **21**, 4128-4140, doi:10.1111/gcb.13021 (2015).

50 Hulme, P. E. Trade, transport and trouble: managing invasive species pathways in an era of globalization. *Journal of Applied Ecology* **46**, 10-18, doi:10.1111/j.1365-2664.2008.01600.x (2009).

51 Waters, C. N. *et al.* The Anthropocene is functionally and stratigraphically distinct from the Holocene. *Science* **351**, 137-+, doi:10.1126/science.aad2622 (2016).

52   Zorner, R. J., Trabucco, A., Bossio, D. A. & Verchot, L. V. Climate change mitigation: A spatial analysis of global land suitability for clean development mechanism afforestation and reforestation. *Agriculture Ecosystems & Environment* **126**, 67-80, doi:10.1016/j.agee.2008.01.014 (2008).
53   Loarie, S. R. *et al.* The velocity of climate change. *Nature* **462**, 1052-U1111, doi:10.1038/nature08649 (2009).
54   Nadeau, C. P. & Fuller, A. K. Accounting for multiple climate components when estimating climate change exposure and velocity. *Methods in Ecology and Evolution* **6**, 697-705, doi:10.1111/2041-210x.12360 (2015).
55   Bjornstad, O. N., Ims, R. A. & Lambin, X. Spatial population dynamics: analyzing patterns and processes of population synchrony. *Trends in Ecology & Evolution* **14**, 427-432, doi:10.1016/s0169-5347(99)01677-8 (1999).
56   cluster: Cluster Analysis Basics and Extensions (R package version 2.0.6. , 2017).

Acknowledgments:

This paper arose from discussions at the sChange Workshop (www.idiv.de/schange) during February 2016, which was supported by and held at sDiv, the synthesis centre of iDiv, the German Centre for Integrative Biodiversity Research Halle-Jena-Leipzig. We thank all other participants of this workshop for the stimulating discussion and compilation of data that directly fed into this project. We also thank Suzanne Fritz, Mary O'Connor and Bob O'Hara for comments on a previous version of the manuscript. DB, JSC & MW were funded by the German Research Foundation (DB: Grant no BO 1221/23-1; JSC & MW: via iDiv: FZT 118). SRS is funded by the National Science Foundation, USA (NSF 1400911). LHA was supported by Fundação para a Ciência e Tecnologia, Portugal (POPH/FSE SFRH/BD/90469/2012). ADB was supported by The Danish Council for Independent Research - Natural Sciences (DFF 4181-00565). MD is funded by the Scottish Funding Council (MASTS, grant reference HR09011) and MD & AEM by the ERC project BioTIME (250189) and BioCHANGE (727440). CW was supported by the Natural Environmental Research Council (grant number NE/L002531/1)

Author contributions:

DB performed the analyses and wrote the first outline of the paper with AEB. All authors

designed the study and helped draft the manuscript.

Competing financial interests:

The authors declare no competing financial interests.

Fig. 1 Strong and positive relationships among anthropogenic drivers of biodiversity change. We find a higher number of correlations between drivers in the terrestrial versus the marine realm. Each link represents a significant and strong positive correlation (with strength >0.7) between two variables across 100 square km grids covering the world (see also Figs S1 and S2). No negative correlations were stronger than -0.7.

Fig. 2 Relationships between climate change and other drivers. Comparing the relationships between each variable and temperature change (air or sea surface temperature – SST) or the velocity of climate change (VOCC), we find weak negative (>-0.3) correlations in the terrestrial realm and weak positive (<0.4) correlations in the marine realm. The length of each bar shows the correlation coefficient between temperature change (upper bar) or VOCC (lower bar) and each variable. * denotes statistical significance after accounting for spatial autocorrelation.

Fig. 3 Regions of the terrestrial and marine realms are exposed to distinct combinations of drivers. The bars show the dominant drivers in each terrestrial and marine region. The length of the bar represents the positive deviation (plus standard deviation) of that region's average + standard deviation (across all variables in each driver – see Table 1) from the average of all regions. Bars with negative average deviations are not shown to simplify the presentation. Regions are presented in declining order of the sum of all their bar lengths. Names of the terrestrial regions were shortened for presentation purposes. Figure S6 shows the full distributions for each individual driver variable in each region as well as gives the full names of the terrestrial regions.

Fig. 4 Geography of the Anthropocene. Different geographic regions of the world are exposed to different Anthropogenic Threat Complexes (numbered I to XI). These regions were obtained by k-medoid clustering based on their similarity of exposure to different drivers of biodiversity change and are colored according to one of the highest ranked variable in each complex (i.e., an most important driver in that region). The bars in the legend show the intensities (between the lower and upper quartiles) of each variable in each complex from 0 (no impact) to 1 (highest impact). Clustered regions are colored to reflect the dominant driver and are harmonized across realms to facilitate comparison. White regions were not included in the analysis of each realm. Fig. S7 provides a larger plot of the legend.

Fig. 5 Regions of the world exposed to high intensities of multiple drivers. The number of the 16 driver variables that each grid cell was in the highest 10% of values within its realm. Regions in the darkest orange are exposed to high intensities of multiple variables, while those in off-white are exposed to lower intensities of all (i.e., still potentially exposed to pressures but the magnitude of the pressures is not in the highest 10% of magnitude values). The same is shown for each of the separate drivers, i.e., the intensity of the color is scaled by the number of variables within each driver (Table 1) with a value in the highest 10%. Note: Greenland was not included in the analysis due to missing data in several of the datasets. Larger versions of the plots are presented in Fig. S8.

Table 1 Anthropogenic drivers of biodiversity change and their respective variables based on available global spatial datasets (Table S1). Time-series datasets are indicated by *. Variables in the same line do not necessarily represent the equivalent variable in each realm.

Anthropogenic driver of biodiversity change	Associated variables	
	Terrestrial	**Marine**
Climate Change	Temperature change*	Temperature change*
	Temperature divergence*	Temperature divergence*
	Change in climate extremes*	Change in climate extremes*
	Velocity of climate change*	Velocity of climate change*
	Aridity change*	Ocean acidification
Human use **(land use or change, resource extraction, exploitation)**	Crop cover	Destructive demersal fishing
	Crop cover trend*	Low by-catch demersal fishing
	Pasture cover trend*	High by-catch demersal fishing
	Forest loss trend*	Low by-catch pelagic fishing
	Urban cover	High by-catch pelagic fishing
	Urban cover trend*	Artisanal fishing
Human population density	Population density	Coastal population density
Pollution	Atmospheric nitrogen deposition	Ocean pollution
	Nitrogen fertilizer application	Inorganic coastal pollution
	Pesticide application	Fertilizer coastal pollution
Invasions **(~ connectivity)**	Travel time to major city ("Accessibility")	Port cargo volume