Genome-wide meta-analysis of depression in 807,553 individuals identifies 102 independent variants with replication in a further 1,507,153 individuals

David M. Howard*[1], Mark J. Adams[1], Toni-Kim Clarke[1], Jonathan D. Hafferty[1], Jude Gibson[1],

Masoud Shirali[1], Jonathan R. I. Coleman[2,3], Joey Ward[4], Eleanor M. Wigmore[1], Clara Alloza[1], Xueyi

Shen[1], Miruna C. Barbu[1], Eileen Y. Xu[1], Heather C. Whalley[1], Riccardo E. Marioni[5,6], David J.

Porteous[5,6], Gail Davies[5,7], Ian J. Deary[5,7], Gibran Hemani[8], Chao Tian[9], David A. Hinds[9], 23andMe

Research Team[9], Major Depressive Disorder Working Group of the Psychiatric Genomics

Consortium†, Maciej Trzaskowski[10], Enda M. Byrne[10], Stephan Ripke[11,12,13], Daniel J. Smith[4], Patrick

F. Sullivan[14,15,16], Naomi R. Wray[10], Gerome Breen[2,3], Cathryn M. Lewis[2,3], and Andrew M.

McIntosh[1,7]

Affiliations:
[1] Division of Psychiatry, University of Edinburgh, Royal Edinburgh Hospital, Edinburgh, UK
[2] Social Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology &
Neuroscience, King's College London, UK
[3] NIHR Biomedical Research Centre for Mental Health, South London and Maudsley NHS Trust,
London, UK
[4] Institute of Health and Wellbeing, University of Glasgow, UK
[5] Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK
[6] Centre for Genomic and Experimental Medicine, Institute of Genetics and Molecular Medicine,
University of Edinburgh, Edinburgh, UK
[7] Department of Psychology, University of Edinburgh, Edinburgh, UK
[8] Medical Research Council (MRC) Integrative Epidemiology Unit, Population Health
Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom
[9] 23andMe, Inc., Mountain View, CA, USA
[10] Queensland Brain Institute, University of Queensland, Brisbane, Queensland, Australia
[11] Department of Psychiatry, Charite Universitatsmedizin Berlin Campus Benjamin Franklin, Berlin,
Germany
[12] Medical and Population Genetics, Broad Institute, Cambridge, MA, USA
[13] Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA
[14] Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
[15] Department of Genetics, University of North Carolina, Chapel Hill, NC, USA
[16] Department of Psychiatry, University of North Carolina, Chapel Hill, NC, USA
† Consortium members and their affiliations are listed in Supplementary Information

*Corresponding author: David M. Howard
Division of Psychiatry, University of Edinburgh, Royal Edinburgh Hospital, Edinburgh, UK
+44 131 537 6288
(E-mail: D.Howard@ed.ac.uk)

Abstract

Major depression is a debilitating psychiatric illness that is typically associated with low mood, anhedonia and a range of comorbidities. Depression has a heritable component that has remained difficult to elucidate with current sample sizes due to the polygenic nature of the disorder. To maximise sample size, we meta-analysed data on 807,553 individuals (246,363 cases and 561,190 controls) from the three largest genome-wide association studies of depression. We identified 102 independent variants, 269 genes, and 15 gene-sets associated with depression, including both genes and gene-pathways associated with synaptic structure and neurotransmission. Further evidence of the importance of prefrontal brain regions in depression was provided by an enrichment analysis. In an independent replication sample of 1,507,153 individuals (474,574 cases and 1,032,579 controls), 87 of the 102 associated variants were significant following multiple testing correction. Based on the putative genes associated with depression this work also highlights several potential drug repositioning opportunities. These findings advance our understanding of the complex genetic architecture of depression and provide several future avenues for understanding aetiology and developing new treatment approaches.

Depression is the leading cause of worldwide disability [1] and an estimated 1 in 6 people will develop

the disorder during their lifetime [2]. Twin studies have provided heritability estimates of the disease of

approximately 30-40% [3], however depression is a polygenic trait influenced by many genetic variants

each of small effect [4]. Therefore, to enable the detection of causal genetic variants associated with

depression there is a need to study very large numbers of individuals. However, obtaining detailed

clinical diagnoses of major depressive disorder in larger cohorts is both time consuming and

expensive. The results of Howard, et al. [5] showed that there is a strong genetic correlation ($r_G = 0.86$,

s.e. = 0.05) between broader self-declared definitions of depression and clinically diagnosed major

depressive disorder (MDD) within a hospital setting. Therefore, analysing larger samples, which have

used different approaches to diagnosis, may provide advances in our understanding of the genetics of

depression.

Major efforts to identify genetic variants associated with depression have included a mega-analysis of

9 cohorts (total n = 18,759; 9240 cases and 9519 controls) for MDD [4] and a meta-analysis of 17

cohorts (total n = 34,549) using a broader diagnostic scale that includes depressive symptoms [6].

However, both of these studies failed to find any replicated variants associated with depression. The

first study to report replicable genetic variants for depression found two significant loci associated

with severe, recurrent MDD (85% enriched for melancholia) in a sample of Han Chinese women

(total n = 10,640; 5,303 cases and 5,337 controls) [7]. A later study conducted by Hyde, et al. [8],

examining research participants from the personal genetics company 23andMe, Inc., used a self-

reported clinical diagnosis of depression as the phenotype and identified 15 associated loci (total n =

459,481; 121,380 cases and 338,101 controls). More recently, a genome-wide association analysis of

UK Biobank by Howard, et al. [5] identified 17 variants associated across three depression phenotypes

(maximum total n = 322,580; 113,769 cases and 208,811 controls). These three depression

phenotypes ranged from self-reported help-seeking for problems with nerves, anxiety, tension or

depression (termed "broad depression"), probable MDD based on self-reported depressive symptoms

with associated impairment, and MDD identified from hospital admission records. Finally, a meta-

analysis of 35 cohorts (total n = 461,134; 130,664 cases and 330,470 controls) conducted by Wray, et

al. [9] (PGC) found 44 loci that were significantly associated with a spectrum of depression phenotypes,

some obtained from structured clinical interview and others based on broader criteria.

To maximise the power of the present study we conducted a genome-wide meta-analysis of

depression using 807,553 individuals (246,363 cases and 561,190 controls, after excluding

overlapping samples) from the three largest studies noted above [5, 8, 9]. From the Hyde, et al. [8] analysis,

our meta-analysis included only the 23andMe discovery cohort (termed "23andMe_307k"; 75,607

cases and 231,747 controls). From the Howard, et al. [5] analysis of UK Biobank, the "broad

depression" phenotype was included, with 4-means clustering of genomic principal components used

to derive a larger sample (127,552 cases and 233,763 controls) than studied previously. The PGC

analysis [9] included the 23andMe_307k discovery cohort and an earlier data release of the UK Biobank

cohort (n = 29,740, 14,260 cases and 15480 controls); we therefore obtained results from the PGC that

excluded both of these cohorts (termed "PGC_139k"; 43,204 cases and 95,680 controls).

We sought replication of the variants associated with depression within a set of 23andMe participants

independent of the 23andMe_307k cohort included in the meta-analysis (474,574 cases of self-

reported clinical diagnosis of depression and 1,032,579 controls). The results from the meta-analysis

were used to calculate genetic correlations and conduct Mendelian randomization to identify

potentially pleiotropic and causal relationships between depression and other diseases and behavioural

traits. The meta-analysis results were also used to identify a set of associated genes and gene-

pathways, as well as enrichment of functional annotations associated with depression. Combining

evidence of enrichment in biological pathways with information on traits correlated with depression

allows for additional inferences about shared aetiological mechanisms, thereby increasing the utility

of the standard association analysis approach. Interactions between associated genes and available

drug treatments were also examined to identify novel drug treatments for depression.

Results

102 independent genetic variants associated with depression

We conducted a genome-wide association meta-analysis of depression using 807,553 individuals

(Table 1; 246,363 cases and 561,190 controls) from three previous studies of depression, after

removing sample overlap; the previous studies were Hyde, et al. [8] Howard, et al. [5], and Wray, et al. [9].

We tested the effects of 8,098,588 genetic variants on depression and identified 9,744 associated

variants ($P < 5 \times 10^{-8}$) of which 102 variants in 101 loci were independently segregating

(Supplementary Table 1). The basepair positions of these loci were identified by clumping all

associated variants (linkage disequilibrium $r^2 < 0.1$ across a 3 Mb window) and then merging any

overlapping clumps. Independent variants in each locus were identified through conditional analysis [10]

using all variants in that locus.

A Manhattan plot of our meta-analysis results is provided in Figure 1 with a quantile-quantile plot

provided in Supplementary Figure 1. Linkage Disequilibrium Score (LDSC) regression [11] produced a

genomic inflation factor (λ_{GC}) estimate of 1.63 with an intercept of 1.015 (0.011) prior to inflation

correction, indicating that the inflation was due to polygenic signal and unlikely to be confounded by

population structure. All of the 102 associated variants had the same direction of effect on depression

across the three contributing studies and also within an independent replication sample of 1,507,153

individuals (Table 1; 474,574 cases and 1,032,579 controls). In the replication sample, 97 out of the

102 associated variants were nominally significant ($P < 0.05$) and 87 were significant after Bonferroni

correction ($\alpha = 0.05 / 102$; $P < 4.90 \times 10^{-4}$). Further examination of the general directionality

agreement of associated variants found in the contributing studies to the meta-analysis is provided in

Supplementary Table 2 and the Supplementary Information. In summary, the direction of effect of

depression variants in the previous studies was consistent with the current meta-analysis.

Polygenic risk scores (PRS), calculated using PRSice-2 software [12], were used to assess the predictive

ability of the current genome-wide meta-analysis of depression within the clinically diagnosed MDD

Generation Scotland (GS) cohort [13]. PRS were also calculated using summary statistics from Wray, et

al. [9] (Wray PRS) with the predictive ability within GS also examined. Both the meta-analysis PRS and

the Wray PRS were significantly associated with MDD in GS; however, the current meta-analysis

PRS explained more of the variance in MDD in GS ($P = 5.6 \times 10^{-50}$, $r^2 = 1.2\%$) than the Wray PRS (P

$= 1.9 \times 10^{-36}$, $r^2 = 0.9\%$). Using the first decile as the reference group, in the current meta-analysis

PRS the odds ratio for depression in the top decile of PRS was 2.3 (95% C.I. = 2.1-2.6) compared to

an odds ratio of 2.1 (95% C.I. = 1.8-2.3) using the Wray PRS (Supplementary Figure 2).

Genetic correlations with depression

LDSC regression [11] was used to calculate pairwise genetic correlations (r_G) to determine the extent of

overlap in the genetic architectures across the three non-overlapping cohorts that contributed to our

meta-analysis, i.e. between the UK Biobank, PGC_139k and 23andMe_307k depression analyses.

There was a strong genetic correlation between each of these cohorts. The r_G between PGC_139k and

23andMe_307k was 0.85 (s.e = 0.03). The UK Biobank had a r_G of 0.87 (s.e = 0.04) with PGC_139k.

A similar r_G was also found between UK Biobank and 23andMe_307k (0.85, s.e = 0.03). This was

despite UK Biobank using a broader phenotype based on self-reported help-seeking behaviour

compared to the self-declared clinical depression phenotype of 23andMe_307k and the primarily

clinically obtained MDD phenotype of PGC_139k.

Depression is known to be comorbid with a wide range of other diseases and disorders[14]. To assess the

shared genetic architecture between depression and many other traits, genetic correlations were

calculated between our meta-analysed summary statistics of all three cohorts for depression and 234

behavioural and disease traits available via LD Hub [15] which implements LDSC regression. Of these

behavioural and disease traits, 41 were significantly genetically correlated ($P_{FDR} < 0.01$) with our

meta-analysis results after applying false discovery rate correction, see Supplementary Figure 3 and

Supplementary Table 3. Significant genetic correlations with depression included schizophrenia ($r_G =$

0.32, s.e = 0.02), bipolar disorder ($r_G = 0.33$, s.e = 0.03), college completion ($r_G = -0.19$, s.e = 0.03),

coronary artery disease ($r_G = 0.13$, s.e = 0.02), triglycerides ($r_G = 0.14$, s.e = 0.02), body fat ($r_G = 0.16$,

s.e = 0.03), and waist-to-hip ratio ($r_G = 0.12$, s.e = 0.02). Many of these genetic correlations are

similar to those reported by Wray, et al. [9] and Howard, et al. [5], including earlier age at menarche ($r_G =$

-0.12, s.e = 0.02). However, a novel genetic correlation was observed for age at menopause ($r_G = -$

0.11, s.e. 0.03), potentially indicating a shared genetic architecture between depression and earlier

female reproductive life events. Additionally, novel genetic correlations were observed between

depression and Crohn's disease (r_G = 0.09, s.e. 0.03) and depression and an earlier age of smoking

initiation (r_G = -0.21, s.e. 0.06).

Causal relationships between depression and other traits

The genetic correlations between depression and other traits/disorders reported in the previous section

may arise from genes with pleiotropic effects and biological influences across both traits.

Alternatively, there may be a causal effect of depression on other traits (e.g. depression influencing

triglyceride level) or from other traits causally influencing depression (e.g. triglyceride level leading

to depression). To determine whether causal relationships exist between depression and the 41

genetically correlated traits in Supplementary Table 3, we used a bi-directional, two-sample

Mendelian randomisation (MR) approach using MR-Base v0.4.9 [16] in R and an inverse-variance

weighted (IVW) regression analysis. Where there was also evidence of variant heterogeneity

additional sensitivity tests were conducted using an MR Egger test and a weighted median test

(Supplementary Table 4).

A number of causal relationships were not tested because of sample overlap, which can lead to biased

effect size estimates[17], or where there were an insufficient number of instrumental genetic variables

($n_{variables}$ < 30). A total of 33 causal effects were tested, of which 24 were for a causal effect of

depression on another trait, and nine tests for a causal effect of another trait on depression. Directional

horizontal pleiotropy, where a genetic variant has an effect on both traits but via differing biological

pathways, can bias the estimates of causal effects between traits[18]. Using the MR Egger intercept test,

there was no evidence ($P \geq 0.05$) of directional horizontal pleiotropy for any of the 33 causal effects

examined.

A putative causal effect of depression on neuroticism[19] was detected using the IVW regression

analysis MR test at the 1% significance threshold after false discovery rate (FDR) correction (beta =

0.146, s.e. = 0.039; P_{FDR} = 2.29 × 10^{-3}; Supplementary Figure 4). However, there was also evidence of

variant heterogeneity (P = 6.01 × 10^{-3}), due to global horizontal pleiotropy, requiring additional

sensitivity tests to examine the consistency of the effect. The additional sensitivity tests both had

effects in the same direction as the IVW test (weighted median beta = 0.119, s.e. = 0.047; MR Egger

beta = 0.050, s.e. = 0.235). There was also evidence of a putative causal effect of depression on ever

vs. never smoked[20] (beta = 0.285, s.e. = 0.077; P_{FDR} = 2.29 × 10^{-3}; Supplementary Figure 5), with no

evidence of variant heterogeneity (P = 0.14). Both the putative causal effects of depression on

neuroticism and depression on ever vs. never smoked remained consistent (P < 6.46 × 10^{-4}) in the

'leave one variant out' IVW analysis indicating that the observed effect was not driven by a single

outlying variant.

Neuroticism was the only trait with a putative causal effect on depression using the IVW regression

analysis MR test (beta = 0.366, s.e = 0.037; P_{FDR} = 2.63 × 10^{-21}; Supplementary Figure 6). However,

there was also evidence of variant heterogeneity (P = 9.62 × 10^{-21}) requiring additional MR sensitivity

testing. The weighted median test produced similar effect size and P-value to the IVW test (beta =

0.337, s.e. = 0.038; P = 1.94 × 10^{-18}), but the MR Egger test had a large standard error and was in the

opposite direction (beta = -0.128, s.e. = 0.271; P = 0.64). This putative causal effect remained

consistent (P = 1.58 × 10^{-21}) in the 'leave one variant out', IVW analysis indicating that the observed

effect was not driven by a single variant.

The observed bi-directional relationship between depression and neuroticism could be confounded by

non-independent instrumental variants across both tests, i.e. a region containing variants associated

with depression was used to test for a causal effect on neuroticism and then variants in that same

region were also used to test for a causal effect of neuroticism on depression. To account for any

overlap, we identified and removed 15 instrumental variants from each bi-directional test where there

was evidence of linkage disequilibrium (LD r^2 > 0.1). The effect size for the IVW regression analysis

MR test for depression on neuroticism was attenuated from 0.146 (s.e. = 0.039) to 0.112 (s.e. = 0.042)

and the P-value was no longer significant after false discovery rate correction (P_{FDR} = 0.037). The

effect size for the IVW regression analysis MR test for neuroticism on depression was attenuated from

0.366 (s.e. = 0.037) to 0.289 (s.e. = 0.042) and the P-value remained significant after false discovery

rate correction (P_{FDR} = 1.34 × 10^{-10}).

Partitioning of the heritability component of depression

The estimate of the SNP-based heritability of depression within our meta-analysis was 0.089 (0.003)

on the liability scale using LDSC regression [11]. Heritability was then partitioned by calculating the

proportion of heritability assigned to 24 functional categories divided by the proportion of variants in

that category [21]. This partitioning showed significant enrichment within conserved, intronic and

H3K4me1 regions of the genome for the heritable component of depression ($P_{corrected} < 0.05$), see

Supplementary Figure 7 and Supplementary Table 5. However, the estimates of enrichment for

intronic (1.16×, s.e. = 0.05) and H3K4me1 (1.41×, s.e. = 0.13) regions were much smaller compared

to the conserved regions (17.49×, s.e. = 1.68) of the genome.

Partitioning the heritability estimate by cell type enrichment (Figure 2A, and Supplementary Table 6)

revealed that central nervous system (CNS) and skeletal muscle tissues were enriched ($P_{corrected} < 0.05$)

for genetic variants contributing to the heritability of depression. Studies have reported altered histone

modifications of skeletal muscle in response to exercise[22] and suggested roles for skeletal muscle

PGC-1α1[23] in depression. The prominence of CNS enrichment led us to examine both brain regions

(Figure 2B, Figure 2C, Supplementary Figure 8, and Supplementary Table 7) and brain cell types

(Supplementary Figure 9, and Supplementary Table 8). There was significant enrichment ($P_{corrected} <$

0.05) of the anterior cingulate cortex, frontal cortex and cortex brain regions and neuron brain cells.

The pseudo-coloring used in Figure 2C and Supplementary Figure 8 highlight in red the significance

and effects sizes of the enriched regions of the brain associated with depression, respectively.

Genes and gene-sets putatively associated with depression

The MAGMA [24] package was used to assess the aggregated genetic effects from the meta-analysis on

17,842 genes. This technique identified a total of 269 putative genes that were associated ($P < 2.80 \times$

10^{-6}) with depression, Supplementary Table 9. The most significant gene ($P = 2.27 \times 10^{-19}$) was

Sortilin related VPS10 domain containing receptor 3 (*SORCS3*) on chromosome 10 (Supplementary

Figure 10). The Neuronal Growth Regulator 1 (*NEGR1*) gene was associated with depression ($P =$

3.55×10^{-15}) and there were two nearby independently-associated variants (rs2568958 and

rs10890020) located in separate loci (Supplementary Figure 11 A-B). rs10890020 overlapped with the

Long Intergenic Non-Protein Coding RNA 1360 (*LINC01306*) coding region which was not available

for analysis in MAGMA. A further short (1.2 Mb) region along 18q.21.2 contained three

independently-associated variants across two loci (rs62091461, rs12966052 and rs12967143;

Supplementary Figure 12 A-C). These variants were close to the Transcription Factor 4 (*TCF4*) and

the *RAB27B* genes which were both putatively associated with depression ($P = 4.55 \times 10^{-16}$ and $P =$

1.39×10^{-9}, respectively). Further consideration of the genes putatively associated with depression is

provided in the discussion.

To identify the biological pathways that are influenced by the putative genes associated with

depression, gene-set analyses were performed using MAGMA [24]. This method identifies the genes

involved in each biological pathway and assesses whether there is evidence of enrichment for each

pathway in depression using the *P*-values of each gene. We identified 14 significant putative gene-sets

($P_{corrected} < 0.05$) for depression (Table 2) using data from the Gene Ontology Consortium. Eight of

these gene-sets were cellular components (areas where the genes were active) and were located in the

nervous system: GO_POSTSYNAPSE, GO_SYNAPSE, GO_NEURON_SPINE,

GO_EXCITATORY_SYNAPSE, GO_SYNAPSE_PART, GO_NEURON_PROJECTION,

GO_NEURON_PART, and GO_DENDRITE. The cellular component gene-sets intimate a role in

synapse function and excitatory mechanisms. The other six associated gene-sets were biological

processes: GO_BEHAVIOR, CO_COGNITION,

GO_MODULATION_OF_SYNAPTIC_TRANSMISSION,

GO_REGULATION_OF_SYNAPSE_STRUCTURE_OR_ACTIVITY,

GO_MODULATION_OF_SYNAPTIC_PLASTICITY, and

GO_SINGLE_ORGANISM_BEHAVIOR. The gene overlap between these gene-sets (Supplementary

Table 10) suggests that the gene-sets generally fall within two clusters. One gene-set cluster typically

relates to synaptic structure and activity while the other gene-set cluster relates to the response or

behaviour to stimuli. We also identified a significant gene-set ($P = 3.87 \times 10^{-5}$) containing

somatosensory pyramidal neurons using brain cell-type data from Skene, et al. [25] (Table 2).

Drug - gene interactions

The 269 putative genes that were identified as being significantly associated ($P < 2.80 \times 10^{-6}$) with

depression using MAGMA [24] were examined for known interactions (including agonistic, partial

agonistic, antagonistic, modulating, binding and blocking interactions) with prescribed medications in

the Drug Gene Interaction Database (dgidb.genome.wustl.edu) v3.0 [26]. A total of 560 interactions

were identified between 57 genes and 514 drugs (see Supplementary Table 11). Anatomical

Therapeutic Chemical (ATC) classifications were available for 220 of these drugs which belonged to

54 different second level ATC classes and which interacted with 37 genes, Figure 3. The greatest

number of drug - gene interactions ($n_{interactions} = 47$) were observed between psycholeptics (N05, which

includes antipsychotics and anxiolytics) and dopamine receptor D2 (*DRD2*).

Discussion

In this study we report a meta-analysis of 807,553 individuals (246,363 cases and 561,190 controls)

using summary statistics from three independent studies of depression, Hyde, et al. [8]

(23andMe_307k), Howard, et al. [5] (UK Biobank) and Wray, et al. [9] (PGC_139k). The meta-analysis

examined 8,098,588 variants and identified 102 independently segregating variants associated ($P < 5$

$\times 10^{-8}$) with depression. These 102 variants were assessed in an independent replication sample of

1,507,153 individuals (474,574 cases and 1,032,579 controls) and 87 variants were significant in that

sample after multiple testing correction. The estimate of the genome-wide SNP-based heritability on

the liability scale was 0.089 (s.e. = 0.003), indicating that the analysed depression phenotype had a

significant genetic component. All of the 102 variants associated with depression had an equivalent

direction of allelic effect across the three studies[5, 8, 9] that contributed to the meta-analysis and the

replication sample (Supplementary Table 1), suggesting that these variants represent robust

associations with depression. Our gene-based analyses revealed pathways relating to

neurotransmission and response to stimuli, with the central nervous system also found to be enriched

when performing heritability partitioning. The partitioning of the heritability for depression also

highlighted the importance of the cortical regions of the brain and further research focussed on the

regions and biological pathways detected is warranted. We detected 269 genes putatively associated

with depression and demonstrated the utility of investigating their interaction with pharmacological

treatments.

The three studies[5, 8, 9] contributing to the meta-analysis were based on different depression

phenotypes; nevertheless, there were strong genetic correlations (> 0.85) between them. PGC_139k

used a predominately clinically ascertained diagnosis of major depression and the observed genetic

correlations indicate that each study was likely to be capturing a similar underlying genetic

architecture, despite the use of different diagnostic instruments. Therefore, larger population-based

samples, where the timescales and costs of obtaining a clinically diagnosis would be prohibitive, can

contribute to our understanding of the genetic architecture of the disease. The meta-analysis also

provides the opportunity to assess the general directionality agreement of variants that have

previously been associated with depression (Supplementary Information and Supplementary Table 2).

The examination of previous findings between studies and the current meta-analysis suggests that

there is good degree of directionality agreement of causal effects of depression when using studies

with over 100,000 participants. However, there is likely to be a need to continue to ascertain clinically

ascertained MDD cohorts to ensure validity of the larger cohorts with broader phenotyping.

There is an ever growing body of evidence that there are shared genetic components between

behavioural and psychiatric disorders[27-29]. Using the meta-analysed results in the current study this

was evidenced by significant genetic correlations for depression with neuroticism, anorexia nervosa,

attention deficit hyperactivity disorder (ADHD), schizophrenia, and bipolar disorder. The MR

analysis also identified a putative bi-directional casual effect between depression and neuroticism.

Removing regions which were used to test the effect in both directions indicated that a putative effect

remained only for neuroticism on depression. This uni-directional effect does make more sense

intuitively as neuroticism is a stable trait whereas depression is a more episodic condition.

Examining significant genes that overlap between the current meta-analysis of depression and the

neuroticism studies of both Luciano, et al. [30] and Okbay, et al. [31] revealed putative associations with

DRD2, CUGBP Elav-like family member 4 (*CELF4*) and ELAV like RNA binding protein 2

(*ELAVL2*). Dopamine acts as a neurotransmitter in the brain with *DRD2* encoding the dopamine

receptor D2 subtype. Genetic variation around the *DRD2* gene has been linked to differences in

structural connectivity between the basal ganglia structure with the frontal cortices[32]. Supporting

evidence of the importance of the cortical brain regions was provided by the heritability partitioning

of the brain regions (Figures 2B and 2C) with significant enrichment of the anterior cingulate cortex,

frontal cortex and cortex brain regions. *DRD2* is also associated with mood modulation and emotion

processing[33] and is also commonly reported in association studies of schizophrenia[34, 35]. *DRD2* was

included in all 14 of the putative Gene Ontology Consortium gene-sets identified (but not the

somatosensory pyramidal neurons gene-set) and has been identified in previous studies of depression[5,

9, 36]. *CELF4* plays a key role in coordinating the synaptic function in excitatory neurons[37] with

dynamic changes in expression during brain development[38] and deletions in the surrounding region

(18q.12.2) associated with autism spectrum disorder[39] and developmental and behavioural disorders[40].

ELAVL2 potentially aids in the regulation of gene expression pathways in human neurodevelopment [41]

and disruption of related pathways may be a factor in neurodevelopmental disorders[42]. Improving our

understanding of the genetic similarities and differences between neuroticism and depression may

provide a route to determining the biological aspects that underpin a more permanent personality trait

or a depressive state. These genetic similarities and differences also provide opportunities for

phenotypic stratification and warrants further investigation in future studies.

The current study further reaffirms the genetic correlation between schizophrenia and depression

observed by Wray, et al. [9]. The current gene-based analysis identified the vaccinia-related kinase 2

(*VRK2*) gene as putatively associated with depression. Extensive research has been conducted

examining the link between *VRK2* and schizophrenia, with the association replicated in both European

[43, 44] and Asian [45] populations. Additionally, associations have been found between schizophrenia and

the arginine and serine rich coiled-coil 1 (*RSRC1*) [46] and myocyte enhancer factor 2C (*MEF2C*) [34, 47]

genes, both of which we found to be putatively associated with depression. Hyde et al. [8] reported that

RSRC1 and *MEF2C* were also close to the genetic variants that they detected, and they highlighted

MEF2C's role in regulation of synaptic function and central nervous system phenotypes. *MEF2C* was

also included in ten of the 15 putative gene-sets identified for depression. The *TCF4* gene coding

region (Supplementary Figure 12) is also noteworthy as it contained two independently-associated

variants for depression and has been identified in previous studies of depression [5, 9, 48]. *TCF4* is

involved in the synaptic plasticity [49] and the excitability of prefrontal neurons [50] and has been

implicated in other psychiatric disorders[51], including schizophrenia [34].

ADHD is a neurodevelopmental disorder, typically diagnosed during early childhood (age 4 to 6) and

is associated with an increased risk of depression during adolescence [52]. The current study

demonstrated that there was a significant genetic correlation between ADHD and depression. The

cadherin 13 (*CDH13*) gene, which codes for cell adhesion molecules, was found to be putatively

associated with depression and has also been implicated in ADHD [53, 54] and specifically hyperactive

and impulsive symptoms [55]. *CDH13* was included in the putative GO_NEURON_PART and

GO_NEURON_PROJECTION gene-sets. A further gene involved in cell adhesion, astrotactin 2

(*ASTN2*), which we found to be putatively associated with depression is also involved in ADHD

aetiology [54, 56] and plays a role in neuronal development in the brain [57]. Dopamine transmission may

also underlie ADHD and depression with both the *DRD2* and ankyrin repeat and kinase domain

containing 1 (*ANKK1*) genes implicated in our analysis of depression and in studies of ADHD [58, 59].

Age of menarche has been found to be phenotypically and genetically correlated with depression [9, 60,

61], with a causal effect of earlier age at menarche on depressive symptoms also reported [62]. The

current meta-analysis identified a significant genetic correlation between age of menarche and

depression, and nominal evidence of a bi-directional causal effect. The lin-28 homolog b (*LIN28B*)

gene was significant in our meta-analysis and has frequently been associated with age of menarche [63-

65]. The biological mechanisms that underlie the association of *LIN28B* with both age of menarche and

depression remain unclear; however, *LIN28B* has been shown to be involved in regulating cell

pluripotency [66] and developmental timing [67], and through its mediation of Lethal-7 miRNA has been

implicated in inflammation and immune response [68]. The current analysis also identified a novel

genetic correlation between age of menopause and depression which may share a similar biological

mechanism to that of age of menarche.

A number of studies, using different methodologies, have examined causal relationships between

depression and smoking with smoking increasing the risk of depression [69, 70], a bi-directional effect [71]

and no effect [72-75] reported. In the current analysis there was an insufficient number of instrumental

variants to test the effect of smoking on depression, but there was evidence of a genetic correlation

between depression and cigarette smoking as well as a causal effect of depression on smoking.

Smoking has been reported to have an anxiolytic and antidepressant effect [76, 77] and this may explain

why we observe a causal effect of depression on smoking; however, as reported by Munafò, Hitsman,

Rende, Metcalfe and Niaura [78] the relationship is likely to be complex and requires further

investigation.

Drug - gene interactions

Examining the number of interactions between the genes associated with depression and the second

level ATC drug classifications demonstrates that there are currently available pharmaceutical

treatments that may target the genetic component of depression. Most notable were the large number

of interactions between the *DRD2* gene and the N05 drug classification, which primarily comprises

typical and atypical antipsychotics. The dopaminergic system has been previously implicated in

depression, particularly the symptoms of anhedonia and decreased motivation, and antidepressant

action has been reported for dopamine reuptake inhibitors and D2-receptor agonists in animal models

of depression [79]. Furthermore, the dopamine and noradrenaline reuptake inhibitor bupropion is

licensed for the treatment of depression [80].

We also identified genes with associated medications which, although not aimed at treating

depression, may provide unpredicted drug benefits or adverse effects for those with the disorder. The

Neuregulin 1 (*NRG1*) receptor ErbB4, found on GABAergic neurones, has been identified as a

potentially druggable target for depression, anxiety and schizophrenia [81]. There is interest in

developing agents, such as basimglurant and fenobam, as antidepressants and anxiolytics that target

glutamatergic receptors [82]. Estrogen Receptor 2 (*ESR2*) has previously been implicated in

antidepressant action through up-regulation by 3β-hydroxysteroid dehydrogenase [83]. Estrogenic

compounds have also been found to have antidepressant effects in rodent models of depression [84].

Further, our work highlights other potential druggable genes associated with depression which are not,

to our knowledge, currently associated with antidepressant treatment or mood-associated adverse

effects. These include the R-type calcium channel gene, Calcium Voltage-Gated Channel Subunit

Alpha1 E (*CACNA1E*), and the nucleosome associated gene, Lysine Methyltransferase 2A (*KMT2A*)

[85].

An intriguing omission among the depression-associated genes identified in our study are genes

linked with the serotonergic system, such as the serotonin transporter *SLC6A4*; the G protein subunit

GNB3; the serotonin receptor *HTR2A* and tryptophan hydroxylase (*TPH2*). This is surprising, as

interaction with the serotonergic system forms the basis of most antidepressant treatments. Our

finding could indicate a functional separation between genetic pathways of depressive disease and

pathways of antidepressant treatment. Thus, serotonin-associated genes, while potentially relevant for

predicting efficacy and adverse effects of serotonergic antidepressants, may not be directly associated

with the aetiology of depression itself (or at least that which is determined by common genetic

variation identified in GWAS). Indeed, a recent review of the genetics studies of depression has

remarked upon the failure to demonstrate association with depression for serotonergic and other

popular candidate genes [86]. It may also be that the pathways of depression and serotonergic

antidepressant effect are separate but entwine through common intermediary genes. One such

candidate is *NRG1*, identified here in depression, and also in a recent meta-analysis of antidepressant

response [87]. These findings suggest that there is potentially a need to concurrently model a range of

'omics data, including genomics, epigenomics, and transcriptomics to gain further understanding of

depression pharmacology.

The principal strength of this study is the increased power obtained from the analysis of three large

independent cohorts. This has allowed the examination of the effects of variants and regions that have

been identified previously to determine whether they maintain an effect on depression. We have used

variant-based analyses to calculate heritability enrichment across different brain regions and then also

applied a complimentary approach using the genes assigned to functional gene-sets, with both

methods highlighting the importance of the prefrontal brain regions. Our study highlights a number of

potential gene targets for drug repositioning; however, due to the causal heterogeneity of depression

[88], it is likely that stratifying depression will lead to clearer distinction between depression subtypes

and potential treatments.

Conclusion

This study describes a meta-analysis of the three largest depression cohorts (total n = 807,553, with

246,363 cases and 561,190 controls) currently available. The meta-analysis identified 102

independently-segregating genetic variants associated with depression in 101 loci and demonstrated a

consistency of effect directionality in a large replication sample (total n = 1,507,153, with 474,574

cases and 1,032,579 controls) and across the three contributing studies allowing greater confidence in

the findings. The heritability enrichments and gene-set analysis both provided evidence for the role of

prefrontal brain regions in depression and the genes identified contribute to our understanding of

biological mechanisms and potential drug targeting opportunities. These findings advance our

understanding of the underlying genetic architecture of depression and provide novel avenues for

future research.

Online Methods

To conduct our analyses we included data from three previous studies of depression [5, 8, 9]. For the UK

Biobank study [5] slightly different quality control was applied and the data were reanalysed. For the

other two studies the summary statistics were obtained directly from the respective authors. Further

information regarding each of these cohorts is provided below.

UK Biobank

The population-based UK Biobank cohort [89] consists of 501,726 individuals with genome-wide data

for 93,095,623 autosomal genetic variants imputed using the HRC and UK10K reference panels [90].

We used the variants from the HRC reference panel to conduct a 4-means clustering approach and

used the first two principal components to identify a genetically homogenous subgroup of 462,065

individuals. We next removed 131,790 individuals that had a shared relatedness up to the third degree

that were identified by UK Biobank based on kinship coefficients (> 0.044) calculated using the

KING toolset [91]. For these 131,790 removed individuals we then calculated a genomic relationship

matrix and identified one individual to be reinstated from within each related group that had a genetic

relatedness less than 0.025 with all other participants which allowed us to add an additional 55,745

individuals back into our sample. We then used a checksum based approach [92] to identify and exclude

954 individuals from within the UK Biobank cohort that overlapped with the Major Depressive

Disorder Working Group of the Psychiatric Genomics Consortium (PGC) cohorts analysed by Wray,

et al. [9]. This was possible for a total of 30 out of 33 cohorts that make up the PGC analysis due to the

availability of genetic data. We also removed those UK Biobank individuals with a variant call rate $<$

90% or that were outliers based on heterozygosity, or variants with a call rate $< 98\%$, a minor allele

frequency < 0.005, those that deviated from Hardy-Weinberg equilibrium ($P < 10^{-6}$), or had an

imputation accuracy score < 0.1, leaving a total of 10,612,809 variants for 371,437 individuals.

Within UK Biobank we used the broad definition of depression [5] with more detailed phenotypic

information available in that paper. In summary, case and control status of broad depression was

defined by the participants' response to the questions 'Have you ever seen a general practitioner for

nerves, anxiety, tension or depression?' or 'Have you ever seen a psychiatrist for nerves, anxiety,

tension or depression?'. Exclusions were applied to participants who were identified with bipolar

disorder, schizophrenia, or personality disorder using self-declared data following the approach of

Smith, et al. [93] as well as prescriptions for antipsychotic medications. This provided a total of 127,552

cases and 233,763 controls (n = 361,315, prevalence = 0.353) for analysis.

Statistical Analysis of UK Biobank

To conduct the association analysis within UK Biobank we used BGENIE v1.1 [89] to assess the effect

of each genetic variant using a linear association test:

486
$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}_1$$

487
$$\hat{\mathbf{y}} = \mathbf{X}\boldsymbol{\beta}$$

488
$$(\mathbf{y} - \hat{\mathbf{y}}) = \mathbf{G}\,b + \boldsymbol{\varepsilon}_2$$

where $\mathbf{y}$ was the vector of binary observations for each phenotype (controls coded as 0 and cases

coded as 1). $\boldsymbol{\beta}$ was the matrix of fixed effects, including sex, age, genotyping array, and the first 8

principal components and $\mathbf{X}$ was the corresponding incidence matrices. $(\mathbf{y} - \hat{\mathbf{y}})$ was a vector of

phenotypes residualized on the fixed effect predictors, $\mathbf{G}$ was a vector of expected genotype counts of

the effect allele (dosages), b was the effect of the genotype on residualized phenotypes, and $\boldsymbol{\varepsilon}_1$ and

$\boldsymbol{\varepsilon}_2$ were vectors of normally distributed errors. The effect sizes and standard errors were transformed

to the logistic scale by dividing each value by $\rho(1-\rho)$, where ρ is the trait prevalence (0.353).

23andMe

We obtained the genome-wide association study results from the discovery 23andMe subset

(23andMe_307k) from the Hyde, et al. [8] analysis. Phenotypic status was based on responses to web-

based surveys with individuals that self-reported as having received a clinical diagnosis or treatment

for depression classified as cases. This provided a total of 75,607 cases and 231,747 controls (n =

307,354, prevalence = 0.25). We excluded variants with an imputation accuracy threshold < 0.6 and

with a minor allele frequency < 0.005, which left a total of 8,995,180 variants.

Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium (PGC)

Wray et al. conducted the largest meta-analysis of MDD to date [9], utilising European-ancestry PGC

cohorts with an emphasis placed on obtaining clinically-derived phenotypes for MDD. Their meta-

analysis included the 23andMe_307k discovery cohort [8] and a previous release of the UK Biobank

data [94]. We therefore obtained the summary statistics from their meta-analysis of major depression

with the 23andMe_307k and the previous UK Biobank cohorts removed (PGC_139k). This provided

a total of 12,149,399 variant calls for 43,204 cases and 95,680 controls (n = 138,884, prevalence =

0.31). We excluded variants with an imputation accuracy threshold < 0.6 or a minor allele frequency

< 0.005, which left a total of 10,365,555 variants.

Meta-analysis of genome-wide association study summary statistics

We used Metal [95] to conduct an inverse variance-weighted meta-analysis of the summary statistics

from the three studies (using the log of the odds ratios and the standard errors of the log of the odds

ratio), conditional on each variant being available in all three of the contributing cohorts. This

provided 8,098,588 genetic variants and up to 246,363 cases and 561,190 controls (n = 807,553) for

the meta-analysis. Linkage disequilibrium score (LDSC) regression intercepts [11] were used for

genomic inflation control of the three contributing cohorts and the final meta-analysis results.

Clumping and merging were used to identify the basepair positions of loci containing depression-

associated variants. Clumping of the meta-analysis results was conducted using Plink v1.90b4 [96]

applying the following parameters: --clump-p1 1e-4 --clump-p2 1e-4 --clump-r2 0.1 --clump-kb 3000,

with merging of the clumped loci conducted using bedtools [97]. A conditional analysis [10] was used to

identify independently-segregating variants in each of the merged loci that was genome-wide

significant, using the linkage disequilibrium structure of UK Biobank as a reference panel. Genome-

wide statistical significance was defined as $P < 5 \times 10^{-8}$, with the meta-analysis results from Metal

reported. Due to the complexity of the major histocompatibility complex region only the most

significant variant in that region is reported.

23andMe replication sample

The variants that were significant ($P < 5 \times 10^{-8}$) in the meta-analysis, described above, were replicated

in an independent sample of 1,507,153 unrelated individuals (474,574 cases and 1,032,579 controls)

from 23andMe, Inc. Individuals within this replication sample were unrelated to individuals within

23andMe_307k. Detailed information regarding this replication sample is provided in the

Supplementary Information. In summary, imputed genetic data was obtained for an unrelated

European sample and the variants identified as genome-wide significant ($P < 5 \times 10^{-8}$) in the meta-

analysis were analysed using logistic regression. The phenotype used in the 23andMe replication

sample was the same used within the 23andMe_307k cohort, described above, with the responses to

web-based surveys used to classify individuals that self-reported as having received a clinical

diagnosis or treatment for depression classified as cases. We used Metal [95] to conduct an inverse

variance-weighted meta-analysis of the significant variants ($P < 5 \times 10^{-8}$) identified in the previous

paragraph (using UK Biobank, 23andMe_307 and PGC_139k) and the 23andMe replication cohort.

Polygenic risk score analysis

Polygenic risk scores (PRS) were created using PRCise-2 [12] and evaluated in Generation Scotland: the

Scottish Family Health Study (GS) [13]. Two polygenic risk scores were created: one using weightings

from the current meta-analysis and one using weightings from a previous association study of major

depression conducted by Wray, et al. [9]. For both of these sets of summary statistics, GS was removed

from the samples prior to calculated single nucleotide polymorphism (SNP) weights which were

filtered so that they only contained SNPs that overlapped between the current meta-analysis PRS and

the Wray, et al. [9] dataset (Wray PRS) (N SNPs = 8,084,638). To create independent SNP sets in

PRCise-2 a linkage disequilibrium r^2 of < 0.1 and a 250kb sliding window was used. PRS were

calculated starting with a P-value threshold of ≤ 0.0001 increasing in increments of 0.00005 up to a P-

value threshold of ≤ 0.5, and also calculating the full model including all SNPs ($P \leq 1$). The best

thresholds for the meta-analysis PRS ($P \leq 0.249$; 46,905 SNPs) and Wray PRS ($P \leq 0.0588$; 18,257

SNPs) were each used to test for an association with MDD in GS.

GS is a family and population-based cohort of 20,195 participants genotyped on the Illumina

OmniExpress BeadChip (706,786 SNPs). The raw genotype data in GS underwent quality control so

that individuals with a call rate $< 98\%$, SNPs with a genotyping rate $< 98\%$, a minor allele frequency

$< 1\%$ and a Hardy-Weinberg equilibrium P-value $< 10^{-6}$ were removed from the dataset, leaving a

total of 19,997 GS individuals and 561,125 SNPs. Participants underwent a clinical diagnosis of MDD

using the Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders

[98] with further information on the phenotype used provided in Fernandez-Pujals, et al. [99]. GS

comprised of 2648 MDD cases and 17,158 controls with genotype data.

The association between MDD in GS and PRS was analysed using ASREML-R (www.vsni.co.uk),

fitting family structure in GS as a random effect using an inverse pedigree matrix. Wald's conditional

F-test was used to calculate *P*-values for fixed effects which were age, sex, four multidimensional

scaling components (to control for population stratification), and the PRS score. A Taylor series

transformation[100] was used to convert effect sizes and their standard errors from the linear scale to the

liability scale and to calculate odds ratios and confidence intervals. The variance explained by the

PRS was calculated by multiplying the PRS by the effect size on the linear scale and dividing by the

variance of the phenotype, to calculate a co-efficient of determination between 0 and 1 (Nakagawa

and Schielzeth [101]). PRS were split into deciles and the odds ratio for MDD in each decile calculated

using the 1st decile as the reference group.

Genetic Correlations

We calculated genetic correlations using LDSC regression [11] and the online software LD hub

(http://ldsc.broadinstitute.org/) [15]. LDSC regression leverages linkage disequilibrium (LD)

information for each genetic variant such that the χ^2 statistic for that variant includes the effects of all

loci in LD with that variant. This can be extended to the analysis of genetic correlations between traits

if the χ^2 statistic is replaced with the product of 2 z-scores from 2 traits of interest [102]. Genetic

correlations were calculated between the three datasets that contributed to the meta-analysis. Genetic

correlations were also calculated between our meta-analysed results (23andMe_307k, UK Biobank

and PGC_139k) for depression and 234 behavioural and disease related traits. *P*-values were false

discovery rate (FDR) adjusted[103] and correlations reported if $P_{FDR} < 0.01$.

Mendelian randomization

We used Mendelian randomization (MR) to assess whether causal effects exist between depression

and a number of other traits and disorders. MR uses genetic variants as a proxy for environmental

exposures, assuming that: i) the genetic variants are associated with the exposure; ii) the genetic

variants are independent of confounders in the exposure-outcome association; iii) the genetic variants

are associated with the outcome only via their effect on the exposure, i.e. there is no horizontal

pleiotropy whereby the variants affect both exposure and outcome independently. Individual genetic

variants may be weak instruments for assessing causality, particularly if they have only small effect

sizes. Using multiple genetic variants can increase the strength of the instrument, but also increases

the risk of violating the MR assumptions.

We performed bidirectional, two-sample MR between our meta-analysis results for depression and all

available traits which had a significant genetic correlation with depression (identified in the previous

section). Traits directly related to or including depression (major depressive disorder, depressive

symptoms and PGC cross disorder) were excluded due to potential bias. The genetic instruments for

depression consisted of the independent, genome-wide significant variants, their effect sizes and

standard errors, as estimated in our genome-wide meta-analysis. Summary statistics from genome-

wide association studies for the other traits were sourced from either publicly available datasets or

from the MR-Base database [16]. Overlapping datasets for the exposure and the outcome can lead to

bias and inflation of causal estimates [17]. To mitigate this, when the source of the other trait included

UK Biobank, 23andMe or any of the studies that contributed to PGC_139k, these studies were

removed from our meta-analysis of depression; for example, the neuroticism trait from van den Berg,

et al. [19] was assessed against a meta-analysis of depression using UK Biobank and 23andMe_307k

only. Where UK Biobank, 23andMe and PGC_139k were all included in the genome-wide

association study of the other trait then an alternative study was sought for that other trait.

All analyses were performed using the MR-Base "TwoSampleMR" v0.4.9 package[16] in R. To avoid

bias in the MR estimates due to linkage disequilibrium (r^2), clumping was applied using the

"clump_data" function with an $r^2 < 0.001$. Genetic variants were required to be available in both the

exposure and outcome traits and were harmonised using the default parameters within the

TwoSampleMR package. Following this harmonisation, we only examined causal relationships where

there were at least 30 instrumental genetic variables.

Directional horizontal pleiotropy, where a genetic instrument has an effect on an outcome

independent of its influence on the exposure, can be a problem in MR analysis, particularly when

multiple genetic variants of unknown function are used. We therefore firstly tested for directional

horizontal pleiotropy using the MR Egger intercept test, as previously described by Hagenaars, Gale,

Deary and Harris [104]. If the MR Egger intercept test had a significant *P*-value (*P* < 0.05) then it was

excluded from the analysis. However, no tests were excluded due to directional horizontal pleiotropy.

The second analysis conducted was a variant heterogeneity test for global horizontal pleiotropy.

Variant heterogeneity is an important metric, but high heterogeneity doesn't necessarily mean bias or

unreliable results; for example, every instrumental variable could have horizontal pleiotropic effects

but if they have a mean effect of 0 then there will be no bias, just larger standard errors due to more

noise. For analyses that had evidence of high variant heterogeneity (*P* < 0.05), additional sensitivity

MR tests were conducted. The sensitivity tests that were used were the MR Egger test and the

weighted median test to examine whether the effect estimate was consistent.

The principal MR test of a causal effect was conducted using inverse-variance weighted (IVW)

regression. This method is based on a regression of the exposure and the outcome which assumes the

intercept is constrained to zero, and produces a causal estimate of the exposure-outcome association.

Where there was no evidence of global horizontal pleiotropy (*P* ≥ 0.05) from the second analysis (see

previous paragraph), a FDR adjusted[103] *P*-value < 0.01 from the IVW test was required for evidence

of a causal effect. Where there was evidence of global horizontal pleiotropy (P < 0.05) from the

second analysis, additional evidence was also sought from the sensitivity tests (MR Egger test and the

weighted median test). To ensure that a causal effect was not driven by a single variant a 'leave one

variant out' IVW regression analysis was conducted with the least significant observed *P*-value used

to assess whether significance was maintained. We tested the causal effect of depression on 24 other

traits, and the causal effect of 9 other traits on depression.

Partitioned heritability analyses

We used stratified LDSC regression to estimate the SNP-based heritability of depression, using the

sample prevalence as the population prevalence (0.302), and then examined the heritability of

partitioned functional categories [21]. This method assigns variants into 24 functional categories and

then evaluates the contribution of each category to the overall heritability of a trait. A category is

enriched for heritability if the variants with high LD to that category have elevated χ^2 statistics. The 24

categories are described in full in the Finucane et al. paper [21]. Briefly, they include genetic annotations

from ReqSeq gene models [105], transcription factor binding sites from ENCODE [105, 106], Roadmap

epigenomics annotations [107], super-enhancers [108], evolutionarily conserved regions [109] and FANTOM5

enhancers [110]. Heritability enrichment is defined as the proportion of heritability assigned to a

functional category divided by the proportion of variants in that category. Cell-type specific

annotations were also analysed for variants in four histone marks: H3K4me1, H3K4me3, H3K9ac and

H3K27ac and these were grouped into 9 cell type groups: central nervous system (CNS),

cardiovascular, connective tissue or bone, gastrointestinal, immune or hematopoietic, kidney, liver,

skeletal muscle, or other. Given the strong contribution to depression heritability from the CNS we

used LDSC applied to specifically expressed genes (LDSC-SEG) [111] across 13 brain regions from the

GTEx gene-expression database and brain cell-types (neuronal, astrocyte, oligodendrocyte) using

specifically expressed genes derived from mouse-forebrain [112].

Gene and gene-set analyses

A gene-based analysis was applied to the results from our meta-analysis using Multi-marker Analysis

of GenoMic Annotation (MAGMA) [24] to assess the simultaneous effect of multiple genetic variants

on 17,842 genes. To account for LD the European panel of the 1,000 Genomes data (phase 3) [113] was

used as a reference panel. Genetic variants were assigned to genes based on their position according to

the NCBI 37.3 build. To identify those genes that were genome-wide significant a *P*-value threshold

was calculated by applying a Bonferroni correction ($\alpha = 0.05 / 17,842$; $P < 2.80 \times 10^{-6}$). Regional

visualisation plots were produced using the online LocusZoom platform [114].

A gene-set analysis was then performed on our gene-based results using gene annotation files from

the Gene Ontology Consortium (http://geneontology.org/) [115] and the Molecular Signatures Database

v5.2 [116]. The annotation file includes gene-sets covering three ontologies; molecular function, cellular

components, and biological function and consisted of 5,917 gene-sets. To correct for multiple testing,

we used the MAGMA default setting of 10,000 permutations, and applied a Bonferroni correction (α

$= 0.05 / 5,917$; $P < 8.45 \times 10^{-6}$). Additional gene-sets were obtained from Skene, et al. [25] providing

expression-weighted enrichment for seven brain cell-types (astrocytes ependymal, endothelial mural,

interneurons, microglia, oligodendrocytes, somatosensory pyramidal neurons, and hippocampus CA1

pyramidal neurons). These brain cell-types were assessed in MAGMA using the default setting of

10,000 permutations with a Bonferroni correction ($\alpha = 0.05 / 7$; $P < 7.14 \times 10^{-3}$) used to assess

significance.

Drug - gene interactions

We examined genes that were identified as significantly associated ($P < 2.80 \times 10^{-6}$) with depression

using MAGMA [24] for interactions with prescribed medications using the Drug Gene Interaction

Database (dgidb.genome.wustl.edu) v3.0 [26]. The Anatomical Therapeutic Chemical (ATC)

classification system was used to determine the second level classification of each drug identified.

ATC classifications for each drug were obtained from the Kyoto Encyclopaedia of Genes and

Genomics (https://www.genome.jp/kegg/drug/) as the primary source of information with additional

classifications obtained from the World Health Organisation Collaborating Centre for Drug Statistics

Methodology (https://www.whocc.no/atc_ddd_index/). Visualisation of the number of interactions

between each gene associated with depression and each second level ATC classification was

undertaken using the R package circlize v0.4.1 [117].

Acknowledgements

This research has been conducted using the UK Biobank resource – application number 4844. We are

grateful to the UK Biobank and all its voluntary participants. The UK Biobank study was conducted

under generic approval from the NHS National Research Ethics Service (approval letter dated 17th

June 2011, Ref 11/NW/0382). All participants gave full informed written consent. The authors

acknowledge the help, advice, and support from all members of the UK Biobank Psychiatric Genetics

Group.

We are grateful to the participants and research teams behind the Psychiatric Genomics Consortium,

UK Biobank and 23andMe. We thank the following members of the 23andMe Research Team:

Michelle Agee, Babak Alipanahi, Adam Auton, Robert K. Bell, Katarzyna Bryc, Sarah L. Elson,

Pierre Fontanillas, Nicholas A. Furlotte, Barry Hicks, Karen E. Huber, Ethan M. Jewett, Yunxuan

Jiang, Aaron Kleinman, Keng-Han Lin, Nadia K. Litterman, Matthew H. McIntyre, Joanna L.

Mountain, Elizabeth S. Noblin, Carrie A.M. Northover, Steven J. Pitts, G. David Poznik, J. Fah

Sathirapongsasuti, Olga V. Sazonova, Janie F. Shelton, Suyash Shringarpure, Joyce Y. Tung,

Vladimir Vacic, Xin Wang, and Catherine H. Wilson. We would like to thank Dr. Nathan Skene for

his advice on analysing the expression-weighted enrichment for brain cell-types.

AMMcI, and IJD acknowledge support from the Wellcome Trust (Wellcome Trust Strategic Award

"STratifying Resilience and Depression Longitudinally" (STRADL) Reference 104036/Z/14/Z and

the Dr Mortimer and Theresa Sackler Foundation. IJD is supported by the Centre for Cognitive

Ageing and Cognitive Epidemiology, which is funded by the Medical Research Council and the

Biotechnology and Biological Sciences Research Council (MR/K026992/1). This investigation

represents independent research part-funded by the National Institute for Health Research (NIHR)

Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's

College London. The views expressed are those of the authors and not necessarily those of the NHS,

the NIHR or the Department of Health. GH is supported by the Wellcome Trust [208806/Z/17/Z].

DJS supported by Lister Institute Prize Fellowship 2016-2021. NRW acknowledges NMHRC grants

1078901 and 1087889.

The PGC has received major funding from the US National Institute of Mental Health and the US

National Institute of Drug Abuse (U01 MH109528 and U01 MH1095320).

Author Contributions

D.M.H., D.J.S., G.B., C.M.L., and A.M.McI. conceived the research project. M.J.A., J.R.I.C., J.W.,

D.J.S., G.B., and A.M.McI. determined the variables that formed the depression phenotypes within

UK Biobank. D.M.H., M.J.A., J.R.I.C., R.E.M., and G.D. applied quality control to the UK Biobank

data. M.J.A. ran the association analysis in UK Biobank. 23andMe R.T. provided the summary

statistics from the Hyde, et al. [8] analysis. M.D.D.W.G.P.G.C., M.T., S.R., P.F.S., N.R.W., and C.M.L.

provided the summary statistics from the Wray, et al. [9] analysis with additional removal of

overlapping cohorts. D.M.H. ran the meta-analysis of the three contributing cohorts. C.T. conducted

the replication analysis in 23andMe for the significant variants in the meta-analysis. T-K.C. and

E.Y.X. conducted the polygenic risk score analysis. T-K.C. and D.M.H calculated the genetic

correlations and partitioned the heritability component of depression. D.M.H. and G.H. performed the

Mendelian randomization analyses. D.M.H. ran the MAGMA-based analyses with J.W., M.S.,

E.M.W., C.A., X.S., and M.C.B. examining the genes and gene-sets identified. J.D.H., D.M.H., and

A.M.McI. conducted the gene x drug interaction analysis. I.J.D., D.J.P., H.C.W., C.S.H., S.R., D.J.S.,

P.F.S., N.R.W., E.M.B., G.B., C.M.L., and A.M.McI. provided expertise of association study

methodology and statistical analysis. D.M.H. oversaw the research project and serves as the primary

contact for all communication. All authors commented on the manuscript.

Competing interests

IJD is a participant in UK Biobank. Chao Tian, David A. Hinds, and Members of the 23andMe

Research Team are employees of 23andMe, Inc. The authors report that no other conflicts of interest

exist.

Data availability

References

1.   World Health Organization. Depression and other common mental disorders. (2017).
2.   Kessler, R.C., *et al.* Lifetime prevalence and age-of-onset distributions of dsm-iv disorders in
the national comorbidity survey replication. *Archives of General Psychiatry* **62**, 593-602 (2005).

3.   Sullivan, P.F., Neale, M.C. & Kendler, K.S. Genetic epidemiology of major depression: review
and meta-analysis. *American Journal of Psychiatry* **157**, 1552-1562 (2000).
4.   Major Depressive Disorder Working Group of the Psychiatric GWAS Consortium. A mega-
analysis of genome-wide association studies for major depressive disorder. *Molecular Psychiatry* **18**,
497-511 (2013).
5.   Howard, D.M.*, et al.* Genome-wide association study of depression phenotypes in UK
Biobank identifies variants in excitatory synaptic pathways. *Nature Communications* **9**, 1470 (2018).
6.   Hek, K.*, et al.* A genome-wide association study of depressive symptoms. *Biological*
Psychiatry **73**, 10.1016/j.biopsych.2012.1009.1033 (2013).
7.   Converge Consortium. Sparse whole-genome sequencing identifies two loci for major
depressive disorder. *Nature* **523**, 588-591 (2015).
8.   Hyde, C.L.*, et al.* Identification of 15 genetic loci associated with risk of major depression in
individuals of European descent. *Nature Genetics* **48**, 1031-1036 (2016).
9.   Wray, N.R.*, et al.* Genome-wide association analyses identify 44 risk variants and refine the
genetic architecture of major depression. *Nature Genetics* **50**, 668-681 (2018).
10.   Yang, J.*, et al.* Conditional and joint multiple-SNP analysis of GWAS summary statistics
identifies additional variants influencing complex traits. *Nature Genetics* **44**, 369 (2012).
11.   Bulik-Sullivan, B.K.*, et al.* LD Score regression distinguishes confounding from polygenicity in
genome-wide association studies. *Nature Genetics* **47**, 291-295 (2015).
12.   Euesden, J., Lewis, C.M. & O'Reilly, P.F. PRSice: Polygenic Risk Score software. *Bioinformatics*
31, 1466-1468 (2015).
13.   Smith, B.H.*, et al.* Cohort profile: Generation Scotland: Scottish Family Health Study
(GS:SFHS). The study, its participants and their potential for genetic research on health and illness.
International Journal of Epidemiology **42**, 689-700 (2013).
14.   Smith, D.*, et al. Depression and multimorbidity: a cross-sectional study of 1,751,841 patients*
in primary care (2014).
15.   Zheng, J.*, et al.* LD Hub: a centralized database and web interface to perform LD score
regression that maximizes the potential of summary level GWAS data for SNP heritability and genetic
correlation analysis. *Bioinformatics* **33**, 272-279 (2017).
16.   Hemani, G.*, et al.* The MR-Base platform supports systematic causal inference across the
human phenome. *eLife* **7**, e34408 (2018).
17.   Burgess, S., Davies, N.M. & Thompson, S.G. Bias due to participant overlap in two-sample
Mendelian randomization. *Genetic Epidemiology* **40**, 597-608 (2016).
18.   Bowden, J., Davey Smith, G. & Burgess, S. Mendelian randomization with invalid
instruments: effect estimation and bias detection through Egger regression. *International Journal of*
Epidemiology **44**, 512-525 (2015).
19.   van den Berg, S.M.*, et al.* Harmonization of neuroticism and extraversion phenotypes across
inventories and cohorts in the Genetics of Personality Consortium: an application of item response
theory. *Behavior Genetics* **44**, 295-313 (2014).
20.   The Tobacco Genetics Consortium. Genome-wide meta-analyses identify multiple loci
associated with smoking behavior. *Nature Genetics* **42**, 441-447 (2010).
21.   Finucane, H.K.*, et al.* Partitioning heritability by functional annotation using genome-wide
association summary statistics. *Nat Genet* **47**, 1228-1235 (2015).
22.   McGee, S.L., Fairlie, E., Garnham, A.P. & Hargreaves, M. Exercise-induced histone
modifications in human skeletal muscle. *The Journal of Physiology* **587**, 5951-5958 (2009).
23.   Agudelo, Leandro Z.*, et al.* Skeletal muscle PGC-1α1 modulates kynurenine metabolism and
mediates resilience to stress-induced depression. *Cell* **159**, 33-45 (2014).
24.   de Leeuw, C.A., Mooij, J.M., Heskes, T. & Posthuma, D. MAGMA: generalized gene-set
analysis of GWAS data. *PLOS Computational Biology* **11**, e1004219 (2015).
25.   Skene, N.G.*, et al.* Genetic identification of brain cell types underlying schizophrenia. *Nature*
Genetics **50**, 825-833 (2018).

26.   Griffith, M., *et al.* DGIdb: mining the druggable genome. *Nature Methods* **10**, 1209-1210
(2013).
27.   Cross-Disorder Group of the Psychiatric Genomics Consortium. Identification of risk loci with
shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet* **381**, 1371-1379
(2013).
28.   Anttila, V., *et al.* Analysis of shared heritability in common disorders of the brain. *Science* **360**
(2018).
29.   The International Schizophrenia Consortium. Common polygenic variation contributes to risk
of schizophrenia and bipolar disorder. *Nature* **460**, 748 (2009).
30.   Luciano, M., *et al.* Association analysis in over 329,000 individuals identifies 116
independent variants influencing neuroticism. *Nature Genetics* **50**, 6-11 (2018).
31.   Okbay, A., *et al.* Genetic variants associated with subjective well-being, depressive
symptoms, and neuroticism identified through genome-wide analyses. *Nature Genetics* **48**, 624-633
(2016).
32.   Markett, S., *et al.* Variation on the dopamine D2 receptor gene (DRD2) is associated with
basal ganglia-to-frontal structural connectivity. *NeuroImage* **155**, 473-479 (2017).
33.   Quarto, T., *et al.* Interaction between DRD2 variation and sound environment on mood and
emotion-related brain activity. *Neuroscience* **341**, 9-17 (2017).
34.   Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights
from 108 schizophrenia-associated genetic loci. *Nature* **511**, 421-427 (2014).
35.   Pergola, G., *et al.* DRD2 co-expression network and a related polygenic index predict
imaging, behavioral and clinical phenotypes linked to schizophrenia. *Translational Psychiatry* **7**,
e1006 (2017).
36.   Whitmer, A.J. & Gotlib, I.H. Depressive rumination and the C957T polymorphism of the DRD2
gene. *Cognitive, Affective, & Behavioral Neuroscience* **12**, 741-747 (2012).
37.   Wagnon, J.L., *et al.* CELF4 regulates translation and local abundance of a vast set of mRNAs,
including genes associated with regulation of synaptic function. *PLOS Genetics* **8**, e1003067 (2012).
38.   Ladd, A.N., Charlet-B, N. & Cooper, T.A. The CELF family of RNA binding proteins is
implicated in cell-specific and developmentally regulated alternative splicing. *Molecular and Cellular*
Biology **21**, 1285-1296 (2001).
39.   Barone, R., *et al.* Familial 18q12.2 deletion supports the role of RNA-binding protein CELF4 in
autism spectrum disorders. *American Journal of Medical Genetics Part A* **173**, 1649-1655 (2017).
40.   Halgren, C., *et al.* Haploinsufficiency of CELF4 at 18q12.2 is associated with developmental
and behavioral disorders, seizures, eye manifestations, and obesity. *European Journal of Human*
Genetics **20**, 1315-1319 (2012).
41.   Fogel, B.L., *et al.* RBFOX1 regulates both splicing and transcriptional networks in human
neuronal development. *Human Molecular Genetics* **21**, 4171-4186 (2012).
42.   Berto, S., Usui, N., Konopka, G. & Fogel, B.L. ELAVL2-regulated transcriptional and splicing
networks in human neurons link neurodevelopment and autism. *Human Molecular Genetics* **25**,
2451-2464 (2016).
43.   Chang, H., *et al.* Further evidence of VRK2 rs2312147 associated with schizophrenia. *The*
World Journal of Biological Psychiatry **17**, 457-466 (2016).
44.   Steinberg, S., *et al.* Common variants at VRK2 and TCF4 conferring risk of schizophrenia.
Human Molecular Genetics **20**, 4076-4081 (2011).
45.   Li, M., *et al.* Meta-analysis and brain imaging data support the involvement of VRK2
(rs2312147) in schizophrenia susceptibility. *Schizophrenia Research* **142**, 200-205 (2012).
46.   Potkin, S.G., *et al.* Gene discovery through imaging genetics: identification of two novel
genes associated with schizophrenia. *Molecular Psychiatry* **14**, 416-428 (2009).
47.   Mitchell, A.C., *et al.* MEF2C transcription factor is associated with the genetic and epigenetic
risk architecture of schizophrenia and improves cognition in mice. *Molecular Psychiatry* (2017).

48.   Mossakowska-Wójcik, J., Orzechowska, A., Talarowska, M., Szemraj, J. & Gałecki, P. The
importance of TCF4 gene in the etiology of recurrent depressive disorders. *Progress in Neuro-*
Psychopharmacology and Biological Psychiatry **80**, 304-308 (2018).
49.   Kennedy, Andrew J., *et al.* TCF4 regulates synaptic plasticity, DNA methylation, and memory
function. *Cell Reports* **16**, 2666-2685 (2016).
50.   Rannals, Matthew D., *et al.* Psychiatric risk gene transcription factor 4 regulates intrinsic
excitability of prefrontal neurons via repression of SCN10a and KCNQ1. *Neuron* **90**, 43-55 (2016).
51.   Forrest, M.P., *et al.* The psychiatric risk gene Transcription Factor 4 (TCF4) regulates
neurodevelopmental pathways associated with schizophrenia, autism, and intellectual disability.
Schizophrenia Bulletin, sbx164-sbx164 (2017).
52.   Chronis-Tuscano, A., Molina, B.G., Pelham, W.E. & et al. Very early predictors of adolescent
depression and suicide attempts in children with attention-deficit/hyperactivity disorder. *Archives of*
General Psychiatry **67**, 1044-1051 (2010).
53.   Neale, B.M., *et al.* Case-control genome-wide association study of attention-
deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry* **49**,
906-920 (2010).
54.   Lesch, K.-P., *et al.* Molecular genetics of adult ADHD: converging evidence from genome-
wide association and extended pedigree linkage studies. *Journal of Neural Transmission* **115**, 1573-
1585 (2008).
55.   Salatino-Oliveira, A., *et al.* Cadherin-13 gene is associated with hyperactive/impulsive
symptoms in attention/deficit hyperactivity disorder. *American Journal of Medical Genetics Part B:*
Neuropsychiatric Genetics **168**, 162-169 (2015).
56.   Lionel, A.C., *et al.* Disruption of the ASTN2/TRIM32 locus at 9q33.1 is a risk factor in males
for autism spectrum disorders, ADHD and other neurodevelopmental phenotypes. *Human Molecular*
Genetics **23**, 2752-2768 (2014).
57.   Wilson, P.M., Fryer, R.H., Fang, Y. & Hatten, M.E. ASTN2, a novel member of the astrotactin
gene family, regulates the trafficking of ASTN1 during glial-guided neuronal migration. *The Journal of*
Neuroscience **30**, 8529-8540 (2010).
58.   Mota, N.R., *et al.* NCAM1-TTC12-ANKK1-DRD2 gene cluster and the clinical and genetic
heterogeneity of adults with ADHD. *American Journal of Medical Genetics Part B: Neuropsychiatric*
Genetics **168**, 433-444 (2015).
59.   Heinrich, H., *et al.* Attention, cognitive control and motivation in ADHD: Linking event-
related brain potentials and DNA methylation patterns in boys at early school age. *Scientific Reports*
7, 3823 (2017).
60.   Kaltiala-Heino, R., Kosunen, E. & Rimpelä, M. Pubertal timing, sexual behaviour and self-
reported depression in middle adolescence. *Journal of Adolescence* **26**, 531-545 (2003).
61.   Joinson, C., Heron, J., Araya, R. & Lewis, G. Early Menarche and Depressive Symptoms From
Adolescence to Young Adulthood in a UK Cohort. *Journal of the American Academy of Child &*
Adolescent Psychiatry **52**, 591-598.e592 (2013).
62.   Sequeira, M.-E., Lewis, S.J., Bonilla, C., Smith, G.D. & Joinson, C. Association of timing of
menarche with depressive symptoms and depression in adolescence: Mendelian randomisation
study. *The British Journal of Psychiatry* **210**, 39-46 (2017).
63.   Ong, K.K., *et al.* Genetic variation in LIN28B is associated with the timing of puberty. *Nature*
Genetics **41**, 729-733 (2009).
64.   Perry, J.R.B., *et al.* Meta-analysis of genome-wide association data identifies two loci
influencing age at menarche. *Nature Genetics* **41**, 648-650 (2009).
65.   Sulem, P., *et al.* Genome-wide association study identifies sequence variants on 6q21
associated with age at menarche. *Nature Genetics* **41**, 734-738 (2009).
66.   Yu, J., *et al.* Induced pluripotent stem cell lines derived from human somatic cells. *Science*
318, 1917-1920 (2007).

67.   Moss, E.G., Lee, R.C. & Ambros, V. The cold shock domain protein LIN-28 controls
developmental timing in C. elegans and is regulated by the lin-4 RNA. *Cell* **88**, 637-646 (1997).
68.   Wei, Y.B., *et al.* Elevation of Il6 is associated with disturbed let-7 biogenesis in a genetic
model of depression. *Translational Psychiatry* **6**, e869 (2016).
69.   Boden, J.M., Fergusson, D.M. & Horwood, L.J. Cigarette smoking and depression: tests of
causal linkages using a longitudinal birth cohort. *British Journal of Psychiatry* **196**, 440-446 (2010).
70.   Kang, E. & Lee, J. A longitudinal study on the causal association between smoking and
depression. *J Prev Med Public Health* **43**, 193-204 (2010).
71.   Wootton, R.E., *et al.* Causal effects of lifetime smoking on risk for depression and
schizophrenia: Evidence from a Mendelian randomisation study. *bioRxiv*, 381301 (2018).
72.   Lewis, S.J., *et al.* Smoking is associated with, but does not cause, depressed mood in
pregnancy – A Mendelian randomization study. *PLoS ONE* **6**, e21689 (2011).
73.   Bjørngaard, J.H., *et al.* The causal role of smoking in anxiety and depression: a Mendelian
randomization analysis of the HUNT study. *Psychological Medicine* **43**, 711-719 (2012).
74.   Taylor, A.E., *et al.* Investigating the possible causal association of smoking with depression
and anxiety using Mendelian randomisation meta-analysis: the CARTA consortium. *BMJ Open* **4**
(2014).
75.   Wium-Andersen, M.K., Ørsted, D.D. & Nordestgaard, B.G. Tobacco smoking is causally
associated with antipsychotic medication use and schizophrenia, but not with antidepressant
medication use or depression. *International Journal of Epidemiology* **44**, 566-577 (2015).
76.   Munafò, M.R. & Araya, R. Cigarette smoking and depression: a question of causation. *British
Journal of Psychiatry* **196**, 425-426 (2018).
77.   Hughes, J.R., Stead, L.F. & Lancaster, T. Anxiolytics for smoking cessation. *Cochrane
Database of Systematic Reviews* (2000).
78.   Munafò, M.R., Hitsman, B., Rende, R., Metcalfe, C. & Niaura, R. Effects of progression to
cigarette smoking on depressed mood in adolescents: evidence from the National Longitudinal Study
of Adolescent Health. *Addiction* **103**, 162-171 (2008).
79.   Bonci, A. & Hopf, F.W. The dopamine D2 receptor: new surprises from an old friend. *Neuron*
47, 335-338 (2005).
80.   Dailly, E., Chenu, F., Renard, C.E. & Bourin, M. Dopamine, depression and antidepressants.
Fundamental & Clinical Pharmacology **18**, 601-607 (2004).
81.   Bi, L.-L., *et al.* Amygdala NRG1-ErbB4 is critical for the modulation of anxiety-like behaviors.
Neuropsychopharmacology **40**, 974-986 (2015).
82.   Fuxe, K. & Borroto-Escuela, D.O. Basimglurant for treatment of major depressive disorder: a
novel negative allosteric modulator of metabotropic glutamate receptor 5. *Expert Opinion on
Investigational Drugs* **24**, 1247-1260 (2015).
83.   Espallergues, J., Temsamani, J., Laruelle, C., Urani, A. & Maurice, T. The antidepressant-like
effect of the 3β-hydroxysteroid dehydrogenase inhibitor trilostane involves a regulation of β-type
estrogen receptors. *Psychopharmacology* **214**, 455-463 (2011).
84.   Estrada-Camarena, E., Fernández-Guasti, A. & López-Rubalcava, C. Antidepressant-like effect
of different estrogenic compounds in the forced swimming test. *Neuropsychopharmacology* **28**, 830
(2002).
85.   Shen, E.Y., *et al.* Neuronal deletion of Kmt2a/Mll1 histone methyltransferase in ventral
striatum is associated with defective spike-timing-dependent striatal synaptic plasticity, altered
response to dopaminergic drugs, and increased anxiety. *Neuropsychopharmacology* **41**, 3103-3113
(2016).
86.   Dunn, E.C., *et al.* Genetic determinants of depression: Recent findings and future directions.
Harvard Review of Psychiatry **23**, 1-18 (2015).
87.   Biernacka, J.M., *et al.* The International SSRI Pharmacogenomics Consortium (ISPC): a
genome-wide association study of antidepressant treatment response. *Translational Psychiatry* **5**,
e553 (2015).

88.   Milaneschi, Y., *et al.* Polygenic dissection of major depression clinical heterogeneity.
Molecular Psychiatry **21**, 516-522 (2016).
89.   Bycroft, C., *et al.* Genome-wide genetic data on ~500,000 UK Biobank participants. *bioRxiv*,
166298 (2017).
90.   McCarthy, S., *et al.* A reference panel of 64,976 haplotypes for genotype imputation. *Nature*
Genetics **48**, 1279-1283 (2016).
91.   Manichaikul, A., *et al.* Robust relationship inference in genome-wide association studies.
Bioinformatics **26**, 2867-2873 (2010).
92.   Ripke, S. GWAS genotypic overlap test without sharing genotypes.
https://personal.broadinstitute.org/sripke/share_links/checksums_download/ (2017).
93.   Smith, D.J., *et al.* Prevalence and characteristics of probable major depression and bipolar
disorder within UK Biobank: cross-sectional study of 172,751 participants. *PLoS ONE* **8**, e75362
(2013).
94.   Allen, N.E., Sudlow, C., Peakman, T. & Collins, R. UK biobank data: come and get it. *Science*
Translational Medicine **6**, 224ed224 (2014).
95.   Willer, C.J., Li, Y. & Abecasis, G.R. METAL: fast and efficient meta-analysis of genomewide
association scans. *Bioinformatics* **26**, 2190-2191 (2010).
96.   Chang, C.C., *et al.* Second-generation PLINK: rising to the challenge of larger and richer
datasets. *GigaScience* **4**, 7 (2015).
97.   Quinlan, A.R. BEDTools: the Swiss-army tool for genome feature analysis. *Current protocols*
in bioinformatics / editoral board, Andreas D. Baxevanis ... [et al.] **47**, 11.12.11-11.12.34 (2014).
98.   First, M.B., Spitzer, R.L., Gibbon Miriam. & Williams, J.B.W. Structured Clinical Interview for
DSM-IV-TR Axis I Disorders, Research Version, Patient Edition. (SCID-I/P) (2002).
99.   Fernandez-Pujals, A.M., *et al.* Epidemiology and heritability of major depressive disorder,
stratified by age of onset, sex, and illness course in generation scotland: scottish family health study
(GS:SFHS). *PLoS ONE* **10**, e0142197 (2015).
100.   International Genetics of Ankylosing Spondylitis Consortium, *et al.* Identification of multiple
risk variants for ankylosing spondylitis through high-density genotyping of immune-related loci.
Nature Genetics **45**, 730 (2013).
101.   Nakagawa, S. & Schielzeth, H. A general and simple method for obtaining R2 from
generalized linear mixed-effects models. *Methods in Ecology and Evolution* **4**, 133-142 (2013).
102.   Bulik-Sullivan, B., *et al.* An atlas of genetic correlations across human diseases and traits.
Nature Genetics **47**, 1236-1241 (2015).
103.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and powerful
approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**,
289-300 (1995).
104.   Hagenaars, S.P., Gale, C.R., Deary, I.J. & Harris, S.E. Cognitive ability and physical health: a
Mendelian randomization study. *Scientific Reports* **7**, 2651 (2017).
105.   Gusev, A., *et al.* Partitioning heritability of regulatory and cell-type-specific variants across 11
common diseases. *Am J Hum Genet* **95**, 535-552 (2014).
106.   Encode Project Consortium. An integrated encyclopedia of DNA elements in the human
genome. *Nature* **489**, 57-74 (2012).
107.   Roadmap Epigenomics Consortium., *et al.* Integrative analysis of 111 reference human
epigenomes. *Nature* **518**, 317-330 (2015).
108.   Hnisz, D., *et al.* Super-enhancers in the control of cell identity and disease. *Cell* **155**, 934-947
(2013).
109.   Lindblad-Toh, K., *et al.* A high-resolution map of human evolutionary constraint using 29
mammals. *Nature* **478**, 476-482 (2011).
110.   Andersson, R., *et al.* An atlas of active enhancers across human cell types and tissues. *Nature*
507, 455-461 (2014).

111. Finucane, H.K., *et al.* Heritability enrichment of specifically expressed genes identifies
disease-relevant tissues and cell types. *Nature Genetics* **50**, 621-629 (2018).
112. Cahoy, J.D., *et al.* A transcriptome database for astrocytes, neurons, and oligodendrocytes: a
new resource for understanding brain development and function. *The Journal of Neuroscience* **28**,
264-278 (2008).
113. The 1000 Genomes Project Consortium. A global reference for human genetic variation.
Nature **526**, 68-74 (2015).
114. Pruim, R.J., *et al.* LocusZoom: regional visualization of genome-wide association scan results.
Bioinformatics **26**, 2336-2337 (2010).
115. The Gene Ontology Consortium. Expansion of the Gene Ontology knowledgebase and
resources. *Nucleic Acids Res* **45**, D331-D338 (2017).
116. Subramanian, A., *et al.* Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences* **102**,
15545-15550 (2005).
117. Gu, Z., Gu, L., Eils, R., Schlesner, M. & Brors, B. circlize implements and enhances circular
visualization in R. *Bioinformatics* **30**, 2811-2812 (2014).

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053 **Figure Titles**

1054 Figure. 1
1055 (A) Manhattan plot of the observed $-\log_{10}$ P-values of each variant for an association with depression
1056 in the meta-analysis (n = 807,553; 246,363 cases and 561,190 controls). Variants are positioned
1057 according to the GRCh37 assembly

1058

Figure. 2
(A) Significance of cell type enrichment using a partitioned heritability approach. The dashed line
represents statistical significance after Bonferroni correction (-log$_{10}P$ = 2.36) and * indicates
significant enrichment for that cell type
(B) Significance of enrichment estimates, based on genetic summary statistics, for brain regions using
GTEx. The dashed line represents the Bonferroni cut-off for statistical significance (-log$_{10}P$ = 2.41)
and * indicates significant enrichment for that brain region
(C) Significant enrichment P-values, based on genetic summary statistics, for brain cell regions using
GTEx overlaid on a physical representation of brain anatomy

The pseudo-coloring used in Figure 2C highlights the P-values of the regions of the brain in red that
were significantly enriched ($P < 0.05$) for depression variants

1070

Figure. 3
Chord diagram of genes significantly associated ($P < 2.80 \times 10^{-6}$) with depression and the second
level Anatomical Therapeutic Chemical classifications to which interacting drugs have been assigned.
The width of each line is determined by the number of drugs known to interact with each gene. The
genes are ordered by significance with depression with those most significantly associated located at
the top of the diagram

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091   **Tables**

1092   Table 1
1093   Sample sizes and the proportion of males and females of the depression cohorts used in the meta-
1094   analysis and the replication cohort

Cohort	Study	Cases (Prop. Male, Female)	Controls (Prop. Male, Female)	Total (Prop. Male, Female)
23andMe_307k	Hyde, et al. [8]	75,607 (0.38, 0.62)	231,747 (0.56, 0.44)	307,354 (0.52, 0.48)
UK Biobank	Howard, et al. [5]	127,552 (0.35, 0.65)	233,763 (0.52, 0.48)	361,315 (0.46, 0.54)
PGC_139k†	Wray, et al. [9]	43,204	95,680	138,884
Meta-analysis		**246,363**	**561,190**	**807,553**
Replication	Unpublished	474,574 (0.30, 0.70)	1,032,579 (0.50, 0.50)	1,507,153 (0.44, 0.56)

1095

1096   † Proportion of males and females not reported for PGC_139k

1097

1098   Table 2
1099   Gene pathways with a significant effect ($P_{corrected} < 0.05$) on depression identified through gene-set
1100   enrichment analysis in MAGMA [24]

Pathway	Number of genes	Beta	Standard error	*P*-value	$P_{Corrected}$
GO_POSTSYNAPSE	351	0.34	0.056	8.15×10^{-10}	4.82×10^{-6}
GO_SYNAPSE	707	0.219	0.040	2.47×10^{-8}	1.46×10^{-4}
GO_NEURON_SPINE	114	0.509	0.096	6.13×10^{-8}	3.63×10^{-4}
GO_EXCITATORY_SYNAPSE	181	0.405	0.077	8.29×10^{-8}	4.90×10^{-4}
GO_SYNAPSE_PART	571	0.225	0.044	1.24×10^{-7}	7.32×10^{-4}
GO_BEHAVIOR	489	0.245	0.049	2.32×10^{-7}	1.38×10^{-3}
GO_COGNITION	238	0.346	0.069	2.77×10^{-7}	1.64×10^{-3}
GO_NEURON_PROJECTION	883	0.18	0.036	3.39×10^{-7}	2.01×10^{-3}
GO_MODULATION_OF_SYNAPTIC_TRANSMISSION	283	0.298	0.061	4.25×10^{-7}	2.52×10^{-3}
GO_REGULATION_OF_SYNAPSE_STRUCTURE_OR_ACTIVITY	219	0.335	0.071	1.19×10^{-6}	7.02×10^{-3}
GO_REGULATION_OF_SYNAPTIC_PLASTICITY	133	0.412	0.089	1.95×10^{-6}	0.012
GO_NEURON_PART	1183	0.142	0.031	3.17×10^{-6}	0.019
GO_DENDRITE	420	0.229	0.052	4.76×10^{-6}	0.028
GO_SINGLE_ORGANISM_BEHAVIOR	367	0.244	0.056	5.86×10^{-6}	0.035
Somatosensory Pyramidal Neurons	1217	0.121	0.031	3.87×10^{-5}	2.71×10^{-4}

1101

1102   Pathways with a "GO" prefix were obtained from the Gene Ontology Consortium [115]. The Somatosensory
1103   Pyramidal Neurons gene-set was obtained from Skene, et al. [25].

1104

1105

1106

1107   **Supplementary Information Titles and Legends**

1108

1109   Supplementary Information

1110   General directionality agreement of variants that have previously been associated with depression

1111   Cohort information for the 23andMe replication dataset

1112   Members of the Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium
1113   and their affiliations

1114

1115   Supplementary Figure 1

1116   Quantile-quantile plot of the observed *P*-values on those expected for each genetic variants
1117   association with depression in our meta-analysis

1118

1119   Supplementary Figure 2

1120   Odds ratios and 95% confidence intervals for Major Depressive Disorder (MDD) in Generation
1121   Scotland based on polygenic risk score (PRS) deciles calculated from the current meta-analysis of
1122   depression and from the summary statistics from the genome-wide association study of major
1123   depression conducted by Wray, et al. [9].

1124

Supplementary Figure 3

Significant genetic correlations (r_G; $P < 0.01$, after false discovery rate correction) between depression
and other behavioural and disease related traits using LD score regression implemented in LD Hub
software (http://ldsc.broadinstitute.org/).

A negative r_G indicates that an earlier or lower value of a continuous trait (i.e. earlier father's age of
death or lower subjective well being was associated with depression. A positive r_G indicates that a
later or higher value of a continuous trait (i.e. higher triglyceride level) was associated with
depression. Where multiple studies have examined a single trait the pubmed number of the study is
given in brackets.

1134

1135   Supplementary Figure 4

1136   Mendelian randomisation test for a putative causal effect of depression on neuroticism using inverse
1137   weighted regression, MR Egger and a weighted median test

1138

1139   Supplementary Figure 5

1140   Mendelian randomisation analysis for a putative causal effect of depression on ever vs. never smoked
1141   using inverse weighted regression

1142

1143   Supplementary Figure 6

1144   Mendelian randomisation test for a putative causal effect of neuroticism on depression using inverse
1145   weighted regression, MR Egger and a weighted median test

1146

Supplementary Figure 7

Contribution of functional annotation categories to the heritability of depression based on the variants
within each category

Error bars represent jackknife standard errors for each the estimate of enrichment, and an asterisk
indicates significance ($P < 0.0021$) after Bonferroni correction. The dashed line represents the
threshold for no enrichment.

1153

1154 Supplementary Figure 8

1155 Coefficients (β) of significantly enriched brain cell regions using GTEx overlaid on physical
1156 representation of brain anatomy

1157 The pseudo-coloring highlights the coefficients of the brain regions in red that were significantly
1158 enriched ($P < 0.05$) for depression variants

1159

1160 Supplementary Figure 9

1161 Stratified LD score regression analyses showing significance of enrichment estimates for 3 brain cell
1162 types in depression.

1163 The dashed line represents the Bonferroni threshold for significance ($P < 0.0167$) and * indicates
1164 significant enrichment for that brain cell type
1165

1166 Supplementary Figure 10

1167 Regional visualization plot centred on the independently-associated variant, rs1021363, close to the
1168 Sortilin related VPS10 domain containing receptor 3 (*SORCS3*) gene on chromosome 10.
1169 Recombination rates used in the plots are based on the European 1000 Genomes panel from
1170 November 2014

1171

1172 Supplementary Figure 11

1173 Regional visualization plots centred on independently-associated variants (A. rs2568958 and B.
1174 rs10890020) close to the Neuronal Growth Regulator 1 (*NEGR1*) gene on chromosome 1.
1175 Recombination rates used in the plots are based on the European 1000 Genomes panel from
1176 November 2014

1177

1178 Supplementary Figure 12

1179 Regional visualization plots centred on independently-associated variants (A. rs62091461, B.
1180 rs12966052, and C. rs12967143) close to the Transcription Factor 4 (*TCF4*) and *RAB27B* genes on
1181 chromosome 18. Recombination rates used in the plots are based on the European 1000 Genomes
1182 panel from November 2014

1183

1184 **Supplementary Table Titles and Legends**

1185   Supplementary Table 1

1186   Variants with a P-value $< 5 \times 10^{-8}$ for an association with depression in the meta-analysis of
1187   PGC_139k, 23andMe_307k and UK Biobank

1188

1189   Supplementary Table 2

1190   The direction of effect of previously reported significant variants for depression across the studies
1191   contributing to the meta-analysis

1192

1193   Supplementary Table 3

1194   Genetic correlations between depression and other behavioural and disease related traits using LD
1195   score regression implemented in LD Hub software (http://ldsc.broadinstitute.org/)

1196

1197   Supplementary Table 4

1198   Mendelian randomization analysis between depression and other traits using MR Egger test for
1199   directional horizontal pleiotropy, inverse variance weighted (IVW) test for variant heterogeneity and
1200   IVW regression, weighted median and MR Egger tests for a causal effect

1201

1202   Supplementary Table 5

1203   Heritability partitioned by functional annotation enrichment. The asterisk indicates significance after
1204   Bonferroni correction for multiple testing ($P < 0.0021$)

1205

1206   Supplementary Table 6

1207   Partitioning of the heritability estimate by cell type enrichment. The asterisk indicates significance
1208   after Bonferroni correction for multiple testing ($P < 0.0056$)

1209

1210   Supplementary Table 7

1211   Enrichment estimates for brain regions using GTEx. The asterisk indicates significance after
1212   Bonferroni correction for multiple testing ($P < 0.0038$)

1213

1214   Supplementary Table 8

1215   Enrichment estimates for brain cell types. The asterisk indicates significance after Bonferroni
1216   correction for multiple testing ($P < 0.0167$)

1217

1218   Supplementary Table 9

1219   Genome-wide significant gene-based hits ($P < 2.80 \times 10^{-6}$) in the meta-analysis of depression using

1220   MAGMA. NSNPS is the number of SNPs in the gene; NiSNPs is the number of independent SNPs in

1221   the gene

1222

Supplementary Table 10

Number and proportion of gene overlap within the Gene Ontology Consortium gene-sets associated

($P_{corrected} < 0.05$) with depression

Values on the lower diagonal are the number of overlapping genes between gene sets. Values on the

upper diagonal are the proportion of overlapping genes within the gene set containing the lower

number of genes

1229

1230   Supplementary Table 11

1231   Drug x gene interactions for the genes identified as significantly associated with depression with

1232   interactions obtained from the drug gene interaction database (http://dgidb.genome.wustl.edu/). The

1233   Anatomical Therapeutic Chemical (ATC) classification for each drug is provided along with the type

1234   of interaction and its source

Figure. 1

Figure. 2

ATC Code	Classification	ATC Code	Classification
A01	Stomatological preparations	J07	Vaccines
A02	Drugs for acid related disorders	L01	Antineoplastic agents
A03	Drugs for functional gastrointestinal disorders	L02	Endocrine therapy
A05	Bile and liver therapy	L03	Immunostimulants
A06	Drugs for constipation	L04	Immunosuppressants
A10	Drugs used in diabetes	M01	Antiinflammatory and antirheumatic products
A11	Vitamins	M03	Muscle relaxants
A12	Mineral Supplements	M04	Antigout preparations
A14	Anabolic agents for systemic use	M05	Drugs for treatment of bone diseases
B05	Blood substitutes and perfusion solutions	N01	Anesthetics
B06	Other hematological agents	N02	Analgesics
C01	Cardiac therapy	N03	Antiepileptics
C03	Diuretics	N04	Anti-Parkinson drugs
C04	Peripheral vasodilators	N05	Psycholeptics
C05	Vasoprotectives	N06	Psychoanaleptics
C07	Beta blocking agents	N07	Other nervous system drugs
C08	Calcium channel blockers	P01	Antiprotozoals
D07	Corticosteroids, dermatological preparations	P03	Ectoparasiticides, incl. scabicides, insecticides and repellents
D10	Anti-acne preparations	R01	Nasal preparations
D11	Other dermatological preparations	R03	Drugs for obstructive airway diseases
G02	Other gynecologicals	R06	Antihistamines for systemic use
G03	Sex hormones and modulators of the genital system	S01	Ophthalmologicals
G04	Urologicals	S02	Otologicals
H01	Pituitary and hypothalamic hormones and analogues	S03	Ophthalmological and otological preparations
H02	Corticosteroids for systemic use	V03	All other therapeutic products
J01	Antibacterials for systemic use	V04	Diagnostic agents
J05	Antivirals for systemic use	V09	Diagnostic radiopharmaceuticals

Figure. 3