Response outcomes gate the impact of expectations on perceptual decisions

Ainhoa Hermoso-Mendizabal[1] *, Alexandre Hyafil [1,2] *, Pavel E. Rueda-Orozco[3], Santiago Jaramillo[4], David Robbe[5] and Jaime de la Rocha[1]

[1] IDIBAPS, Barcelona, Spain
[2] Center for Brain and Cognition,Universitat Pompeu Fabra, Barcelona, Spain
[3] Universidad Nacional Autónoma de México, México
[4] University of Oregon, Oregon,USA
[5] Institut de Neurobiologie de la Méditerranée, Marseille, France

*These authors contributed equally to this work

ABSTRACT

Perceptual decisions are not only determined by current sensory information but are also influenced by recent experience. Can the magnitude of expectation-based decision biases be flexibly modulated based on the outcome of previous decisions? We trained rats in several two-alternative forced choice auditory tasks, where the probability to repeat the previous stimulus category was varied in blocks of trials. All rats capitalized on the serial correlations of the stimulus sequence by consistently exploiting a *transition bias:* a tendency to repeat or alternate their previous response using an internal trial-by-trial estimate of the sequence repeating probability. Surprisingly, this expectation bias was null in trials immediately following an error. The internal estimate however was not reset and it became effective again causing a bias after the next correct response. This ability to rapidly activate and inactivate the bias was captured by a non-linear generative model of rat behavior, whereby a reward-driven modulatory signal gated the use of the latent estimate of the environment statistics on the current decision. These results demonstrate that rats use behavioral outcomes to flexibly modulate how expectations derived from the recent history influence their decisions.

INTRODUCTION

Imagine Rafa Nadal returning Roger Federer's serve in the decisive game of a Grand Slam final. Serving at 190 km per hour, Nadal has few hundred milliseconds to visually estimate the ball trajectory, prepare the motor plan including where he aims to return the ball and execute it. In such speeded decisions based on partial or ambiguous sensory information, the anticipation provided by an informed prior expectation can be decisive because subjects can respond faster. Based on past games bringing the two players together, and on the pattern of the last serves executed by Federer, Nadal inevitably forms an expectation about where the next ball will arrive. Combined with the visual motion of the ball, this expectation may allow him to gain some decisive tens of milliseconds in the return of the serve(Vernon et al., 2018). However, if his prediction fails and he concedes an ace, does he need to choose between trashing his prior model on Federer's serve or sticking to it in the subsequent point? Or can Nadal transiently downplay the weight of his prediction on the next serve without modifying his prior?

Normative theories describe how prior expectations and ambiguous stimulus evidence should be combined in order to maximize categorization performance (Ernst and Banks, 2002; Stocker

and Simoncelli, 2006). In dynamical environments, in which the statistics of the sensory information varies with time, subjects must be constantly updating their internal model by accumulating past stimuli, actions and outcomes (Yu and Cohen, 2008). The updating of the prior based on the actions occurring in each trial typically introduces sequential effects, which are systematic history-dependent choice biases reflecting the impact of the trial-to-trial variations in expectation (Abrahamyan et al., 2016; Akaishi et al., 2014; Akrami et al., 2018; Ashourian and Loewenstein, 2011; Braun et al., 2018; Busse et al., 2011; Cho et al., 2002; Fischer and Whitney, 2014; Fründ et al., 2014; Hwang et al., 2017; Meyniel et al., 2016; Nogueira et al., 2017). There are circumstances however, where subjects seem able to quickly and flexibly modulate the impact of prior expectations in driving their choices. One of such examples is the switch between (1) exploiting choices which, according of their current statistical model of the environment, are more likely to yield reward and (2) exploring alternative choices that are not aimed to maximize reward given that internal model, but to reduce environmental uncertainty and eventually refine the current model (Ebitz et al., 2018; Karlsson et al., 2012). In particular, when the task design potentiates stochastic exploration, rats are able to operate in an expectation-free mode in which choices did not depend on previous history (Tervo et al., 2014). In other tasks, the updating of the internal prior is not done in a continuous manner as new information is presented, but subjects update their internal estimates abruptly and intermittently when they feel there has been a change-point in the environment (Gallistel et al., 2004). Recent studies have shown that, in the absence of feedback, the magnitude of the expectation bias on current choice is smaller after a low confidence response (Braun et al., 2018; Samaha et al., 2018; Urai et al., 2017). Despite these findings, we lack a conceptual framework that could explain both how expectations are formed and which are the factors that regulate their use in a moment to moment basis.

Here we investigate the extent to which the combination of expectation and sensory evidence can be dynamically modulated. Moreover, we aim to develop a unified model that jointly describes the dynamics of expectation build-up and its modulatory variables on a trial by trial basis. Behavioral experiments yielding tens of thousands of trials per animal allowed us to tease apart the different types of history biases. In particular, rats accumulated evidence over previous choice transitions in order to predict the next rewarded response. Crucially, this transition bias disappeared after an error, reflecting a fast switch into an expectation-free categorization mode. This switch did not imply however the reset of the accumulated expectation which resumed its influence on behavior as soon as the animal obtained a new reward. This ubiquitous behavior across animals was readily captured by a non-linear dynamical model in which previous outcome acted as a gate for the impact of past transitions on future choices.

RESULTS

A reaction time auditory discrimination task promoting serial biases

To study how the recent history of stimuli, responses and outcomes influence perceptual choices, we trained rats in a two-alternative forced choice (2AFC) auditory discrimination task in which serial correlations were introduced in stimulus trial sequences (Fig.1a-c) (Braun et al., 2018; Goldfarb et al., 2012; Jones et al., 2013; Kim et al., 2017). This design mimicked the temporal regularities of ecological environments and allowed us to probe the trial-by-trial expectations that animals formed about upcoming stimuli based on the changing statistics of the

stimulus sequence. The serial correlations between trials were created using a two-state Markov chain (Fig. 1b) parameterized by the probability to repeat the previous stimulus category P_{rep} (the unconditioned probabilities for each of the two categories were equal). We varied P_{rep} between Repeating blocks, in which $P_{rep}= 0.7$ and Alternating blocks in which $P_{rep}= 0.2$ (Fig. 1c). By poking into the center port, rats triggered the presentation of the stimuli, which lasted until they freely withdrew from the port. Each acoustic stimulus was a superposition of a high-frequency and a low-frequency amplitude-modulated tones and animals were rewarded for correctly discriminating the tone with the higher average amplitude. The discrimination difficulty of each stimulus, termed stimulus strength s, was randomly and independently determined in each trial, and set the relative amplitude of each tone (Fig.1b,d). When stimulus strength s was null, i.e. contained no net evidence in favor of either alternative, the rewarded side was determined by the stimulus category sequence (Fig. 1b).

Across-trial dynamics of history-dependent choice biases

Animals in Group 1 (n=10 animals) completed an average of 508 trials per session (range 284 - 772 average trials), gathering an average of 56,242 trials in total per animal (range 15,911 - 81,654 trials). For each rat, psychometric curves showing the proportion of Right responses as a function of the stimulus evidence indicated that neither sensitivity nor fixed side bias, the history-independent fixed preference to choose one side, varied with block type (Fig. 2a left). To estimate the impact of stimulus serial correlations, we also analyzed the repeating psychometric curves, showing the proportion of trials where the animals repeated the previous choice as a function of the sensory evidence in favor of the repeating choice. This analysis showed that all animals developed a block-dependent repeating bias b (Fig. 2a right, b left): after correct trials, b was positive in the Repetitive Block, and negative in the Alternating Block (i.e. tendency to repeat or alternate, respectively). Interestingly, the side bias, which was heterogeneous across animals, was similar across blocks for each animal (Fig. 2c left), showing that animals side preference was independent of the changes in repeating bias caused by block switching. Surprisingly, in trials following an error, b almost vanished in both block types (Fig. 2b-c right). Thus, after errors rats seemed unable to use previous history to guide their decision (e.g. in the Repetitive block, after an incorrect Right, the Left response is more likely to be rewarded). In contrast, the side bias was analogous after correct and error trials (Supplementary Fig. 1b).

Rats used history information by tracking several trials over short windows into the past: the magnitude of the repeating bias b built up with the number of consecutive correct past repetitions or alternations n until it plateaued after $n = 5$-10 trials (Fig. 2e blue and red line). This plateau was greater after repetitive patterns rather than alternating patterns. Importantly however, irrespective of n, the repeating bias b vanished almost completely with a single incorrect response for all rats (Fig. 2e). The vanishing occurred independently of the strength of the incorrectly categorized stimulus (not shown). To control that the vanishing was not caused by forgetting due to the 5s time-out imposed after errors, we trained a subset of rats using random time-out durations (range 1-5 s) and found that the bias extinction was maintained independently of time-out duration (Supplementary Fig.1c). The vanishing was only observed after errors and not after correct but unexpected responses, e.g. one alternation after several repetitions (Supplementary Fig. 2a). Accordingly, performance was higher for trials following a correct trial than for trials following an error, and it increased with n (Supplementary Fig. 3a-b). Moreover, the repeating bias increased performance when it was consistent with the block

tendency but it decreased performance when it was inconsistent with it (Supplementary Fig.3c-d). The impact on performance was largest for low stimulus strength, when the sensory evidence was weak and animals relied more strongly in their *a priori* belief, and it vanished to zero as the stimulus strength increased and the classification became easier. Together, these observations suggest that rats update their beliefs about the environment in a trial-by-trial basis and that this update crucially relies on the outcome of the preceding trials: longer sequences of rewarded repetitions/alternations lead to stronger response prior, but one error was sufficient to make the animals abandon this prior.

A GLM analysis of integration of sensory evidence and recent history information

Having identified that rats used previous responses and outcomes to guide decisions, we aimed to identify the specific factors in trial history generating this repeating choice bias. These factors could be classified as: (1) *sensory after-effects* yielding a changed sensitivity to the stimuli that were recently presented; (2) a *lateral bias* that creates an attraction or repulsion towards the Left or Right side depending on previous responses; (3) a *transition bias* that creates an attraction towards repeating or alternating depending on the history of previous repetitions and alterations (i.e. a rule bias). Although only the transition bias is adaptive in the task, since it allows to flexibly capitalize the sequence correlations in both types of blocks, the three factors could in principle contribute to the repeating bias *b* described above. We thus used a generalized linear model (GLM) to estimate the impact onto current decisions of each event in recent history including stimuli, sides of response, transitions and outcomes in the last ten trials (Fig. 3; see Methods for details). Given the effect of error responses suppressing the use of previous history (Fig. 2b-d right), we fitted for each rat the weights of the GLM separately for trials following correct and error responses. A consistent pattern for the regressor weights emerged (Fig. 4 and Supplementary Fig. 4a). First, as expected, the current stimulus, and in particular the first acoustic frames in the stimulus, had a strong impact on decisions (Fig. 4a). Second, choices were biased away from the sides associated with previously presented stimuli. This repulsive bias was consistent with an after-effect caused by sensory adaptation in which a strong acoustic power at a given frequency would reduce the likelihood to perceive that frequency in subsequent trials (Fig. 4b). Third, the impact of Left or Right responses on future choices depended on their outcome following a win-stay-lose-switch pattern: while rats displayed a tendency to opt again for the side of previously rewarded responses, sometimes called *reinforcers (Corrado et al., 2005; Lau and Glimcher, 2005)*, they tended to opt away from previously non-rewarded responses (Fig. 4c). We termed this first-order sequential effect the *lateral bias.*

Finally, the most influent history-dependent regressor in biasing choices were previous transitions *T*, meaning repetitions (*T*=+1) and alternations (*T*= -1) (Fig. 4d). Positive weighting of recent transitions implied that repetitions increased the transition bias γ_T (i.e. increased the tendency to repeat), while alternations had a negative impact on γ_T (i.e. increased the probability of alternating). We separated the effect of transitions between two consecutive correct responses (*T*++) and transitions between two responses where either the first (*T*-+), the second (*T*+-) or both responses (*T*--) were incorrect. While transitions between two correct responses *T*++ were strongly reinforced in subsequent trials, the transitions *T*-+, *T*+- and *T*-- with at least one error barely influenced subsequent trials (Fig. 4d). In other words, rats lacked the adaptive behavior to use non-rewarded trials for re-evaluating the current probability of repetition vs alternation. After incorrect trials, consistently with previous analyses, the impact of *T*++

transitions completely vanished (Figure 4d, black points). Thus, following an error, rats behavior was completely blind to the history of previous repetitions and alternations, and was driven only by sensory information and lateral biases. The reset of the transition bias was not an idiosyncratic strategy followed by some of our animals but it was found in every animal we trained (Supplementary Fig. 5). In fact the magnitude of transition kernel was much more homogenous across animals than the lateral kernel (Supplementary Fig. 5). The reset effect was however not observed neither in the after-effect bias nor in the lateral bias that were not affected or moderately affected by errors, respectively (Fig. 4b-c). Thus, the bias reset following errors was specific of the transition term but extremely reliable across subjects.

The impact of the current sensory stimulus was an order of magnitude larger than the transition bias, which was itself an order of magnitude larger than lateral bias (Fig. 4e). The after-effect bias was the weakest determinant of behaviour. A modified analysis separating the effects of repetitions and alternations showed that they had largely symmetrical effects suggesting that animals summarized both types of transition into a single rule that could take positive or negative values (Supplementary Fig. 4d). Importantly, the weights were identical when computed separately in repetition and alternation blocks (Supplementary Fig. 6). This suggests that rats adopted a single strategy across all blocks, and the different repeating choice bias found in each block (Fig. 2b-e) simply reflected the different statistics of the stimulus sequence used in that block (Fig. 1c). Because the impact of transitions decayed in around 5 trials (Fig. 4d left), the strategy allowed animals to switch the sign of their repeating bias relatively fast when switching between blocks (Supplementary Fig. 1a) at the cost of suffering relatively large fluctuations in the repeating bias within each block. Model comparisons further confirmed that the full model fitted separately for trials following correct trials and errors provided a better fit to rats decisions than the full model fitted to all trials, or alternative models where the lateral and/or transition module were removed (Supplementary Fig. 4c). Importantly, the GLM with only lateral biases yielded an non-monotonic kernel for the Lateral responses, a result that could lead to spurious interpretations when the effect of previous transitions was not considered (Supplementary Fig. 4d).

To test the extent to which these findings depended on the task design, we trained a new group of rats (Group 2, n= 6) in a different level discrimination 2AFC task in which noise stimuli had to be classified according to the intensity difference between the two lateral speakers (Pardo-Vazquez et al.; 2018). The stimulus sequence followed the same pattern as before with repeating and alternating blocks (Fig. 1b-c). Performing the same GLM analysis in this task yielded quantitatively similar results, including the reset of the transition bias after errors (Supplementary Fig. 7). Finally, we found that the presence of a transition bias and its reset after errors was not contingent on the presence of serial correlations in the stimulus sequence. A third group of rats (Group 3, n = 9) exposed to only an uncorrelated stimulus sequence, exhibited the same qualitative pattern for the impact of previous transitions, although with a smaller magnitude (Supplementary Fig. 8c). Once the sessions started containing serial correlations (Fig. 1b-c), the magnitude of the transition weights increased (Supplementary Fig. 8c) suggesting that the transition bias is an intrinsic behavior of the animals, but its magnitude can be adapted to the statistics of the environment. In total, these analyses show that the dependence on previous outcome of history-dependent biases is a general result across animals and across different tasks. It represents a novel non-linearity that cannot be captured by standard regression methods.

Transition evidence is blocked but not reset after an error

We then asked whether the system monitoring transitions (i) underwent a *complete reset* after error trials, meaning that the build-up of the transition bias γ_T started back from zero following errors (Fig. 5a), or whether, in contrast, (ii) information about previous transitions was maintained in memory at the bias reset and was later used to compute γ_T following new correct responses. The latter scenario, which we termed the *gating hypothesis* (Fig. 5b), was formulated by defining the accumulated transition evidence z_T, a variable that maintained an estimate of the transition probabilities and which, on trial by trial basis, was used or not (i.e. gated) to generate the transition bias γ_T. To test these two hypotheses, we quantified how the value of the bias γ_T in trial t could predict the bias at trial $t+1$, $t+2$ and further, depending on the outcome of each of these trials. When trial t was correct, the bias was passed on to $t+1$ with a discounting decay that mirrored the shape of the transition kernel in the GLM analysis (Figure 5c dark orange dots). The same discounting occured going from $t+1$ to $t+2$ when trial $t+1$ was correct. By contrast, if t was incorrect, because of the bias reset after errors, the value of γ_T was not predictive of the decision at trial $t+1$, nor at trial $t+2$ if $t+1$ was also incorrect (Fig. 5c black dots). Crucially though, the bias γ_T in trial t strongly influenced choices at trial $t+2$ if trial t was incorrect but trial $t+1$ was correct (Fig. 5c light orange dots). Its impact was significantly larger than zero for all rats ($p<10^{-6}$) and close in magnitude to the impact when both trials t and $t+1$ were correct. This rebound in choice predictability was even observed at $t+3$ after two incorrect responses followed by a correct one ($p<10^{-4}$). These results are consistent with the gating hypothesis (Fig. 5b) in which errors do not cause a reset of the accumulated transition evidence z_T but do cause a transient cut off in the influence of z_T on choice measured as a reset in γ_T. This influence became effective again once the animal made a new correct response giving rise to the measured correlation between the values of the bias before and after the reset (Fig. 5b gray vertical arrows). An equivalent analysis on the lateral bias γ_L showed that the bias transferred to the subsequent trials with a rapid decay, which was moderately affected by the outcome of the trials and showed no evidence of reset-and-rebound dynamics (Fig. 5d).
The rebound of the transition bias was observed more directly when we extended our GLM analysis by conditioning on the outcome of the last two trials, i.e. fitting separately for choices following a '++', '+-', '-+' or '-- ' sequence. When the last two trials were correct (++), the correct transitions T++ occuring in the last ten trials had a strong influence (Fig. 5e dark orange dots) whereas when the last trial was an error (+- or --) they had no influence (Fig. 5e gray and black dots). The influence of T++ transitions was again strong, however, when the choice followed a -+ sequence (Fig. 5e light orange dots) confirming the ability of the animals to rapidly recover the history-dependent transition bias based on the outcome of the last trial.

A dynamical model of history-dependent outcome-gated biases

Having found that the transition bias underwent reset-and-rebound dynamics, we built a generative model that implemented the gating hypothesis. One latent variable in the model was the accumulated transition evidence z_T, which was updated in each trial depending on whether the last choice was a repetition or an alternation and therefore maintained a running estimate of the transition statistics (Braun et al., 2018; Busse et al., 2011; Cho et al., 2002; Fründ et al., 2014) (Fig. 6a). A second modulatory variable c_T, modulated the influence of the transition evidence onto the current decision by setting the transition bias equal to $\gamma_T = c_T * z_T$. Importantly c_T was updated after each trial only based on the trial's outcome. In addition to the transition

bias, the model had also an accumulated lateral evidence z_L that directly resulted in a lateral bias (i.e. $\gamma_L = z_L$).

We fitted the model parameters to the series of choices made by each rat (Fig. 6b-g and Supplementary Fig. 9) and obtained results in agreement with our previous analyses: first, correct transitions (++) lead to a strong changes in the transition evidence z_T, while the other transitions (+-,-+,--) did not lead to any consistent pattern (Fig. 6d). Second, the update parameters for c_T corresponded to an almost complete reset after errors for at least 7 rats out of 10, and a very strong recovery after any correct trial (Fig. 6f). This effectively converted the variable c_T into a gating variable that was able to completely block the use of the transition accumulated evidence after a single error (Fig. 6g). By contrast, the leak of z_T was not significantly different after correct trials and after errors (p>0.6), providing further evidence that the reset transition bias after errors did not correspond to a loss of the accumulated evidence. Third, correct Right (Left) responses increased the lateral bias in favor of the Right (Left) response (Fig. 6c). Fourth, adding a modulatory variable c_L to the lateral module only had a marginal impact on model performance (Supplementary Fig. 10). Finally, model comparison concluded that this canonical dynamical model gave a better account than versions where either c_T or the lateral bias z_L were omitted, as well as of the GLM described in the previous section (Supplementary Fig. 11).

Generative model simulation versus experimental data

Finally we assessed the capacity of the compact dynamical model to account for the dynamics of the previously reported repeating bias *b* (Fig. 2d-e) by comparing model simulations to actual rat data. The model very closely reproduced the build-up dynamics of *b* in series of correct repetitions and alternations (Fig. 7a). Moreover, the model allowed to partition the value of *b* into the lateral and transition biases showing that, while the transition bias was perfectly symmetric in series of repetitions and alternations, the lateral bias behaved very differently: it only built up during series of repetitions, in which all the responses were on the same side, while it oscillated around zero in series of alternations, in which the contribution of each response was partially cancelled by the next one. Thus the dissection of the repeating bias into the lateral and transition biases explained the overall asymmetry found between the two blocks. Furthermore, the block asymmetry in the accumulation of the lateral bias also explained asymmetries in *b* found after correct unexpected responses (Supplementary Fig. 2). Model simulations also reproduced the reset of repeating bias when a series of correct repetitions/alternations was interrupted by an error (Fig. 7b), and the subsequent rebound when the rat performed correctly again (Fig. 7c). Impressively, the model replicated the asymmetry in the magnitude of this rebound between the Repeating and Alternating blocks by summing (Fig. 7c top) or subtracting (Fig. 7c bottom), respectively, contributions of transition and lateral biases. Furthermore, the model provided a very good fit (Pearson's r=0.96) to *b* for all possible sequences of 2-6 correct trials (Fig. 7d). In sum, by factorising the transition bias into accumulated transition evidence z_T and the modulatory signal c_T, the model captured the non-linear across-trial dynamics of history-dependent biases pointing towards possible modulatory circuit mechanisms that could implement this computation (see Discussion).

DISCUSSION

We employed a standard acoustic 2AFC task to characterize how rats' perceptual categorizations are affected by expectations derived from the history of past stimuli, choices and outcomes and how these expectations can be captured by a simple dynamical model. A thorough analysis of the behavior isolated three different sequential effects. First, because in our task the stimuli in each category varied strongly in magnitude, we could disentangle the impact of past physical stimuli from the stimulus category revealing a repulsive after-effect that lasted for at least ten trials (Fig. 3b). Although this after-effect resonates with the predominant response adaptation found in several stages of the auditory pathway, its time scale, spanning tens of seconds (Fig. 4b), exceeds the previously reported recovery period from adaptation (Asari and Zador, 2009; Natan et al., 2015). Second, we identified a sequential lateral effect that biased choices towards or away from the recently rewarded or unrewarded targets, respectively (Fig. 3c). This win-stay-lose-switch strategy has been extensively characterized both in humans (Abrahamyan et al., 2016; Braun et al., 2018; Fründ et al., 2014) and rodents (Akrami et al., 2018; Hwang et al., 2017). Finally, we identified the sequential transition bias, a form of rule bias that had been previously shown to impact human reaction times (Cho et al., 2002; Kirby, 1976; Soetens et al., 1985), choices (Maloney et al., 2005) and neural responses (Jones et al., 2013; Sommer et al., 1999). Our results however go beyond from previous reports in several important aspects regarding error responses: first, repetitions or alternations did not influence subsequent choices whenever one of the two trials of the transition was unrewarded, meaning that the running estimate of the transition probabilities only accumulated evidence from repetitions and alternations of two rewarded trials. Second, the transition bias was absent after an error trial, i.e. animal responses temporarily ignored the recent history of rewarded repetitions and alternations. However, this reset did not imply the reset of the accumulated transition evidence, i.e. the tallie keeping track of the number of recent repetition vs. alternations, whose influence over behavior was restored as soon as the animal obtained a reward (Fig. 5c,e).

The probability of a subject to repeat the previous response (Fig. 7a) is a common measure to characterize history effects (Abrahamyan et al., 2016; Urai et al., 2018). By dissecting the distinct contribution of both first- and second-order serial biases (Gokaydin et al., 2011; Jones et al., 2013; Meyniel et al., 2016; Wilder et al., 2009), i.e. the lateral and the transition biases, respectively, to the repeating bias we were able to understand , the asymmetry in its magnitude between the Repeating and Alternating blocks (Fig. 2c-e): in series of correct repetitions, both transition and lateral bias add up and yield a strong tendency of the animals to repeat the last rewarded response (Fig. 7a top). In contrast, in alternating environments, the lateral bias does not build up and the negative repeating bias (tendency to switch) is solely given by the transition bias (Fig. 7a bottom). In sum, the first-order lateral bias favors repetition over alternation; the second-order transition bias has a symmetric effect. In fact, our analysis provides indirect evidence that animals recapitulated previous repetitions and alternations into a single and symmetric transition bias and not into separate variables (Supplementary Fig. 4). A recent modeling study has proposed that estimating first and second-order rates is part of the same core computation that the brain performs when analyzing binary sequences. This computation comes down to estimate the two independent transition probabilities $P(L_t|R_{t-1})$ and $P(R_t|L_{t-1})$ between consecutive trials t -1 and t (Meyniel et al., 2016). Our findings seem at odds with this hypothesis because the dependence of each type of bias on the response outcome was very different: whereas incorrect responses L^- tended to cause a negative switch effect (Fig. 4c),

incorrect transitions (T^{+-}, T^{-+}, and T^{--}) had no effect (Fig. 4d). Furthermore, only the transition bias showed a reset-and-rebound dynamics caused by error responses (Fig. 5c-d). An alternative hypothesis, based on analysis of response evoked potentials (ERP), proposes that the lateral bias is generated by the *processing of the response* whereas the transition bias from the *stimulus processing* (Jones et al., 2013; Wilder et al., 2009). Preliminary data obtained in the same task in the absence of any stimuli (speakers were turned off) seems to indicate that the transition bias is still present and thus does not seem to be contingent on any sensory processing.

Several of our findings, together with previous literature (Cho et al., 2002; Jones et al., 2013; Kirby, 1976; Soetens et al., 1985; Sommer et al., 1999), suggest that the transition bias is a fundamental aspect of sequence processing preserved across subjects, species and conditions and which does not seem particularly adaptive to the details of the experiment. First, the transition bias was the same in both Repeating and Alternating blocks (Supplementary Fig. 6) reflecting the use of a single fixed strategy that could switch from generating a net positive repeating bias in a Repeating block to generating a negative bias in the alternating block (Fig. 2e). Interestingly, this invariance of the transition bias across the repetitive and alternating blocks has also been found in humans performing a 2AFC task (Jones et al., 2013). Second, the transition bias was also present when sequences are uncorrelated and the bias can only hinder performance (Supplementary Fig. 8a; (Cho et al., 2002; Jones et al., 2013; Kirby, 1976; Soetens et al., 1985; Sommer et al., 1999). Third, the trial integration window over which animals estimated the repetition rate (~3-5 trials; Fig. 4d) does not seem adapted to the block length (200 trials). This short-span estimate allowed to reverse the repetition bias rapidly after a block switch (Supplementary Fig. 1a) at the cost of a noisier estimate of the repetition rate (Gallistel et al., 2004; Nassar et al., 2010; Sutton and Barto, 2018). Quantification of this integration window in human subjects performing different 2AFC tasks yields numbers in the range of 2-10 trials, despite the use of very long trial-blocks with constant sequence correlations (Jones et al 2013). Thus, rather than an overestimation of the environment's volatility (Behrens et al., 2007; Nassar et al., 2010), the short fixed windows might reflect structural adaptation to the statistics of natural environments (Seriès and Seitz, 2013) or a capacity limitation of the system. Fourth, the sophisticated outcome-dependent across-trial dynamics of the transition bias were found systematically in every animal we tested (Supplementary Fig.5) showing that they do not reflect idiosyncratic strategies but the action of an unknown basic cognitive process. Finally, there was one aspect of the mechanism that seemed adaptive: the magnitude of the transition kernel gradually increased when animals trained using uncorrelated sequences were presented with correlated sequences (Supplementary Fig. 8). Rats in fact were previously shown to suppress sequential biases when those can be turned up against them (Tervo et al., 2014). Thus, the transition bias can be adapted to the temporal structure of the environment, if not in nature, at least in magnitude (Abrahamyan et al., 2016) (Supplementary Fig. 8).

Why does the transition bias reset after errors? Previous studies have shown that an uncued change in stimulus-outcome contingencies leading to an unexpected number of unrewarded choices can trigger an abrupt behavioral change in rats, switching from the exploitation of a statistical model of the environment to an exploration mode in which they sample the environment in an unbiased way in order to build new beliefs (Karlsson et al., 2012). This suggests that the reset-and-rebound dynamics of the transition bias could be interpreted as a fast switching between the exploitation of their internal model, represented by their estimate of the transition probability, to a mode that relies almost exclusively on sensory information. This

expectation-free mode, however, is different from the standard exploration mode in which animals guide their choices aiming to reduce the uncertainty of the environment. In contrast, our animals, perhaps unable to use their prior after not obtaining the reward (i.e. not knowing *what* they must repeat after an error), guide their choices based on the sensory evidence alone. To capture the reset-and-rebound dynamics, we built a generative-sufficient novel model that could jointly describe the latent trial-to-trial dynamics of (1) the expectation formation following standard reinforcement learning updating rules (Behrens et al., 2007) (Fig. 6a and b) and (2) a modulatory signal c_T that had a multiplicative effect on the impact of the transition evidence in biasing choices. The fitting of the model parameters revealed that c_T reset to zero after errors and then increased progressively with a series of correct trials (Fig. 6c). This modulatory variable may reflect subjects' confidence in their internal model of the environment statistics or, alternatively, the probability that the subject operated in the exploitation mode versus the expectation-free mode. Furthermore, in this state in which the prior is not used, it seems that it also cannot be updated, as can be concluded from the finding that only ++ transitions impacted subsequent choices (Fig. 4d).

Recent studies found that, in the absence of feedback, the impact of a choice on the subsequent trial was weaker if subject was unsure of her choice (Braun et al., 2018; Samaha et al., 2018; Urai et al., 2017). The explanation provided in two of these studies was that, according to a normative theory describing how to accumulate noisy evidence in the face of uncued changes (Glaze et al 2015), low confidence choices should have a weaker contribution on the belief about what will be the next stimulus category (Braun et al., 2018). In our latent variable model, this is is indeed true because unrewarded transitions, *T+-* and *T--*, supposedly generating the lowest confidence about what the true transition was, have a weaker contribution to the accumulated evidence z_T (see fitted values of Δ_T in Fig. 6c). However, a bias reset after incorrect or low confidence trials was not reported in that study, i.e. errors without feedback did not seem to modulate retroactively the impact of previous trials onto the next choice, unlike what was observed in our rats. Also, in (Braun et al., 2018) subjects were cued about the value of transition probabilities in each block, whereas our animals were constantly estimating the transition probability. A fair assessment on the existence of the expectation bias reset in humans would necessitate of an experiment in which subjects are blind about the sequence correlations but receive feedback in every trial. The activation of noradrenergic inputs onto the anterior cingulate cortex have been shown to control the switching into a behavioral mode in which beliefs based on previous experience do not guide choices (Tervo et al., 2014). Because in quoted study the experimental condition was a free choice task, removing the impact of history-effects resulted in stochastic exploration (Tervo et al., 2014). This prompts the question of whether the activation of this very same modulatory pathway underlies the after-error switch into the expectation-free sensory-based mode observed in our task. Future pharmacological an electrophysiological experiments could shed light into the brain regions encoding the expectation signals, the modulatory variables as well the circuit mechanism underlying their combination and their integration with the afferent sensory information.

FIGURES:

Figure 1. Auditory discrimination task and stimulus sequence statistics:
(a) Sketch of one trial of the task: cued by center port LED, rats poke in the center port to trigger the presentation of a mixture of two AM tones, each of which is associated with reward in the Left (L) or Right (R) port. Correct responses are rewarded with water and incorrect responses are punished with a light plus a 5 s time-out. **(b-c)** Serial correlations in the sequence of stimuli were introduced by setting the probability of repeating the previous stimulus category P_{rep} (top in b) in trial blocks named Repetitive Block and Alternating Block (c). The stimulus strength s_k was randomly drawn in each trial (bottom in b) to yield the stimulus evidence e_k, that determined the distance to the categorization boundary, i.e. the discrimination difficulty of the stimulus (right in b). **(d)** The stimulus evidence e determined the distribution (top) from which the instantaneous evidence was drawn in each cycle of the sound envelope (see color match). An instantaneous evidence trace (middle) and the AM modulated tones that result (bottom) are shown for an example stimulus with e = -0.48 (asterisks in b and d).

Figure 2. Build up and reset dynamics of the Repeating Bias:

(a) Psychometric curves for an example animal showing the proportion of Right responses vs. stimulus evidence (left) or of Repeated responses vs. Repeating stimulus evidence (right) computed in the Repetitive (blue dots) or Alternating blocks (red dots; color code applies for all panels)**.** This animal shows a block independent fixed side bias B>0 towards the Right (left), and a block-dependent repeating bias b matching the tendency of each block. Curves show fits using a probit function. **(b)** Proportion of Repeated responses (median across n=10 animals) computed in trials following a correct (left) or an incorrect response (right). **(c)** Repeating bias b versus fixed side bias B in the two blocks after a correct (left) or an incorrect response (right). Each pair of connected dots represents one animal. **(d)** *Left:* Fits of the proportion of Repeated responses following trial sequences made of a different number of correct repetitions (blue gradient) or alternations (red gradient; see insets for color code). X^+ and Y^+ represent either Right or Left correct responses. E represents an error. Time in the sequences progresses from left to right. *Right:* same curves obtained when the sequence of correct repetitions is terminated by an error. **(e)** Repeating bias versus the sequence length of correct repetitions (left, blue) or alternations (right, red). Sequences terminated by an error are shown in black. Dark traces show median across animals while light traces show individual animals. Error bars show Std. Dev. (a) or first and third quartiles (b, e).

Figure 3. GLM assessing the impact of sensory evidence and recent history onto rats choices. **(a)**, Exemplar series of recent history trials used to model rat decisions at current trial *t*. The Stimulus series depicts the amplitude of each acoustic frame for the low (green bars) and high frequency tones (purple bars). The Response series shows the rat decisions (green for Left and purple for Right). The Transition series shows the relation of two consecutive responses (blue for Repeat, red for Alternation) and the Outcome series indicates whether the responses were rewarded (orange) o not (black). These series are combined to generate the regressors that are grouped into the Sensory (b), Lateral (c) and Transition (d) modules. **(b)**, The weighted sum of the amplitude difference in each acoustic frame of the current trial provides the Stimulus evidence γ^S_t. The weighted sum of the net stimulus evidence in each of the previous trials (*t-1*, *t-2*, etc) provides the After-effect bias γ^A_t. **(c)**, The Lateral module weights separately previous rewarded *L+* and unrewarded responses *L-* that take the values -1 (Left), +1 (Right) or 0. The sum of these two series gives rise to the lateral bias γ^L_t. **(d)**, Transitions are considered separately depending on the outcome of the two trials in the transition: *T++* (rewarded-rewarded), *T-+* (error-rewarded), *T+-* (rewarded-error) and *T—*(error-error) which take the values -1 (alternation), +1 (repetition) and 0. The weighted sum of transition regressors yields the Transition bias γ^T which is multiplied by the previous response r_{t-1} in order to be combined with variables providing Right vs Left evidence. **(e)**, The sum $\gamma^S_t + \gamma^A_t + \gamma^L_t + r_{t-1}\,\gamma^T_t$ is then passed through a sigmoid function to yield the trial-by-trial probability of selecting a Right response. For each rat, the weights were fitted to maximize the model's probability to generate the actual animal choices.

Figure 4. Fitted weights quantifying the impact onto animals decisions of the different modules. Average coefficients (n=10 animals) obtained in the GLM when separately fitting the choices in trials after a correct (orange) and error response (black). **(a)** Frame-by-frame influence of the current stimulus onto rats decisions. **(b)** Influence of the sensory after-effect caused by the net sensory evidence of previous trials as a function of trial lag (i.e. number of trials back from current trial). **(c)** Influence of the response side (Left vs Right) from previously rewarded (*L+*, left panel) and unrewarded (*L-*, right panel) trials. **(d).** Influence of previous transitions (repetition vs. alternation) computed separately for *T++* (a rewarded trial followed by a rewarded trial), *T-+* (error-rewarded), *T+-* (rewarded-error) and *T--* (error-error). **(e)** Normalized variance averaged over animals (see Methods) quantifies the overall relative impact of each module (current stimulus, sensory after-effect, lateral bias and transition bias) on the rats choices. (a-e) Error bars indicate first and third quartile.

Figure 5. Transition bias is reset after errors but accumulated transition evidence is maintained.
(a-b) Schematics showing three example traces of the transition accumulated evidence z^T_t (top) and transition bias on current response γ^T_t (bottom) in two hypothetical scenarios. **(a)** Complete reset hypothesis: after an error at *t*, both variables reset $z^T_{t+1} \simeq \gamma^T_{t+1} \simeq 0$. Evidence z^T_{t+2} is then built-up *de novo*, implying that biases before (γ^T_t) and after (γ^T_{t+2}) the reset are uncorrelated (i.e. γ^T_t traces do not maintain the sorting). **(b)** Gating hypothesis: after an error, evidence z^T_{t+1} is maintained but it does not convert into a bias, leading to the reset $\gamma^T_{t+1} \simeq 0$. After a correct response at *t*+1 the conversion is recovered and the value γ^T_{t+2} correlates with γ^T_t (i.e. γ^T_t traces maintain the sorting). **(c)** Transfer coefficient $\gamma^T_t \rightarrow \gamma^T_{t+k}$ versus trial lag *k*, quantifying the degree to which the transition bias at trial *t* affects the bias on subsequent trials, is calculated separately depending on the outcome of each trial (black lines show rewarded choices and colored lines error choices). While the transfer coefficient vanishes after errors (i.e. reset of the bias; black dots), a correct response following an error (orange) brings it close to the value obtained when there are no errors (red dots). This implies that the information about the value of the bias γ^T_t must be maintained when the bias is reset (i.e. gating hypothesis). **(d)** Same as in **c** but computed for lateral bias γ^L_t. **(e)** Average coefficients for transitions *T*++ (n=10 animals) obtained when the GLM is fitted separately

depending on the outcome of last two trials (see inset; e.g. -+ represents that *t-2* was incorrect and *t-1* was correct). Error bars indicate the first and third quartiles.

Figure 6. Dynamic generative model of history-dependent perceptual decisions. (**a**) Architecture of the model. Decisions integrate the evidence from the sensory, lateral and transition modules. The Sensory module accumulates the instantaneous evidence of each stimulus frame of the current trial. The lateral module maintains a representation of the lateral bias z_L which is updated depending on the side and outcome of each trial response (weights Δ_L), and has an outcome-dependent leak λ_L. The transition module maintains the accumulated transition evidence z_T, is updated depending on the last transition (e.g. repetition ++, alternation +-, etc; weights Δ_T) and has an outcome-dependent leak λ_T. . Evidence z_T is multiplied at each trial by the modulatory signal c_T, updated by the outcome of each response (weights Δ_C), to yield the transition bias $\gamma_T = z_T \, c_T$. The sum of sensory evidence and the lateral and transition biases determined the probability to choose either response at the current trial. Parameters were fitted to the choices of each rat separately . (**b-e**). Best-fitting values of the update parameters in the generative model (median across rats). **b)** update of the lateral bias Δ_L depending on trial outcome **c)** outcome-dependent leak of the lateral bias λ_L. **d)** update of the transition bias Δ_T depending on outcome of the last two trials (++, -+, +-,--). **e)** outcome-dependent leak of the transition bias λ_T. **f)** outcome-dependent update of the transition gating signal Γ_C. A value of -1 correspond to an extinction of

the confidence signal on the subsequent trial (i.e. a full blockade of the corresponding bias), while +1 correspond to full recovery of the bias (i.e. confidence equal to its maximum value of 1). **(g)** Example of the model dynamics across 25 trials switching from a repetition to an alternation block (model parameters were fitted to an example animal). Traces from top to bottom depict the stimulus evidence,lateral bias z_L, transition bias z_T, transition modulatory signal c_T and overall probability to choose a Right response. The inputs to the variables z_L, z_T and c_T are shown as a sequence of symbols on the corresponding trial axis: Left (green) vs. Right (purple) responses, repeating (blue) vs. alternating (red) transitions and rewarded (orange) vs. error (black) outcomes. Symbols shape represent different outcome combinations (see inset). Notice the reset of c_T after errors and the maintenance of z_T afterwards (asterisks).

Figure 7. Generative model simulation compared to experimental data:
Comparison between experimental data (dots) and model simulation (black curves) showing the Repeating bias b for different trial sequences. In the model, b was decomposed into the transition bias (blue curves) and lateral bias (green curves). **(a)** Similar to Fig 2e. Repeating bias b versus number n of correct repetitions (top) or alternations (bottom). **(b)** Similar to Fig 2e. Repeating bias after a repetitive (top) or alternating sequence (bottom) terminated by an error. **(c)** Repeating bias for sequences with an error E flanked by correct repetitions (top) or alternations (bottom). **(d)** Repeating bias for all sequences made of $n \leq 8$ repetitions (R) and alternations (A). Top panel shows correct sequences while bottom panel shows correct sequences terminated by an error. In all panels, data and model show median across n=10 rats. Error bars and shaded areas show first and third quartiles.

ONLINE METHODS

All experimental procedures were approved by the local ethics committee (Comité d'Experimentació Animal, Universitat de Barcelona, Spain, Ref 390/14).

Animal Subjects

Animals were male Long-Evans rats (n=25, 350-650g; Charles River), pair-housed during behavioral training and kept on stable conditions of temperature (23ºC) and humidity (60%) with a constant light-dark cycle (12h:12h, experiments were conducted during the light phase). Rats had free access to food, but water was restricted to behavioral sessions. Free water during a limited period was provided on days with no experimental sessions.

Task description

The two tasks performed were auditory reaction-time two-alternative forced choice procedures: an LED on the center port indicated that the rat could start the trial by poking in (Fig. 1a). After a fixation period of 300 ms, an acoustic stimulus consisting in a superposition of two amplitude-modulated sounds (see details below) was presented. The rats had to discriminate the dominant sound and seek reward in the associated port. Animals could respond any time after stimulus onset. Withdrawal from the center port during the stimulus immediately stopped the stimulus. Correct responses were rewarded with a 24 µl drop of water and incorrect responses were punished with a bright light and a 5 s time-out. Trials in which the rat did not make a side poke response within 4 seconds after leaving the center port were considered invalid trials and were excluded from the analysis (on average, only 0.4% of the trials were invalid). Behavioral setup (Island Motion, NY) was controlled by a custom software developed in Matlab (Mathworks, Natick, MA), based on the open-source BControl framework (http://brodylab.princeton.edu/bcontrol). Rats performed an average of 694 trials per session (range: 335 - 1188), one session per day lasting 60-90 min, 6 days per week, during 9 months. Rats were trained using an automated training protocol that had several stages and lasted between 2 and 3 months (depending of the animal). The data presented in this study was taken from the period after training yielding an average of 56,506 valid trials per rat. A first group of ten rats were trained in the frequency discrimination version of the task (see below) in which the correlated sequence of trials was present from the training. A subset of three rats from this group were also trained in a random time-out version of the task where the duration of the after-error time-out was randomly chosen between 1, 3 or 5 s. A second group of n=9 rats were trained in the same frequency discrimination version of the task but starting with uncorrelated stimulus sequences and only after several weeks, introducing the correlated sequences used in the first group of animals. A third group of n=6 rats were trained in a level discrimination version of the task using the same correlated sequence than the first group.

Acoustic stimulus

In the two acoustic tasks used, the stimulus $S(t)$ in each trial was created by simultaneously playing two amplitude modulated (AM) sounds $s_R(t)$ and $s_L(t)$:

$$S(t) = [1 + sin(f_{AM}t + \varphi)][a_L(t)\, s_L(t)\ + a_R(t)\, s_R(t)\,] \tag{1}$$

The AM frequency was f_{AM}=20 Hz and the phase delay $\varphi = 3\pi/2$ made the envelope zero at t=0. In the frequency discrimination task, s_L(t) and s_R(t) were pure tones with frequencies 6.5 kHz and 31 kHz, respectively, played simultaneously in the two speakers. In the level discrimination task (Supplementary Fig. 7) they were broadband noise played either from the Left or the Right speaker, respectively. The amplitudes of the sounds s_L(t) and s_R(t) were separately calibrated at 70 dB. Sounds were delivered through generic electromagnetic dynamic speakers (STAX, SRS-2170) located on each side of the chamber, and calibrated using a free-field microphone (Med Associates Inc, ANL-940-1).

Stimulus Sequence

The Markov chain generated the sequence of stimulus category c_k= {-1,1}, that determined whether the reward in the k-th trial was available in the Left or the Right port respectively (Fig. 1b top). The stimulus category set which of the two sounds s_L(t) and s_R(t) composing each stimulus was dominant, which ultimately determined the statistics of the sound amplitudes a_1(t) and a_2(t) (Eq. 1) as described below. In each trial, independently of c_k, the stimulus strength s_k was also randomly generated (Fig. 1b bottom). Stimulus strength defined the relative weights of the dominant and non-dominant sounds: for example, when s_k =1 only the dominant sound was played (i.e. easiest trials) whereas when s_k =0 the two sounds had on average the same amplitude (i.e. hardest trials). We used four possible values for s = 0, 0.23, 0.48 and 1. The stimulus evidence was defined in each trial as the combination e_k = c_k*s_k . The value of e_k determined the p.d.f. from which the instantaneous evidence x(t) was drawn in each frame t (i.e. in each 50 ms AM-envelope cycle; Fig. 1d top): when e_k:=±1 the p.d.f. was $f(x) = \delta(x\mp 1)$ (i.e. a Dirac delta p.d.f.) whereas when $e_k \in$ (-1,1), it was a stretched Beta distribution with support [-1,1], mean equal to e_k and variance equal to 0.06 (Fig. 1d top). Finally, the amplitudes a_L(t) and a_R(t) of the two AM envelopes (Eq. 1) were obtained using a_L(t)=[1+x(t)]/2 and a_R(t)=[1-x(t)])/2 (see example in Fig. 1d). With this choice the sum of the two envelopes was constant in all frames a_L(t)+a_R(t)=1.

Psychometric curve analysis

We computed two types of psychometric curves for each animal, by pooling together trials across all sessions for each type of block and for each of the 7 different stimulus evidences (e= 0, ±0.23, ±0.48, ±1). We calculated (1) the proportion of Right responses vs. stimulus evidence e (Fig. 1a left) and (2) the Proportion of Repeated responses as function of Repeating Stimulus Evidence (Fig. 1b), where positive or negative Repeating Stimulus Evidence denote trials in which the animals had evidence to repeat their previous choice (e.g. a Right stimulus with evidence e = +0.23 after a Left response implied a repeating stimulus evidence equal to - 0.23). Both psychometric curves were separately fitted to a 2-parameter probit function (using Matlab function *nlinfit*):

$$f(x) = \tfrac{1}{2}\left(1 + erf(\tfrac{\beta x + \alpha}{\sqrt{2}})\right) \tag{2}$$

The sensitivity β quantified the stimulus discrimination ability, while the bias α in the Proportion of Right responses captures the animal Fixed side preference for the Left or Right port, and the bias α in the Proportion of Repeated responses arose from the expectation generated by the history of recent choices, i.e. it showed the animal's tendency to repeat or alternate their

previous choice. We termed these two biases the Fixed side bias *B* and the Repeating bias *b*, respectively. Within-subject error bars were estimated by one standard deviation of a non-parametric bootstrap (n=1000). Across-subject error bars, corresponded to the first and third quartiles.

REFERENCES

Abrahamyan, A., Silva, L.L., Dakin, S.C., Carandini, M., and Gardner, J.L. (2016). Adaptable history biases in human perceptual decisions. Proc. Natl. Acad. Sci. U. S. A. *113*, E3548–E3557.

Akaishi, R., Umeda, K., Nagase, A., and Sakai, K. (2014). Autonomous mechanism of internal choice estimate underlies decision inertia. Neuron *81*, 195–206.

Akrami, A., Kopec, C.D., Diamond, M.E., and Brody, C.D. (2018). Posterior parietal cortex represents sensory history and mediates its effects on behaviour. Nature *554*, 368–372.

Asari, H., and Zador, A.M. (2009). Long-lasting context dependence constrains neural encoding models in rodent auditory cortex. J. Neurophysiol. *102*, 2638–2656.

Ashourian, P., and Loewenstein, Y. (2011). Bayesian inference underlies the contraction bias in delayed comparison tasks. PLoS One *6*, e19551.

Behrens, T.E.J., Woolrich, M.W., Walton, M.E., and Rushworth, M.F.S. (2007). Learning the value of information in an uncertain world. Nat. Neurosci. *10*, 1214–1221.

Braun, A., Urai, A.E., and Donner, T.H. (2018). Adaptive History Biases Result from Confidence-weighted Accumulation of Past Choices. J. Neurosci.

Busse, L., Ayaz, A., Dhruv, N.T., Katzner, S., Saleem, A.B., Schölvinck, M.L., Zaharia, A.D., and Carandini, M. (2011). The detection of visual contrast in the behaving mouse. J. Neurosci. *31*, 11351–11361.

Cho, R.Y., Nystrom, L.E., Brown, E.T., Jones, A.D., Braver, T.S., Holmes, P.J., and Cohen, J.D. (2002). Mechanisms underlying dependencies of performance on stimulus history in a two-alternative forced-choice task. Cogn. Affect. Behav. Neurosci. *2*, 283–299.

Corrado, G.S., Sugrue, L.P., Seung, H.S., and Newsome, W.T. (2005). Linear-Nonlinear-Poisson models of primate choice dynamics. J. Exp. Anal. Behav. *84*, 581–617.

Ebitz, R.B., Albarran, E., and Moore, T. (2018). Exploration Disrupts Choice-Predictive Signals and Alters Dynamics in Prefrontal Cortex. Neuron *97*, 475.

Ernst, M.O., and Banks, M.S. (2002). Humans integrate visual and haptic information in a statistically optimal fashion. Nature *415*, 429–433.

Fischer, J., and Whitney, D. (2014). Serial dependence in visual perception. Nat. Neurosci. *17*, 738–743.

Fründ, I., Wichmann, F.A., and Macke, J.H. (2014). Quantifying the effect of intertrial dependence on perceptual decisions. J. Vis. *14*.

Gallistel, C.R., Fairhurst, S., and Balsam, P. (2004). The learning curve: implications of a quantitative analysis. Proc. Natl. Acad. Sci. U. S. A. *101*, 13124–13131.

Gokaydin, D., Ma-Wyatt, A., Navarro, D., and Perfors, A. (2011). Humans use different statistics for sequence analysis depending on the task. (Cognitive Science Society),.

Goldfarb, S., Wong-Lin, K., Schwemmer, M., Leonard, N.E., and Holmes, P. (2012). Can post-error dynamics explain sequential reaction time patterns? Front. Psychol. *3*, 213.

Hwang, E.J., Dahlen, J.E., Mukundan, M., and Komiyama, T. (2017). History-based action selection bias in posterior parietal cortex. Nat. Commun. *8*, 1242.

Jones, M., Curran, T., Mozer, M.C., and Wilder, M.H. (2013). Sequential effects in response time reveal learning mechanisms and event representations. Psychol. Rev. *120*, 628–666.

Karlsson, M.P., Tervo, D.G.R., and Karpova, A.Y. (2012). Network resets in medial prefrontal cortex mark the onset of behavioral uncertainty. Science *338*, 135–139.

Kim, T.D., Kabir, M., and Gold, J.I. (2017). Coupled Decision Processes Update and Maintain Saccadic Priors in a Dynamic Environment. J. Neurosci. *37*, 3632–3645.

Kirby, N.H. (1976). Sequential effects in two-choice reaction time: automatic facilitation or subjective expectancy? J. Exp. Psychol. Hum. Percept. Perform. *2*, 567–577.

Lau, B., and Glimcher, P.W. (2005). Dynamic response-by-response models of matching behavior in rhesus monkeys. J. Exp. Anal. Behav. *84*, 555–579.

Maloney, L.T., Dal Martello, M.F., Sahm, C., and Spillmann, L. (2005). Past trials influence perception of ambiguous motion quartets through pattern completion. Proc. Natl. Acad. Sci. U. S. A. *102*, 3164–3169.

Meyniel, F., Maheu, M., and Dehaene, S. (2016). Human Inferences about Sequences: A Minimal Transition Probability Model. PLoS Comput. Biol. *12*, e1005260.

Nassar, M.R., Wilson, R.C., Heasly, B., and Gold, J.I. (2010). An approximately Bayesian delta-rule model explains the dynamics of belief updating in a changing environment. J. Neurosci. *30*, 12366–12378.

Natan, R.G., Briguglio, J.J., Mwilambwe-Tshilobo, L., Jones, S.I., Aizenberg, M., Goldberg, E.M., and Geffen, M.N. (2015). Complementary control of sensory adaptation by two types of cortical interneurons. Elife *4*.

Nogueira, R., Abolafia, J.M., Drugowitsch, J., Balaguer-Ballester, E., Sanchez-Vives, M.V., and Moreno-Bote, R. (2017). Lateral orbitofrontal cortex anticipates choices and integrates prior with current information. Nat. Commun. *8*, 14823.

Samaha, J., Switzky, M., and Postle, B.R. (2018). Confidence boosts serial dependence in orientation estimation. bioRxiv.

Seriès, P., and Seitz, A.R. (2013). Learning what to expect (in visual perception). Front. Hum. Neurosci. *7*, 668.

Soetens, E., Boer, L.C., and Hueting, J.E. (1985). Expectancy or automatic facilitation?

Separating sequential effects in two-choice reaction time. J. Exp. Psychol.

Sommer, W., Leuthold, H., and Soetens, E. (1999). Covert signs of expectancy in serial reaction time tasks revealed by event-related potentials. Percept. Psychophys. *61*, 342–353.

Stocker, A.A., and Simoncelli, E.P. (2006). Noise characteristics and prior expectations in human visual speed perception. Nat. Neurosci. *9*, 578–585.

Sutton, R.S., and Barto, A.G. (2018). Reinforcement Learning: An Introduction (MIT Press).

Tervo, D.G.R., Proskurin, M., Manakov, M., Kabra, M., Vollmer, A., Branson, K., and Karpova, A.Y. (2014). Behavioral variability through stochastic choice and its gating by anterior cingulate cortex. Cell *159*, 21–32.

Urai, A.E., Braun, A., and Donner, T.H. (2017). Pupil-linked arousal is driven by decision uncertainty and alters serial choice bias. Nat. Commun. *8*, 14637.

Urai, A.E., de Gee, J.W., Tsetsos, K., and Donner, T.H. (2018). Choice history biases subsequent evidence accumulation.

Vernon, G., Farrow, D., and Reid, M. (2018). Returning Serve in Tennis: A Qualitative Examination of the Interaction of Anticipatory Information Sources Used by Professional Tennis Players. Front. Psychol. *9*, 895.

Wilder, M., Jones, M., and Mozer, M.C. (2009). Sequential effects reflect parallel learning of multiple environmental regularities. In Advances in Neural Information Processing Systems 22, Y. Bengio, D. Schuurmans, J.D. Lafferty, C.K.I. Williams, and A. Culotta, eds. (Curran Associates, Inc.), pp. 2053–2061.

Yu, A.J., and Cohen, J.D. (2008). Sequential effects: Superstition or rational behavior? Adv. Neural Inf. Process. Syst. *21*, 1873–1880.