1   Genetic analysis of the *Komagataella phaffii* centromeres by a color-based

2                               plasmid stability assay

Luiza Cesca Piva[1], Janice Lisboa De Marco[1], Lidia Maria Pepe de Moraes[1], Viviane

Castelo Branco Reis[1], Fernando Araripe Gonçalves Torres[1]*

[1]Laboratório de Biologia Molecular, Universidade de Brasília, Brasília, DF, Brazil,

70910-900.

*Corresponding author

E-mail: ftorres@unb.br

## Abstract

The yeast *Komagataella phaffii* is widely used as a microbial host for heterologous protein production. However, molecular tools for this yeast are basically restricted to a few integrative and replicative plasmids. Four sequences that have recently been proposed as the *K. phaffii* centromeres could be used to develop a new class of mitotically stable vectors. In this work we designed a color-based genetic assay to investigate genetic stability in *K. phaffii*. Plasmids bearing each centromere and the *ADE3* marker were evaluated in terms of mitotic stability in an *ade2/ade3* auxotrophic strain which allows plasmid screening through colony color. Plasmid copy number was verified through qPCR. Our results confirmed that the centromeric plasmids were maintained at low copy number as a result of typical chromosome-like segregation during cell division. These features, combined with high transformation efficiency and *in vivo* assembly possibilities, prompt these plasmids as a new addition to the *K. phaffii* genetic toolbox.

## Author summary

The methylotrophic yeast *Komagataella phaffii* is considered as one of the most important platforms for the production of proteins and metabolites. We sought in this study to develop a color-based genetic system widely used in other yeasts to assess mitotically stability of vectors carrying the proposed *K. phaffii* centromeres. First, we constructed a *K. phaffii* strain (LA3) mutant for *ADE2* and *ADE3*; this resulted in a strain that forms white colonies and when transformed with a vector (pPICH-ADE3) carrying *ADE3* turns red. Next, the four *K. phaffii* centromeres were cloned into pPICH-

ADE3 and tested in LA3 for copy number and plasmid stability. Centromeres are responsible for proper chromosome segregation during cell division, hence guaranteeing that both daughter cells receive one copy of the duplicated DNA. Our results show that three *K. phaffii* centromeres behaved as expected conferring extra stability to the replicative plasmids and maintaining them at low copy number. Once characterized, centromeres can be used as parts in the construction of advanced genetic manipulation tools, thus allowing the construction of strains capable of expressing large metabolic pathways for the production of complex biochemicals.

**Introduction**

*Komagataella phaffii* is a methylotrophic yeast of great industrial importance which has been used for more than 30 years as a heterologous protein production platform [1]. Its genome was first published in 2009 and has since then been refined and thoroughly studied [2,3]. As a result, in addition to a protein factory, *K. phaffii* has also been widely considered as a platform for the production of chemicals, biopharmaceuticals, vitamins and other molecules. However, the construction and regulation of new pathways demand complex molecular biology tools which are not readily available for this yeast [4].

*K. phaffii* genetic manipulation traditionally involves the use of shuttle vectors assembled in *Escherichia coli* and subsequently integrated into the yeast´s genome [5]. Recent studies have described the development of a wide range of genetic parts for use in this yeast, as well as new methods of plasmid assembly and transformation [6]. An alternative to integrative strategies is the use of replicative plasmids, which are usually

based on the well-known ARS1 sequence [1]. These plasmids may overcome some

drawbacks such as genetic instability in multi-copy strains and non-specific integration

[7,8]. In addition, they present higher transformation efficiency when compared to

integrative vectors and can be assembled by *in vivo* recombination, which eliminates the

need for bacterial transformation [9,10]. However, replicative plasmids show low

mitotic stability when compared to integrative vectors and few vector options are

available for use [11]. Stability problems can be circumvented by the creation of

centromeric plasmids, which may provide proper segregation during mitosis. A greater

mitotic stability as well as low copy number allow stable and constant protein

expression [12]. Centromeric plasmids can be constructed *in vivo,* allowing the

assembly and cloning of large sequences including whole metabolic pathways and

regulatory regions [13]. Therefore, the construction of such vectors would be of great

value for *K. phaffii* strain development in the context of synthetic biology.

Centromeres are typically surrounded by large heterochromatin sections in most

organisms [14]. Their structure ranges from simple "point" centromeres of only ~125 bp

in *Saccharomyces cerevisiae* to epigenetic, sequence-independent centromeres, such as

those present in plants and animals. The reason for this phenomenon is that, for most

eukaryotes, centromeres are maintained epigenetically and not genetically. Sequence

homologies are rare in and between species, hampering the definition of a consensus

sequence. In addition, some DNA regions can be centromeric or not depending on its

function in previous cell cycles, which highlights the epigenetic nature of the

centromere [15].

As for non-conventional yeasts there are wide variations in centromere size and

structure. *Candida glabrata* has centromeres that show some homology to the CDEI and

CDEIII regions of *S. cerevisiae* while *Kuraishia capsulata* centromeres have 200-bp

conserved sequences [16,17]. On the other hand, *Candida tropicalis,*

*Schizosaccharomyces pombe* and *Candida albicans* have regional centromeres named

after their sizes which range from 3 to 110 kb [18–20].

*K. phaffii* centromeres have recently been identified, bearing no sequence

similarities to those of any other yeast [3]. Since centromere function relies strongly on its

structure rather than on its sequence, a centromere-specific histone H3 variant (CSE4)

was used in the search for centromeric regions in *K. phaffii*. A CSE4 homolog was

identified in chromosome 2 and tagged with a fluorescence marker. The corresponding

nuclear localization of the histone-DNA complex indicated a centromere pattern typical

of budding yeasts [3]. Tridimensional conformation analysis followed the centromere

clustering pattern observed in yeasts and narrowed down all four *K. phaffii* centromere

locations to 20 kb windows [21].

Considering that a low transcription rate is typical of centromeric regions, RNA-

seq analysis allowed to pinpoint the putative centromeric locations for all four *K. phaffii*

centromeres [3]. Similarly to *C. tropicalis* and *S*. pombe, *K. phaffii* centromeres are

formed by inverted repeats. All four sequences have two inverted repeats of ~2,5 kb,

separated by a central segment of 800 to 1300 bp. Chromatin immunoprecipitation

sequencing analysis showed that the CSE4 histone binds preferably to the central

region, but also along the inverted repeats [22].

*K. phaffii* centromeric sequences contain early replication peaks with

autonomously replicating sequences, characteristics that are also observed in

centromeres of other yeasts [23,24]. According to recently published studies, there are

native ARS sequences contained within centromeres 2 and 4. These comprise regions

within the inverted repeats, as well as unique adjacent sequences [22,25].

111            In order to expand the functional analysis of the *K. phaffii* centromeres we

112     sought in this study to develop a genetic system based on an *ade2/ade3* auxotrophic

113     strain and a replicative vector carrying the wild-type *ADE3*. Vectors carrying each

114     individual centromere were used to assess plasmid copy number and mitotic stability.

116     **Results and Discussion**

118            In yeasts, adenine synthesis pathway is used as a tool for auxotrophic selection,

gene copy number indicator and for plasmid stability analysis [26]. Many genes from

this pathway have been deleted in *S. cerevisiae* in order to create auxotrophic strains,

while in *K. phaffii* studies have only focused on *ADE1* and *ADE2*[ 27,28]. *K. phaffii*

LA2, a strain mutant for *ADE2* [29], was used as a starting point for the construction of

a strain that would allow plasmid stability verification. Deletion of *ADE2* results in cells

auxotrophic for adenine which accumulate a red pigment [26] while deletion of genes

located upstream, such as *ADE1* or *ADE3*, should prevent the formation of such

pigment [27]. As expected, the deletion of *ADE3* in LA3 strain results in white colonies

(Fig 1). Deletion of *ADE3* in *S. cerevisiae* has regulatory effects in the histidine

synthesis pathway [30]. Consequently, *ade2 ade3* strains are not only auxotrophic for

adenine, but also for histidine. In order to verify if this phenotype is applicable to *K.*

*phaffii*, we plated strains X-33, LA2 and LA3 on MD medium without supplementation,

comparing growth and colony color to cells plated on MD medium with adenine and

histidine (Fig 1). LA3 strain displayed the expected histidine auxotrophy phenotype,

showing that the adenine-histidine pathways in *K. phaffii* and *S. cerevisiae* have

common characteristics.

In order to assess plasmid stability, we first constructed plasmid pPICH-ADE3

bearing the *ADE3* gene (Fig 2). When transformed with pPICH-ADE3, LA3 cells

should return to being red and any changes on colony color would allow a simple

screening of plasmid loss [26]. Although adenine auxotrophy has been explored for

other purposes in *K. phaffii* [28], this particular color-based system has not yet been

used for measuring plasmid stability in this yeast.

pPICH-ADE3 was used for cloning of all four *K. phaffii* centromeres. Since it

revealed extremely difficult to amplify entire centromeric regions we designed a

strategy to amplify centromeres in halves in order to reduce fragment size and to avoid

primer annealing inside the inverted repeats (Fig 3). Amplified fragments exhibited in

their ends overlapping regions that would allow recombination between each other and

with vector pPICH-ADE3.

LA3 strain was individually transformed with pPICH-ADE3 and all four

constructed centromeric plasmids (pPICH-CEN1-4). Plasmids were verified for

autonomous replication by PCR analysis which confirmed their circular conformation

(data not shown) and by plasmid rescue in *E. coli*. In this case, only pPICH-CEN3 could

not be recovered, probably because CEN3 was the only centromere that could not be

cloned directly in *E. coli*.

Plasmid stability was firstly verified through colony color in non-selective

medium (Fig 4). When plated on YPD non-selective medium, colonies transformed with

pPICH-ADE3 lost their color rapidly and presented a red center with large white edges,

a result consistent with plasmid instability. In contrast, strains transformed with all

centromeric plasmids presented a uniform red coloration throughout the colony.

Further stability examination of the centromeric plasmids was performed by

growing cells in liquid YPD medium for 144 hours. After diluting and plating cultures

on non-selective medium, red and white colonies were counted and compared between

each construction (Fig 5). LA3 strain transformed with pPICH-ADE3 did not yield red

colonies in any growth period, indicating that the plasmid was mitotically unstable.

Conversely, centromeric plasmids presented a higher mitotic stability than pPICH-

ADE3. After 96 hours of growth, cells with pPICH-CEN1 started to present white

colonies, while all other centromeric plasmids remained stable. After 144 hours,

pPICH-CEN1 was lost in most colonies while the other centromeric plasmids were lost

in <10% cells. The reason for the instability of pPICH-CEN1 is still unclear but it could

be related to the absence of important regulatory regions.

Yeast centromeric plasmids knowingly have a higher mitotic stability under non-

selective conditions than common replicative vectors since they are equally segregated

between daughter cells and therefore provide a uniform culture of cells containing the

plasmid [26]. A centromeric vector containing *K. phaffii* CEN2 has been constructed

and it presented an enhanced stability when compared to a replicative plasmid [25]. In

addition to *K. phaffii* and *S. cerevisiae*, centromeric plasmids have been developed for

other yeasts such as *S. pombe, C. glabrata* and *Scheffersomyces stipitis* and in all cases

enhanced plasmid stability under non-selective conditions was verified [12,17].

Yeast replicative plasmids are normally replicated but are unevenly distributed

between daughter cells, which creates both multi-copy and plasmidless cells [26]. Under

selective conditions, cells lacking the plasmid are unable to survive and the result is a

population of multi-copy plasmid-containing cells. The construction of centromeric

plasmids should provide better plasmid segregation and stability and cells should

maintain a low and stable plasmid copy number during yeast growth [31]. Plasmid copy

number was assessed by qPCR after strains were grown in YPD medium containing

zeocin in order to ensure that all cells assayed were harboring the centromeric plasmids.

Results from qPCR (Fig 6) indicate that the strain transformed with centromeric

plasmids carried 1-2 copies per cell while the replicative plasmid was present at

approximately 25 copies per cell. The difference between plasmid copy number for the

replicative vector and all centromeric vectors was significant according to a t-test

($p < 0.05$). This result illustrates the expected segregation pattern described above for

growth in selective conditions and, together with the mitotic stability analysis, provides

a clear picture of *K. phaffii* genetic manipulation using centromeric plasmids.

In a previous study, *K. phaffii* transformed with a centromeric plasmid was

analyzed regarding plasmid copy number and compared to both a replicative plasmid

and an integrative strategy [25]. Results exhibited an equal low number of plasmids per

cell in all strategies, which does not correspond to our observation. This can be

explained by the fact that their analysis was performed in non-selective medium, a

condition in which replicative plasmids are readily lost as we have shown in Fig 5A

(pPICH-ADE3).

Overall, our results indicate that centromeric plasmids could be employed as a

new tool for genetic manipulation of *K. phaffii*. Plasmids were maintained for long

periods in non-selective medium, indicating that growth can be performed without the

addition of antibiotics or any form of selective pressure. The centromeric plasmids' low

copy numbers per cell characterize a stable and homogeneous culture that can provide

reliable expression results. Finally, their structure as a circular molecule allows *in vivo*

plasmid assembly with relatively short homologous sequences when compared to

genomic integration techniques where sequences have to be much longer for directed

homologous recombination. Simpler assembly may also facilitate the construction of

larger and more sophisticated vectors such as yeast artificial chromosomes (YAC)

whose stability features may be also analysed by the color-based assay described in this

work.

**Methods**

**Strains and Media**

DNA cloning was performed using chemically competent *Escherichia coli* XL-

10 Gold (Agilent Technologies) grown in LB medium (5 g $L^{-1}$ yeast extract, 10 g $L^{-1}$

peptone and 10 g $L^{-1}$ NaCl, pH 7,2). When needed, agar was added to a final

concentration of 1,5%. When zeocin (25 µg $mL^{-1}$) was used for bacterial antibiotic

selection, NaCl concentration was reduced to 5 g $L^{-1}$.

*K. phaffii* strains were derived from X-33 (Invitrogen). LA2 strain (*amd2 ade2*)

was described in a previous work [29]. Yeast was routinely grown in YPD medium (10 g

$L^{-1}$ yeast extract, 20 g $L^{-1}$ peptone and 20 g $L^{-1}$ glucose). Solid medium used 2% agar.

Zeocin and geneticin, when used, were added at 100 µg $mL^{-1}$ and 500 µg $mL^{-1}$,

respectively. Hygromycin B was used to a final concentration of 50 µg $mL^{-1}$. Minimal

medium (MD) used 0,34% Yeast Nitrogen Base, 1% $(NH_4)_2SO_4$, 2% glucose,

0,00004% biotin and 0,0002% adenine or 0,004% histidine, when needed.

**PCR**

DNA was amplified using Invitrogen Platinum Taq DNA Polymerase (High

Fidelity), Promega GoTaq Colorless Master Mix or Sigma-Aldrich Accutaq LA DNA

Polymerase. All primers used in this work are shown in Table 1.

**Table 1. Primers used in this work.**

| Primer | Sequence | Enzyme |
|---|---|---|
| ADE3up-F | GATAAGCTTGATATCGAATTCCTGCAGCCC<u>CCC GGG</u>ACGTAATGGAATAACTGCTGAC | *Sma*I |
| ADE3up-R | GAAGTTATGGATCCTACGAGGTAATTGAAGGCT CAC | |
| ADE3lox-F | CTTCAATTACCTCGTAGGATCCATAACTTCGTAT AATG | |
| ADE3lox-R | CAATCTCTCCCTTGTCATCGGATCCATAACTTCG TATAG | |
| ADE3dw-F | GAAGTTATGGATCCGATGACAAGGGAGAGATT GAAG | |
| ADE3dw-R | GGTGGCGGCCGCTCTAGAACTAGTGGATCCCCC <u>CCCGGG</u>TGCAATGTACTGTTGAGTAGG | *Sma*I |
| ADE3conF | GGGGACCGGAGGTAAAAGAC | |
| ADE3conR | GTTGGAATAATTGCATGGTCTG | |
| MUT1(Hpa) | CCATTGACATGGTAAACAGTTGGA | |
| MUT2(Bam) | GAACACTGGGATCTTGGTTGAGG | |
| Cen1-F | GAGTTTAAAC<u>GGATCC</u>ACGAAGCAATGGATAG GCACT | *Bam*HI |
| Cen1-R | CCGCGAATTC<u>GGATCC</u>TGAAGTCTTTCAGAGAG GAGCA | *Bam*HI |
| Cen1c-F | CAAGTATGCGTGATCCCAGGT | |
| Cen1c-R | TACGAATTGTGGGGCTCTGT | |
| Cen2-F | GAGTTTAAAC<u>GGATCC</u>ATCTCCGTTGATACTCC CAAC | *Bam*HI |
| Cen2-R | CCGCGAATTC<u>GGATCC</u>ATCGACAAGCAGAACA CTAAG | *Bam*HI |

| | | |
|---|---|---|
| Cen2c-F | GAATGGAGGTGCTGGTGGTTA | |
| Cen2c-R | TGTAATGCTCGCTGGTGAGT | |
| Cen3-F | GAGTTTAAAC<u>GGATCC</u>AAGTGGTACACCAGTCAGCG | *Bam*HI |
| Cen3-R | CCGCGAATTC<u>GGATCC</u>TCAGTATTCAACTGCAACTGC | *Bam*HI |
| Cen3c-F | TCAGCCGAATACCCACACTT | |
| Cen3c-R | TCAGCCGTCAGCGAAATGAT | |
| Cen4-F | GAGTTTAAAC<u>GGATCC</u>CAAACGCACCGTCTTGTTCA | *Bam*HI |
| Cen4-R | CCGCGAATTC<u>GGATCC</u>AATTGATGTAGACGAGCAGC | *Bam*HI |
| Cen4c-F | TCAAGAATCGTACTGGCACCT | |
| Cen4c-R | CAAGCTCGTGAGATGGGATGT | |
| Cen370-F | CGCTCAGTGGAACGAAAACTCACGTTAAGGGATTTGGTCATGAGATCAGATCTAACATCCAAAGACGAAAGGTTGAATGAGTTTAAACGGATCCAAGTGG | |
| Cen370-R | GCTGGCCCTCTCTTCCCAGCTCACGAATCAGATCCTAAGTCCTACTCAACAGTACATTGCAGCGGCCGCGTTTAAACGAATTCGGATCCTCAGTATTCAA | |
| qZEO-F | CGACGTGACCCTGTTCATCA | |
| qZEO-R | TGGACACGACCTCCGACCA | |
| qHIS-F | GTGTATCCTGGCTTGGCATCT | |
| qHIS-R | GCCAAGTACGGTGTGACGTT | |

234 Restriction sites are underlined.

236 **DNA manipulation**

237   All basic DNA manipulation and analysis were performed as previously

described [32]. Restriction digestion was performed in accordance to the manufacturer

instructions (New England Biolabs), as well as vector dephosphorylation with Shrimp

Alkaline Phosphatase (Promega) and ligation with T4 DNA ligase (USB). In-Fusion

Cloning Kit (Clontech) was used for in-vitro assembly of plasmids. Site-directed

mutagenesis was performed using the Transformer Site-Directed Mutagenesis kit

(Clontech). PCR and gel purification used Promega Wizard SV Gel and PCR Clean-Up

System.

**Quantitative PCR (qPCR)**

Cells were grown to an O.D. of 1 in 10 mL YPD containing zeocin and then

collected by centrifugation at 2000 x g for 5 minutes. Cell pellet was resuspended with

1 mL 0,25% SDS and incubated at 98ºC for 8 minutes according to Nakamura (2018).

Finally, cell debris was removed by centrifugation and DNA was diluted 10-fold in

water before qPCR reactions.

Quantitative PCR reactions used primers qZEO-F and qZEO-R for plasmid

quantification and qHIS-F and qHIS-R as an internal single-copy control. Assays were

carried out with iTaq Universal SYBR Green Supermix (Bio-Rad) in a Rotor-Gene Q

(Qiagen) thermal cycler. Analysis used the absolute quantification method and standard

curves that ranged from $1x10^4$ to $1x10^8$ copies of the gene of interest. pPIC9

(Invitrogen) and pPICH linearized plasmids were used for construction of the standard

curves.

**Yeast transformation**

*K. phaffii* was electroporated following two different protocols. For integrative

cassettes, we followed the *Pichia* Expression Kit protocol (Invitrogen) and when using

replicative plasmids, we proceeded as described previously [33].

**Construction of an *ade2 ade3* strain for color-based stability assays**

266      Strain LA2 [29] was transformed with an *ADE3* deletion cassette and had the

marker recycled before moving on with the centromeric plasmid transformations.

Construction of the deletion cassette used PCR reactions assembled by an "In-Fusion"

cloning reaction. Briefly, primers ADE3up-F and R; ADE3dw-F and R were used for

PCR amplification of 491 bp and 582 bp, respectively, from *K. phaffii* genome. These

reactions amplified sequences used for directing homologous recombination and

substitution of the complete *ADE3* coding sequence. Meanwhile, primers ADE3lox-F

and ADE3lox-R amplified the *kanR* geneticin resistance cassette from plasmid pGKL

[34]. PCR fragments were assembled and cloned into pBluescript II SK$^+$ linearized with

*Sma*I. A final PCR reaction using primers ADE3up-F and ADE3dw-R amplified the

whole deletion cassette which was used for transformation of *K. phaffii* LA2. Cells were

selected in YPD containing geneticin.

278      The resulting *ade2 ade3* strain was later transformed with pYRCre2 [35] and

selected in YPD supplied with hygromycin B. This step promoted a Cre-mediated

excision of the *kanR* cassette eliminating geneticin resistance. After PCR confirmation

of marker recycling using primers ADE3conF and ADE3conR, the resulting strain was

plated in non-selective YPD medium, causing loss of the pYRCre2 plasmid. The

resulting strain was named LA3.

**Construction of centromeric plasmids containing *ADE3***

286      Plasmid pPICH [29], which is derived from pPICHOLI (MoBiTec), was digested

with *Not*I for cloning of the *K. phaffii* native *ADE3*. The complete gene was amplified

from X-33 DNA using primers ADE3up-F and ADE3dw-R following digestion with

*Not*I. After vector dephosphorylation, fragments were ligated and transformed into *E.*

*coli* XL-10 Gold. One positive clone was then submitted to site-directed mutagenesis

using primers Mut1(Hpa) and Mut2(Bam) for removal of the *Bam*HI restriction site

present within the *ADE3* coding sequence. The final plasmid containing ARS1, the *Sh*

*ble* resistance marker and *ADE3* was named pPICH-ADE3.

pPICH-ADE3 was digested with *Bam*HI for cloning of all four *K. phaffii*

centromeres. These were amplified from *K. phaffii* X-33 genomic DNA using two PCR

reactions for each centromeric sequence. Primers Cen1/2/3/4-F and Cen1/2/3/4c-R

amplified the first inverted repeat of each centromere while primers Cen1/2/3/4c-F and

Cen1/2/3/4-R amplified the other half of the sequences. In order to promote *in vitro*/*in*

*vivo* assembly amplicons had approximately 80 bp homology between each other and

15 bp with pPICH-ADE3. Firstly, we attempted an "In-Fusion" cloning reaction for

each of the four centromeres using linearized pPICH-ADE3 and the two PCR

fragments. Plasmids were extracted and analyzed by restriction digestion. Centromeres

1, 2 and 4 were successfully assembled and cloned into the plasmid through this

strategy.

Centromere 3 did not yield any *E. coli* clones following the "In-Fusion"

reaction; therefore, we proceeded to an *in vivo* assembly strategy. Primers Cen370-F

and Cen3c-R; Cen3c-F and Cen370-R amplified both inverted repeats adding 70 bp of

homologous sequences between the fragments and pPICH-ADE3. Finally, we

transformed *K. phaffii* LA3 using the linearized vector and both centromeric fragments,

using 85 bp of homology for directing recombination. Clones were selected in YPD

supplied with zeocin.

The resulting plasmids were named pPICH-CEN1, pPICH-CEN2, pPICH-CEN3

and pPICH-CEN4. All plasmids were transformed into *K. phaffii* LA3 for subsequent

stability and quantification assays.

**Stability analysis**

LA3 strain transformed with each of the four centromeric plasmids was grown in

20 mL YPD for 16 hours at 28ºC and 200 rpm. This culture was inoculated to 20 mL

YPD to an initial optical density of 0,1. After 24 h of growth under the same conditions

the culture was used as inoculum for another flask containing 20 mL YPD  to an $OD_{600}$

of 0,1. This procedure was repeated every 24 h until a total of 144 hours. At 96 and at

144 hours of growth, a small amount of cells was diluted $10^6$-fold and 100 μL of this

dilution were plated on YPD. Plates were incubated at 30 ºC for 72 h.

**References**

1. Cregg, J. M., Barringer, K. J., Hessler,  a Y., Madden, K. R. *Pichia pastoris* as a host

system for transformations. Mol. Cell. Biol. 1985; 5(12): 3376–85.

2. De Schutter, K., Lin, Y.-C., Tiels, P., Van Hecke, A., Glinka, S., Weber-Lehmann, J.,

et al. Genome sequence of the recombinant protein production host *Pichia pastoris*. Nat.

Biotechnol. 2009;27(6): 561–566.

3. Sturmberger, L., Chappell, T., Geier, M., Krainer, F., Day, K. J., Vide, U., et al.

Refined *Pichia pastoris* reference genome sequence. J. Biotechnol. 2016; 235: 121–131.

4. Schwarzhans, J. P., Luttermann, T., Geier, M., Kalinowski, J., Friehs, K. Towards

systems metabolic engineering in *Pichia pastoris*. Biotechnol. Adv. 2017;35(6): 681–

710.

5. Cereghino, J. L., Cregg, J. M. Heterologous protein expression in the methylotrophic

yeast *Pichia pastoris*. FEMS Microbiol. Rev. 2000; 24(1): 45–66.

6. Obst, U., Lu, T. K., Sieber, V. A modular toolkit for generating *Pichia pastoris*

secretion libraries. *ACS Synth. Biol.* 2017; 6 (6): 1016-1025.

7. Camattari, A., Goh, A., Yip, L. Y., Tan, A. H. M., Ng, S. W., Tran, A., et al.

Characterization of a panARS-based episomal vector in the methylotrophic yeast *Pichia*

*pastoris* for recombinant protein production and synthetic biology applications. Microb.

Cell Fact. 2016;15(139). doi: 10.1186/s12934-016-0540-5

8. Schwarzhans, J.-P., Wibberg, D., Winkler, A., Luttermann, T., Kalinowski, J., Friehs,

K. Integration event induced changes in recombinant protein productivity in *Pichia*

*pastoris* discovered by whole genome sequencing and derived vector optimization.

Microb. Cell Fact. 2016;15( 84). doi: 10.1186/s12934-016-0486-7

9. Yu, X., Wang, R., Zhang, M., Xu, Y., Xiao, R. Enhanced thermostability of a

Rhizopus chinensis lipase by in vivo recombination in *Pichia pastoris*. Microb. Cell

Fact. 2012;11(102). doi : 10.1186/1475-2859-11-102

10. Lee, C. C., Williams, T. G., Wong, D. W. S., Robertson, G. H. An episomal

expression vector for screening mutant gene libraries in *Pichia pastoris*. Plasmid

2005;54: 80–85.

11. Liachko, I., Youngblood, R. A., Tsui, K., Bubb, K. L., Queitsch, C., Raghuraman,

M. K.,  et al. GC-Rich DNA elements enable replication origin activity in the

methylotrophic yeast *Pichia pastoris*. PLOS Genet. 2014;10:3. doi:

10.1371/journal.pgen.1004169

12. Cao, M., Gao, M., Lopez-garcia, C. L., Wu, Y., Seetharam, A. S., Severin, A. J., et

al. Centromeric DNA facilitates nonconventional yeast genetic engineering. ACS Synth.

Biol. 2017; 6(8): 1545-1553.

13. Kouprina, N., Larionov, V. Transformation-associated recombination (TAR)

cloning for genomics studies and synthetic biology. Chromosoma 2016; 125(4): 621-

632.

14. Lyn Chan, F., Wong, L. H. Transcription in the maintenance of centromere

chromatin identity. Nucleic Acids Res. 2012;40(22): 11178–11188.

15. Malik, H. S., Henikoff, S. Major evolutionary transitions in centromere complexity.

Cell 2009;138(6): 1067–1082.

16. Morales, L., Noel, B., Porcel, B., Marcet-Houben, M., Hullo, M. F., Sacerdot, C., et

al. Complete DNA sequence of *Kuraishia capsulata* illustrates novel genomic features

among budding yeasts (Saccharomycotina). Genome Biol. Evol. 2013;5(12):  2524–

2539.

17. Kitada, K., Yamaguchi, E., Arisawa, M. Isolation of a *Candida glabrata* centromere

and its use in construction of plasmid vectors. Gene 1996;175(1-2): 105–108.

18. Ketel, C., Wang, H. S. W., McClellan, M., Bouchonville, K., Selmecki, A., Lahav,

T., et al. Neocentromeres form efficiently at multiple possible loci in *Candida albicans*.

PLOS Genet. 2009;5(3). doi: 10.1371/journal.pgen.1000400

19. Chatterjee, G., Sankaranarayanan, S. R., Guin, K., Thattikota, Y., Padmanabhan, S.,

Siddharthan, R., et al. Repeat-associated fission yeast-like regional centromeres in the

ascomycetous budding yeast *Candida tropicalis*. PLOS Genet. 2016;12(2). doi:

10.1371/journal.pgen.1005839

20. Smirnova, J. B., McFarlane, R. J. The unique centromeric chromatin structure of

*Schizosaccharomyces pombe* is maintained during meiosis. J. Biol. Chem. 2002;

277(22); 19817–19822.

21. Varoquaux, N., Liachko, I., Ay, F., Burton, J. N., Shendure, J., Dunham, M. J., et al.

Accurate identification of centromere locations in yeast genomes using Hi-C. Nucleic

Acids Res. 2015;43(11): 5331–5339.

22. Coughlan, A. Y., Hanson, S. J., Byrne, K. P., Wolfe, K. H. Centromeres of the yeast

*Komagataella phaffii* (*Pichia pastoris*) have a simple inverted-repeat structure. Genome

Biol. Evol. 2016;8(8): 2482–2492.

23. McCarroll, R. M., Fangman, W. L. Time of replication of yeast centromeres and

telomeres. Cell 1988;54(4): 505–513.

24. Koren, A., Tsai, H., Tirosh, I., Burrack, L. S., Barkai, N., Berman, J. Epigenetically-

inherited centromere and neocentromere DNA replicates earliest in s-phase. PLOS

Genet. 2010;6(8). doi: 10.1371/journal.pgen.1001068

25. Nakamura, Y., Nishi, T., Noguchi, R., Ito, Y., Watanabe, T., Nishiyama, T., et al.

Construction of a new stable, autonomously replicating plasmid vector containing

*Pichia pastoris* centromeric DNA. Appl. Environ. Microbiol. 2018;84(15). doi:

10.1128/AEM.02882-17

26. Hieter, P., Mann, C., Snyder, M., Davis, R. W. Mitotic stability of yeast

chromosomes : a colony color assay that measures nondisjunction and chromosome

loss. Cell 1985;40(2): 381–392.

27. Ugolini, S., Bruschi, C. V. The red/white colony color assay in the yeast

*Saccharomyces cerevisiae*: epistatic growth advantage of white ade8-18, ade2 cells over

red ade2 cells. Curr. Genet. 1996;30(6): 485–492.

28. Du, M., Battles, M. B., Nett, J. H. A color-based stable multi-copy integrant

selection system for *Pichia pastoris* using the attenuated *ADE1* and *ADE2* genes as

auxotrophic markers. Bioeng. Bugs 2012;3(1): 32–37.

29. Piva, L. C., De Marco, J. L., de Moraes, L. M. P., Reis, V. C. B., Torres, F. A. G.

Acetamidase as a dominant recyclable marker for *Komagataella phaffii* strain

engineering. Appl. Microbiol. Biotechnol. 2018;102(6): 2753–2761.

30. Rébora, K., Laloo, B., Daignan-Fornier, B. Revisiting purine-histidine cross-

pathway regulation in *Saccharomyces cerevisiae*: a central role for a small molecule.

Genetics 2005;170(1): 61–70.

31. Steensels, J., Snoek, T., Meersman, E., Nicolino, M. P., Voordeckers, K.,

Verstrepen, K. J. Improving industrial yeast strains: exploiting natural and artificial

diversity. FEMS Microbiol. Rev. 2014;38(5): 947–995.

32. Sambrook, J., Fritsch, E., Maniatis, T. Molecular cloning: a laboratory manual. 2nd

ed. New York: Cold Spring Harbor Laboratory Press; 1989.

33. Wu, S., Letchworth, G. J. High efficiency transformation by electroporation of

*Pichia pastoris* pretreated with lithium acetate and dithiothreitol. Biotechniques

2004;36(1): 152–154.

34. Reis, V. C. B., Nicola, A. M., de Souza Oliveira Neto, O., Batista, V. D. F., de

Moraes, L. M. P., Torres, F. A. G. Genetic characterization and construction of an

auxotrophic strain of *Saccharomyces cerevisiae* JP1, a Brazilian industrial yeast strain

for bioethanol production. J. Ind. Microbiol. Biotechnol. 2012;39(11): 1673–1683.

35. Betancur, M. O., Reis, V. C. B., Nicola, A. M., De Marco, J. L., de Moraes, L. M.

P., Torres, F. A. G. Multicopy plasmid integration in *Komagataella phaffii* mediated by

a defective auxotrophic marker. Microb. Cell Fact. 2017;16(99). doi: 10.1186/s12934-

017-0715-8

**Fig 1. Strain phenotypic analysis on defined media.** *K. phaffii* X-33 (wild-type), LA2

and LA3 cultures were spotted on MD medium with or without supplementation.

**Fig 2. Map of vector pPICH-ADE3.** The *Not*I site was used for cloning *K. phaffii*

*ADE3* gene.

**Fig 3. Strategy for amplification of *K. phaffii* centromeres.** Schematic representation

of a typical *K. phaffii* centromere. Inverted repeats are represented by green arrows.

Primer annealing regions are shown by small arrows.

**Fig 4. Plasmid stability analysis.** LA3 strain transformed with pPICH-ADE3 and the

four centromeric plasmids was grown on YPD medium for 3 days until color

development.

**Fig 5. Plasmid stability test.** (A) Aliquots of the liquid cultures were collected at after

96 and 144 hours of growth and plated on YPD medium. Red colonies represent cells

that maintained the *ADE3*-containing plasmid, while white colonies have lost it. Red

portions of the bars represent red colonies; light pink bars represent white colonies. (B)

A plate representing a typical result after 144 h growth.

**Fig 6. Plasmid copy number determination.** The number of plasmids in each cell was

estimated by qPCR. LA3 was used as negative control. Statistical analysis comparing

each of the centromeric plasmids with the replicative vector was performed through a t-

test using GraphPad Prism 5. ($p < 0.05$). Error bars depict the standard deviation of the

mean (n = 3).

X-33  LA2  LA3

MD

MD
+Adenine

MD
+Adenine
+Histidine

**pPICH-CEN1**  **pPICH-CEN2**  **pPICH-CEN3**  **pPICH-CEN4**

**pPICH-ADE3**