G-Protein signaling accelerates stem cell divisions in *Drosophila* males

Manashree Malpe[1,*], Leon F. McSwain[2,*], Karl Kudyba[1,*], Chun L. Ng[3], Jennie

Nicholson[1], Maximilian Brady[1], Yue Qian[4], Vinay Choksi[5], Alicia G. Hudson[1], Benjamin

B. Parrott[6], and Cordula Schulz[1,#]

1 Department of Cellular Biology, University of Georgia, Athens GA 30602, USA

[2] Winship Cancer Institute, Emory University, Atlanta, GA 30322, USA

[3] University of Texas Southwestern Medical Center, Dallas, TX

[4] University of North Georgia, Department of Biology, Oakwood, GA 30566

[5] School of Medicine, Duke University, Durham, NC 27708, USA

[6] Odum School of Ecology, University of Georgia, Athens, GA 30602, USA

* These authors contributed equally

Corresponding author: cschulz@uga.edu

Abstract

Adult stem cells divide to renew the stem cell pool and replenish specialized cells that

are lost due to death or usage. However, little is known about the mechanisms

regulating how stem cells adjust to a demand for specialized cells. A failure of the stem

cells to respond to this demand can have serious consequences, such as tissue loss, or

prolonged recovery post injury.

Here, we challenge the male germline stem cells (GSCs) of *Drosophila*

melanogaster for the production of specialized cells using mating experiments. We

show that repeated mating reduced the sperm pool and accelerated germline stem cell

(GSC) divisions. The increase in GSC divisions depended on the activity of the highly

conserved G-proteins. Germline expression of RNA-Interference (RNA-*i*) constructs

against G-proteins or a dominant negative G-protein eliminated the increase in GSC

divisions in mated males. Consistent with a role for the G-proteins in the regulation of

GSC divisions, RNA-*i* against seven out of 35 G-protein coupled receptors (GPCRs)

within the germline cells also eliminated the capability of males to accelerate their GSC

divisions in response to mating. Our data show that GSCs are receptive to GPCR

stimulus, potentially through a network of interactions among multiple signaling

pathways.

Introduction

Metazoan tissues undergo homeostasis wherein stem cells divide and their daughter

cells proliferate and differentiate to replace lost cells. The human hematopoietic stem

cells, for example, renew a remarkable number of about one trillion blood cells per day

[1, 2]. Stem cells have to maintain a baseline mitotic activity for the production of daughter

cells that account for the daily turnover of differentiated cells. However, whether stem

cells can modulate their mitotic activity in response to demands that challenge the

system is not fully explored. In some instances, stem cells respond to physiological

cues; for example, murine hematopoietic stem cells divide more frequently during

pregnancy due to increased oestrogen levels [3]. In *Drosophila melanogaster*, intestinal

stem cells initiate extra cell divisions upon ablation of differentiated gut cells. *Drosophila*

GSCs modulate their mitotic activity in response to environmental conditions, such as

nutrient availability and temperature [4-7].

Drosophila is an excellent model for identifying the molecules and mechanisms

that regulate and fine-tune tissue homeostasis. A plethora of genetic tools are available

for manipulating and monitoring dividing adult stem cells in *Drosophila*. The small size

of the fly, the short generation cycle, and the fairly low costs covering their maintenance

allow for high throughput screens. Here, we subjected several thousand male and

several million virgin female flies to mating experiments, a task challenging to perform

with vertebrate model organisms. We discovered that repeated mating caused a

reproducible and significant increase in GSC division frequency in *Drosophila wild-type*

(*wt*) males. Our analysis revealed that this response to mating was dependent on the

activity G-proteins. Impairing G-protein activity from the germline cells eliminated the

ability of the GSCs to increase their division frequency in response to mating.

G-proteins are highly conserved molecules that associate with GPCRs. GPCRs

constitute a large family of cell surface receptors that mediate the cell's response to a

wide range of external stimuli, including odors, pheromones, hormones, and

neurotransmitters. Loss of GPCR signaling affects countless developmental and neural

processes in humans, as well as vertebrate and invertebrate model organisms [8-10]. Here

we show that reducing the expression of seven out of 35 GPCRs via RNA-*i* from the

germline cells eliminated the capability of males to accelerate their GSC divisions when

mated. These were the Serotonin (5-HT) Receptors 1A, 1B and 7, Metuselah (Mth),

Metuselah-like5 (Mth-l5), Octopamineβ2-Receptor (Octβ2R), and a predicted GPCR

encoded by *CG12290*.

A role for any of these GPCRs in regulating GSC division frequency is novel. No

previous study has identified any functional role for Mth-l5 or CG12290. Serotonin,

Octopamine, and Mth signaling play opposing roles in life-span, locomotion, and sleep

[11-16]. Mth signaling also regulates vesicle trafficking at the synapse, Octopamine

signaling regulates ovulation, and Serotonin signaling plays essential roles in memory

formation and learning [17-19].

Results

Mating increased the percentage of GSCs in mitosis

As is typical for many stem cells, the *Drosophila* GSCs are found in a specific cellular

microenvironment. They are located at the tip of the gonad, where they are attached to

somatic hub cells (Figure 1A, A'). Upon GSC division, one of the daughter cells, called

gonialblast, undergoes four rounds of stem cell daughter characteristic transit amplifying

divisions, resulting in 16 spermatogonia. Subsequently, spermatogonia enter a tissue-

specific differentiation process. They grow in size, undergo the two rounds of meiosis,

and develop through extensive morphological changes into elongated spermatids [20].

According to this tightly controlled homeostasis program, each GSC division can only

produce 64 spermatids (Figure 1A). Thus, an increase in sperm production is reliant on

the GSCs.

We investigate division frequency using an established immuno-fluorescence

protocol [7]. In this approach, Vasa-positive GSCs are identified based on their position

adjacent to FasciclinIII (FasIII)-positive hub cells (Figure 1A'). The percentage of GSCs

in mitosis, the M-phase index (MI), is investigated by staining against phosphorylated

Histone-H3 (pHH3). The MI of the GSCs (MI^{GSC}) is calculated by dividing the number of

pHH3-positive GSCs by the total number of GSCs.

To investigate if stem cells can modulate their division frequency in response to a

demand for specialized cells, we challenged *Drosophila* males in mating experiments.

For each experiment, 80-100 males were exposed individually to virgin females. An

equal number of male siblings were each kept in solitude and served as the non-mated

controls. To keep experimental variation to a minimum, we employed a three-day

mating protocol for all experiments, kept the animals under the same conditions,

dissected the testes at the same time of the same day, and dissected experimental

groups in tandem. Using *wt* males, we obtained robust and reproducible increases in

MI^{GSC} in response to mating. The box-plot in Figure 1B shows the observed difference

in MI^{GSC} between mated and non-mated populations of isogenized *wt*, *Oregon R* (*OR*),

males from 17 independent mating experiments. Interestingly, we observed variability in

MI^{GSC} among males of each condition. The MI^{GSC} of non-mated males ranged from six

to nine percent, with a median at seven percent. The MI^{GSC} of mated males ranged from

11 to 18 percent, with a median at 16.5 percent. We hypothesize that this variability in

MI^{GSC} within each condition is due to naturally occurring physiological differences within

the flies. Likewise, the increase in MI^{GSC} in response to mating varied among the

different experiments, but, in each of the experiments, the increase was biologically and

statistically significant.

We next investigated if only a few males within a population contributed to the

increase in MI^{GSC} or whether the effect of mating is reflected by changes in the MI^{GSC}

across a population. These data are displayed in frequency distribution graphs (FDGs).

FDGs show how often a particular value is represented within a population. When the

distribution of the MI^{GSC} for testes within one population of *OR* flies was plotted, the

resulting FDG revealed that mated males had significantly fewer testes with an MI^{GSC} of

zero and more testes with higher MI^{GSC} compared to non-mated siblings (Figure 1C).

We observed the same result for another isogenized *wt* strain, *Canton S* (*CS*, Figure

1D). We conclude that mating affected the MI^{GSC} of many males within one mated

population.

Finally, we asked how long or frequently we had to mate the males to see an

increase in MI^{GSC}. For this, we mated *OR* males to varying numbers of virgin females

and subsequently analyzed how many of their GSCs were in mitotic division. When we

exposed *OR* males for 24 hours to one (1F, 24 hrs), two (2F, 24 hrs), or three (3F,

24hrs) female virgins, no significant difference in MI^{GSC} between non-mated and mated

males was apparent (Figure S1A). Robust and reproducible increases were seen in *OR*

males that were exposed to three virgin females on each of two (2x3F, 48 hrs) or three

(3x3F, 72 hrs) days of mating (Figure S1A). We conclude that males have to mate

repeatedly for an increase in MI^{GSC} to occur. The increase in MI^{GSC} in mated males was

reversible, showing that the response to mating was dynamic. Moving males back into

solitude after the three-day mating experiment for 48 hours (3x3F, 120 hrs) eliminated

the increase in MI^{GSC} (Figure S1A). Control males mated for 120 hours (5x3F, 120 hrs),

in contrast, still had a significant increase in MI^{GSC} (Figure S1A).

Mating increased GSC division frequency

As another measure of cell divisions, we investigated the percentage of GSCs in

synthesis phase (S-phase) of the cell cycle. Testes were labeled with 5-ethynyl-2'-

deoxyuridine (EdU) and the S-phase index of the GSCs (SI^{GSC}) was calculated by

dividing the number of EdU-positive GSCs by the total number of GSCs. Using pulse-

labeling experiments, we observed that mated *OR* males displayed significant higher

SI^{GSC} compared to their non-mated siblings (Figure 1E). Together with the increase in

the MI^{GSC} this suggests that mating accelerates stem cell divisions.

To test this hypothesis, the lengths of the cell cycle were measured using EdU

feeding experiments. In this approach, *OR* animals were fed EdU during the mating

experiment. We then calculated how many GSCs had been in S-phase at different time

points. Our EdU-incorporation experiment revealed that the number of EdU-positive

GSCs increased rapidly after 24 hours of feeding and reached 80% at 60 hours of

feeding (Figure 1F). Prolonged feeding further increased the numbers of EdU-positive

GSCs but this data was excluded from the study as the majority of males that were fed

EdU while mating had died by 72 hours of the experiment. The response curve we

obtained in this time-course experiment is different from the response curves reported

by other groups that used bromo-deoxy-uridine (BrDU) as the thymidine analog instead

of EdU. For example, the non-mated males in our experiment had about 70% of EdU-

positive GSCs after 48 hours of feeding. A study using *white* (*w*) mutant animals fed the

same concentration of the thymidine homologue had a steeper response curve, in which

85% of the GSCs were BrDU-marked after 48 hours of feeding [21]. Another study using

yellow, vermillion (y, v) flies showed even steeper response curves where 100% of the

GSCs were BrDU-labeled after 24 hours. However, in this study, animals were fed a 30

times higher concentration of the thymidine homologue than used in our study [22]. We

propose that the different response curves are due to the different genetic backgrounds,

chemicals and doses.

Most importantly, mated males had significantly more EdU-positive GSCs at 36

and 48 hours of mating compared to their non-mated siblings (Figure 1F). This

experiment shows that, in mated males, more GSCs had entered S-phase of the cell

cycle. We conclude that mated males had accelerated GSC divisions.

To further investigate how mating affects the cell cycle, we employed the Fly-

Fucci technology in combination with the UAS-Ga4 expression system (Duffy, 2002

#321)(Phelps, 1998 #34)(Zielke, 2014 #1122). With Fly-Fucci, the coding regions of

fluorescent proteins are fused to the destruction boxes of cell cycle regulators, allowing

the marking of different cell cycle stages. These artificial proteins are expressed under

control of the Yeast Upstream Activating Sequences [23] (Zielke, 2014 #1122). UAS-

controlled target genes can be expressed under spatial control using tissue-specific

Gal4-transactivators. In addition, temporal control can be applied to their expression by

exposing the flies to different temperatures (Phelps, 1998 #34)(Duffy, 2002 #321). For

our experiments, we used a *nanos-Gal4*-transactivator (NG4) with a reported

expression in GSCs, gonialblasts, and spermatogonia (Van Doren, 1998 #55). Using

two independent Fucci-lines, we observed that MI^{GSC} did not significantly increase in

mated *Fucci/NG4* males while mated controls animals (*Fucci/wt*) increased their MI^{GSC}

compared to non-mated siblings (Figure S1B). We conclude that expressing FUCCI-

constructs from these fly lines within the GSCs interfered with their ability to significantly

increase MI^{GSC} in response to mating. One possible explanation for this could be that

the expression of proteins with destruction boxes could overload the cell cycle

machinery of male GSCs.

Mating reduced the sperm pool

To confirm that our mating experiments created a demand for sperm, we explored

differences in the sperm pool of the seminal vesicles between non-mated and mated

males. For this, we used two different transgenic constructs that label the sperm. A Don

Juan-Green Fluorescent Protein (DJ-GFP) reporter labels the sperm bodies and allows

to assess the overall amount of sperm within the seminal vesicles [24]. A ProtamineB-

GFP (Mst35B-GFP) line, on the other hand, only labels the sperm heads and can be

used to count the sperm within the seminal vesicles [25]. With each of these reporters,

individualized mature sperm was normally seen within the seminal vesicle of the male

reproductive tract.

According to the literature, the total number of sperm within one seminal vesicle

varies among different *Drosophila* species and among genetic backgrounds [25-27]. To

keep the genetic background consistent among our experiments, we crossed each of

the reporter lines to *OR* females and used their male progeny for our mating

experiments. The seminal vesicles were then analyzed at days one to three of the

experiment. Based on the size and the fluorescence of the seminal vesicles, we first

sorted them into three classes. Class 1 and class 2 seminal vesicles were completely

filled with GFP-positive sperm heads. However, class 1 seminal vesicles were very wide

(Figure 2A), while class 2 seminal vesicles were thinner (Figure 2B). Class 3 seminal

vesicles contained only few GFP-positive sperm heads and had areas that were not

filled with GFP (Figure 2C, arrows). A quantification revealed that non-mated males had

mostly class 1 and 2 seminal vesicles, while mated males had mostly class 3 seminal

vesicles. While we still detected class 1 and 2 seminal vesicles in males that had mated

for only one day, their numbers were severely reduced in males after two and three

days of mating (Figure 2D-F).

To further validate our observation that mating reduces the amount of sperm, we

developed an automated procedure that calculates the volume occupied by Mst35B-

GFP-positive sperm heads per seminal vesicle in all focal planes (voxels in Figure 2G).

This allowed us to investigate larger numbers of seminal vesicles compared to a

previously reported method, in which images through the seminal vesicles were

flattened and sperm heads counted by eye [25]. Furthermore, a computer-based

calculation eliminates subjective bias introduced by the investigator. Based on our

computer calculation, the sperm heads of mated males occupied significantly less

volume within the seminal vesicles than the sperm in non-mated males (Figure 2G).

Notably, the total volume occupied by sperm became more reduced with every day of

mating. By days two and three of mating it ranged from 0.1 to 0.4 x 10^6 voxels per

seminal vesicle. The non-mated sibling controls, in contrast, maintained a large GFP-

occupied volume in their seminal vesicles, with an average of 1.2 x 10^6 voxels per

seminal vesicle. The computer program estimated the numbers of sperm per seminal

vesicle of non-mated males around 2000, while males that were mated for two or three

days had less than 500 sperm in their seminal vesicles. As our mated males showed a

drastic reduction in sperm, we argue that we have created a demand for sperm.

Mating had no effect on GSC numbers

It was previously reported that females significantly increased the numbers of their

GSCs upon mating [28]. According to the literature, an adult male gonad contains up to

twelve GSCs per testis, but the exact number of GSCs per testes appears to vary

among different strains and laboratories. One study using a *wt* strain of males reported

six to ten GSCs per testis, while other studies using transgenic males in a *w* mutant

genetic background reported 8.94 and 12.3 GSCs per testis, respectively [29-31]. Among

our fly lines, we found variation in GSC numbers as well. The distribution of GSCs

ranged from one to 14 per testis, with an average of seven GSCs per testis. Males from

an isogenized *OR* stock had the lowest average number of GSCs, having only four to

five GSCs per testis (Figure 3A). Males from an isogenized *CS* stock had an average of

six GSCs per testis (Figure 3B). Animals mutant for *w* alleles, w^{1118} and w^1, had on

average eight and seven GSCs per testis, respectively (Figures 3C, 3D). Males from a

v^1, y^1 stock, which serves as the genetic background for many RNA-*i* lines, had the

highest average number of GSCs, at 11 GSCs per testis (Figure 3E). We believe that

the obtained GSC numbers are specific to the fly lines in our laboratory and do not

necessarily reflect the numbers of GSCs in fly stocks of other laboratories.

Importantly, we did not observe a significant difference in the numbers of GSCs

between non-mated and mated siblings in any of these fly lines (Figure 3A-E). We

concluded that mating did not affect the numbers of GSCs in our fly stocks. However,

the observed variation in GSC numbers prompted us to perform our experiments in

animals from as similar genetic backgrounds as possible. All males reported in the

following of this manuscript carried the X-chromosome from our isogenized *OR* line.

The increase in MI^{GSC} upon mating required G-protein signaling

Drosophila mating is a complex and genetically controlled behavior that is dependent on

neural circuits [32]. This implicates a possible neuronal control in regulating GSC divisions

during mating. Therefore, we wanted to focus on the type of signaling pathway

commonly stimulated during neural activity, G-protein signaling [33, 34]. In a non-stimulated

cell, a trimeric complex of G-proteins, G_α, G_β, and G_γ is associated with classical

GPCRs (Figure 4A, step 1). When ligand binds to the GPCR, a guanidyl exchange

factor within the GPCR becomes activated that exchanges GDP for GTP in the G_α

subunit. The exchange leads to the dissociation of G_α and the $G_{\beta/\gamma}$ complex from each

other and from the GPCR. Remaining attached to the membrane, G_α and $G_{\beta/\gamma}$ diffuse

along it and activate downstream signal transducers (Figure 4A, step 2) [35, 36]. Most

organisms have multiple genes that encode for each of the G-protein subunits.

Drosophila has six G_α, three G_β, and two G_γ proteins, yet only a few examples are

available in the literature associating a specific *Drosophila* G-protein with an upstream

GPCR [35, 37].

Animals mutant for G-protein subunits are often lethal, making it problematic to

investigate their roles in the adult. Furthermore, studying G-protein signaling in animals

lacking their function throughout the whole body may could affect behavior and

physiology of the fly, leading to confounding effects on mating and GSC divisions.

Fortunately, large collections of RNA-*i*-lines are available that are expressed under

control of UAS. To reduce G-protein signaling we employed two separate *nanos*-Gal4-

transactivators (*NG4*), *NG4-1* and *NG4-2.* When RNA-*i* against the different G-protein

subunit was expressed within the germline cells via *NG4-1*, several of the mated males

displayed only a weak increase in MI^{GSC} compared to their non-mated siblings (Table

1). We focused on an RNA-*i*-line that is directed against the subunit $G_{\alpha}i$ as animals

expressing this construct within the germline did not show any increase in MI^{GSC} in

response to mating (Table 1). For reproducibility, we conducted each of the following

experiments in triplicates. We used progeny from transgenic Gal4 and UAS-flies that

had been crossed to *wt* as positive controls. As expected, each population of positive

control males displayed a significant increase in MI^{GSC} when mated (Figure 4B).

Experimental flies expressing $G_{\alpha}i$-*i* via *NG4-1* or *NG4-2*, however, failed to increase

MI^{GSC} (Figure 4C).

We next sought to validate the role for G-protein signaling in GSC division

frequency by an alternative approach. A dominant negative version of *Drosophila* $G_{\gamma}1$

(dnG$_{\gamma}$1) is available that serves as a reliable tool to abolish G-protein signaling [38]. Males

expressing dnG$_{\gamma}$1 via either *NG4-1* or *NG4-2* did not show an increase in MI^{GSC} in

response to mating (Figure 3C). Control dnG$_{\gamma}$1/*wt* animals, on the other hand, had

increased MI^{GSC} upon mating (Figure 3B). These data clearly show that signaling via G-

proteins is required for the increase in MI^{GSC}. Plotting the results in FDGs confirmed that

mated control animals had significantly fewer testes with an MI^{GSC} of zero and more

testes with higher MI^{GSC} compared to non-mated males (Figures S3A-D), and that this

response to mating was eliminated in experimental males (Figures S3E-H).

In mammalian cells, three major G-protein-dependent signaling cascades have

been described (Figure 3A, steps 3a, b, c) [33, 39]. For *Drosophila*, the literature provides

little information on the signaling cascades downstream of GPCRs but it is generally

assumed that the mammalian signal transducers are conserved in flies. To further

validate that an increase in MI^{GSC} upon mating is regulated by G-protein signaling we

expressed RNA-*i* and mis-expression constructs for conserved signal transducers via

NG4 and found that males expressing RNA-*i*-lines for one of the *Drosophila Protein*

Kinase C (*PKC*) proteins, *PKC98E*, for *Inositol-triphosphate 3-Kinase* (*IP3K),* and for

Ca2+/Calmodulin-dependent protein kinase II (CaMKII) indeed failed to increase MI^{GSC}

in response to mating (Table 1).

314   RNA-i against seven distinct GPCRs blocked the increase in MI^{GSC} upon mating

To further confirm that G-protein signaling regulates the increase in MI^{GSC} we aimed

towards identifying the upstream GPCRs. Next Generation Sequencing (NGS) of RNA

from *wt* testis tips revealed the expression of 140 receptors, including 35 classical

GPCRs (Figure 5 and Table 2). The functions of many of these GPCRs have not been

studied yet and mutant animals are only available in rare cases. Expressing RNA-*i*-

constructs against most GPCRs in the germline had little to no effect on the ability of the

GSCs to increase their MI^{GSC} in response to mating (Table 2). RNA-*i* against three

Serotonin Receptors (5HT-1A, 5HT-1B and 5HT-7), Mth, Mth-l5, Octβ2R, and a

predicted GPCR encoded by CG12290, clearly and reproducibly eliminated this ability.

Animals carrying UAS-controlled RNA-*i*-constructs against these GPCRs (*GPCR-i*) were

crossed to *wt*, *NG4-1* and *NG4-2*, and MI^{GSC} of their progeny was investigated. Each of

the controls (*GPCR-i/wt*) increased their MI^{GSC} when repeatedly mated to females in

each of the triplicate experiments (Figure 5A and Figures S3A-G). However, when the

GPCR-i-animals were crossed with either *NG4-1* (Figure 5B and Figures S3H-N) or

NG4-2 (Figure 5C and Figures S3O-U) the MI^{GSC} of their non-mated and mated progeny

did not significantly differ. Confirming the necessity of the GPCRs in increasing MI^{GSC},

we investigated alternative RNA-*i*-lines. A second RNA-*i*-line for Mth blocked the

increase in MI^{GSC} in mated males and a second RNA-*i*-line for 5HT-1A displayed only a

weak response to mating (Table 3).

Mating success was evaluated by two criteria: visual observation and the

appearance of progeny. When flies were anesthetized to exchange the females for

fresh virgins, several copulating pairs of males and females were always observed.

Furthermore, 100 single females that had been exposed to males on day one of the

experiment were placed into one food vial each and mating success evaluated a few

days later by counting the percentage of vials with progeny. Most males in this study

sired 60-90% of the females. Specifically, each of the *GPCR-i/NG4-1* males produced

offspring (Table S1), showing that a block in the increase in MI^{GSC} is not caused by a

failure to mate but by a lack of GPCR signaling. Viable alleles of 5HT-1A and 5HT-1B

were not pursued as alternative strategies because they displayed only a weak mating

success rate (Table S1).

Finally, we wanted to assure that male age had no effect on the increase in

MI^{GSC}. We performed a time-course experiment of one, two, three, and four-week old

OR males. We found that mated males of all ages showed robust increases in MI^{GSC}

compared to their non-mated siblings (Figure S1C). We conclude that aging animals for

up to four weeks had little to no effect on the ability of *wt* GSCs to increase their MI^{GSC}

in response to mating, and that the age of the transgenic animals used in this study

(three weeks of age at the time of testes dissection) had no impact on the obtained

results.

Discussion

Here, we show that repeated mating reduced the sperm pool and increased GSC

division frequency. Using highly controlled experiments, we demonstrate that mated

males had more GSCs in M-phase and S-phase of the cell cycle compared to non-

mated males. Mated males also showed faster incorporation of EdU indicating that their

GSCs progressed faster through the cell cycle. Our findings demonstrate that GSCs can

respond to a demand for sperm by accelerating their mitotic activity. Based on RNA-*i*

targeting G-proteins and a dominant negative construct against G$_\gamma$1, the increase in

MI^{GSC} of mated males is dependent on G-protein signaling. Furthermore, signal

transducers predicted to act downstream of G-proteins and GPCRs predicted to act

upstream of G-proteins also appeared to be required for the response to mating.

Due to the lack of mutants and a potential interference of whole animal knock-

down in the behavior of the flies, we used tissue-specific expression of RNA-*i*-

constructs. It is surprising that our studies revealed potential roles for seven instead of a

single GPCR in the increase of MI^{GSC} in response to mating. A possible explanation is

that some of the RNA-*i*-lines have off-target effects. RNA-*i*-hairpins can cause the

down-regulation of unintended targets due to stretches of sequence homologies,

especially when long hairpins are used [40, 41]. However, with the exception of the RNA-*i*-

line directed against 5HT-7, all lines that produced a phenotype contain second

generation vectors with a short, 21 nucleotide hairpin predicted to have no off-target

effects [42]. We hypothesize that multiple GPCRs regulate the increase in MI^{GSC} in

response to mating. Consistent with this, expression of second RNA-*i*-line directed

against Mth or 5HT-1A interfered with the increase in MI^{GSC} in mated males.

Our finding that RNA-*i* against several GPCRs blocked the increase in MI^{GSC} in

mated males suggests a high level of complexity in the regulation of GSC divisions. In

the literature, increasing evidence has emerged that GPCRs can form dimers and

oligomers and that these physical associations have a variety of functional roles,

ranging from GPCR trafficking to modification of G-protein mediated signaling [43-45]. In *C.*

elegans, two Octopamine receptors, SER-3 and SER-6, additively regulate the same

signal transducers for food-deprived-mediated signaling. One possible explanation for

the non-redundant function of the two receptors was the idea that they form a functional

dimer [46]. In mammalian cells, 5-HT receptors can form homo-dimers and hetero-dimers

and, dependent on this, have different effects on G-protein signaling [47-49]. In cultured

fibroblast cells, for example, G-protein coupling is more efficient when both receptors

within a 5-HT4 homo-dimer bind to agonist instead of only one [50]. In cultured

hippocampal neurons, hetero-dimerization of 5-HT1A with 5-HT7 reduces G-protein

activation and decreases the opening of a potassium channel compared to 5-HT1A

homo-dimers [51]. The formation of hetero-dimers of GPCRs with other types of receptors

plays a role in depression and in the response to hallucinogens in rodents [52, 53].

Alternatively, or in addition to the possibility that some or all of the seven GPCRs

form physical complexes, a role for several distinct GPCRs in regulating GSC division

frequency could be explained by crosstalk among the downstream signaling cascades.

One signaling cascade could, for example, lead to the expression of a kinase that is

activated by another cascade. Similarly, one signaling cascade could open an ion

channel necessary for the activity of a protein within another cascade. Unfortunately,

the literature provides little information on *Drosophila* GPCR signal transduction

cascades and only very few mutants have been identified that affect a process

downstream of GPCR stimulation. Thus, it remains to be explored how stimulation of

the GPCRs and G-proteins increase GSC divisions.

The role for G-protein signaling in regulating the frequency of stem cell divisions

is novel. Our data suggest that the increase in MI^{GSC} in response to mating is regulated

by external signals, potentially arising from the nervous system, that stimulate G-protein

signaling within the GSCs. Based on the nature of the GPCRs, the activating signal

could be Serotonin, the Mth ligand, Stunted, Octopamine, or two other, yet unknown,

signals that activate Mth-I5, and CG12290 [54-56]. It will be interesting to address which of

these ligands are sufficient to increase MI^{GSC}, in what concentrations they act, by which

tissues they are released, and whether they also affect other stem cell populations.

Methods

Fly husbandry

Flies were raised on a standard cornmeal/agar diet and maintained in temperature-,

light-, and humidity-controlled incubators. Unless otherwise noted, all mutations,

markers, and transgenic lines are described in the *Drosophila* database and were

obtained from the Bloomington stock center (Consortium, 2003 #132).

UAS/Gal4-expression studies

Two separate X; UAS-*dicer*; nanos-*Gal4* (NG4-1 and NG4-2) fly lines were used as

transactivators. Females from the transactivator line or *wt* females were crossed with

males carrying target genes under the control of UAS in egg lay containers with fresh

apple juice-agar and yeast paste to generate either experimental or control flies. The

progeny were transferred into food bottles, raised to adulthood at 18°C, males collected

and then shifted to 29°C for seven days to induce high activity of Gal4 prior to the

mating experiment. Note that the males were not collected as virgins as to avoid any

potential developmental or learning effects on our experiments.

Mating experiments

Unless otherwise noted, mating experiments were performed at 29°C. Males and virgin

females were placed on separate apple juice-agar plates with yeast paste overnight to

assure they were well fed prior to their transfer into mating chambers. Single males

were placed into each mating slot either by themselves (non-mated) or with three virgin

females (mated) and the chambers closed with apple juice-agar lids with yeast paste.

Females were replaced by virgin females on each of the following two days and apple

juice-agar lids with yeast paste were replaced on a daily basis for both non-mated and

mated animals. In most instances, females from the stock $X\square X, y, w, f$ / Y / *shi*[ts] were

used as virgins. When raised at 29°C, only females hatch from this stock. For fertility

tests, *OR* virgins were used. Note that 10-20% of the mated males died during the

experiment while only 5% of the non-mated siblings died.

Immuno-fluorescence and microscopy

Animals were placed on ice to immobilize them. Gonads were dissected in Tissue

Isolation Buffer (TIB) and collected in a 1.5 ml tube with TIB buffer on ice for no more

than 30 minutes. Gonads were then fixed, followed by immuno-fluorescence staining

and imaging as previously described [7]. The mouse anti-FasciclinIII (FasIII) antibody

(1:10) developed by C. Goodman was obtained from the Developmental Studies

Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of

Iowa, Department of Biology, Iowa City, IA 52242. Goat anti-Vasa antibody (1:50 to

1:300) was obtained from Santa Cruz Biotechnology Inc. (sc26877), anti-

phosphorylated Histone H3 (pHH3) antibodies (1:100 to 1:1000) were obtained from

Fisher (PA5-17869), Milllipore (06-570), and Santa Cruz Biotechnology Inc. (sc8656-R).

Secondary Alexa 488, 568, and 647-coupled antibodies (1:1000) and Slow Fade Gold

embedding medium with DAPI were obtained from Life Technologies. Images were

taken with a Zeiss Axiophot, equipped with a digital camera, an apotome, and

Axiovision Rel. software. Statistical relevance was determined using the two-tailed

Graphpad student's t-test.

EdU-labeling experiments

The EdU-labeling kit was obtained from Invitrogen and the procedure performed

following manufacturer's instructions. For EdU-pulse labeling experiments, animals

were mated as described above, and the dissected testes incubated with 10mM EdU in

PBS for 30 minutes at room temperature prior to fixation. For EdU-feeding experiments,

OR males were fed 10 mM EdU in liquid yeast provided on paper towels. These animals

were mated at room temperature (21°C) because the paper towels easily dried out at

higher temperatures, causing the flies to dehydrate and die.

Sperm head volumetric calculations

In order to easily evaluate sperm numbers, we turned to computer analysis in Python.

By quantifying the volume of GFP signal we generated estimates to the amount of

sperm in each seminal vesicle. Image stacks were taken of individual seminal vesicles.

After masking relevant regions, each image set was normalized by mean subtraction

and division by the standard deviation, followed by rescaling image intensity to

encompass the range of the image. To remove signal noise, a median filter was applied

and the mask refined by Otso thresholding. We determined signal volume by hysteresis

thresholding. This approach initially thresholds an image at an upper limit, and then

expands the region by adjacent pixels satisfying the lower threshold. We set the lower

bound at the value generated from a triangle threshold and the upper threshold as the

median value above the lower limit. The number of signal voxels was calculated and

normalized to an expected size of a single sperm head. Our analysis utilized OpenCV

3.4.2, Scipy1.2.1, Scikit-image 0.14.2, Numpy 1.16.2, Matplotlib 3.0.3, Seaborn 0.9.0,

as well as built in Python 3.7.3 modules [57-61].

References

1.   Dancey, J.T., Deubelbeiss, K.A., Harker, L.A. & Finch, C.A. Neutrophil kinetics in

man. *J Clin Invest* **58**, 705-715 (1976).

2.   Erslev, A. Production of erythrocytes, in *Hematology*. (ed. B.E. William WJ,

Erslev AJ, Lichtman MA) 365-376 (

Mc-Graw-Hill, New York, NY; 1983).

3.   Nakada, D. *et al.* Oestrogen increases haematopoietic stem-cell self-renewal in

females and during pregnancy. *Nature* **505**, 555-558 (2014).

4.   Hsu, H.J., LaFever, L. & Drummond-Barbosa, D. Diet controls normal and

tumorous germline stem cells via insulin-dependent and -independent

mechanisms in Drosophila. *Dev Biol* **313**, 700-712 (2008).

5.   Amcheslavsky, A., Jiang, J. & Ip, Y.T. Tissue damage-induced intestinal stem

cell division in Drosophila. *Cell Stem Cell* **4**, 49-61 (2009).

6.   McLeod, C.J., Wang, L., Wong, C. & Jones, D.L. Stem cell dynamics in response

to nutrient availability. *Curr Biol* **20**, 2100-2105 (2010).

7.   Parrott, B.B., Hudson, A., Brady, R. & Schulz, C. Control of germline stem cell

division frequency--a novel, developmentally regulated role for epidermal growth

factor signaling. *PLoS One* **7**, e36460 (2012).

8.   Schoneberg, T. *et al.* Mutant G-protein-coupled receptors as a cause of human

diseases. *Pharmacol Ther* **104**, 173-206 (2004).

9.   Wettschureck, N. & Offermanns, S. Mammalian G proteins and their cell type

506   specific functions. *Physiol Rev* **85**, 1159-1204 (2005).

507   10.   Langenhan, T. *et al.* Model Organisms in G Protein-Coupled Receptor Research.

508   *Mol Pharmacol* **88**, 596-603 (2015).

509   11.   Lin, Y.J., Seroude, L. & Benzer, S. Extended life-span and stress resistance in

510   the Drosophila mutant methuselah. *Science* **282**, 943-946 (1998).

511   12.   Selcho, M., Pauls, D., El Jundi, B., Stocker, R.F. & Thum, A.S. The role of

512   octopamine and tyramine in Drosophila larval locomotion. *J Comp Neurol* **520**,

513   3764-3785 (2012).

514   13.   Silva, B., Goles, N.I., Varas, R. & Campusano, J.M. Serotonin receptors

515   expressed in Drosophila mushroom bodies differentially modulate larval

516   locomotion. *PLoS One* **9**, e89641 (2014).

517   14.   Crocker, A. & Sehgal, A. Octopamine regulates sleep in drosophila through

518   protein kinase A-dependent mechanisms. *J Neurosci* **28**, 9377-9385 (2008).

519   15.   Yuan, Q., Joiner, W.J. & Sehgal, A. A sleep-promoting role for the Drosophila

520   serotonin receptor 1A. *Curr Biol* **16**, 1051-1062 (2006).

521   16.   Li, Y. *et al.* Octopamine controls starvation resistance, life span and metabolic

522   traits in Drosophila. *Sci Rep* **6**, 35359 (2016).

523   17.   Song, W. *et al.* Presynaptic regulation of neurotransmission in Drosophila by the

524   g protein-coupled receptor methuselah. *Neuron* **36**, 105-119 (2002).

525   18.   Lee, H.G., Seong, C.S., Kim, Y.C., Davis, R.L. & Han, K.A. Octopamine receptor

526   OAMB is required for ovulation in Drosophila melanogaster. *Dev Biol* **264**, 179-

527   190 (2003).

19.   Sitaraman, D. *et al.* Serotonin is necessary for place memory in Drosophila. *Proc*

Natl Acad Sci U S A **105**, 5579-5584 (2008).

20.   Fuller, M.T. Spermatogenesis, in *The development of Drosophila melanogaster*,

Vol. 1. (ed. M. Bate, Martinez-Arias, A) 71-147 (Cold Spring Harbor Press, Cold

Spring Harbor; 1993).

21.   Wallenfang, M.R., Nayak, R. & DiNardo, S. Dynamics of the male germline stem

cell population during aging of Drosophila melanogaster. *Aging Cell* **5**, 297-304

(2006).

22.   Yang, H. & Yamashita, Y.M. The regulated elimination of transit-amplifying cells

preserves tissue homeostasis during protein starvation in Drosophila testis.

Development **142**, 1756-1766 (2015).

23.   Abdouh, M., Albert, P.R., Drobetsky, E., Filep, J.G. & Kouassi, E. 5-HT1A-

mediated promotion of mitogen-activated T and B cell survival and proliferation is

associated with increased translocation of NF-kappaB to the nucleus. *Brain*

Behav Immun **18**, 24-34 (2004).

24.   Santel, A., Blumer, N., Kampfer, M. & Renkawitz-Pohl, R. Flagellar mitochondrial

association of the male-specific Don Juan protein in Drosophila spermatozoa. *J*

Cell Sci **111 (Pt 22)**, 3299-3309 (1998).

25.   Tirmarche, S. *et al.* Drosophila protamine-like Mst35Ba and Mst35Bb are

required for proper sperm nuclear morphology but are dispensable for male

fertility. *G3 (Bethesda)* **4**, 2241-2245 (2014).

26.   Pitnick, S., Markow, T.A. Male gametic Strategies: Sperm Size, Testes Size, and

the Allocation of Ejaculate among Successive Mates by the Sperm-Limited Fly

Drosophila Pachea and its Relatives. *The American Naturalist* **143**, 785-819

(1994).

27.   Kubrak, O.I., Kucerova, L., Theopold, U., Nylin, S. & Nassel, D.R.

Characterization of Reproductive Dormancy in Male Drosophila melanogaster.

Front Physiol **7**, 572 (2016).

28.   Ameku, T. & Niwa, R. Mating-Induced Increase in Germline Stem Cells via the

Neuroendocrine System in Female Drosophila. *PLoS Genet* **12**, e1006123

(2016).

29.   Chen, D. *et al.* Gilgamesh is required for the maintenance of germline stem cells

in Drosophila testis. *Sci Rep* **7**, 5737 (2017).

30.   Yamashita, Y.M., Jones, D.L. & Fuller, M.T. Orientation of asymmetric stem cell

division by the APC tumor suppressor and centrosome. *Science* **301**, 1547-1550

(2003).

31.   Sheng, X.R. & Matunis, E. Live imaging of the Drosophila spermatogonial stem

cell niche reveals novel mechanisms regulating germline stem cell output.

Development **138**, 3367-3376 (2011).

32.   Manoli, D.S., Fan, P., Fraser, E.J. & Shah, N.M. Neural control of sexually

dimorphic behaviors. *Curr Opin Neurobiol* **23**, 330-338 (2013).

33.   Geppetti, P., Veldhuis, N.A., Lieu, T. & Bunnett, N.W. G Protein-Coupled

Receptors: Dynamic Machines for Signaling Pain and Itch. *Neuron* **88**, 635-649

(2015).

34.   Lee, D. Global and local missions of cAMP signaling in neural plasticity, learning,

and memory. *Front Pharmacol* **6**, 161 (2015).

35. McCudden, C.R., Hains, M.D., Kimple, R.J., Siderovski, D.P. & Willard, F.S. G-

protein signaling: back to the future. *Cell Mol Life Sci* **62**, 551-577 (2005).

36. Oldham, W.M. & Hamm, H.E. Heterotrimeric G protein activation by G-protein-

coupled receptors. *Nat Rev Mol Cell Biol* **9**, 60-71 (2008).

37. Boto, T., Gomez-Diaz, C. & Alcorta, E. Expression analysis of the 3 G-protein

subunits, Galpha, Gbeta, and Ggamma, in the olfactory receptor organs of adult

Drosophila melanogaster. *Chem Senses* **35**, 183-193 (2010).

38. Deshpande, G., Godishala, A. & Schedl, P. Ggamma1, a downstream target for

the hmgcr-isoprenoid biosynthetic pathway, is required for releasing the

Hedgehog ligand and directing germ cell migration. *PLoS Genet* **5**, e1000333

(2009).

39. Moolenaar, W.H. G-protein-coupled receptors, phosphoinositide hydrolysis, and

cell proliferation. *Cell Growth Differ* **2**, 359-364 (1991).

40. Kulkarni, M.M. *et al.* Evidence of off-target effects associated with long dsRNAs

in Drosophila melanogaster cell-based assays. *Nat Methods* **3**, 833-838 (2006).

41. Moffat, J., Reiling, J.H. & Sabatini, D.M. Off-target effects associated with long

dsRNAs in Drosophila RNAi screens. *Trends Pharmacol Sci* **28**, 149-151 (2007).

42. Perkins, L.A. *et al.* The Transgenic RNAi Project at Harvard Medical School:

Resources and Validation. *Genetics* **201**, 843-852 (2015).

43. Filizola, M. & Weinstein, H. The study of G-protein coupled receptor

oligomerization with computational modeling and bioinformatics. *FEBS J* **272**,

2926-2938 (2005).

44.   Milligan, G. G protein-coupled receptor dimerisation: molecular basis and

597   relevance to function. *Biochim Biophys Acta* **1768**, 825-835 (2007).

598   45.   Terrillon, S. & Bouvier, M. Roles of G-protein-coupled receptor dimerization.

599   *EMBO Rep* **5**, 30-34 (2004).

600   46.   Yoshida, M., Oami, E., Wang, M., Ishiura, S. & Suo, S. Nonredundant function of

601   two highly homologous octopamine receptors in food-deprivation-mediated

602   signaling in Caenorhabditis elegans. *J Neurosci Res* **92**, 671-678 (2014).

603   47.   Lukasiewicz, S. *et al.* Hetero-dimerization of serotonin 5-HT(2A) and dopamine

604   D(2) receptors. *Biochim Biophys Acta* **1803**, 1347-1358 (2010).

605   48.   Herrick-Davis, K. Functional significance of serotonin receptor dimerization. *Exp*

606   *Brain Res* **230**, 375-386 (2013).

607   49.   Xie, Z., Lee, S.P., O'Dowd, B.F. & George, S.R. Serotonin 5-HT1B and 5-HT1D

608   receptors form homodimers when expressed alone and heterodimers when co-

609   expressed. *FEBS Lett* **456**, 63-67 (1999).

610   50.   Pellissier, L.P. *et al.* G protein activation by serotonin type 4 receptor dimers:

611   evidence that turning on two protomers is more efficient. *J Biol Chem* **286**, 9985-

612   9997 (2011).

613   51.   Renner, U. *et al.* Heterodimerization of serotonin receptors 5-HT1A and 5-HT7

614   differentially regulates receptor signalling and trafficking. *J Cell Sci* **125**, 2486-

615   2499 (2012).

616   52.   Borroto-Escuela, D.O., Tarakanov, A.O. & Fuxe, K. FGFR1-5-HT1A

617   Heteroreceptor Complexes: Implications for Understanding and Treating Major

618   Depression. *Trends Neurosci* **39**, 5-15 (2016).

53. Moreno, J.L., Holloway, T., Albizu, L., Sealfon, S.C. & Gonzalez-Maeso, J.

Metabotropic glutamate mGlu2 receptor is necessary for the pharmacological

and behavioral effects induced by hallucinogenic 5-HT2A receptor agonists.

Neurosci Lett **493**, 76-79 (2011).

54. Saudou, F., Boschert, U., Amlaiky, N., Plassat, J.L. & Hen, R. A family of

Drosophila serotonin receptors with distinct intracellular signalling properties and

expression patterns. *EMBO J* **11**, 7-17 (1992).

55. Cvejic, S., Zhu, Z., Felice, S.J., Berman, Y. & Huang, X.Y. The endogenous

ligand Stunted of the GPCR Methuselah extends lifespan in Drosophila. *Nat Cell*

Biol **6**, 540-546 (2004).

56. Maqueira, B., Chatwin, H. & Evans, P.D. Identification and characterization of a

novel family of Drosophila beta-adrenergic-like octopamine G-protein coupled

receptors. *J Neurochem* **94**, 547-560 (2005).

57. van der Walt, S., Colbert, C., Varoquaux, G. The NumPy Array: A Structure for

Efficient Numerical Vomputation. *Computing in Science and Engineering*, 22-30

(2011).

58. van der Walt, S., Schoenberger, J.L., Nunez-Iglesias, J., Boulogne, F., Warner,

J.D., Yager, N., Gouillart, E., Yu, T., and the scikit-image contributors scikit-

image: Image processing in Python. *PerJ:e453* (2014).

59. Travis, E., Oliphant, E. A guide to NumPy. (2006).

60. Hunter, J.D. Matplotib: A 2D Graphics Environment. *Computing in Science and*

Engineering, 90-95 (2007). https://archive.org/details/NumPyBook.

61.   Jones, E., Oliphant, E., Peterson, P., *et al* SciPy: Open Source Scientific Tools

for Python. (2001-). https://www.scipy.org.

Acknowledgements

The authors dedicate this manuscript to Bruce Baker, who was one of the foremost

scientists in the field, and a great colleague and friend. The authors are grateful to

Richard Zoller, Yue Qian, Megan Aarnio, Heather Kudyba, Jacqueline Uribe,

Chidemman Ihenacho, Stefani Moore, Sampreet Reddy, Amanda Cameron, Chederli

Belongilot, Dylan Ricke, Kenneth Burgess, Amanda Redding, Erin Guillebeau, Jennifer

Murphy, Chantel McCarty, Sarah Murphy, Haley Grable, Mitch Hanson, Haein Kim, and

Sarah Rupert for technical assistance. We thank Bruce Baker, Carmen Robinett,

Edward Kravitz, Matthew Freeman, Mark Brown, Erika Matunis, Steve DiNardo,

Margaret Fuller, Alan Spradling, Hannele Rahuola-Baker, Eric Bohman, Celeste Berg,

Wolfgang Lukowitz, Patricia Moore, Jim Lauderdale, Michael Tiemeyr, Scott Dougan,

Rachel Roberts-Galbraith, and Rhonda Snook for helpful discussions, and Heath Aston

Zachary Letts for comments on the manuscript. We are especially grateful to Barry

Ganetzky for the $X \square X, shi^{ts}$ fly stock and to Wolfgang Lukowitz for the use of his

microscope. This work was supported by NSF grants #0841419 and #1355009, and

UGA bridge funds given to CS.

Author contributions

M.M, B.B.P, and C.S developed and supervised the project, L.F.M coordinated the

mating experiments, K.K. identified the GPCRs expressed in testes tips and developed

the computer analysis for the sperm counts, all authors performed the experiments,

M.M, K.K., and CS wrote the manuscript.

Competing interests

The authors declare no competing interests.

Tables

Table 1. MI^{GSC} from control, RNA-*i* and overexpression lines directed against G-

protein subunits and other signal transducers

UAS-driven expression for the listed genes in the germline via *NG4-1*. BL #:

Bloomington stock number, Single and Mated: number of pHH3-positive GSCs/total

number of GSCs = MI^{GSC}, Diff: MI^{GSC} of mated males minus MI^{GSC} of non-mated males.

For RNA-*i*-lines marked by asterisks siblings outcrossed to *wt* did not show a strong

response to mating either, suggesting leakiness of the lines.

Genotype	BL#	Crossed to:	MI^{GSC} Single	MI^{GSC} Mated	Diff.
UAS-G$_\alpha$f-i	43201	NG4-1	19/448=4.2%	52/452=11.5%	7.3
	25930*	NG4-1	54/1031=5.2%	76/1183=6.1%	0.9
UAS-G$_\alpha$i-i	34924	NG4-1	22/335=6.6%	49/322=15.2%	8.6
	40890	NG4-1	33/597=5.5%	59/536=11%	5.5
	31133	NG4-1	5/269=1.9%	19/285=6.7%	4.8
UAS-G$_\alpha$o-i	34653*	NG4-1	23/313=7.3%	26/295=8.8%	1.5
	28010	NG4-1	18/240=7.5%	27/231=11.7%	4.2

UAS-G$_\alpha$q-i	36820	NG4-1	24/403=6.0%	32/268=11.9%	5.9
	33765	NG4-1	33/320=10%	50/298=16.8%	6.8
	36775	NG4-1	9/153=5.9%	21/255=8.6%	2.7
	31268	OR	17/233=7.3	15/169=8.9	1.6
		OR	14/335=4.2	12/164=7.3	3.1
		OR	31/568=5.5	27/333=8.1	2.6
		NG4-1	25/556=4.5	8/296=2.7	-2.8
		NG4-1	23/332=6.9	23/291=7.9	1
		NG4-1	23/542=4.2	19/577=3.3	-0.9
		NG4-1	71/1430=5.0	50/1164=4.3	-0.7
	30735	NG4-1	14/318=4.4%	23/293=7.8%	3.4
UAS-G$_\alpha$s-i	29576	NG4-1	49/605=8.1%	60/615=9.7%	1.6
	50704	NG4-1	82/1137=7.3%	121/1527=7.9%	0.6
UAS-G$_\beta$5-i	28310	NG4-1	10/306=3.3%	20/292=6.9%	3.6
UAS-G$_\beta$13F-i	35041	NG4-1	38/752=4.8%	46/785=5.7%	0.9
	31134	NG4-1	12/198=6%	31/221=14%	8.0
UAS-G$_\beta$76C-i	28507	NG4-1	14/219=6.4%	26/226=11.5%	5.1
UAS-G$_\gamma$1-i	25934	NG4-1	21/283=7.4%	45/311=14.4%	7.0
	34372	NG4-1	21/434=4.8%	46/400=11.5%	6.7
UAS-G$_\gamma$30A-i	25932	NG4-1	16/319=5.0%	18/286=6.3%	1.3
	34484	NG4-1	9/320=2.8%	31/323=9.6%	6.8
UAS-CaMKI-i	41900	NG4-1	17/337=5.0%	31/328=9.4%	4.4
	35362	NG4-1	10/222=4.5%	17/212=8.0%	3.5
	26726	NG4-1	16/301=5.3%	22/282=7.8%	2.5
UAS-CaMKII-i	35330	NG4-1	49/784=6.2%	91/858=10.6%	4.4
	29401	NG4-1	38/332=11.4%	43/342=12.6%	1.2
UAS-CrebA-i	42526	NG4-1	13/301=4.3%	29/298=9.7%	5.4

UAS-Gprk1-i	35246	NG4-1	13/323=4.0%	15/207=7.4%	3.4
	28354	NG4-1	13/304=4.3%	24/289=12%	7.7
UAS-Gprk2-i	41933	NG4-1	3/218=1.4%	24/228=10.5%	9.1
	35326	NG4-1	12/268=4.5%	35/267=13.1%	8.6
UAS-IP3K-i	35296	NG4-1	10/225=4.4%	14/152=9.2%	4.8
	31733	OR	25/336=7.4%	31/331=9.4%	2.0
	31733	NG4-1	67/572=11.7%	57/555=10.3%	-1.4
UAS-PKC53E-i	34716	NG4-1	18/304=5.9%	22/289=7.6%	1.7
UAS-PKC98E-i	29311	NG4-1	14/293=4.8%	15/284=5.3%	0.5
	35275	OR	9/266=3.4%	25/288=8.4%	5.0
		NG4-1	49/657=7.5%	45/603=7.5%	0.0
	44074	OR	30/597=5.1	36/346=10.4	5.3
		NG4-1	31/289=10.7	38/318=11.9	1.2
UAS-PLC2-i	33719	NG4-1	23/264=8.7%	37/311=11.9%	3.2
UAS-bsk-i	53310	NG4-1	21/380=5.5%	32/346=9.3%	3.8
UAS-Ira-i	31595	NG4-1	6/272=2.2%	8/190=4.2%	2.0
UAS-kay-i	27722	NG4-1	11/256=4.3%	25/259=9.6%	5.3
	31322	NG4-1	22/384=5.7%	61/334=18.3%	12.6
	31391	NG4-1	24/291=8.2%	29/218-13.3%	5.1
UAS-rl-i	36059	NG4-1	21/269=7.8%	32/297=10.8%	3.0
UAS-wt-5-HT1A	27630	NG4-1	16/335=4.8%	30/302=10%	5.2
	27631	NG4-1	10/240=4.2%	24/271=8.9%	4.7
UAS-wt-CrebB17A	7220	NG4-1	51/636=8.0%	98/628=15.6%	7.6
	9232	NG4-1	32/637=5.0%	84/592=14.2%	9.2
UAS-wt-CaMK2R3	29662	NG4-1	20/292=6.8%	28/289=10.0%	3.2
UAS- CaMKII.T287A/	29663	NG4-1	16/287=5.6%	25/283=8.8%	3.2
UAS-wt-Gαs	6489	NG4-1	32/289=11.1%	39/300=13.0%	1.9

	6489	NG4-1	40/610=6.6%	54/652=8.3%	1.7
UAS-wt-Ira	7216	NG4-1	15/342=4.4%	44/378=11.6%	7.2
UAS-wt-Kay	7213	NG4-1	24/341=7.4%	70/350=20.0%	12.6

Table 2. MI^{GSC} from select RNA-*i*-lines directed against GPCRs

UAS-driven expression of RNA-*i* for the listed GPCRs i via *NG4-1* did not block the

increase in MI^{GSC} in response to mating. BL #: Bloomington stock number, Single and

Mated: number of pHH3-positive GSCs/total number of GSCs = MI^{GSC}, Diff: MI^{GSC} of

mated males minus MI^{GSC} of non-mated males. Note the variability in MI^{GSC} among the

different genotypes. GPCRs marked by asterisks were excluded from further studies

because their siblings outcrossed to *wt* did not show a stronger response to mating than

the experimental (GPCR-*i*/*NG4-1*) flies.

GPCR	BL #	Single	Mated	Diff.
UAS-5-HT2A-i	31882	25/490=5.1	36/465=7.3	2.2
	56870	19/582=3.3	32/553=5.8	2.5
UAS-5-HT2B-i	60488	6/261=2.3	21/272=7.7	5.4
	25874	4/228=1.7	34/236=14.4	12.7
UAS-Ado-R-i	27536	11/276=4.0	20/209=9.6	5.6
UAS-AKHR-i	29577	23/492=4.7	54/574=9.4	4.7
UAS-AR-2-i	25935	13/363=3.6	25/336=7.4	3.6
UAS-CG13229-i	29519	9/285=3.2	33/297= 11.1	7.9
UAS-CG14539-i	25855	21/318=6.6	31/307=10.1	3.5

UAS-CG15556-i	44574	28/425=6.6	43/401=10.7	4.1
UAS-CG15744-i	28516	18/279=6.4	27/252=10.7	4.3
	42497	23/323=7.1	36/236=11.0	3.9
UAS-CG30106-i	27669	10/244=4.1	34/273=12.4	8.3
UAS-CG33639-i	28614	32/300=10.7	47/371=12.7	2.0
*UAS-CCHaR1-i**	51168	27/407=6.6	27/323=8.4	1.8
UAS-Cry-i	43217	43/389=11.0	75/521=14.4	3.4
UAS-CrzR-i	52751	14/337=4.2	31/333=9.3	5.1
UAS-Dop1R1-i	62193	12/352=3.4	25/308=8.1	4.6
	55239	11/300=3.7	44/267=16.5	12.8
UAS-GABA BR2-i	50608	6/176=3.4	35/207=16.9	13.5
	27699	7/291=2.4	18/282=6.4	4.0
UAS-GABA BR3-i	42725	10/190=5.3	28/243=11.5	6.2
UAS-Moody-i	36821	5/301=1.7	16/234=6.8	5.1
UAS-Mth-l1-i	41930	11/279=4.0	42/300=14.0	10
UAS-Mth-l3-i	41877	54/817=6.6	81/850=11.8	5.2
	36822	19/231=8.2	39/241=16.2	8.0
UAS-Mth-l8-i	36886	48/933=5.1	70/915 =7.6	2.5
UAS-Mth-l9-i	51695	61/985=6.2	82/910 =9.0	2.8
UAS-Mth-l15-i	28017	14/349=4.0	25/337=7.1	3.1
UAS-PK1R-i	27539	28/478=5.9	37/375=9.9	4.0
UAS-Smo-i	27037	12/263=4.6	33/288=11.4	6.8
	43134	19/274=7.0	15/147=10.2	3.2
UAS-Tre1-i	34956	5/234=2.1	28/252=11.1	9.0
*UAS-TKR86D-i**	31884	31/564=5.5	28/394=7.1	1.6

UAS-TKR99D-i	55732	30/506=5.9	49/467=10.5	4.6
	27513	4/240=1.7	30/294=10.2	8.5

Table 3. MI^{GSC} from additional RNA*i*-lines with modified expression of the GPCRs

blocking the increase in MI^{GSC} in mated males

BL #: Bloomington stock number, Single and Mated: number of pHH3-positive

GSCs/total number of GSCs = MI^{GSC}, Diff: MI^{GSC} of mated males minus MI^{GSC} of non-

mated males.

GPCR	BL #	Single	Mated	Diff.
5HT-1A-i/NG4-1	25834	64/841=7.6	67/777=8.6	1
5HT-1B-i/NG4-1	25833	16/256=6.2	26/268=9.7	3.5
	27635	3/304=1.0	27/317=8.5	7.5
	51842	20/381=5.2	32/375-8.5	3.3
	54006	13/405=3.2	23/351=6.5	3.3
5HT-7-i/NG4-1	32471	8/238=3.4	17/229=7.4	4.0
CG12290-i/NG4-1	42520	4/260=1.5	31/246=12.6	11.1
Mth-i/NG4-1	27495	14/352=4	15/336=4.5	0.5

Figure legends

Figure 1. Mating increased male stem cell division frequency.

A) Cartoon depicting the stages of *Drosophila* spermatogenesis. Note that every GSC

division produces exactly 64 spermatids. GB: gonialblast, SG: spermatogonia, SC:

spermatocytes, SP: spermatids.

A') The apical tip of a *wt* testis. The FasIII-positive hub (asterix) is surrounded by seven

Vasa-positive GSCs (green), one of which is in mitosis based on anti-pHH3-staining

(arrowhead). Scale bar: 10µm.

B-F) Blue: non-mated condition, red: mated condition, ***: P-value < 0.001, numbers of

GSCs and number of gonads (n=) as indicated.

B) Box plots showing the range of MI^{GSC}. Lines within boxes represent medians,

whiskers represent outliers.

C, D) FDGs showing bin of MI^{GSC} (bin width=10) across a population of c) *OR* and D)

CS males on the X-axis and the percentage of testes with each MI^{GSC} on the Y-axis.

E) Bar graph showing SI^{GSC} of *OR* males from three independent experiments.

F) Graph showing the percentage of EdU-marked *OR* GSCs on the Y-axis and hours of

feeding and mating on the X-axis.

Figure 2. Mating reduced the mature sperm pool.

A-C) Class 1,2 and 3 seminal vesicles from Mst35B-GFP males. Scale bars: 0.1 mm;

arrows point to GFP-negative regions.

D-G) Numbers of seminal vesicles (n=) as indicated, n-m: non-mated, m: mated, ***: P-

value < 0.001.

D-F) Bar graphs showing the distribution of Class 1 to 3 seminal vesicles in non-mated

and mated males at days one to three of the mating experiment. Three fly lines that

carry GFP-marked sperm were used: one that carries Dj-GFP (BL#5417), one that

carries the Mst35B-GFP (BL#58408), and one line that carries both constructs

(BL#58406).

G) Box plot showing sperm head volume (based on MST35B-GFP) per seminal vesicle

in non-mated and mated males on days one to three of the experiment.

Figure 3. Mating did not affect GSC numbers.

A-E) Blue: non-mated condition, red: mated condition, numbers of gonads (n=) as

indicated, genotypes as indicated.

A-E) FDGs showing numbers of GSCs on the X-axis and percentage of testes with the

number of GSCs on the Y-axis. No difference in GSC numbers was observed between

non-mated and mated males from different genetic backgrounds.

Figure 4. G-proteins were required for the increase in MI^{GSC} in response to

mating.

A) Cartoon depicting the activation of G-proteins upon GPCR stimulation by ligand. 1:

G-protein association before GPCR stimulation, 2: G-protein distribution after GPCR

stimulation, 3a-c: downstream signaling cascades. AC: Adenylyl Cyclase, cAMP: cyclic

Adenosine Monophosphate, PKA: Protein Kinase A, CREB: cAMP responsive element-

binding protein, PLC: Phospho Lipase C, DAG: Diacylglycerol, PKC: Protein Kinase C,

MAPK: Map Kinase, IP3: Inositol Triphosphate, CaMK: Calcium$^{2+/}$calmodulin-dependent

protein kinase.

B, C) Bar graphs showing MI^{GSC}. Blue: non-mated condition, red: mated condition, ***:

P-value < 0.001, numbers of GSCs as indicated, genotypes as indicated.

B) Control animals increased MI^{GSC} in response to mating.

C) Males expressing *G$_\alpha$i-i* or dnG$_\gamma$1 in the germline did not increase MI^{GSC} after mating.

Figure 5. Expression of RNA-*i* against seven distinct GPCRs blocked the increase

in MI^{GSC} in response to mating.

A-C) Bar graphs showing MI^{GSC}. Blue: non-mated condition, red: mated condition, ***:

P-value < 0.001, numbers of GSCs as indicated, genotypes as indicated.

A) Control males have significantly higher MI^{GSC} than their non-mated siblings.

B, C) Mated (B) *GPCR-i*/NG4-1 and (C) *GPCR-i*/NG4-2 males did not have significantly

higher MI^{GSC} compared to their non-mated siblings.