Factors Affecting Electroconvulsive Therapy Ictal Outcomes:
Duration and Postictal Suppression

Wendy Marie Ingram, PhD*[1, 2], Cody Weston, MD, PhD [2], Wei Dar Lu, MD[1], Caleb Hodge, MD[1], S. Mark Poler, MD[1], Fatin Nahi, MD[1], Sharon Larson, PhD[1, 3]

[1]Geisinger Health System, Danville, Pennsylvania; [2]Johns Hopkins University, Baltimore, Maryland; [3]Lankenau Institute for Medical Research, Wynnewood, Pennsylvania
* Corresponding author: E-mail address – wingram5@jhu.edu

Abstract

Electroconvulsive therapy (ECT) is an effective and rapid treatment for severe depression, however predictors of therapeutic outcomes remain insufficiently understood. Ictal duration and postictal suppression are two outcomes that may be correlated with patient response, yet patient and treatment variables which may influence these outcomes have not been thoroughly explored. We collected ECT stimulus metrics, EEG parameters, patient demographics, primary diagnosis, and anesthesia type for retrospective ECTs. Univariate and multivariate mixed-effects linear regression models were used to identify variables associated with ictal duration and postictal suppression index. For both outcomes, multivariate models which included all variables resulted in the best fit, reflecting the complex influences of a variety of factors on the ictal response. These results are an important step forward in elucidating patterns in retrospective ECT clinical data which may lead to new clinical knowledge of modifiable factors to optimize ECT treatment outcomes.

Key words: Secondary use of EHR data; Precision medicine; Outcomes research, clinical epidemiology, population health

Introduction

Electroconvulsive Therapy, or ECT, is one of the most effective and safe treatments available to psychiatrists for treating severe mood disorders and psychosis. ECT emerged as a therapy due to the observation that epilepsy seemed to be rare in people with schizophrenia. Seizures were originally induced using chemical methods including camphor injection. Later, in 1938, it was found that electrical current could more safely and reliably induce seizures, resulting in improvements in patient symptoms. However, the mechanism of how a controlled seizure leads to reduced depressive symptoms remains poorly understood(1). Despite popular perception that ECT is antiquated, it can be superior to medication and serves as a vital hope for remission in treatment-resistant cases of depression, bipolar disorder, suicidality and catatonia(2,3).

Unlike pharmaceutical interventions, ECT requires that practitioners consistently generate a "therapeutic seizure" under continually varying circumstances. For example, a given patient's seizure threshold is influenced by their age, sex, hydration status, the number of recent seizures that they have had, as well as intrinsic unobservable factors. In turn, the ECT metrics that are involved in stimulating a seizure are numerous. These include percent energy set, stimulus duration, frequency, pulse width, and current. Thus, determining what constitutes "dosage" is not immediately clear in ECT(4). Consequently, there is considerable variability in the practice of ECT in the United States and around the world.

Optimizing for clinical outcomes requires consideration of the many factors involved in inducing an "adequate" seizure. Early ECT optimization work focused on the prediction of ictal duration, or the

duration of the induced seizure(5,6). More recently it has been determined that ictal duration does not reliably predict clinical outcome, so long as it is between 25 and 180 seconds(7,8). Many clinicians continue to optimize for this ictal duration window in their practice today. More recent evidence, however, points to postictal suppression, or the extent to which EEG activity is diminished after the seizure terminates, as being the best predictor of therapeutic outcome(9–12). As mentioned above, there are many variables that can affect ictal properties and, by extension, clinical outcomes. The identification of individual or combinations of parameters which predict ictal duration and postictal suppression could feasibly improve our ability to reliably produce optimal seizures and thus patient outcomes.

Here we utilize univariate and multivariate linear regression mixed-effects models to determine which patient, ECT, and EEG parameters predict ictal duration and postictal suppression in a retrospective study.

Methods

ECT Data. All ECT sessions were performed in outpatient surgery at Geisinger Medical Center, located in Danville, Pennsylvania between June 1st, 2013 and May 31st, 2017. Patients were included in the study retrospectively if they were 18 years or older, had a primary diagnosis of depression (ICD-9 codes: 296.2x, 296.3x, 311; ICD-10 codes: F32.x, F33.x, F43l.x) or bipolar disorder (ICD-9 codes: 296.4x, 296.5x, 296.6x, 296.8x; ICD-10 codes: F30.x, F31.x), and had received an acute series of ECT sessions. An acute series is defined as 6 or more sequential ECTs, each no more than 3 days apart. ECTs were performed using a Thymatron device (Somatics, LLC. Lake Bluff, IL, USA). Part way through the study period, the machine was changed from a Thymatron DGx with a pulse width of 1.0 mSec to a Thymatron IV with a pulse width of 0.5 mSec. Machine type and pulse width are perfectly correlated variables, thus we included only pulse width in the analysis. ECTs were performed in the mornings on Mondays, Wednesdays, and Fridays. Anesthesia and analgesic agents were chosen based on physician experience and preference and included methohexital, propofol, and ketamine which were administered either alone or in combination. For each seizure, the input percent energy set, current, stimulus duration, and frequency were recorded. The ictal outcomes of duration and postictal suppression index were measured and summarized by the Thymatron machines. The procedure start time, the sex and age of the patient, and the number of ECTs in the acute series were also collected from the electronic health record. Bilateral ECT initial dose was based on patient sex (females: 10% energy, 50 mcoul charge; males: 15% energy, 76 mcoul charge) and increased if patients did not respond clinically or if seizures were insufficient during the ECT course (i.e. ictal duration <25 Sec).

Statistical analysis. Univariate and multivariate mixed-effects linear regression analysis were performed in R (2017, R Core Team, Vienna, Austria) using package lme4(13). Visualizations used GraphPad Prism 6 (La Jolla, CA). Patient-specific variability was modeled as a random variable allowing for random patient intercept in all models. Coefficients and restricted maximum likelihood (REML) values for each model are reported and p-values were calculated from associated t-values.

Results

We evaluated data from 84 patients which underwent a total of 724 ECT sessions. Patient demographics, pulse width, and anesthesia numbers and percentages are shown in Table 1 for patients and ECTs. All patients were non-Hispanic white. Females were slightly more common than males, and most patients had a primary diagnosis of depression. A third of ECTs were performed on the second machine, and two thirds of patients received methohexital at least once during an ECT session.

Table 1. Demographic Data and ECT parameters. Number and percentages provided. Anesthesia groups are not mutually exclusive.

	Patients	**%**	**ECTs**	**%**
	N=84	-	N=724	-
Males	36	43	295	41
Females	48	57	429	59
Bipolar	16	19	116	16
Depression	68	81	608	84
Pulse width 1.0	59	70	476	66
Pulse width 0.5	25	30	248	34
Methohexital	66	79	555	77
Propofol	28	33	172	24
Ketamine	12	14	37	5

Ictal duration. Univariate mixed-effects linear regression coefficients and p-values are presented in Table 2. Being administered methohexital was significantly associated with a longer ictal duration while propofol was associated with a shorter duration. The number of ECTs a patient had received in a series, the percent energy, stimulus duration, and frequency were all significantly associated with very minor decreases in ictal duration. The fitness of each mixed-effects model, as estimated by the restricted maximum likelihood (REML) (Figure 1), only changes modestly between each univariate model. Three of the six statistically significant univariate models had the lowest REML values amongst them, indicating the best fit. When these six significant parameters along with patient age and sex are included in a selective multivariate model, the REML value of the model is dramatically reduced, indicating improved fit. Stimulus duration and frequency remain statistically significant, while the other four variables do not. When including all variables in a complete multivariate model, the REML value decreases further, but with only the number of ECTs and the frequency remaining statistically significant predictors of ictal duration.

Postictal suppression index. Univariate models for postictal suppression index result in pulse width, ketamine administration, and stimulus duration being positively predictive of higher values and statistically significant. These are presented in Table 2. The hour of day during which the ECT is performed is statistically significant and associated with a negative effect on postictal suppression. When these four variables are included with age and sex in a selective multivariate model, only ketamine remains statistically significant with a large positive effect. By including all variables in a complete multivariate model, six variables become statistically significant including those that were significant as univariates (pulse width, ketamine, and stimulus duration) and those that were not (number of ECTs, percent energy, and frequency). Postictal suppression index model REML values were similar in trend to those of ictal duration (Figure 1).

Discussion

Here we present and compare univariate and multivariate linear mixed-effects models predicting two critical outcome measures of ECT. Ictal duration has been studied extensively due to its previously believed association with therapeutic outcomes and known association with cognitive and other adverse effects. Therefore, its modifiable input parameters have been relatively well studied. Univariate modeling identified expected independent variables predictive of ictal duration. Anesthesia type affected

Table 2. Mixed-effects model coefficients and p-values for univariate and multivariate linear regressions of ictal duration and postictal suppression index. Significance: $p<0.05*$, $p<0.01**$, $p<0.001***$. Univar coef = Univariate mixed model coefficients; Selective Multivar coef = multivariate model coefficients including age, sex, and significant univariate variables; Complete Multivar coef = multivariate model coefficients including all univariate model variables.

Independent variables	Factors	Univar coef	p-value	Selective Multivar coef	p-value	Complete Multivar coef	p-value
			Ictal Duration				
Sex	Female (ref)	-	-	-	-	-	-
	Male	-1.70	0.63	0.95	0.77	0.65	0.84
Age (years)		-0.20	0.11	-0.17	0.14	-0.17	0.15
Primary dx code	Bipolar (ref)	-	-	-	-	-	-
	Depression	2.32	0.61	-	-	4.38	0.28
Pulse-width (mSec)	1.0 (ref)	-	-	-	-	-	-
	0.5	-2.99	0.34	-	-	-2.28	0.80
Ketamine		-3.06	0.51	-	-	3.21	0.58
Methohexital		8.66	0.008**	9.05	0.16	10.27	0.13
Propofol		-7.14	0.02**	-2.67	0.68	-1.66	0.80
Hour of ECT		-0.05	0.94	-	-	0.79	0.33
Number of ECTs		-0.73	<0.001***	-0.37	0.06	-0.39	0.05*
Percent energy set (%)		-0.24	<0.001***	0.08	0.53	0.08	0.59
Current (A)		65.35	0.46	-	-	-89.87	0.35
Stimulus duration (Sec)		-1.40	0.02**	-1.94	0.04*	-1.87	0.40
Frequency (Hz)		-0.35	<0.001***	-0.36	0.04*	-0.39	0.03*
			Postictal Suppression Index				
Sex	Female (ref)	-	-	-	-	-	-
	Male	0.83	0.77	1.08	0.71	0.57	0.06
Age (years)		-0.07	0.48	-0.11	0.28	-0.13	0.18
Primary dx code	Bipolar (ref)	-	-	-	-	-	-
	Depression	6.23	0.09	5.82	0.11	6.62	0.06
Pulse-width (mSec)	1.0 (ref)	-	-	-	-	-	-
	0.5	6.84	0.01**	-2.71	0.69	-19.55	0.04*
Ketamine		15.18	0.001***	11.17	0.02*	11.64	0.02*
Methohexital		2.52	0.40	-	-	0.94	0.89
Propofol		-3.15	0.28	-	-	-0.13	0.98
Hour of ECT		-1.83	0.03*	-1.45	0.09	-1.60	0.06
Number of ECTs		-0.15	0.39	-	-	-0.40	0.05*
Percent energy set (%)		0.02	0.63	-	-	-0.43	0.008**
Current (A)		7.28	0.93	-	-	104.85	0.31
Stimulus duration (Sec)		1.65	0.003**	1.70	0.23	7.54	0.002**
Frequency (Hz)		-0.01	0.91	-	-	0.50	0.01**

ictal duration, which is consistent with previous studies that show that methohexital is associated with longer seizure time(14–16). However, we found that the multivariate model that included all independent variables modeled the data best. In this model, ECT number and frequency were significant predictors of ictal duration. Frequency is not surprising because it is one of the principle components that can be varied to increase "dosage" and overcome seizure threshold(17,18). ECT number as a predictor of ictal duration is also consistent with existing literature, as it is well-known that seizure threshold increases with each subsequent ECT administration in an acute series(19). It will be of interest to explore variable interactions in future mixed-effects modeling and explore the structure of the random effects in this data.

Figure 1. Restricted Maximum Likelihood (REML) of univariate and multivariate mixed-effects models for ictal duration (A) and postictal suppression (B), rank ordered. REML is the unbiased estimate of variance and covariance parameters for mixed-effects models, with lower REML relating to better fit of the data.

Multivariate modeling of postictal suppression indicates that pulse width, ketamine use, stimulus duration, ECT number, percent energy, and frequency are all significantly predictive of postictal suppression. With the current understanding that postictal suppression is a good predictor of clinical outcome, recognition of these factors has potential to improve practice. The detection of anesthesia choice in this study is consistent with other preliminary studies(15,16). This is useful information as the standard of care in anesthesia and analgesia for ECT continues to evolve with our understanding of its interplay with clinical outcomes. Previous studies have demonstrated that ketamine may be associated with improved outcomes in the early ECT course(20). The findings of the previous study are provocative since the field currently recommends methohexital as the standard anesthesia method and has not yet produced consistent evidence to recommend ketamine as a primary or adjunct agent(20). Interestingly, ketamine infusion is emerging as a promising standalone antidepressant treatment(21), so perhaps ketamine and ECT are interacting in a combinatorial way via postictal suppression.

Perhaps the most surprising finding in the present study is that time of day significantly negatively affects postictal suppression when modeled in a univariate model. While this variable loses significance in the multivariate model, the size of the effect is sizeable and has a near significant p-value of 0.06. Others have recently found that the timing of anesthesia and time of ECT delivery may affect seizure quality, but did not look at postictal suppression specifically(22). This effect may be due to factors including length of time fasting, time since last medication administration, or circadian effects. While the underlying

reasons remain unclear and warrant future studies, the recognition of the effect alone might merit the adjustment of patient procedure timing in particularly difficult cases. The key caveat is that it is not clear that time of day was random. It is possible that cases were already stratified according to clinical severity, inpatient/outpatient status, or some other clinical history feature without this information being recorded.

There are several limitations of this study. The study population was ethnically homogenous and based out of a single center located in rural Pennsylvania. This may limit generalizability to other parts of the country and the world. By performing a retrospective study, many factors are not able to be controlled for and confounders could exist due to a lack of randomization. In addition, the data recorded by the ECT machine is generally considered to not be as reliable as expert clinical review of the EEG trace. To estimate the extent of this, future studies may benefit from clinical review of a random sampling to estimate machine/clinician agreement. Another key caveat to be mindful of with clinical interpretation is that the pulse width is correlated perfectly with a change in machine. Thus, it is possible that idiosyncrasies related to the individual machines are being misinterpreted as attributable to pulse width. Future studies at multiple sites as well as larger sample sizes will help to address some of these concerns.

There are also many strengths of this study. Retrospective studies such as this examine ECT sessions that were performed under clinical conditions on a clinical population, rather than in an artificial setting with selection bias of enrolled participants. The findings presented here are thus immediately relevant to the clinical practice at the investigation site. The unbiased nature of retrospective observational studies can be powerful for hypothesis generation. This repurposing of clinical data is low cost in comparison to randomized controlled trials, thus, there is reasonable cause to thoroughly investigate trends and identify possible predictive factors in existing clinical data prior to conducting prospective studies. Finally, use of mixed-effects linear regression modeling of many variables both in univariate and multivariate analyses ultimately allows us to begin to unravel the complex interactions and effects that multiple factors have on the ictal outcomes of duration and postictal suppression.

Conclusions

ECT is just one example of a medical procedure with a multitude of patient, clinician, and treatment mediated variables. For decades, many adjustments of ECT parameters have been driven largely by clinical judgment and preference. In the current era of information technology, it is possible to use historical data to model outcomes of interest and improve care guidelines without having to prospectively randomize patients. The approach demonstrated here could be applied to procedures in many medical domains and represents one high-yield area for informatics techniques to directly improve the health of communities and improve patient centered care.

Acknowledgements

We would like to thank Jason Brown from the Geisinger Phenomics and Clinical Data Analytics Core for his tireless work refining the data pull, Sarah Robishaw for her help with chart review, and Christopher Bauer, PhD for his excellent comments and suggestions on the manuscript.

References

1.   Fosse R, Read J. Electroconvulsive treatment: Hypotheses about mechanisms of action. Front Psychiatry. 2013;4(AUG):1–10.

2.   Ross EL, Zivin K, Maixner DF. Cost-effectiveness of Electroconvulsive Therapy vs Pharmacotherapy/Psychotherapy for Treatment-Resistant Depression in the United States. JAMA

psychiatry [Internet]. 2018 Jul 1;75(7):713–22. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29800956

3. Schoeyen HK, Kessler U, Andreassen OA, Auestad BH, Bergsholm P, Malt UF, et al. Treatment-resistant bipolar depression: A randomized controlled trial of electroconvulsive therapy versus algorithm-based pharmacological treatment. Am J Psychiatry. 2015;172(1):41–51.

4. Peterchev A V., Rosa MA, Deng Z-D, Prudic J, Lisanby SH. Electroconvulsive therapy stimulus parameters: rethinking dosage. J ECT [Internet]. 2010 Sep;26(3):159–74. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20805726

5. Sackeim HA, Decina P, Kanzler M, Kerr B, Malitz S. Effects of electrode placement on the efficacy of titrated, low-dose ECT. Am J Psychiatry. 1987;144(11):1449–55.

6. Miller AL, Faber RA, Hatch JP, Alexander HE. Factors affecting amnesia, seizure duration, and efficacy in ECT. Am J Psychiatry. 1985;142(6):692–6.

7. Folkerts H. The ictal electroencephalogram as a marker for the efficacy of electroconvulsive therapy. Eur Arch Psychiatry Clin Neurosci [Internet]. 1996;246(3):155–64. Available from: http://www.ncbi.nlm.nih.gov/pubmed/8739401

8. Rasimas JJ, Stevens SR, Rasmussen KG. Seizure length in electroconvulsive therapy as a function of age, sex, and treatment number. J ECT [Internet]. 2007 Mar;23(1):14–6. Available from: http://www.ncbi.nlm.nih.gov/pubmed/17435566

9. Sackeim HA, Devanand DP, Prudic J. Stimulus intensity, seizure threshold, and seizure duration: impact on the efficacy and safety of electroconvulsive therapy. Psychiatr Clin North Am [Internet]. 1991 Dec;14(4):803–43. Available from: http://www.ncbi.nlm.nih.gov/pubmed/1771150

10. Suppes T, Webb A, Carmody T, Gordon E, Gutierrez-Esteinou R, Hudson JI, et al. Is postictal electrical silence a predictor of response to electroconvulsive therapy? J Affect Disord [Internet]. 1996 Nov 4;41(1):55–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/8938205

11. Azuma H, Fujita A, Sato K, Arahata K, Otsuki K, Hori M, et al. Postictal suppression correlates with therapeutic efficacy for depression in bilateral sine and pulse wave electroconvulsive therapy. Psychiatry Clin Neurosci [Internet]. 2007 Apr;61(2):168–73. Available from: http://doi.wiley.com/10.1111/j.1440-1819.2007.01632.x

12. Gangadhar BN, Subbakrishna DK, Janakiramaiah N, Motreja S, Narayana Dutt D, Paramehwara G. Post-seizure EEG fractal dimension of first ECT predicts antidepressant response at two weeks. J Affect Disord. 1999;52(1–3):235–8.

13. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. J Stat Softw [Internet]. 2015;67(1). Available from: http://arxiv.org/abs/1406.5823

14. Andersen FA, Arsland D, Holst-Larsen H. Effects of combined methohexitone-remifentanil anaesthesia in electroconvulsive therapy. Acta Anaesthesiol Scand [Internet]. 2001 Aug;45(7):830–3. Available from: http://www.ncbi.nlm.nih.gov/pubmed/11472282

15. Vaidya P V, Anderson EL, Bobb A, Pulia K, Jayaram G, Reti I. A within-subject comparison of propofol and methohexital anesthesia for electroconvulsive therapy. J ECT [Internet]. 2012 Mar;28(1):14–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22330701

16. Wojdacz R, Święcicki Ł, Antosik-Wójcińska A. Comparison of the effect of intravenous anesthetics used for anesthesia during electroconvulsive therapy on the hemodynamic safety and

the course of ECT. Psychiatr Pol [Internet]. 2017 Dec 30;51(6):1039–58. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29432502

17. Grunhaus L, Zelnik T, Albala AA, Rabin D, Haskett RF, Zis AP, et al. Serial dexamethasone suppression tests in depressed patients treated only with electroconvulsive therapy. J Affect Disord [Internet]. 13(3):233–40. Available from: http://www.ncbi.nlm.nih.gov/pubmed/2960715

18. Pande AC, Krugler T, Haskett RF, Greden JF, Grunhaus LJ. Predictors of response to electroconvulsive therapy in major depressive disorder. Biol Psychiatry [Internet]. 1988 May;24(1):91–3. Available from: http://www.ncbi.nlm.nih.gov/pubmed/3370280

19. Scott AI, Boddy H. The effect of repeated bilateral electroconvulsive therapy on seizure threshold. J ECT [Internet]. 2000 Sep;16(3):244–51. Available from: http://www.ncbi.nlm.nih.gov/pubmed/11005045

20. Gálvez V, McGuirk L, Loo CK. The use of ketamine in ECT anaesthesia: A systematic review and critical commentary on efficacy, cognitive, safety and seizure outcomes. World J Biol Psychiatry [Internet]. 2017;18(6):424–44. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27892759

21. Fond G, Loundou A, Rabu C, Macgregor A, Lançon C, Brittner M, et al. Ketamine administration in depressive disorders: a systematic review and meta-analysis. Psychopharmacology (Berl) [Internet]. 2014 Sep;231(18):3663–76. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25038867

22. Gálvez V, Hadzi-Pavlovic D, Wark H, Harper S, Leyden J, Loo CK. The Anaesthetic-ECT Time Interval in Electroconvulsive Therapy Practice--Is It Time to Time? Brain Stimul [Internet]. 9(1):72–7. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26452698