1 Classification: Biological Sciences: Microbiology

3 Hfq-assisted RsmA regulation is central to *Pseudomonas*

4 *aeruginosa* biofilm and motility

Yasuhiko Irie[1,2]*, Jessica E. Geyer[1] and Victoria Shingler[2]

[1] Department of Biology, University of Dayton, Dayton, OH 45469, USA

[2] Department of Molecular Biology, Umeå University, Umeå, 901 87, Sweden

Short Title: *Pseudomonas aeruginosa* RsmA regulates Vfr

Keywords: *Pseudomonas aeruginosa*, RsmA, Vfr, FleQ, Hfq

*Corresponding author, current mailing address: Department of Biology, University of

Dayton, Science Center 234, 300 College Park, Dayton, OH 45469-2320, USA. Tel: +1-

937-229-3061. E-mail: yirie1@udayton.edu

ORCID ID for Yasuhiko Irie: 0000-0002-7162-4772

18   ABSTRACT

Expression of biofilm and motility genes is controlled by multiple regulatory elements,

allowing bacteria to appropriately adopt a sessile or motile lifestyle. In *Pseudomonas*

aeruginosa, the post-transcriptional regulator RsmA has been implicated in the control of

various biofilm- and motility-associated genes, but much of the evidence for these links is

limited to transcriptomic and phenotypic studies. RsmA binds to target mRNAs to

modulate translation by affecting ribosomal access and/or mRNA stability. Here we trace a

global regulatory role of RsmA to the inhibition of Vfr – a transcription factor required for

efficient production of two other transcriptional regulators, namely FleQ and AlgR. FleQ

and AlgR, in turn, directly control flagella and pili genes, respectively. FleQ also directly

controls biofilm-associated genes that encode the PEL polysaccharide biosynthesis

machinery. Furthermore, we show that RsmA cannot bind *vfr* mRNA alone, but requires

the RNA chaperone protein Hfq. This is the first example where a RsmA protein family

member is demonstrated to require another protein for RNA binding.

33   SIGNIFICANCE STATEMENT

Microorganisms are subjected to dynamic changes in their environment and rapidly adapt

their gene expressions accordingly. Such environmental cues influence planktonic/biofilm

and acute/chronic infection modes of growth. In the opportunistic pathogen *Pseudomonas*

aeruginosa, RsmA facilitates this switch by controlling gene expression on the RNA level.

In this study, we present evidence that RsmA lies central to several downstream regulatory

cascades that govern biofilm formation, flagellar motility, and Type IV pilus-mediated

motility – all previously described as part of the RsmA regulon. Importantly, we show that

RsmA-binding to the *vfr* mRNA requires Hfq. These findings have implications for the

global regulatory role of RsmA and necessitates a genetic and biochemical re-evaluation of

what is currently thought to be the RsmA regulon.

 INTRODUCTION

The opportunistic pathogen *Pseudomonas aeruginosa* can be isolated from a wide range of

environmental niches, in large part owing to its versatile metabolic capabilities. It is also

proficient in colonizing various eukaryotic organisms and can cause both acute and chronic

infections, with the latter often associated with biofilm-like modes of growth at the sites of

infection (1). *P. aeruginosa* is also highly competitive against other microbial species, with

its adaptability and competitive fitness being aided by the production of various secondary

metabolites and virulence factors (2, 3). Many of these cellular processes are inter-regulated

by multiple regulatory pathways, presumably to provide appropriate response mechanisms

to a wide range of environmental cues. One of the central regulators that determine *P.*

aeruginosa behavior is the post-transcription factor RsmA (<u>R</u>egulator of <u>S</u>econdary

<u>M</u>etabolite) as evidenced by its null mutant phenotypes linked to motility (4), virulence (5-

7), biofilm (5, 8), growth (8), and secreted products (4, 9).

RsmA belongs to the RsmA/CsrA family of dimeric RNA-binding proteins (10-12),

homologues of which are found across a wide range of both Gram-negative and Gram-

positive bacterial species (13). RsmA/CsrA proteins generally function as translation

inhibitors by binding to the ribosome binding sites of target mRNAs, typically overlapping

either the Shine-Dalgarno or the start codon (14). However, in a few cases described, they

can also serve as positive regulators by altering the secondary structures of bound RNAs

(15, 16) or protecting RNAs from degradation (17).

Many RsmA/CsrA-targeted mRNAs are reported to have altered RNA decay rates. Several

studies attempted to exploit this property to determine the global *P. aeruginosa* RsmA

regulon using microarrays (18-20). However, this approach cannot distinguish directly

regulated genes from indirectly regulated gene sets that result from cascade regulation, nor

can it identify target RNAs that do not exhibit changes in their stability (21, 22). For

example, direct RsmA-mediated translation inhibition has been demonstrated for the *P.*

aeruginosa biofilm polysaccharide operon *psl* without any effect on the level of

transcription (8). Hence, the *psl* operon was not prominently featured in transcriptomic

studies despite a strong biofilm phenotype (18, 19), raising concerns of what is currently

considered to be 'the RsmA regulon' of *P. aeruginosa*.

Here, we present evidence that RsmA serves a master regulatory role through several

intermediate transcription factors that are known to control different cellular processes.

Through cascade regulation, originating from a direct effect on *vfr* mRNA, RsmA indirectly

affects both bacterial motility and biofilm formation. Furthermore, we discovered an

unexpected regulatory interplay between RsmA and the RNA chaperone protein Hfq on the

target mRNA, whereby RsmA could only bind in the presence of Hfq *in vitro*. To our

knowledge, this is the first example where RsmA/CsrA is unable to bind to its target alone.

Because Hfq is known to facilitate RNA degradation (23), these results may provide a

mechanistic insight into RsmA-associated mRNA stability changes.

 RESULTS

RsmA is a post-transcriptional regulator of *vfr* through which *fleQ* and *pel* transcripts are indirectly regulated.

P. aeruginosa produces at least three different extracellular biofilm polysaccharides: alginate, PEL, and PSL (24). PEL and PSL are co-regulated by several factors including the secondary messenger molecule c-di-GMP via the c-di-GMP-binding transcriptional regulator FleQ (25) and possibly also quorum sensing (26, 27). Similar to PSL, PEL has long been considered to be regulated by RsmA (28) based on a model originally proposed from a microarray study (29). However, unlike the *psl* operon transcript, which was genetically and biochemically demonstrated to be directly regulated by RsmA (8), no direct evidence for the *pel* operon has been presented.

Consistent with a previous microarray study (18), using a chromosomal transcriptional reporter, we found that *pel* was up-regulated in the Δ*rsmA* mutant as compared to the wild type (WT) background (Fig. S1A). However, the PSL polysaccharides are also over-expressed in a Δ*rsmA* strain (8), and as a direct consequence of this over-overexpression, intracellular levels of c-di-GMP are elevated (30). Given that the *pel* genes are known to be transcriptionally up-regulated by c-di-GMP (25, 31) and Δ*rsmA* has elevated c-di-GMP, it appeared plausible that the up-regulation of *pel* could be caused by changes in c-di-GMP levels. To uncouple *pel* expression from elevated c-di-GMP, we also measured *pel* transcriptional reporter activities when introduced into a Δ*rsmA* Δ*pel* Δ*psl* triple mutant background, which has low c-di-GMP (30). In this strain, *pel* was still up-regulated (Fig. S1A), indicating that the up-regulation was caused by lack of RsmA rather than changes in intracellular c-di-GMP concentration.

Genes that are directly regulated by RsmA/CsrA often manifest differences that are experimentally observed at the transcriptional reporter level, since bound RNAs frequently have altered half-lives (14). Therefore, we next determined whether the apparent *pel*

transcriptional changes between the WT and Δ*rsmA* strains were actually a consequence of

post-transcriptional RNA stability changes. However, quantification of *pelA* transcripts at

progressive time points after blocking *de novo* transcription revealed no differences in

stability between the WT and Δ*rsmA* strains (Fig. S1B). These results suggest that the

observed differences in *pel* expression likely occur at the promoter activity level. Because

the RNA-binding protein RsmA is not a transcription factor, we next turned our attention to

the possibility of RsmA affecting expression of a known transcriptional regulator of *pel*,

namely FleQ. As a well-characterized transcriptional activator of *pel* (25, 31), potential

RsmA-mediated alterations in expression of FleQ would be anticipated to affect

transcription of the *pel* genes. Using a similar approach as described for *pel* above, we

found that a transcriptional reporter fusion of *fleQ* gave lower levels in the absence of

RsmA (Fig. S2A). This is consistent with previous data showing *pel* transcript levels are

up-regulated in Δ*fleQ* backgrounds (25).

The data above could lend themselves to the interpretation that RsmA serves as a positive

regulator of *fleQ*. While RsmA is more commonly known to be a translational repressor,

this possibility is not without precedence. Examples of RsmA/CsrA proteins stimulating

translation of target mRNAs include *phz2* and *moaA* in *P. aeruginosa* (15, 16) and *flhDC* in

E. coli (17). While *P. aeruginosa* FleQ and *E. coli* FlhD$_4$C$_2$ share no sequence or

mechanistic similarities, they are functional counterparts, with both being class I master

regulators of flagellar biosynthesis in their respective hosts (32). In addition, flagellar

motility was previously found to be positively controlled by RsmA in *P. aeruginosa* (4),

while in *E. coli,* CsrA has been shown to up-regulate flagellar gene expression through

protecting *flhDC* mRNA from degradation (17). It follows that if RsmA regulated *fleQ* in

an analogous manner, *fleQ* mRNAs would be hyper-stabilized by RsmA and thus, less

stable in its absence. However, no evidence of altered mRNA turnover rate was found

between WT and Δ*rsmA* strains (Fig. S2B). Therefore, we reasoned that the direct action of

RsmA may lie even further upstream in a regulatory cascade that involved control of *fleQ*

by Vfr.

Vfr acts as a transcriptional repressor of the *fleQ* promoter (33). Therefore, to be consistent

with the above phenotypes, RsmA would have to serve as a repressor of *vfr*. In line with

this idea, comparison of *vfr* transcriptional reporter activities in WT and Δ*rsmA* strains

revealed up-regulation in the absence of RsmA (Fig. 1A). To ensure that the regulatory

cascade functioned as anticipated, we also tested the prediction that over-expression of Vfr

should ultimately result in increased activity of the *pel* transcriptional reporter. As shown in

Fig. 1B, this is indeed the case. In contrast to *pel* (Fig. S1B) and *fleQ* (Fig. S2B), we found

that the transcript stability of *vfr* was significantly different between WT and Δ*rsmA*

strains, such that *vfr* mRNAs were more stable in the absence of RsmA (Fig. 1C). These

results indicated that RsmA affects the *vfr* mRNA levels by mediating changes in turnover

rates.

RsmA regulation of Vfr alters swimming and twitching motility.

P. aeruginosa uses at least two major modes of motility: flagellar-driven swimming

motility through liquid and surface 'twitching' motility using Type IV pilus (34, 35).

Bacterial migration is an important aspect during *P. aeruginosa* biofilm development as

initial attachment to a surface is thought to require flagella and mature microcolony

formation has been attributed to Type IV pili functions (36, 37). Thus, fine-tuned regulatory

control of motility and biofilm genes is a necessity for effective *P. aeruginosa* adaptation,

particularly when switching between motile and sessile lifestyles.

Because FleQ is a class I master regulator of flagellar genes (32) and RsmA and Vfr lie

upstream of FleQ in the regulatory cascade, lack of either protein should predictively

impact flagellar motility, but in opposite ways. As would be predicted, Δ*rsmA* has reduced

motility while Δ*vfr* is hyper-motile (Fig. 2A). Mutants in *vfr* have previously been

documented to be defective in Type IV pilus-dependent twitching motility (38, 39) due to

Vfr's positive regulation of the AlgZR two-component system (40, 41) required for

expression of the Type IV pili genes (42, 43). Based on the results that RsmA serves as a

repressor of *vfr* (Fig. 1), we predicted that the over-expression of RsmA and lack of Vfr

should have epistatically similar phenotypic effect on twitching motility. As shown in Fig.

2, Δ*vfr* and RsmA over-expressing strains were both defective in twitching motility.

Hfq is required for RsmA to bind to *vfr* mRNA.

Given the evidence that RsmA may post-transcriptionally regulate *vfr*, we next examined

the binding properties of RsmA to *vfr* mRNA. *P. aeruginosa* RsmA has a binding

specificity for the CANGGAYG consensus sequence of its target mRNA (44), which is

similar to the *E. coli* CsrA consensus sequence RUACARGGAUGU (45). The

transcriptional start site for *vfr* lies 153 bases upstream of the translational start site (46).

Inspection of the *vfr* leader sequence identified one putative RsmA-binding site overlapping

the Shine-Dalgarno sequence (Fig. 3A).

RsmA, purified as used here, directly binds *psl* mRNA (8) in RNA electrophoretic mobility

shift assays (EMSA). However, no RsmA-binding was observed in similar assays with *vfr*

RNA spanning the leader sequence (Fig. 3B). This initially perplexing result led us to

consider that RsmA may require another factor to bind the *vfr* RNA. Our attention was

drawn to Hfq for three reasons. Firstly, the small RNA RsmY, which binds multiple RsmA

proteins at high affinity to sequester and relieve RsmA-bound mRNAs (47), also has the

capacity to be bound by Hfq (48), although it was unclear whether co-binding of both

proteins occurred. Secondly, Hfq was recently identified to bind to *vfr* mRNA by global

ChIP-Seq analyses (49). Thirdly, *in silico* analysis of the *vfr* leader sequence revealed two

potential Hfq binding sites upstream of the putative RsmA-binding site (Fig. 3A).

Biochemical studies, primarily of *E. coli* Hfq, have shown that this hexameric protein

complex has four regions which can all be involved in RNA-binding: the proximal face, the

distal face, the rim, and the C-terminal tail (50). *P. aeruginosa* and *P. putida* Hfq have

shortened C-termini compared to *E. coli* Hfq and may lack C-terminal tail-binding

altogether (51). The proximal site preferentially binds to U-rich RNA sequence (52), the

distal site binds to A-rich $(ARN)_n$ triplet repeats (53), while the rim associates with UA-rich

regions (54). The two potential Hfq-binding sites found within the *vfr* leader sequence (Fig.

3A) represent one A-rich ARN repeat region and one U-rich region.

RNA EMSA analyses showed that *P. aeruginosa* Hfq bound *vfr* RNA with high efficiency,

resulting in three distinct band shifts (Fig. 3C) indicative of two or more binding sites.

Because wild-type *P. putida* Hfq showed identical binding patterns to *P. aeruginosa* Hfq on

vfr RNA (Fig. S3A), we took advantage of previously characterized derivatives of *P. putida*

Hfq, namely, a distal site mutant (Hfq_{Y25D}; A-rich ARN-binding deficient) and a proximal

site mutant (Hfq_{K56A}; U-rich-binding deficient) (55) to further analyze Hfq-binding to the

vfr RNA. As seen in Fig. S3B, Hfq_{Y25D} only recapitulated the second band shift while

Hfq_{K56A} only recapitulated the first band shift. Notably, the third band shift is absent with

both mutant derivatives of Hfq. Based on this set of data, we conclude that the first shift is

caused by Hfq-binding to the U-rich region of the *vfr* leader sequence (Hfq-site 2 in Fig.

3A), the second shift is caused when the ARN repeats are bound by Hfq (Hfq-site 1 in Fig.

3A), while the third shift occurs when both sites are occupied simultaneously.

Having ascertained that Hfq does bind *vfr* RNA, we next added RsmA and Hfq

simultaneously to the *vfr* RNA. As seen in Fig. 3D, this resulted in super-shifting of all

three Hfq band shifts in the presence of RsmA. Because the GG doublet within the

consensus sequences for RsmA and CsrA binding are essential for their abilities to bind

cognate target RNAs (8, 45), we made a CC substitution of the GG doublet within the

predicted RsmA-binding region of *vfr* RNA. No RsmA-mediated super-shifting of Hfq

bands shifts was detected with the CC substituted RNA (Fig. 3E).

We infer from the findings above that Hfq-binding is a pre-requisite for RsmA to bind to its

target within *vfr* mRNA. Together with the data in preceding sections, these results lead us

to conclude: 1) that the direct binding of RsmA on *vfr* RNA requires the RNA chaperone

protein Hfq and 2) that RsmA is an indirect regulator of biofilm polysaccharide locus *pel*,

flagellar motility and Type IV pili-mediated twitching motility, through its direct action on

vfr propagated through a regulatory cascade from Vfr to FleQ and AlgR.

238	DISCUSSION

In this study, we report that *P. aeruginosa* RsmA requires the RNA chaperone Hfq to assist

its binding to *vfr* mRNA to initiate a regulatory cascade that ultimately impacts both

motility- and biofilm-associated genes. The requirement for Hfq was unexpected, since all

other biochemical analysis of RsmA/CsrA family members document unaided binding to

their target RNAs (e.g.: 8, 15-17, 56-59). We identified two independent Hfq-binding sites

upstream of the RsmA-binding motif, which overlaps the ribosome-binding site of *vfr* (Fig.

3A). Based on a mFOLD-predicted secondary structure, a large hairpin loop base-pairs the

identified U-rich and A-rich Hfq-binding sites, resulting in a dsRNA structure that would

predictively block both RsmA and the ribosome from accessing the mRNA (Fig. 4A). As

depicted in Fig. 4B, the simplest model that arises from our findings is that Hfq-binding is

required to open the stem-loop to expose the RsmA-binding site. Subsequent binding of

RsmA would directly block the Shine-Dalgarno sequence, preventing ribosome access and

thereby inhibiting translation. We also provide evidence that *vfr* mRNA is hyper-stable in

the absence of RsmA (Fig. 1C), indicating that Hfq- and RsmA-bound *vfr* mRNA may

undergo more rapid degradation. It was previously suggested that Hfq may be able to

directly recruit RNase E for the degradation of bound RNA in *E. coli* (60), and it is possible

that a similar mechanism causes *vfr* mRNA instability.

Due to its histidine-rich C-terminus, *E. coli* Hfq is a common contaminant of His-tagged

proteins purified by nickel affinity chromatography after over-expression in *E. coli*, and

visually undetectable Hfq levels can critically influence *in vitro* analysis of RNA-binding

(51, 61). Because the proteins used in this study were purified from either *P. aeruginosa*

(RsmA) or a Hfq null strain of *E. coli* (Hfq proteins), our analyses were not complicated by

this issue. However, a large majority of analyses performed in past publications use C-

terminal His-tagged RsmA/CsrA proteins derived from over-expression in *E. coli,* raising

the possibility of *E. coli Hfq* contamination. Given that Hfq binds *vfr* RNA with apparent

high affinity (Figs. 3C and S3), it is plausible that Hfq-assisted RsmA-binding may be

prevalent. Although such a possibility would not alter major conclusions and remains to be

experimentally verified, it is provocative that comparison of transcriptomic studies done for

RsmA, Hfq, and Vfr regulons in *P. aeruginosa* identify numerous overlaps of genes that

were differentially expressed between WT and the respective null mutants (18, 19, 29, 62,

63).

A diverse range of phenotypes have been associated with RsmA functions in *P. aeruginosa*

(64), but phenotypic observations do not distinguish direct regulatory activities from

indirect effects through regulatory cascades. Our identification of Hfq and RsmA action on

vfr mRNA resulted from an initial aim to determine whether RsmA directly targeted *pel*

RNA, as is the case for *psl* RNA (8). Our back-tracking to trace the ultimate cause of RsmA

effects on *pel* expression revealed a regulatory pathway originating from Vfr to the

transcriptional regulator FleQ and from there to *pel* (Figs. S1, S2, and 1) – a cascade that in

addition to biofilm-associated genes, also impacts bacterial motility (Fig. 2).

As summarized in Fig. 4C, this work places the global post-transcriptional regulators RsmA

and Hfq as a hub at the center of upstream and downstream regulatory pathways and thus

provides a conceptually new framework to evaluate/interpret past (and future) work. The

two component GacS/GacA system lies at the top of the hierarchy that controls RsmA-

mediated regulation. Although the environmental cue(s) for activation is still unknown,

GacS activity is modulated through interactions with RetS and LadS (65-67). GacA, in turn

controls the production of the small RNAs – *rsmY* and *rsmZ* – which possess multiple high

affinity binding sites to titrate RsmA away from target mRNAs (4, 68). In light of a report

indicating that Hfq also binds to *rsmY* (48), it appears plausible that these small RNAs may

also require Hfq to assist their function.

In addition to motility- and biofilm-associated phenotypes, there are numerous other

processes associated with RsmA in *P. aeruginosa*, including acute virulence (5, 69). Many

previously assigned 'RsmA-regulated factors and phenotypes' are likely to be indirectly

regulated through Vfr initiated regulatory cascades. Here we present evidence for

swimming motility (Vfr/FleQ/Flagella cascade, Fig. 4C center), twitching motility

(Vfr/AlgR/Type IV pili, Fig. 4C right) and PEL (Vfr/FleQ/PEL cascade, Fig. 4C left),

Based on our findings, it will be crucial to determine and distinguish direct versus indirect

regulatory routes to gain a greater understanding of the RsmA regulon.

MATERIALS AND METHODS

Bacterial strains and growth conditions

Table S1 lists the bacterial strains used in this study. *E. coli* and *P. aeruginosa* strains were

grown in lysogeny broth (LB) at 37°C unless specified otherwise. VBMM citrate medium

(70) was used for selecting *P. aeruginosa* post-conjugation. For *E. coli*, the following

antibiotics concentrations were used: 50 μg·ml^{-1} carbenicillin, 10 μg·ml^{-1} gentamicin, and

10 μg·ml^{-1} tetracycline. For *P. aeruginosa* strains: 300 μg·ml^{-1} carbenicillin, 100 μg·ml^{-1}

gentamicin, and 100 μg·ml^{-1} tetracycline were used. Sucrose counter-selection for plasmids

carrying the *sacB* gene used in *P. aeruginosa* strain constructions was performed by

streaking colonies on LB agar (no salt) supplemented with 10% w/v sucrose. Plates were

incubated at 30°C for 24 hours, after which the counter-selected colonies were confirmed

for the loss of antibiotic resistance and mutations confirmed by PCR for double-cross-over

genomic mutants.

Strain constructions

Transcriptional fusion constructs were generated in single copy on the chromosome of *P.*

aeruginosa strains via integration into the chromosomal *attB* site as previously published

(8). In brief, promoter regions were PCR amplified using oligonucleotides in Table S2.

PCR products were ligated between the EcoRI and BamHI sites of mini-CTX *lacZ* (Fig.

S4). These plasmids were then introduced into *P. aeruginosa* by conjugation. After double

site recombination, plasmid backbones were removed by FLP recombinase, and then strains

cured of the pFLP2 plasmid by sucrose counter-selection.

Motility assays

Swimming motility assays were performed essentially as previously described (71). LB

plates containing 0.3% Bacto agar were inoculated with overnight cultures with a sterile

inoculation needle, ensuring the needle tip was inserted approximately halfway into the

agar but not to the plastic petri dish bottom, and incubated for 24 hours at 30°C. Swim ring

diameters were measured for quantitation. For twitching motility, we followed the protocol

of Haley *et al.* (72) by using LB plates supplemented with 5 mg/ml porcine gastric mucin

(Sigma-Aldrich). A sterile inoculation needle was inserted through the agar until it touched

the plastic bottom. Plates were incubated for 48 hours at 37°C, then at room temperature for

48 more hours. Subsequently, the agar was peeled off, and the plates were stained with 1%

w/v crystal violet to visualize the twitching zone diameters prior to measurement (73). All

motility experiments were performed in biological quadruplicates.

β-galactosidase assays

Quantitative β-galactosidase activities were assayed using Galacto-Light Plus kit (Thermo-

Fisher) as previously published (74). All *P. aeruginosa* cultures were grown in VBMM

citrate at 37°C to exponential phase, and lysed using chloroform as previously described

(8). β-galactosidase activity units were normalized to total proteins per ml as determined

using Bradford assay reagents (Bio-Rad). Assays were performed in biological triplicates.

Quantitative real-time PCR and RNA stability analyses

Real-time PCR was performed as previously described (30), using the oligonucleotides

listed in Table S2. For RNA stability experiments, exponential phase *P. aeruginosa*

cultured in VBMM citrate at 37°C were treated with 200 μg·ml^{-1} rifampicin (75). RNAs

were extracted from 1 ml of the cultures at various time points as previously described (8)

after the addition of rifampicin. RNA extractions were performed using the RNeasy Mini

Kit (Qiagen) after treating the harvested cells with RNAprotect (Qiagen). Genomic DNA

was removed using DNase I (Promega) and removal confirmed by PCR using primers

designed against the *rplU* gene (Table S2). SuperScript III First-Strand Synthesis

(Invitrogen) was used to synthesize cDNA as per manufacturer's protocol using random

hexamers. Quantitative real-time PCR was performed using SYBR Green PCR Master Mix

(Thermo/Applied Biosystems). The internal control gene used was *ampR*. All experiments

were done in biological quadruplicates.

Purified proteins

Proteins were purified by previously established protocols. Over-expressed RsmA-His$_6$ was

purified from *P. aeruginosa* bearing pUCP18::*rsmA*-His$_6$ (8), while all Hfq-His$_6$ proteins

were purified from a Hfq null derivative of *E. coli* BL21(DE3) (55) carrying P$_{T7}$ expression

plasmids (pVI2344-2346 or pVI2357). All proteins were purified through Ni-NTA affinity

with HisTrap HP 1 ml and further by ion exchange (HiTrap SP HP 5 ml for RsmA and

ResourceQ 1ml for Hfq) by ÄKTA avant FPLC (GE Healthcare). Final protein

concentrations were determined using the BCA Protein Assay Kit (Pierce/Fisher).

RNA synthesis

Wild-type and mutant *vfr* mRNAs were generated using Ambion's MEGAscript kit as

recommended for *in vitro* transcription from the P$_{T7}$ promoter of linearized plasmids.

Reactions (total 40 µl) containing 1 µg of EcoRV-linearized pVI2358 or pVI2359, were

incubated for 6 hours prior to 15 minutes DNase I treatment. Reactions were terminated by

adding 230 µl RNase-free H$_2$O and 30 µl stop solution (5 M ammonium acetate; 100 mM

EDTA). The resulting 233 nt RNAs (encompassing co-ordinates -106 to +116 relative to

the A of the initiation codon of *vfr*) were extracted twice with phenol:chloroform:IAA

(25:24:1) and once with chloroform prior to precipitation and resuspension in 50 µl RNase-

free H$_2$O.

RNA electrophoretic mobility shift assays (EMSAs)

Reactions (total 10 µl) contained 32 nM *in vitro* transcribed RNA with the indicated

concentrations of RsmA and/or Hfq. RsmA and Hfq molarities given here are for the

dimeric and hexameric complexes respectively. RNA was first heated at 80^{o}C for 5 min and

then immediately chilled in an ice-water bath. Additional final reaction mixture

components were as follows: 10 mM HEPES pH 7.9, 2 mM MgCl$_2$, 90 ng yeast total RNA

(Fisher), 4U RNasin, and 35 mM KCl. Binding reactions were performed at 20^{o}C for 40

min prior to adding 2.5 µl loading buffer (40% sucrose). Reactions were placed on ice and

analyzed on 5% TBE Criterion pre-cast gels (Bio-Rad) with electrophoresis at 100V for

~130 min at 4^{o}C. Gels were stained at room temperature with 1:10,000 TBE-diluted SYBR

Gold (Fisher) in a light-protected container with agitation. Images were captured using

Typhoon FLA 9500 (GE Healthcare).

Statistics

Presented data are the mean with standard deviations of the data collected. Student's t-test

using the GraphPad Prism software was used to calculate p-values.

395   ACKNOWLEDGEMENTS

396   The authors would like to thank Victoriia Murina for her assistance with the FPLC. Plasmid

397   pMMBGW:RBS-*vfr* was a kind gift from Joe J. Harrison. YI would like to thank Matt

398   Parsek for his more than a decade of scientific and moral support that led to the culmination

399   of this project. The authors declare no conflict of interests.

401   FUNDING INFORMATION

402   This work was supported by the J C Kempe & S M Kempe Foundation (JCK-1523 to VS).

REFERENCES

1.   Gellatly SL & Hancock RE (2013) *Pseudomonas aeruginosa*: new insights into
pathogenesis and host defenses. *Pathog. Dis.* 67(3):159-173.
2.   Khare A & Tavazoie S (2015) Multifactorial Competition and Resistance in a Two-
Species Bacterial System. *PLoS Genet.* 11(12):e1005715.
3.   Tashiro Y, Yawata Y, Toyofuku M, Uchiyama H, & Nomura N (2013) Interspecies
interaction between *Pseudomonas aeruginosa* and other microorganisms. *Microbes*
Environ. 28(1):13-24.
4.   Heurlier K*, et al.* (2004) Positive control of swarming, rhamnolipid synthesis, and
lipase production by the posttranscriptional RsmA/RsmZ system in *Pseudomonas*
aeruginosa PAO1. *J. Bacteriol.* 186(10):2936-2945.
5.   Mulcahy H, O'Callaghan J, O'Grady EP, Adams C, & O'Gara F (2006) The
posttranscriptional regulator RsmA plays a role in the interaction between
Pseudomonas aeruginosa and human airway epithelial cells by positively regulating
the type III secretion system. *Infect. Immun.* 74(5):3012-3015.
6.   Mulcahy H*, et al.* (2008) *Pseudomonas aeruginosa* RsmA plays an important role
during murine infection by influencing colonization, virulence, persistence, and
pulmonary inflammation. *Infect. Immun.* 76(2):632-638.
7.   O'Grady EP, Mulcahy H, O'Callaghan J, Adams C, & O'Gara F (2006)
Pseudomonas aeruginosa infection of airway epithelial cells modulates expression
of Kruppel-like factors 2 and 6 via RsmA-mediated regulation of type III
exoenzymes S and Y. *Infect. Immun.* 74(10):5893-5902.
8.   Irie Y*, et al.* (2010) *Pseudomonas aeruginosa* biofilm matrix polysaccharide Psl is
regulated transcriptionally by RpoS and post-transcriptionally by RsmA. *Mol.*
Microbiol. 78(1):158-172.
9.   Pessi G*, et al.* (2001) The global posttranscriptional regulator RsmA modulates
production of virulence determinants and *N*-acylhomoserine lactones in
Pseudomonas aeruginosa. *J. Bacteriol.* 183(22):6676-6683.
10.   Gutiérrez P*, et al.* (2005) Solution structure of the carbon storage regulator protein
CsrA from *Escherichia coli*. *J. Bacteriol.* 187(10):3496-3501.
11.   Rife C*, et al.* (2005) Crystal structure of the global regulatory protein CsrA from
Pseudomonas putida at 2.05 A resolution reveals a new fold. *Proteins* 61(2):449-
453.
12.   Schubert M*, et al.* (2007) Molecular basis of messenger RNA recognition by the
specific bacterial repressing clamp RsmA/CsrA. *Nat. Struct. Mol. Biol.* 14(9):807-
813.
13.   White D, Hart ME, & Romeo T (1996) Phylogenetic distribution of the global
regulatory gene *csrA* among eubacteria. *Gene* 182(1-2):221-223.
14.   Romeo T, Vakulskas CA, & Babitzke P (2013) Post-transcriptional regulation on a
global scale: form and function of Csr/Rsm systems. *Environ. Microbiol.* 15(2):313-
324.

15. Patterson-Fortin LM, Vakulskas CA, Yakhnin H, Babitzke P, & Romeo T (2013)
Dual posttranscriptional regulation via a cofactor-responsive mRNA leader. *J. Mol.*
Biol. 425(19):3662-3677.
16. Ren B, Shen H, Lu ZJ, Liu H, & Xu Y (2014) The *phzA2-G2* transcript exhibits
direct RsmA-mediated activation in *Pseudomonas aeruginosa* M18. *PloS one*
9(2):e89653.
17. Yakhnin AV*, et al.* (2013) CsrA activates *flhDC* expression by protecting *flhDC*
mRNA from RNase E-mediated cleavage. *Mol. Microbiol.* 87(4):851-866.
18. Brencic A & Lory S (2009) Determination of the regulon and identification of novel
mRNA targets of *Pseudomonas aeruginosa* RsmA. *Mol. Microbiol.* 72(3):612-632.
19. Burrowes E, Baysse C, Adams C, & O'Gara F (2006) Influence of the regulatory
protein RsmA on cellular functions in *Pseudomonas aeruginosa* PAO1, as revealed
by transcriptome analysis. *Microbiology* 152(Pt 2):405-418.
20. Lawhon SD*, et al.* (2003) Global regulation by CsrA in *Salmonella* typhimurium.
Mol. Microbiol. 48(6):1633-1645.
21. Baker CS*, et al.* (2007) CsrA inhibits translation initiation of *Escherichia coli hfq*
by binding to a single site overlapping the Shine-Dalgarno sequence. *J. Bacteriol.*
189(15):5472-5481.
22. Pannuri A*, et al.* (2012) Translational repression of NhaR, a novel pathway for
multi-tier regulation of biofilm circuitry by CsrA. *J. Bacteriol.* 194(1):79-89.
23. Bandyra KJ & Luisi BF (2013) Licensing and due process in the turnover of
bacterial RNA. *RNA Biol.* 10(4):627-635.
24. Ryder C, Byrd M, & Wozniak DJ (2007) Role of polysaccharides in *Pseudomonas*
aeruginosa biofilm development. *Curr. Opin. Microbiol.* 10(6):644-648.
25. Hickman JW & Harwood CS (2008) Identification of FleQ from *Pseudomonas*
aeruginosa as a c-di-GMP-responsive transcription factor. *Mol. Microbiol.*
69(2):376-389.
26. Gilbert KB, Kim TH, Gupta R, Greenberg EP, & Schuster M (2009) Global position
analysis of the *Pseudomonas aeruginosa* quorum-sensing transcription factor LasR.
Mol. Microbiol. 73(6):1072-1085.
27. Sakuragi Y & Kolter R (2007) Quorum-sensing regulation of the biofilm matrix
genes (*pel*) of *Pseudomonas aeruginosa*. *J. Bacteriol.* 189(14):5383-5386.
28. Gooderham WJ & Hancock RE (2009) Regulation of virulence and antibiotic
resistance by two-component regulatory systems in *Pseudomonas aeruginosa*.
FEMS Microbiol. Rev. 33(2):279-294.
29. Goodman AL*, et al.* (2004) A signaling network reciprocally regulates genes
associated with acute infection and chronic persistence in *Pseudomonas aeruginosa*.
Dev. Cell 7(5):745-754.
30. Irie Y*, et al.* (2012) Self-produced exopolysaccharide is a signal that stimulates
biofilm formation in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. U. S. A.*
109(50):20632-20636.
31. Baraquet C, Murakami K, Parsek MR, & Harwood CS (2012) The FleQ protein
from *Pseudomonas aeruginosa* functions as both a repressor and an activator to

488		control gene expression from the *pel* operon promoter in response to c-di-GMP.
489		*Nucleic Acids Res.* 40(15):7207-7218.
490	32.	Dasgupta N, *et al.* (2003) A four-tiered transcriptional regulatory circuit controls
491		flagellar biogenesis in *Pseudomonas aeruginosa. Mol. Microbiol.* 50(3):809-824.
492	33.	Dasgupta N, Ferrell EP, Kanack KJ, West SE, & Ramphal R (2002) *fleQ*, the gene
493		encoding the major flagellar regulator of *Pseudomonas aeruginosa*, is σ^{70}
494		dependent and is downregulated by Vfr, a homolog of *Escherichia coli* cyclic AMP
495		receptor protein. *J. Bacteriol.* 184(19):5240-5250.
496	34.	Burrows LL (2012) *Pseudomonas aeruginosa* twitching motility: type IV pili in
497		action. *Annu. Rev. Microbiol.* 66:493-520.
498	35.	Kazmierczak BI, Schniederberend M, & Jain R (2015) Cross-regulation of
499		*Pseudomonas* motility systems: the intimate relationship between flagella, pili and
500		virulence. *Curr. Opin. Microbiol.* 28:78-82.
501	36.	Klausen M, *et al.* (2003) Biofilm formation by *Pseudomonas aeruginosa* wild type,
502		flagella and type IV pili mutants. *Mol. Microbiol.* 48(6):1511-1524.
503	37.	O'Toole GA & Kolter R (1998) Flagellar and twitching motility are necessary for
504		*Pseudomonas aeruginosa* biofilm development. *Mol. Microbiol.* 30(2):295-304.
505	38.	Beatson SA, Whitchurch CB, Sargent JL, Levesque RC, & Mattick JS (2002)
506		Differential regulation of twitching motility and elastase production by Vfr in
507		*Pseudomonas aeruginosa. J. Bacteriol.* 184(13):3605-3613.
508	39.	Whitchurch CB, Alm RA, & Mattick JS (1996) The alginate regulator AlgR and an
509		associated sensor FimS are required for twitching motility in *Pseudomonas*
510		*aeruginosa. Proc. Natl. Acad. Sci. U. S. A.* 93(18):9839-9843.
511	40.	Luo Y, *et al.* (2015) A hierarchical cascade of second messengers regulates
512		*Pseudomonas aeruginosa* surface behaviors. *mBio* 6(1).
513	41.	Pritchett CL, *et al.* (2015) Expression analysis of the *Pseudomonas aeruginosa*
514		AlgZR two-component regulatory system. *J. Bacteriol.* 197(4):736-748.
515	42.	Belete B, Lu H, & Wozniak DJ (2008) *Pseudomonas aeruginosa* AlgR regulates
516		type IV pilus biosynthesis by activating transcription of the *fimU-pilVWXY1Y2E*
517		operon. *J. Bacteriol.* 190(6):2023-2030.
518	43.	Lizewski SE, *et al.* (2004) Identification of AlgR-regulated genes in *Pseudomonas*
519		*aeruginosa* by use of microarray analysis. *J. Bacteriol.* 186(17):5672-5684.
520	44.	Schulmeyer KH, *et al.* (2016) Primary and Secondary Sequence Structure
521		Requirements for Recognition and Discrimination of Target RNAs by *Pseudomonas*
522		*aeruginosa* RsmA and RsmF. *J. Bacteriol.* 198(18):2458-2469.
523	45.	Dubey AK, Baker CS, Romeo T, & Babitzke P (2005) RNA sequence and
524		secondary structure participate in high-affinity CsrA-RNA interaction. *Rna*
525		11(10):1579-1587.
526	46.	Fuchs EL, *et al.* (2010) The *Pseudomonas aeruginosa* Vfr regulator controls global
527		virulence factor expression through cyclic AMP-dependent and -independent
528		mechanisms. *J. Bacteriol.* 192(14):3553-3564.
529	47.	Kay E, *et al.* (2006) Two GacA-dependent small RNAs modulate the quorum-
530		sensing response in *Pseudomonas aeruginosa. J. Bacteriol.* 188(16):6026-6033.

48.   Sorger-Domenigg T, Sonnleitner E, Kaberdin VR, & Bläsi U (2007) Distinct and
overlapping binding sites of *Pseudomonas aeruginosa* Hfq and RsmA proteins on
the non-coding RNA RsmY. *Biochem. Biophys. Res. Commun.* 352(3):769-773.
49.   Kambara TK, Ramsey KM, & Dove SL (2018) Pervasive Targeting of Nascent
Transcripts by Hfq. *Cell Rep.* 23(5):1543-1552.
50.   Updegrove TB, Zhang A, & Storz G (2016) Hfq: the flexible RNA matchmaker.
Curr. Opin. Microbiol. 30:133-138.
51.   Milojevic T, Grishkovskaya I, Sonnleitner E, Djinovic-Carugo K, & Bläsi U (2013)
The *Pseudomonas aeruginosa* catabolite repression control protein Crc is devoid of
RNA binding activity. *PloS one* 8(5):e64609.
52.   Møller T*, et al.* (2002) Hfq: a bacterial Sm-like protein that mediates RNA-RNA
interaction. *Mol. Cell.* 9(1):23-30.
53.   Link TM, Valentin-Hansen P, & Brennan RG (2009) Structure of *Escherichia coli*
Hfq bound to polyriboadenylate RNA. *Proc. Natl. Acad. Sci. U. S. A.*
106(46):19292-19297.
54.   Schu DJ, Zhang A, Gottesman S, & Storz G (2015) Alternative Hfq-sRNA
interaction modes dictate alternative mRNA recognition. *EMBO J.* 34(20):2557-
2573.
55.   Madhushani A, Del Peso-Santos T, Moreno R, Rojo F, & Shingler V (2015)
Transcriptional and translational control through the 5'-leader region of the *dmpR*
master regulatory gene of phenol metabolism. *Environ. Microbiol.* 17(1):119-133.
56.   Baker CS, Morozov I, Suzuki K, Romeo T, & Babitzke P (2002) CsrA regulates
glycogen biosynthesis by preventing translation of *glgC* in *Escherichia coli*. *Mol.*
Microbiol. 44(6):1599-1610.
57.   Dubey AK*, et al.* (2003) CsrA regulates translation of the *Escherichia coli* carbon
starvation gene, *cstA*, by blocking ribosome access to the cstA transcript. *J.*
Bacteriol. 185(15):4450-4460.
58.   Lapouge K*, et al.* (2007) Mechanism of *hcnA* mRNA recognition in the Gac/Rsm
signal transduction pathway of *Pseudomonas fluorescens*. *Mol. Microbiol.*
66(2):341-356.
59.   Wang X*, et al.* (2005) CsrA post-transcriptionally represses *pgaABCD*, responsible
for synthesis of a biofilm polysaccharide adhesin of *Escherichia coli*. *Mol.*
Microbiol. 56(6):1648-1663.
60.   Morita T, Maki K, & Aiba H (2005) RNase E-based ribonucleoprotein complexes:
mechanical basis of mRNA destabilization mediated by bacterial noncoding RNAs.
Genes Dev. 19(18):2176-2186.
61.   Moreno R*, et al.* (2015) The Crc and Hfq proteins of *Pseudomonas putida* cooperate
in catabolite repression and formation of ribonucleic acid complexes with specific
target motifs. *Environ. Microbiol.* 17(1):105-118.
62.   Sonnleitner E*, et al.* (2018) Interplay between the catabolite repression control
protein Crc, Hfq and RNA in Hfq-dependent translational regulation in
Pseudomonas aeruginosa. *Nucleic Acids Res.* 46(3):1470-1485.

63.   Wolfgang MC, Lee VT, Gilmore ME, & Lory S (2003) Coordinate regulation of
bacterial virulence genes by a novel adenylate cyclase-dependent signaling
pathway. *Dev. Cell* 4(2):253-263.
64.   Vakulskas CA, Potts AH, Babitzke P, Ahmer BM, & Romeo T (2015) Regulation
of bacterial virulence by Csr (Rsm) systems. *Microbiol. Mol. Biol. Rev.* 79(2):193-
224.
65.   Chambonnier G*, et al.* (2016) The Hybrid Histidine Kinase LadS Forms a
Multicomponent Signal Transduction System with the GacS/GacA Two-Component
System in *Pseudomonas aeruginosa. PLoS Genet.* 12(5):e1006032.
66.   Goodman AL*, et al.* (2009) Direct interaction between sensor kinase proteins
mediates acute and chronic disease phenotypes in a bacterial pathogen. *Genes Dev.*
23(2):249-259.
67.   Kong W*, et al.* (2013) Hybrid sensor kinase PA1611 in *Pseudomonas aeruginosa*
regulates transitions between acute and chronic infection through direct interaction
with RetS. *Mol. Microbiol.* 88(4):784-797.
68.   Sonnleitner E, Schuster M, Sorger-Domenigg T, Greenberg EP, & Bläsi U (2006)
Hfq-dependent alterations of the transcriptome profile and effects on quorum
sensing in *Pseudomonas aeruginosa. Mol. Microbiol.* 59(5):1542-1558.
69.   Intile PJ, Diaz MR, Urbanowski ML, Wolfgang MC, & Yahr TL (2014) The AlgZR
two-component system recalibrates the RsmAYZ posttranscriptional regulatory
system to inhibit expression of the *Pseudomonas aeruginosa* type III secretion
system. *J. Bacteriol.* 196(2):357-366.
70.   Hoang TT & Schweizer HP (1997) Fatty acid biosynthesis in *Pseudomonas*
aeruginosa: cloning and characterization of the *fabAB* operon encoding β-
hydroxyacyl-acyl carrier protein dehydratase (FabA) and β-ketoacyl-acyl carrier
protein synthase I (FabB). *J. Bacteriol.* 179(17):5326-5332.
71.   Shrout JD*, et al.* (2006) The impact of quorum sensing and swarming motility on
Pseudomonas aeruginosa biofilm formation is nutritionally conditional. *Mol.*
Microbiol. 62(5):1264-1277.
72.   Haley CL, Kruczek C, Qaisar U, Colmer-Hamood JA, & Hamood AN (2014)
Mucin inhibits *Pseudomonas aeruginosa* biofilm formation by significantly
enhancing twitching motility. *Can. J. Microbiol.* 60(3):155-166.
73.   Déziel E, Comeau Y, & Villemur R (2001) Initiation of biofilm formation by
Pseudomonas aeruginosa 57RP correlates with emergence of hyperpiliated and
highly adherent phenotypic variants deficient in swimming, swarming, and
twitching motilities. *J. Bacteriol.* 183(4):1195-1204.
74.   Lequette Y, Lee J-H, Ledgham F, Lazdunski A, & Greenberg EP (2006) A distinct
QscR regulon in the *Pseudomonas aeruginosa* quorum-sensing circuit. *J. Bacteriol.*
188(9):3365-3370.
75.   Lory S (1986) Effect of iron on accumulation of exotoxin A-specific mRNA in
Pseudomonas aeruginosa. *J. Bacteriol.* 168(3):1451-1456.
614

Fig. 1   RsmA is a repressor of *vfr*.

A. Single-copy transcriptional *lacZ* fusion constructs reveal up-regulation of *vfr* transcripts

in the Δ*rsmA* background as compared to WT.

B. The single-copy P*pelA*::*lacZ* transcriptional fusion has a higher activity in a Vfr over-

expressing strain compared to the same strain harboring the vector control (VC).

C. Real-time quantitative PCR at different time points after rifampicin treatment show that

vfr mRNA is less stable in WT as compared to the Δ*rsmA* strain.

Fig. 2   RsmA regulation of Vfr indirectly regulates flagellar and Type IV pilus-

mediated motilities.

A. Flagellar-mediated swimming motility is decreased in the absence of RsmA (Δ*rsmA*)

and enhanced in the absence of Vfr (Δ*vfr*) as compared to the WT strain.

B. Type IV pilus-mediated twitching motility is unaffected when *rsmA* is mutated but is

reduced in the Δ*vfr* and RsmA over-expression strains.

Fig. 3   Direct targeting of *vfr* mRNA by RsmA requires Hfq.

A. The leader sequence of the *vfr* transcript (46) possesses two putative Hfq-binding sites

(blue boxes) and a RsmA-binding site (green box) overlapping the predicted Shine-

Dalgarno sequence (red italics) upstream of the translational start site (underlined italics).

B. RNA Electrophoretic mobility shift assay (EMSA) shows that RsmA does not bind to

the *vfr* RNA when RsmA is present alone. Dimeric molar excess relative to RNA; U =

unbound RNA.

C. The *vfr* RNA produces three distinct band shifts (open arrow heads 1, 2, and 3) with

increasing concentrations of Hfq. Hexameric molar excess relative to RNA.

D. The presence of RsmA causes all Hfq-shifted bands to super-shift (filled arrow heads

1+, 2+, and 3+), indicating that RsmA binds to Hfq-bound RNA. Note that the RsmA

concentration used (36× molar excess of RNA) is incapable of producing a shift when

present alone (panel B).

E. A GG→CC substitution within the RsmA-binding site abolishes RsmA super-shifts but

does not alter Hfq-binding.

Fig. 4 Model for Hfq-assisted RsmA regulation of *vfr* and its consequences

A. RNA secondary structure of the leader sequence region of *vfr* as predicted by mFOLD.

The locations of *in silico*-detected Hfq- and RsmA-binding sites are indicated. Note that the

RsmA-binding site located at the base of the stem loop structure overlaps with the Shine-

Dalgarno sequence of the ribosome binding site. The AUG start codon is shown underlined

in italics.

B. Schematic illustration of a model for Hfq-assisted RsmA regulation of *vfr*. Upon Hfq-

binding of *vfr* RNA, the stem loop structure shown in panel A is disrupted, exposing the

RsmA-binding site. RsmA-binding then directly blocks the Shine Dalgarno (SD) sequence.

RsmA-binding is depicted as producing a smaller stem-loop structure (10-12), preventing

ribosome access and thereby inhibiting translation. *In vivo*, the Hfq- and RsmA-bound

mRNAs are presumed to be subsequently processed for degradation.

C. Simplified overview of the Gac/Rsm pathway that controls RsmA levels and pertinent

parts of its downstream regulon.

Fig. 1

Fig. 2

A

*p<0.003 vs WT

B

*p<0.0001 vs WT

Fig. 3

A

CGCGGUUUACUGGCACACUUCCUGAUCUGCCCGGUUGGGGGGAGCUCUCCGCUGAGCUCC

CGGGCCCUGGCAGGUCUCUCCAUAAGAACUC AAAAAAACA UCGCCUUGUACGUCAGGCAU
Hfq-site 1

A UUCUUUAUU C UUUC *GGGA* CUCGGGC *AUG*→
Hfq-site 2   RsmA-binding site
Shine-Dalgarno   *Start codon*

B dimeric molar excess
RsmA: 0 10× 14× 18×
U

C hexameric molar excess
Hfq: 0 1× 2× 3× 4× 6×
◁3
◁2
◁1
U

D
Hfq: 0   0   2× 2× 4× 4×
RsmA: 0 18× 0 18× 0 18×
3+
2+
1+
U
Wild-type RsmA-binding mRNA
–UC UUUC *GGGA* CUCGG GC *AUG*–

E
Hfq: 0   0   2× 2× 4× 4×
RsmA: 0 18× 0 18× 0 18×
◁3
◁2
◁1
U
Mutant RsmA-binding site mRNA
–UC UUUC *G*CC*A* CUCGGGC *AUG*–

Fig. 4