Hfq-assisted RsmA regulation is central to *Pseudomonas aeruginosa* biofilm polysaccharide PEL expression

Yasuhiko Irie[1][#][*], Victoriia Murina[1,2], Vasili Hauryliuk[1,2,3], and Victoria Shingler[1]

[1] Department of Molecular Biology, Umeå University, Umeå, 901 87, Sweden

[2] The Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå, 901 87, Sweden

[3] Institute of Technology, University of Tartu, Nooruse 1, Tartu 50411, Estonia

Running Title: *Pseudomonas aeruginosa* RsmA regulates Vfr

Keywords: *Pseudomonas aeruginosa*, RsmA, Vfr, FleQ, Hfq

*Corresponding author

[#]current address: Institute of Technology, University of Tartu, Nooruse 1, Tartu 50411, Estonia. Tel: +372-737-4800. E-mail: yasuhiko.irie@ut.ee

ORCID ID for Yasuhiko Irie: 0000-0002-7162-4772

ORCID ID for Victoriia Murina: 0000-0003-1523-1816

ORCID ID for Vasili Hauryliuk: 0000-0003-2389-5057

ORCID ID for Victoria Shingler: 0000-0002-7349-1678

24   ABSTRACT

Expression of biofilm-associated genes is controlled by multiple regulatory elements,

allowing bacteria to appropriately switch between sessile and motile lifestyles. In

Pseudomonas aeruginosa, the post-transcriptional regulator RsmA has been implicated in

the control of various genes including those related to biofilms, but much of the evidence

for these links is limited to transcriptomic and phenotypic studies. RsmA binds to target

mRNAs to modulate translation by affecting ribosomal access and/or mRNA stability. Here

we trace a global regulatory role of RsmA to the inhibition of Vfr – a transcription factor

that controls a transcriptional regulator FleQ. FleQ directly controls biofilm-associated

genes that encode the PEL polysaccharide biosynthesis machinery. Furthermore, we show

that RsmA cannot bind *vfr* mRNA alone, but requires the RNA chaperone protein Hfq. This

is the first example where a RsmA protein family member is demonstrated to require

another protein for RNA binding.

                                        INTRODUCTION

The opportunistic pathogen *Pseudomonas aeruginosa* can be isolated from a wide range of

environmental niches, in large part owing to its versatile metabolic capabilities. It is also

proficient in colonizing various eukaryotic organisms and can cause both acute and chronic

infections, with the latter often associated with biofilm-like modes of growth at the sites of

infection (Gellatly and Hancock, 2013). *P. aeruginosa* is also highly competitive against

other microbial species, with its adaptability and competitive fitness being aided by the

production of various secondary metabolites and virulence factors (Tashiro et al.,

2013;Khare and Tavazoie, 2015). Many of these cellular processes are inter-regulated by

multiple regulatory pathways, presumably to provide appropriate response mechanisms to a

wide range of environmental cues. One of the central regulators that determine *P.*

aeruginosa behavior is the post-transcription factor RsmA (Regulator of Secondary

Metabolite) as evidenced by its null mutant phenotypes linked to motility (Heurlier et al.,

2004), virulence (Mulcahy et al., 2006;O'Grady et al., 2006;Mulcahy et al., 2008), biofilm

(Mulcahy et al., 2006;Irie et al., 2010), growth (Irie et al., 2010), and secreted products

(Pessi et al., 2001;Heurlier et al., 2004).

RsmA belongs to the RsmA/CsrA family of dimeric RNA-binding proteins (Gutiérrez et

al., 2005;Rife et al., 2005;Schubert et al., 2007), homologues of which are found across a

wide range of both Gram-negative and Gram-positive bacterial species (White et al., 1996):

in some species including *E. coli*, they are called CsrA, while they are named RsmA in

other species such as *P. aeruginosa*. RsmA/CsrA proteins inhibit translation of specific

target mRNAs by binding to the ribosome binding sites, typically overlapping either the

Shine-Dalgarno or the start codon (Romeo et al., 2013). However, in a few cases, they can

also serve as positive regulators of translation by altering the secondary structures of bound

RNAs (Patterson-Fortin et al., 2013;Ren et al., 2014) or protecting RNAs from degradation

(Yakhnin et al., 2013).

Many RsmA/CsrA-targeted mRNAs are reported to have altered RNA decay rates. Several

studies attempted to exploit this property to determine the global *P. aeruginosa* RsmA

regulon using microarrays (Lawhon et al., 2003;Burrowes et al., 2006;Brencic and Lory,

2009). However, this approach cannot distinguish directly regulated genes from indirectly

regulated gene sets that result from cascade regulation, nor can it identify target RNAs that

do not exhibit changes in their stability (Baker et al., 2007;Pannuri et al., 2012). For

example, in *P. aeruginosa*, direct RsmA-mediated translation inhibition has been

demonstrated for the *P. aeruginosa*-specific biofilm polysaccharide operon *psl* without any

effect on the level of transcription (Irie et al., 2010). Hence, the *psl* operon was not

prominently featured in transcriptomic studies despite a strong biofilm phenotype

(Burrowes et al., 2006;Brencic and Lory, 2009), raising concerns of what is currently

considered to be 'the RsmA regulon' of *P. aeruginosa*. More direct global analyses of CsrA

using techniques such as CLIP-Seq, RIP-Seq, and ribosome profiling have been performed

in other organisms (Holmqvist et al., 2016;Potts et al., 2017;Sahr et al., 2017), but to our

knowledge, such biochemical techniques to analyze the RsmA regulon have not been

performed for *P. aeruginosa*.

Here, we present evidence that RsmA serves a master regulatory role through several

intermediate transcription factors that are known to control different cellular processes.

Through cascade regulation, originating from a direct effect on *vfr* mRNA, RsmA indirectly

affects biofilm polysaccharide expression. Furthermore, we discovered an unexpected

regulatory interplay between RsmA and the RNA chaperone protein Hfq on the target

mRNA, whereby RsmA could only bind in the presence of Hfq. To our knowledge, this is

the first example where RsmA/CsrA is unable to bind to its target alone. Because Hfq is

known to facilitate RNA degradation (Bandyra and Luisi, 2013), these results may provide

a mechanistic insight into RsmA-associated mRNA stability changes.

 RESULTS

93 RsmA is a post-transcriptional regulator of *vfr* through which *fleQ* and *pel* transcripts

94 are indirectly regulated.

P. aeruginosa produces at least three different extracellular biofilm polysaccharides:

alginate, PEL, and PSL (Ryder et al., 2007). PEL and PSL are co-regulated by several

factors including the secondary messenger molecule c-di-GMP via the c-di-GMP-binding

transcriptional regulator FleQ (Hickman and Harwood, 2008) and possibly also quorum

sensing (Sakuragi and Kolter, 2007;Gilbert et al., 2009). Similar to PSL, PEL has long been

considered to be regulated by RsmA (Gooderham and Hancock, 2009) based on a model

originally proposed from a microarray study (Goodman et al., 2004). However, unlike the

psl operon transcript, which was genetically and biochemically demonstrated to be directly

regulated by RsmA (Irie et al., 2010), no direct evidence for the *pel* operon has been

presented.

Consistent with a previous microarray study (Brencic and Lory, 2009), here using a

chromosomal transcriptional reporter, we found that *pel* is up-regulated in the Δ*rsmA*

mutant as compared to the wild type (WT) background (Fig. 1A). However, the PSL

polysaccharides are also over-produced in a Δ*rsmA* strain (Irie et al., 2010) resulting in

elevated intracellular levels of c-di-GMP due to PSL signaling (Irie et al., 2012). Given that

the *pel* genes are transcriptionally up-regulated by c-di-GMP (Hickman and Harwood,

2008;Baraquet et al., 2012) and the Δ*rsmA* strain has elevated c-di-GMP levels, it is

plausible that the up-regulation of *pel* could be due to increased c-di-GMP levels. To

uncouple *pel* expression from elevated c-di-GMP, we measured *pel* transcriptional reporter

activities when introduced into a low c-di-GMP Δ*rsmA* Δ*pel* Δ*psl* triple mutant background

(Irie et al., 2012). In this strain *pel* is still up-regulated (Fig. 1A), indicating that the up-

regulation is caused by lack of RsmA rather than changes in intracellular c-di-GMP

concentration.

Since direct regulation of mRNA levels by RsmA/CsrA often acts via altered mRNA

stability (Romeo et al., 2013), we tested whether the apparent *pel* transcriptional changes

between the WT and Δ*rsmA* strains is due to altered *pel* mRNA stability. This, however, is

not the case (Fig. 1B), suggesting that the differences in *pel* expression are likely due to

altered activity of the promoter. Because the RNA-binding protein RsmA is not a

transcription factor, we tested the possibility of RsmA affecting expression of a known

transcriptional regulator of *pel*, namely FleQ. As a well-characterized transcriptional

activator of *pel* (Hickman and Harwood, 2008;Baraquet et al., 2012), potential RsmA-

mediated alterations in expression of FleQ would be anticipated to affect transcription of

the *pel* genes. Using a similar approach as described for *pel* above, we found that a

transcriptional reporter fusion of *fleQ* gave lower levels in the absence of RsmA (Fig. 1C).

This is consistent with previous data showing *pel* transcript levels are up-regulated in Δ*fleQ*

backgrounds (Hickman and Harwood, 2008).

The data above could lend themselves to the interpretation that RsmA serves as a positive

regulator of *fleQ*. While RsmA is more commonly known to be a translational repressor,

this possibility is not without precedence. Examples of RsmA/CsrA proteins stimulating

translation of target mRNAs include *phz2* and *moaA* in *P. aeruginosa* (Patterson-Fortin et

al., 2013;Ren et al., 2014) and *flhDC* in *E. coli* (Yakhnin et al., 2013). While *P. aeruginosa*

FleQ and *E. coli* FlhD$_4$C$_2$ share no sequence or mechanistic similarities, they are functional

counterparts, with both being class I master regulators of flagellar biosynthesis in their

respective organisms (Dasgupta et al., 2003). In addition, flagellar motility was previously

found to be positively controlled by RsmA in *P. aeruginosa* (Heurlier et al., 2004), while in

E. coli, CsrA has been shown to up-regulate flagellar gene expression through protecting

flhDC mRNA from degradation (Yakhnin et al., 2013). It follows that if RsmA regulated

fleQ in an analogous manner, *fleQ* mRNAs would be hyper-stabilized by RsmA and thus,

less stable in the null mutant. However, no evidence of altered mRNA turnover rate was

found between WT and Δ*rsmA* strains (Fig. 1D). Therefore, we hypothesized that the direct

action of RsmA may lie even further upstream in a regulatory cascade that involved control

of *fleQ* by Vfr.

Vfr acts as a transcriptional repressor of the *fleQ* promoter (Dasgupta et al., 2002).

Therefore, to be consistent with the above phenotypes, RsmA would have to serve as a

repressor of *vfr*. In line with this idea, comparison of *vfr* transcriptional reporter activities in

WT and Δ*rsmA* strains revealed up-regulation of *vfr* transcripts in the absence of RsmA

(Fig. 2A). To ensure that the regulatory cascade functioned as anticipated, we tested the

prediction that over-expression of Vfr should ultimately result in increased activity of the

pel transcriptional reporter. This is, indeed, the case (Fig. 2B). In contrast to *pel* (Fig. 1B)

and *fleQ* (Fig. 1D), the transcript stability of *vfr* is significantly different between WT and

Δ*rsmA* strains, such that *vfr* mRNAs are more stable in the absence of RsmA (Fig. 2C).

These results indicate that RsmA affects *vfr* mRNA levels by mediating changes in

turnover rates.

RsmA regulation of Vfr alters swimming and twitching motility.

P. aeruginosa uses at least two major modes of motility: flagellar-driven swimming

motility through liquid and surface 'twitching' motility using Type IV pilus (Burrows,

2012;Kazmierczak et al., 2015). Bacterial migration is an important aspect during *P.*

aeruginosa biofilm development as initial attachment to a surface is thought to require

flagella and mature microcolony formation has been attributed to Type IV pili functions

(O'Toole and Kolter, 1998;Klausen et al., 2003). Thus, fine-tuned regulatory control of

motility and biofilm genes is a necessity for effective *P. aeruginosa* adaptation, particularly

when switching between motile and sessile lifestyles.

Because FleQ is a class I master regulator of flagellar genes (Dasgupta et al., 2003) and

RsmA and Vfr lie upstream of FleQ in the regulatory cascade, lack of either protein should

predictively impact flagellar motility, but in opposite ways. As would be predicted, Δ*rsmA*

and Vfr over-expressing strains have reduced flagellar (swimming) motilities (Fig. 3A).

Motilities of Δ*vfr* and RsmA over-expressing strain are comparable to the WT and vector

control (VC) strains respectively probably because de-repression of flagellar genes do not

necessarily lead to functional hyper-motility.

Mutants in *vfr* have previously been documented to be defective in Type IV pilus-

dependent twitching motility (Whitchurch et al., 1996;Beatson et al., 2002) due to Vfr's

positive regulation of the AlgZR two-component system (Luo et al., 2015;Pritchett et al.,

2015) required for expression of the Type IV pili genes (Lizewski et al., 2004;Belete et al.,

2008). As shown in Fig. 3B, Δ*vfr* and RsmA over-expressing strains were both defective in

twitching motility. Motilities of Δ*rsmA* and Vfr over-expressing strain are comparable to

the WT and VC strains respectively probably because of similar reasons discussed with

swimming motility above.

Hfq is required for RsmA to bind to *vfr* mRNA.

Given the evidence that RsmA may post-transcriptionally regulate *vfr*, we next examined

the binding properties of RsmA to *vfr* mRNA. *P. aeruginosa* RsmA has a binding

specificity for the CANGGAYG consensus sequence of its target mRNA (Schulmeyer et

al., 2016), which is similar to the *E. coli* CsrA consensus sequences RUACARGGAUGU

generated by SELEX (Dubey et al., 2005) and AUGGAUG generated by CLIP-Seq (Potts

et al., 2017). The transcriptional start site for *vfr* lies 153 bases upstream of the translational

start site (Fuchs et al., 2010). Inspection of the *vfr* leader sequence identified one putative

RsmA-binding site overlapping the likely GGGA Shine-Dalgarno sequence (Fig. 4A).

Purified RsmA protein directly binds *psl* mRNA (Irie et al., 2010) in RNA electrophoretic

mobility shift assays (EMSA). However, no RsmA-binding was observed in similar assays

with *vfr* RNA spanning the leader sequence (Fig. 4B). This result led us to consider the

possibility that RsmA may require another factor to bind the *vfr* mRNA. RNA-binding

chaperone protein Hfq was raised as a candidate for three reasons. First, the small RNA

RsmY, which binds multiple RsmA proteins at high affinity to sequester and relieve RsmA-

bound mRNAs (Kay et al., 2006), also has the capacity to be bound by Hfq (Sorger-

Domenigg et al., 2007), although it was unclear from their study whether co-binding of

both proteins occurred. Second, Hfq was recently identified to bind to *vfr* mRNA by global

ChIP-Seq analyses (Kambara et al., 2018). Third, *in silico* analysis of the *vfr* leader

sequence revealed two potential Hfq binding sites upstream of the putative RsmA-binding

site (Fig. 4A).

Biochemical studies, primarily of *E. coli* Hfq, have shown that this hexameric protein

complex has at least four regions which can all be involved in RNA-binding: the proximal

face, the distal face, the rim, and the C-terminal tail (Updegrove et al., 2016). *P. aeruginosa*

and *P. putida* Hfq have shortened C-termini compared to *E. coli* Hfq and may lack C-

terminal tail-binding altogether (Milojevic et al., 2013). The proximal site preferentially

binds to U-rich RNA sequence (Møller et al., 2002), the distal site binds to A-rich $(ARN)_n$

triplet repeats (Link et al., 2009), while the rim associates with UA-rich regions (Schu et

al., 2015). The two potential Hfq-binding sites found within the *vfr* leader sequence (Fig.

4A) represent one A-rich ARN repeat region and one U-rich region.

RNA EMSA analyses show that *P. aeruginosa* Hfq binds *vfr* RNA, resulting in three

distinct band shifts (Fig. 4C) indicative of two or more binding sites. Because wild-type *P.*

putida Hfq showed identical binding patterns to *P. aeruginosa* Hfq on *vfr* RNA (Fig. S1A),

we took advantage of previously characterized derivatives of *P. putida* Hfq, namely, a

distal site mutant (Hfq$_{Y25D}$; A-rich ARN-binding deficient) and a proximal site mutant

(Hfq$_{K56A}$; U-rich-binding deficient) (Madhushani et al., 2015) to further analyze Hfq-

binding to the *vfr* RNA. As seen in Fig. S1B, Hfq$_{Y25D}$ only recapitulates the second band

shift while Hfq$_{K56A}$ only recapitulated the first band shift. Notably, the third band shift is

absent with both mutant derivatives of Hfq. We conclude that the first shift is caused by

Hfq-binding to the U-rich region of the *vfr* leader sequence (Hfq-site 2 in Fig. 4A), the

second shift is caused when the ARN repeats are bound by Hfq (Hfq-site 1 in Fig. 4A),

while the third shift occurs when both sites are occupied simultaneously.

Having ascertained that Hfq does bind *vfr* RNA, we next added RsmA and Hfq

simultaneously to the *vfr* RNA. As seen in Fig. 4D, this results in super-shifting of all three

Hfq band shifts in the presence of RsmA, suggesting that binding of Hfq to either of its

target sites is sufficient to allow RmsA-binding. Because the GG doublet within the

consensus sequences for RsmA and CsrA binding are essential for their abilities to bind

cognate target RNAs (Dubey et al., 2005;Irie et al., 2010), we made a CC substitution of

the GG doublet within the predicted RsmA-binding region of *vfr* RNA. No RsmA-mediated

super-shifting of Hfq band shifts was detected with the CC substituted RNA (Fig. 4E).

We infer from the findings above that Hfq-binding is a pre-requisite for RsmA to bind to its

target within *vfr* mRNA. Together with the data in preceding sections, these results lead us

to conclude: 1) that the direct binding of RsmA on *vfr* RNA requires the RNA chaperone

protein Hfq and 2) that RsmA is an indirect regulator of biofilm polysaccharide locus *pel*,

through its direct action on *vfr* propagated through a regulatory cascade from Vfr to FleQ.

DISCUSSION

In this study, we report that *P. aeruginosa* RsmA requires the RNA chaperone Hfq to assist

its binding to *vfr* mRNA to initiate a regulatory cascade that ultimately impacts *pel* biofilm

polysaccharide genes. The requirement for Hfq was unexpected, since all other biochemical

analyses of RsmA/CsrA family members document unaided binding to their target RNAs

(Baker et al., 2002;Dubey et al., 2003;Wang et al., 2005;Lapouge et al., 2007;Irie et al.,

2010;Patterson-Fortin et al., 2013;Yakhnin et al., 2013;Ren et al., 2014).

We identified two independent Hfq-binding sites upstream of the RsmA-binding motif,

which overlaps the ribosome-binding site of *vfr* (Fig. 4A). Based on a mFOLD-predicted

secondary structure (Zuker, 1989;2003), a large hairpin loop base-pairs the identified U-

rich and A-rich Hfq-binding sites, resulting in a dsRNA structure that would predictively

block both RsmA and the ribosome from accessing the mRNA (Fig. 5A). As depicted in

Fig. 5B, the simplest model that we propose from our findings is that Hfq-binding is

required to open the stem-loop to expose the RsmA-binding site. Subsequent binding of

RsmA would directly block the Shine-Dalgarno sequence, preventing ribosome access and

thereby inhibiting translation. We also provide evidence that *vfr* mRNA is hyper-stable in

the absence of RsmA (Fig. 2C), indicating that Hfq- and RsmA-bound *vfr* mRNA may

undergo more rapid degradation. It was previously suggested that Hfq may be able to

directly recruit RNase E for the degradation of bound RNA in *E. coli* (Morita et al., 2005),

and it is possible that a similar mechanism causes *vfr* mRNA instability.

Due to its histidine-rich C-terminus, *E. coli* Hfq is a common contaminant of His-tagged

proteins purified by nickel affinity chromatography after over-expression in *E. coli*, and

visually undetectable (on SDS-PAGE) Hfq levels can critically influence *in vitro* analyses

of RNA-binding (Milojevic et al., 2013;Moreno et al., 2015). Because the proteins used in

this study were purified from over-expressing them in either *P. aeruginosa* (RsmA) or a

Hfq null strain of *E. coli* (Hfq proteins), our analyses were not complicated by this issue.

However, a large majority of analyses performed in past publications use C-terminal His-

tagged RsmA/CsrA proteins derived from over-expression in *E. coli,* raising the possibility

of *E. coli* native Hfq contamination. Given that Hfq binds *vfr* RNA with apparent high

affinity (Figs. 4C and S1), it is plausible that Hfq-assisted RsmA-binding may be prevalent.

Although such a possibility would not alter major conclusions and remains to be

experimentally verified, it reveals provocative observations that comparison of

transcriptomic studies done for RsmA, Hfq, and Vfr regulons in *P. aeruginosa* identify

numerous overlaps of genes that were differentially expressed between WT and the

respective null mutants (Wolfgang et al., 2003;Goodman et al., 2004;Burrowes et al.,

2006;Brencic and Lory, 2009;Sonnleitner et al., 2018). Overlaps were also seen in *E. coli*

between the Hfq and CsrA regulons (Potts et al., 2017), suggesting that interplays between

the two RNA-binding regulators is not unique to *P. aeruginosa.*

A diverse range of phenotypes have been associated with RsmA functions in *P. aeruginosa*

(Vakulskas et al., 2015), but phenotypic observations do not distinguish direct regulatory

activities from indirect effects through regulatory cascades. Our identification of Hfq and

RsmA action on *vfr* mRNA resulted from an initial aim to determine whether RsmA

directly targeted *pel* RNA, as is the case for *psl* RNA (Irie et al., 2010). Our backtracking to

trace the ultimate cause of RsmA effects on *pel* expression revealed a regulatory pathway

originating from Vfr to the transcriptional regulator FleQ and from there to *pel* (Fig. 5C).

This work places the global post-transcriptional regulators RsmA and Hfq as a hub at the

center of upstream and downstream regulatory pathways and thus provides a conceptually

new framework to evaluate and interpret past work. The two component GacS/GacA

system lies at the top of the hierarchy that controls RsmA-mediated regulation. Although

the environmental cue(s) for activation is still unknown, GacS activity is modulated

through interactions with RetS and LadS (Goodman et al., 2009;Kong et al.,

2013;Chambonnier et al., 2016). GacA, in turn controls the production of the small RNAs –

rsmY and *rsmZ* – which possess multiple high affinity binding sites to titrate RsmA away

from target mRNAs (Heurlier et al., 2004;Sonnleitner et al., 2006). In light of a report

indicating that Hfq also binds to *rsmY* (Sorger-Domenigg et al., 2007), it appears plausible

that these small RNAs may also require Hfq to assist their functions.

In addition to biofilm-associated phenotypes, there are numerous other processes associated

with RsmA in *P. aeruginosa*, including motility (Fig. 3) and acute virulence (Mulcahy et

al., 2006;Intile et al., 2014). Many of these 'RsmA-regulated factors and phenotypes' are

likely to be indirectly regulated through Vfr- and/or FleQ-initiated regulatory cascades.

Based on our findings, it will be crucial to determine and distinguish direct versus indirect

regulatory routes to gain a greater understanding of the RsmA regulon.

 MATERIALS AND METHODS

Bacterial strains and growth conditions

Table S1 lists the bacterial strains used in this study. *E. coli* and *P. aeruginosa* strains were grown in lysogeny broth (LB) Lennox composition at 37°C unless specified otherwise. VBMM citrate medium (Hoang and Schweizer, 1997) was used for selecting *P. aeruginosa* post-conjugation. For *E. coli*, the following antibiotics concentrations were used: 50 μg·ml^{-1} carbenicillin, 10 μg·ml^{-1} gentamicin, and 10 μg·ml^{-1} tetracycline. For *P. aeruginosa* strains: 300 μg·ml^{-1} carbenicillin, 100 μg·ml^{-1} gentamicin, and 100 μg·ml^{-1} tetracycline were used. Sucrose counter-selection for plasmids carrying the *sacB* gene used in *P. aeruginosa* strain constructions was performed by streaking colonies on LB agar (no salt) supplemented with 10% w/v sucrose. Plates were incubated at 30°C for 24 hours, after which the counter-selected colonies were confirmed for the loss of antibiotic resistance and mutations confirmed by PCR for double-cross-over genomic mutants.

Strain constructions

Transcriptional fusion constructs were generated in single copy on the chromosome of *P. aeruginosa* strains via integration into the chromosomal *attB* site as previously published (Irie et al., 2010). In brief, promoter regions were PCR amplified using oligonucleotides in Table S2. PCR products were ligated between the EcoRI and BamHI sites of mini-CTX *lacZ* (Fig. S2). These plasmids were then introduced into *P. aeruginosa* by conjugation. After double site recombination, plasmid backbones were removed by FLP recombinase, and then strains cured of the pFLP2 plasmid by sucrose counter-selection.

β-galactosidase assays

Quantitative β-galactosidase activities were assayed using Galacto-Light Plus kit (Thermo-Fisher) as previously published (Lequette et al., 2006). All *P. aeruginosa* cultures were grown in VBMM citrate at 37°C to exponential phase, and lysed using chloroform as previously described (Irie et al., 2010). β-galactosidase activity units were normalized to

total proteins per ml as determined using Bradford assay reagents (Bio-Rad). Assays were

performed in biological triplicates.

Quantitative real-time PCR and RNA stability analyses

Real-time PCR was performed as previously described (Irie et al., 2012), using the

oligonucleotides listed in Table S2. For RNA stability experiments, exponential phase *P.*

aeruginosa cultured in VBMM citrate at 37°C were treated with 200 μg·ml^{-1} rifampicin

(Lory, 1986). RNAs were extracted from 1 ml of the cultures at various time points as

previously described (Irie et al., 2010) after the addition of rifampicin. RNA extractions

were performed using the RNeasy Mini Kit (Qiagen) after treating the harvested cells with

RNAprotect (Qiagen). Genomic DNA was removed using DNase I (Promega) and removal

confirmed by PCR using primers designed against the *rplU* gene (Table S2). SuperScript III

First-Strand Synthesis (Invitrogen) was used to synthesize cDNA as per manufacturer's

protocol using random hexamers. Quantitative real-time PCR was performed using SYBR

Green PCR Master Mix (Thermo/Applied Biosystems). The internal control gene used was

ampR. All experiments were done in biological quadruplicates.

Motility assays

Motility assays were performed essentially as previously described (Shrout et al., 2006).

For swimming motility, LB-Lennox plates containing 0.3% Bacto agar were inoculated

with overnight cultures with a sterile inoculation needle, ensuring the needle tip was

inserted approximately halfway into the agar but not to the plastic petri dish bottom, and

incubated for 24 hours at 30°C. Swim ring diameters were measured for quantitation. For

twitching motility, LB-Lennox plates containing 0.5% Bacto agar were inoculated by

inserting a sterile inoculation needle through the agar until it touched the plastic bottom.

Plates were incubated for 72 hours at 30°C. Subsequently, the agar was peeled off, and the

plates were stained with 1% w/v crystal violet to visualize the twitching zone diameters

prior to measurement (Déziel et al., 2001). All motility experiments were performed in at

least biological quadruplicates each in technical quadruplicates.

Protein expression and purification

Proteins were purified by previously established protocols. Briefly, over-expressed RsmA-His$_6$ was purified from *P. aeruginosa* PAO1 bearing pUCP18::*rsmA*-His$_6$ (Irie et al., 2010), while all Hfq-His$_6$ proteins were purified from a Hfq null derivative of *E. coli* BL21(DE3) (Madhushani et al., 2015) carrying P$_{T7}$ expression plasmids (pVI2344-2346 or pVI2357).

RsmA and Hfq over-expressing strains were grown in 2× YT broth to stationary or exponential phases, respectively. Cell pellets were re-suspended in lysis buffer (buffer A: 0.3 M NaCl, 20 mM Tris-HCl pH 8, 5 mM imidazole) supplemented with 1 tablet/L cOmplete^{TM} EDTA-free protease inhibitor tablet (Sigma). Cells were lysed by Stansted Fluid Power SPCH ultra high-pressure cell disrupter/homogenizer. Cell debris were removed by centrifugation (35,000 rpm, 40 minutes, 4^{o}C) and the supernatants loaded onto 1 mL HisTRAP HP (GE Healthcare) columns equilibrated in lysis buffer (buffer A). The column was washed with high salt buffer (buffer B: 5 M NaCl, 20 mM Tris-HCl pH 8, 5 mM imidazole) to remove RNA contamination and the proteins subsequently eluted with the linear gradient of imidazole (25 mM to 750 mM) in buffer C (buffer C: 0.5 M NaCl, 20 mM Tris-HCl pH 8). Fractions containing the desired proteins were pooled. For RsmA-His$_6$, an additional ion exchange step using a 5 ml HiTrap SP HP column (GE Healthcare) equilibrated with buffer D (0.1 M NaCl, 20 mM Tris-HCl pH 8) was performed, with RmsA subsequently being eluted with 1 M NaCl.

All proteins were dialyzed across Slide-a-lyzer 10,000 MWCO cassettes (Thermo Fisher) against 2 lots of 2 L dialysis buffer (40 mM Tris-HCl pH 8, 600 mM NaCl, 2 mM EDTA) at 4^{o}C for 2 days prior concentration using 10 MWCO Centricons (Amicon). Protein concentrations were determined using the BCA Protein Assay Kit (Pierce/Fisher) and purity confirmed by SDS-PAGE. Protein were stored at -20°C in final storage buffer (20 mM Tris-HCl pH 8, 300 mM NaCl, 1 mM EDTA, 40% glycerol) until use.

RNA synthesis

Wild-type and mutant *vfr* mRNAs were generated using Ambion's MEGAscript kit as

recommended for *in vitro* transcription from the P_{T7} promoter of linearized plasmids.

Reactions (total 40 µl) containing 1 µg of EcoRV-linearized pVI2358 or pVI2359, were

incubated for 6 hours prior to 15 minutes DNase I treatment. Reactions were terminated by

adding 230 µl RNase-free H_2O and 30 µl stop solution (5 M ammonium acetate; 100 mM

EDTA). The resulting 233 nt RNAs (encompassing co-ordinates -106 to +116 relative to

the A of the initiation codon of *vfr*) were extracted twice with phenol:chloroform:IAA

(25:24:1) and once with chloroform prior to precipitation and resuspension in 50 µl RNase-

free H_2O.

RNA electrophoretic mobility shift assays (EMSAs)

Reactions (total 10 µl) contained 32 nM *in vitro* transcribed RNA with the indicated

concentrations of RsmA and/or Hfq. RsmA and Hfq molarities given here are for the

dimeric and hexameric complexes respectively. RNA was first heated at $80^{o}C$ for 5 min and

then immediately chilled in an ice-water bath. Additional final reaction mixture

components were as follows: 10 mM HEPES pH 7.9, 2 mM $MgCl_2$, 90 ng yeast total RNA

(Fisher), 4U RNasin, and 35 mM KCl. Binding reactions were performed at $20^{o}C$ for 40

min prior to adding 2.5 µl loading buffer (40% sucrose). Reactions were placed on ice and

analyzed on 5% TBE Criterion pre-cast gels (Bio-Rad) with electrophoresis at 100V for

≈130 min at $4^{o}C$. Gels were stained at room temperature with 1:10,000 TBE-diluted SYBR

Gold (Fisher) in a light-protected container with agitation. Images were captured using

Typhoon FLA 9500 (GE Healthcare).

Statistics

Presented data are the mean with standard deviations of the data collected. Student's *t*-test

using the GraphPad Prism software was used to calculate *p*-values.

ACKNOWLEDGEMENTS

Plasmid pMMBGW:RBS-*vfr* was a kind gift from Joe J. Harrison. YI would like to thank

Matt Parsek for his more than a decade of scientific and moral support that led to the

culmination of this project. The manuscript partially overlaps with a preprint at BioRxiv

(Irie Y *et al.*, 2019).

AUTHOR CONTRIBUTIONS

YI and VS designed and performed the experiments and analyzed the data; VM and VH

assisted in purification of the RsmA and Hfq proteins. YI and VS wrote the manuscript with

input from VH and VM.

FUNDING INFORMATION

This work was supported by the J C Kempe & S M Kempe Foundation (JCK-1523 to VS)

and The Swedish Research Council (grant numbers 2016-02047 to VS).

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interests.

REFERENCES

Baker, C.S., Eöry, L.A., Yakhnin, H., Mercante, J., Romeo, T., and Babitzke, P. (2007). CsrA inhibits translation initiation of *Escherichia coli hfq* by binding to a single site overlapping the Shine-Dalgarno sequence. *J. Bacteriol.* 189**,** 5472-5481.

Baker, C.S., Morozov, I., Suzuki, K., Romeo, T., and Babitzke, P. (2002). CsrA regulates glycogen biosynthesis by preventing translation of *glgC* in *Escherichia coli*. *Mol. Microbiol.* 44**,** 1599-1610.

Bandyra, K.J., and Luisi, B.F. (2013). Licensing and due process in the turnover of bacterial RNA. *RNA Biol.* 10**,** 627-635.

Baraquet, C., Murakami, K., Parsek, M.R., and Harwood, C.S. (2012). The FleQ protein from *Pseudomonas aeruginosa* functions as both a repressor and an activator to control gene expression from the *pel* operon promoter in response to c-di-GMP. *Nucleic Acids Res.* 40**,** 7207-7218.

Beatson, S.A., Whitchurch, C.B., Sargent, J.L., Levesque, R.C., and Mattick, J.S. (2002). Differential regulation of twitching motility and elastase production by Vfr in *Pseudomonas aeruginosa*. *J. Bacteriol.* 184**,** 3605-3613.

Belete, B., Lu, H., and Wozniak, D.J. (2008). *Pseudomonas aeruginosa* AlgR regulates type IV pilus biosynthesis by activating transcription of the *fimU-pilVWXY1Y2E* operon. *J. Bacteriol.* 190**,** 2023-2030.

Brencic, A., and Lory, S. (2009). Determination of the regulon and identification of novel mRNA targets of *Pseudomonas aeruginosa* RsmA. *Mol. Microbiol.* 72**,** 612-632.

Burrowes, E., Baysse, C., Adams, C., and O'gara, F. (2006). Influence of the regulatory protein RsmA on cellular functions in *Pseudomonas aeruginosa* PAO1, as revealed by transcriptome analysis. *Microbiology* 152**,** 405-418.

Burrows, L.L. (2012). *Pseudomonas aeruginosa* twitching motility: type IV pili in action. *Annu. Rev. Microbiol.* 66**,** 493-520.

Chambonnier, G., Roux, L., Redelberger, D., Fadel, F., Filloux, A., Sivaneson, M., De Bentzmann, S., and Bordi, C. (2016). The Hybrid Histidine Kinase LadS Forms a Multicomponent Signal Transduction System with the GacS/GacA Two-Component System in *Pseudomonas aeruginosa*. *PLoS Genet.* 12**,** e1006032.

Dasgupta, N., Ferrell, E.P., Kanack, K.J., West, S.E., and Ramphal, R. (2002). *fleQ*, the gene encoding the major flagellar regulator of *Pseudomonas aeruginosa*, is σ^{70} dependent and is downregulated by Vfr, a homolog of *Escherichia coli* cyclic AMP receptor protein. *J. Bacteriol.* 184**,** 5240-5250.

Dasgupta, N., Wolfgang, M.C., Goodman, A.L., Arora, S.K., Jyot, J., Lory, S., and Ramphal, R. (2003). A four-tiered transcriptional regulatory circuit controls flagellar biogenesis in *Pseudomonas aeruginosa*. *Mol. Microbiol.* 50**,** 809-824.

Déziel, E., Comeau, Y., and Villemur, R. (2001). Initiation of biofilm formation by *Pseudomonas aeruginosa* 57RP correlates with emergence of hyperpiliated and highly adherent phenotypic variants deficient in swimming, swarming, and twitching motilities. *J. Bacteriol.* 183**,** 1195-1204.

Dubey, A.K., Baker, C.S., Romeo, T., and Babitzke, P. (2005). RNA sequence and
secondary structure participate in high-affinity CsrA-RNA interaction. *Rna* 11**,**
1579-1587.
Dubey, A.K., Baker, C.S., Suzuki, K., Jones, A.D., Pandit, P., Romeo, T., and Babitzke, P.
(2003). CsrA regulates translation of the *Escherichia coli* carbon starvation gene,
cstA, by blocking ribosome access to the cstA transcript. *J. Bacteriol.* 185**,** 4450-
4460.
Fuchs, E.L., Brutinel, E.D., Jones, A.K., Fulcher, N.B., Urbanowski, M.L., Yahr, T.L., and
Wolfgang, M.C. (2010). The *Pseudomonas aeruginosa* Vfr regulator controls global
virulence factor expression through cyclic AMP-dependent and -independent
mechanisms. *J. Bacteriol.* 192**,** 3553-3564.
Gellatly, S.L., and Hancock, R.E. (2013). *Pseudomonas aeruginosa*: new insights into
pathogenesis and host defenses. *Pathog. Dis.* 67**,** 159-173.
Gilbert, K.B., Kim, T.H., Gupta, R., Greenberg, E.P., and Schuster, M. (2009). Global
position analysis of the *Pseudomonas aeruginosa* quorum-sensing transcription
factor LasR. *Mol. Microbiol.* 73**,** 1072-1085.
Gooderham, W.J., and Hancock, R.E. (2009). Regulation of virulence and antibiotic
resistance by two-component regulatory systems in *Pseudomonas aeruginosa*.
FEMS Microbiol. Rev. 33**,** 279-294.
Goodman, A.L., Kulasekara, B., Rietsch, A., Boyd, D., Smith, R.S., and Lory, S. (2004). A
signaling network reciprocally regulates genes associated with acute infection and
chronic persistence in *Pseudomonas aeruginosa*. *Dev. Cell* 7**,** 745-754.
Goodman, A.L., Merighi, M., Hyodo, M., Ventre, I., Filloux, A., and Lory, S. (2009).
Direct interaction between sensor kinase proteins mediates acute and chronic
disease phenotypes in a bacterial pathogen. *Genes Dev.* 23**,** 249-259.
Gutiérrez, P., Li, Y., Osborne, M.J., Pomerantseva, E., Liu, Q., and Gehring, K. (2005).
Solution structure of the carbon storage regulator protein CsrA from *Escherichia*
coli. *J. Bacteriol.* 187**,** 3496-3501.
Heurlier, K., Williams, F., Heeb, S., Dormond, C., Pessi, G., Singer, D., Camará, M.,
Williams, P., and Haas, D. (2004). Positive control of swarming, rhamnolipid
synthesis, and lipase production by the posttranscriptional RsmA/RsmZ system in
Pseudomonas aeruginosa PAO1. *J. Bacteriol.* 186**,** 2936-2945.
Hickman, J.W., and Harwood, C.S. (2008). Identification of FleQ from *Pseudomonas*
aeruginosa as a c-di-GMP-responsive transcription factor. *Mol. Microbiol.* 69**,** 376-
389.
Hoang, T.T., and Schweizer, H.P. (1997). Fatty acid biosynthesis in *Pseudomonas*
aeruginosa: cloning and characterization of the *fabAB* operon encoding β-
hydroxyacyl-acyl carrier protein dehydratase (FabA) and β-ketoacyl-acyl carrier
protein synthase I (FabB). *J. Bacteriol.* 179**,** 5326-5332.
Holmqvist, E., Wright, P.R., Li, L., Bischler, T., Barquist, L., Reinhardt, R., Backofen, R.,
and Vogel, J. (2016). Global RNA recognition patterns of post-transcriptional
regulators Hfq and CsrA revealed by UV crosslinking *in vivo*. *EMBO J.* 35**,** 991-
1011.

Intile, P.J., Diaz, M.R., Urbanowski, M.L., Wolfgang, M.C., and Yahr, T.L. (2014). The
AlgZR two-component system recalibrates the RsmAYZ posttranscriptional
regulatory system to inhibit expression of the *Pseudomonas aeruginosa* type III
secretion system. *J. Bacteriol.* 196**,** 357-366.
Irie, Y., Borlee, B.R., O'connor, J.R., Hill, P.J., Harwood, C.S., Wozniak, D.J., and Parsek,
M.R. (2012). Self-produced exopolysaccharide is a signal that stimulates biofilm
formation in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. U. S. A.* 109**,** 20632-
20636.
Irie, Y., Starkey, M., Edwards, A.N., Wozniak, D.J., Romeo, T., and Parsek, M.R. (2010).
Pseudomonas aeruginosa biofilm matrix polysaccharide Psl is regulated
transcriptionally by RpoS and post-transcriptionally by RsmA. *Mol. Microbiol.* 78**,**
158-172.
Kambara, T.K., Ramsey, K.M., and Dove, S.L. (2018). Pervasive Targeting of Nascent
Transcripts by Hfq. *Cell Rep.* 23**,** 1543-1552.
Kay, E., Humair, B., Dénervaud, V., Riedel, K., Spahr, S., Eberl, L., Valverde, C., and
Haas, D. (2006). Two GacA-dependent small RNAs modulate the quorum-sensing
response in *Pseudomonas aeruginosa*. *J. Bacteriol.* 188**,** 6026-6033.
Kazmierczak, B.I., Schniederberend, M., and Jain, R. (2015). Cross-regulation of
Pseudomonas motility systems: the intimate relationship between flagella, pili and
virulence. *Curr. Opin. Microbiol.* 28**,** 78-82.
Khare, A., and Tavazoie, S. (2015). Multifactorial Competition and Resistance in a Two-
Species Bacterial System. *PLoS Genet.* 11**,** e1005715.
Klausen, M., Heydorn, A., Ragas, P., Lambertsen, L., Aaes-Jørgensen, A., Molin, S., and
Tolker-Nielsen, T. (2003). Biofilm formation by *Pseudomonas aeruginosa* wild
type, flagella and type IV pili mutants. *Mol. Microbiol.* 48**,** 1511-1524.
Kong, W., Chen, L., Zhao, J., Shen, T., Surette, M.G., Shen, L., and Duan, K. (2013).
Hybrid sensor kinase PA1611 in *Pseudomonas aeruginosa* regulates transitions
between acute and chronic infection through direct interaction with RetS. *Mol.*
Microbiol. 88**,** 784-797.
Lapouge, K., Sineva, E., Lindell, M., Starke, K., Baker, C.S., Babitzke, P., and Haas, D.
(2007). Mechanism of *hcnA* mRNA recognition in the Gac/Rsm signal transduction
pathway of *Pseudomonas fluorescens*. *Mol. Microbiol.* 66**,** 341-356.
Lawhon, S.D., Frye, J.G., Suyemoto, M., Porwollik, S., Mcclelland, M., and Altier, C.
(2003). Global regulation by CsrA in *Salmonella* typhimurium. *Mol. Microbiol.* 48**,**
1633-1645.
Lequette, Y., Lee, J.-H., Ledgham, F., Lazdunski, A., and Greenberg, E.P. (2006). A
distinct QscR regulon in the *Pseudomonas aeruginosa* quorum-sensing circuit. *J.*
Bacteriol. 188**,** 3365-3370.
Link, T.M., Valentin-Hansen, P., and Brennan, R.G. (2009). Structure of *Escherichia coli*
Hfq bound to polyriboadenylate RNA. *Proc. Natl. Acad. Sci. U. S. A.* 106**,** 19292-
19297.
Lizewski, S.E., Schurr, J.R., Jackson, D.W., Frisk, A., Carterson, A.J., and Schurr, M.J.
(2004). Identification of AlgR-regulated genes in *Pseudomonas aeruginosa* by use
of microarray analysis. *J. Bacteriol.* 186**,** 5672-5684.

Lory, S. (1986). Effect of iron on accumulation of exotoxin A-specific mRNA in *Pseudomonas aeruginosa*. *J. Bacteriol.* 168**,** 1451-1456.

Luo, Y., Zhao, K., Baker, A.E., Kuchma, S.L., Coggan, K.A., Wolfgang, M.C., Wong, G.C., and O'toole, G.A. (2015). A hierarchical cascade of second messengers regulates *Pseudomonas aeruginosa* surface behaviors. *MBio* 6.

Madhushani, A., Del Peso-Santos, T., Moreno, R., Rojo, F., and Shingler, V. (2015). Transcriptional and translational control through the 5'-leader region of the *dmpR* master regulatory gene of phenol metabolism. *Environ. Microbiol.* 17**,** 119-133.

Milojevic, T., Grishkovskaya, I., Sonnleitner, E., Djinovic-Carugo, K., and Bläsi, U. (2013). The *Pseudomonas aeruginosa* catabolite repression control protein Crc is devoid of RNA binding activity. *PLoS One* 8**,** e64609.

Møller, T., Franch, T., Hojrup, P., Keene, D.R., Bachinger, H.P., Brennan, R.G., and Valentin-Hansen, P. (2002). Hfq: a bacterial Sm-like protein that mediates RNA-RNA interaction. *Mol. Cell.* 9**,** 23-30.

Moreno, R., Hernández-Arranz, S., La Rosa, R., Yuste, L., Madhushani, A., Shingler, V., and Rojo, F. (2015). The Crc and Hfq proteins of *Pseudomonas putida* cooperate in catabolite repression and formation of ribonucleic acid complexes with specific target motifs. *Environ. Microbiol.* 17**,** 105-118.

Morita, T., Maki, K., and Aiba, H. (2005). RNase E-based ribonucleoprotein complexes: mechanical basis of mRNA destabilization mediated by bacterial noncoding RNAs. *Genes Dev.* 19**,** 2176-2186.

Mulcahy, H., O'callaghan, J., O'grady, E.P., Adams, C., and O'gara, F. (2006). The posttranscriptional regulator RsmA plays a role in the interaction between *Pseudomonas aeruginosa* and human airway epithelial cells by positively regulating the type III secretion system. *Infect. Immun.* 74**,** 3012-3015.

Mulcahy, H., O'callaghan, J., O'grady, E.P., Maciá, M.D., Borrell, N., Gómez, C., Casey, P.G., Hill, C., Adams, C., Gahan, C.G., Oliver, A., and O'gara, F. (2008). *Pseudomonas aeruginosa* RsmA plays an important role during murine infection by influencing colonization, virulence, persistence, and pulmonary inflammation. *Infect. Immun.* 76**,** 632-638.

O'grady, E.P., Mulcahy, H., O'callaghan, J., Adams, C., and O'gara, F. (2006). *Pseudomonas aeruginosa* infection of airway epithelial cells modulates expression of Kruppel-like factors 2 and 6 via RsmA-mediated regulation of type III exoenzymes S and Y. *Infect. Immun.* 74**,** 5893-5902.

O'toole, G.A., and Kolter, R. (1998). Flagellar and twitching motility are necessary for *Pseudomonas aeruginosa* biofilm development. *Mol. Microbiol.* 30**,** 295-304.

Pannuri, A., Yakhnin, H., Vakulskas, C.A., Edwards, A.N., Babitzke, P., and Romeo, T. (2012). Translational repression of NhaR, a novel pathway for multi-tier regulation of biofilm circuitry by CsrA. *J. Bacteriol.* 194**,** 79-89.

Patterson-Fortin, L.M., Vakulskas, C.A., Yakhnin, H., Babitzke, P., and Romeo, T. (2013). Dual posttranscriptional regulation via a cofactor-responsive mRNA leader. *J. Mol. Biol.* 425**,** 3662-3677.

Pessi, G., Williams, F., Hindle, Z., Heurlier, K., Holden, M.T., Cámara, M., Haas, D., and Williams, P. (2001). The global posttranscriptional regulator RsmA modulates

production of virulence determinants and *N*-acylhomoserine lactones in *Pseudomonas aeruginosa*. *J. Bacteriol.* 183**,** 6676-6683.

Potts, A.H., Vakulskas, C.A., Pannuri, A., Yakhnin, H., Babitzke, P., and Romeo, T. (2017). Global role of the bacterial post-transcriptional regulator CsrA revealed by integrated transcriptomics. *Nat. Commun.* 8**,** 1596.

Pritchett, C.L., Little, A.S., Okkotsu, Y., Frisk, A., Cody, W.L., Covey, C.R., and Schurr, M.J. (2015). Expression analysis of the *Pseudomonas aeruginosa* AlgZR two-component regulatory system. *J. Bacteriol.* 197**,** 736-748.

Ren, B., Shen, H., Lu, Z.J., Liu, H., and Xu, Y. (2014). The *phzA2-G2* transcript exhibits direct RsmA-mediated activation in *Pseudomonas aeruginosa* M18. *PLoS One* 9**,** e89653.

Rife, C., Schwarzenbacher, R., Mcmullan, D., Abdubek, P., Ambing, E., Axelrod, H., Biorac, T., Canaves, J.M., Chiu, H.J., Deacon, A.M., Didonato, M., Elsliger, M.A., Godzik, A., Grittini, C., Grzechnik, S.K., Hale, J., Hampton, E., Han, G.W., Haugen, J., Hornsby, M., Jaroszewski, L., Klock, H.E., Koesema, E., Kreusch, A., Kuhn, P., Lesley, S.A., Miller, M.D., Moy, K., Nigoghossian, E., Paulsen, J., Quijano, K., Reyes, R., Sims, E., Spraggon, G., Stevens, R.C., Van Den Bedem, H., Velasquez, J., Vincent, J., White, A., Wolf, G., Xu, Q., Hodgson, K.O., Wooley, J., and Wilson, I.A. (2005). Crystal structure of the global regulatory protein CsrA from Pseudomonas putida at 2.05 A resolution reveals a new fold. *Proteins* 61**,** 449-453.

Romeo, T., Vakulskas, C.A., and Babitzke, P. (2013). Post-transcriptional regulation on a global scale: form and function of Csr/Rsm systems. *Environ. Microbiol.* 15**,** 313-324.

Ryder, C., Byrd, M., and Wozniak, D.J. (2007). Role of polysaccharides in *Pseudomonas aeruginosa* biofilm development. *Curr. Opin. Microbiol.* 10**,** 644-648.

Sahr, T., Rusniok, C., Impens, F., Oliva, G., Sismeiro, O., Coppee, J.Y., and Buchrieser, C. (2017). The *Legionella pneumophila* genome evolved to accommodate multiple regulatory mechanisms controlled by the CsrA-system. *PLoS Genet.* 13**,** e1006629.

Sakuragi, Y., and Kolter, R. (2007). Quorum-sensing regulation of the biofilm matrix genes (*pel*) of *Pseudomonas aeruginosa*. *J. Bacteriol.* 189**,** 5383-5386.

Schu, D.J., Zhang, A., Gottesman, S., and Storz, G. (2015). Alternative Hfq-sRNA interaction modes dictate alternative mRNA recognition. *EMBO J.* 34**,** 2557-2573.

Schubert, M., Lapouge, K., Duss, O., Oberstrass, F.C., Jelesarov, I., Haas, D., and Allain, F.H. (2007). Molecular basis of messenger RNA recognition by the specific bacterial repressing clamp RsmA/CsrA. *Nat. Struct. Mol. Biol.* 14**,** 807-813.

Schulmeyer, K.H., Diaz, M.R., Bair, T.B., Sanders, W., Gode, C.J., Laederach, A., Wolfgang, M.C., and Yahr, T.L. (2016). Primary and Secondary Sequence Structure Requirements for Recognition and Discrimination of Target RNAs by *Pseudomonas aeruginosa* RsmA and RsmF. *J. Bacteriol.* 198**,** 2458-2469.

Shrout, J.D., Chopp, D.L., Just, C.L., Hentzer, M., Givskov, M., and Parsek, M.R. (2006). The impact of quorum sensing and swarming motility on *Pseudomonas aeruginosa* biofilm formation is nutritionally conditional. *Mol. Microbiol.* 62**,** 1264-1277.

Sonnleitner, E., Schuster, M., Sorger-Domenigg, T., Greenberg, E.P., and Bläsi, U. (2006).
Hfq-dependent alterations of the transcriptome profile and effects on quorum
sensing in *Pseudomonas aeruginosa*. *Mol. Microbiol.* 59**,** 1542-1558.
Sonnleitner, E., Wulf, A., Campagne, S., Pei, X.Y., Wolfinger, M.T., Forlani, G., Prindl,
K., Abdou, L., Resch, A., Allain, F.H., Luisi, B.F., Urlaub, H., and Bläsi, U. (2018).
Interplay between the catabolite repression control protein Crc, Hfq and RNA in
Hfq-dependent translational regulation in *Pseudomonas aeruginosa*. *Nucleic Acids*
Res. 46**,** 1470-1485.
Sorger-Domenigg, T., Sonnleitner, E., Kaberdin, V.R., and Bläsi, U. (2007). Distinct and
overlapping binding sites of *Pseudomonas aeruginosa* Hfq and RsmA proteins on
the non-coding RNA RsmY. *Biochem. Biophys. Res. Commun.* 352**,** 769-773.
Tashiro, Y., Yawata, Y., Toyofuku, M., Uchiyama, H., and Nomura, N. (2013).
Interspecies interaction between *Pseudomonas aeruginosa* and other
microorganisms. *Microbes Environ.* 28**,** 13-24.
Updegrove, T.B., Zhang, A., and Storz, G. (2016). Hfq: the flexible RNA matchmaker.
Curr. Opin. Microbiol. 30**,** 133-138.
Vakulskas, C.A., Potts, A.H., Babitzke, P., Ahmer, B.M., and Romeo, T. (2015).
Regulation of bacterial virulence by Csr (Rsm) systems. *Microbiol. Mol. Biol. Rev.*
79**,** 193-224.
Wang, X., Dubey, A.K., Suzuki, K., Baker, C.S., Babitzke, P., and Romeo, T. (2005). CsrA
post-transcriptionally represses *pgaABCD*, responsible for synthesis of a biofilm
polysaccharide adhesin of *Escherichia coli*. *Mol. Microbiol.* 56**,** 1648-1663.
Whitchurch, C.B., Alm, R.A., and Mattick, J.S. (1996). The alginate regulator AlgR and an
associated sensor FimS are required for twitching motility in *Pseudomonas*
aeruginosa. *Proc. Natl. Acad. Sci. U. S. A.* 93**,** 9839-9843.
White, D., Hart, M.E., and Romeo, T. (1996). Phylogenetic distribution of the global
regulatory gene *csrA* among eubacteria. *Gene* 182**,** 221-223.
Wolfgang, M.C., Lee, V.T., Gilmore, M.E., and Lory, S. (2003). Coordinate regulation of
bacterial virulence genes by a novel adenylate cyclase-dependent signaling
pathway. *Dev. Cell* 4**,** 253-263.
Yakhnin, A.V., Baker, C.S., Vakulskas, C.A., Yakhnin, H., Berezin, I, Romeo, T., and
Babitzke, P. (2013). CsrA activates *flhDC* expression by protecting *flhDC* mRNA
from RNase E-mediated cleavage. *Mol. Microbiol.* 87**,** 851-866.
Zuker, M. (1989). On finding all suboptimal foldings of an RNA molecule. *Science* 244**,**
48-52.
Zuker, M. (2003). Mfold web server for nucleic acid folding and hybridization prediction.
Nucleic Acids Res. 31**,** 3406-3415.
708

 FIGURE LEGENDS

Fig. 1   RsmA does not directly regulate *pel* nor *fleQ*.

A. Single-copy transcriptional *lacZ* fusion constructs reveal up-regulation of *pel* transcripts

in the Δ*rsmA* background as compared to wild type (WT). Δ*rsmA* strain overexpresses the

PSL polysaccharides, which causes an elevation of intercellular concentration of c-di-GMP

(Irie et al., 2012). Thus, we also tested a Δ*rsmA* Δ*pel* Δ*psl* strain, which has low c-di-GMP

due to the absence of PSL (Irie et al., 2012). Note that *pel* is equally up-regulated in the

Δ*rsmA* Δ*pel* Δ*psl* strain, demonstrating that the *pel* transcriptional phenotype of Δ*rsmA* is

PEL-, PSL-, and c-di-GMP-independent *p = 0.0002; **p = 0.0013.

B. Real-time quantitative PCR (qPCR) at different time points after rifampicin treatment

show indistinguishable stabilities of the *pelA* transcript in the WT and Δ*rsmA* strains.

C. Single-copy transcriptional *lacZ* fusion constructs reveal down-regulation of *fleQ*

transcripts in the Δ*rsmA* background as compared to WT.

D. qPCR at different time points after rifampicin treatment shows indistinguishable

stabilities of the *fleQ* transcript in the WT and Δ*rsmA* strains.

Fig. 2   RsmA directly represses *vfr* by altering the mRNA stability.

A. Single-copy transcriptional *lacZ* fusion constructs reveal up-regulation of *vfr* transcripts

in the Δ*rsmA* background as compared to WT.

B. The single-copy P$_{pelA}$::*lacZ* transcriptional fusion has a higher activity in a Vfr over-

expressing strain compared to the same strain harboring the vector control (VC).

C. qPCR at different time points after rifampicin treatment show that *vfr* mRNA is less

stable in WT as compared to the Δ*rsmA* strain.

Fig. 3   RsmA regulatory cascade indirectly affects flagellar and Type IV pilus-

mediated motilities.

A. Flagellar-mediated swimming motility is decreased for Δ*rsmA* and Vfr over-expressing

strains. VC = vector control. Motility assays for VC, RsmA over-expressing strain, and Vfr

over-expressing strains were performed in the presence of 300 μg/ml carbenicillin to ensure

maintenance of the plasmids.

B. Type IV pilus-mediated twitching motility is reduced for Δ*vfr* compared to WT. Δ*rsmA*

motility is comparable to the WT strain probably because WT motility is maximized in this

assay. Motility assays for VC, RsmA over-expressing strain, and Vfr over-expressing

strains were performed in the presence of 300 μg/ml carbenicillin.

Fig. 4   Direct targeting of *vfr* mRNA by RsmA requires Hfq.

A. The leader sequence of the *vfr* transcript (Fuchs et al., 2010) possesses two putative Hfq-

binding sites (blue boxes) and a RsmA-binding site (green box) overlapping the predicted

Shine-Dalgarno sequence (red italics) upstream of the translational start site (underlined

italics).

B. RNA Electrophoretic mobility shift assay (EMSA) shows that RsmA does not bind to

the *vfr* RNA when RsmA is present alone. Dimeric molar excess relative to RNA; U =

unbound RNA.

C. The *vfr* RNA produces three distinct band shifts (open arrow heads 1, 2, and 3) with

increasing concentrations of Hfq. Hexameric molar excess relative to RNA.

D. The presence of RsmA causes all Hfq-shifted bands to super-shift (filled arrow heads

1+, 2+, and 3+), indicating that RsmA binds to Hfq-bound RNA. Note that the RsmA

concentration used (36✕ molar excess of RNA) is incapable of producing a shift when

present alone (panel B).

E. A GG→CC substitution within the RsmA-binding site abolishes RsmA super-shifts but

does not alter Hfq-binding.

Fig. 5 Model for Hfq-assisted RsmA regulation of *vfr* and its consequences

A. RNA secondary structure of the leader sequence region of *vfr* as predicted by mFOLD

(Zuker, 1989;2003). The locations of *in silico*-detected Hfq- and RsmA-binding sites are

indicated. Note that the RsmA-binding site located at the base of the stem loop structure

overlaps with the Shine-Dalgarno sequence of the ribosome binding site. The AUG start

codon is shown underlined in italics.

B. Schematic illustration of a model for Hfq-assisted RsmA regulation of *vfr*. Upon Hfq-

binding of *vfr* RNA, the stem loop structure shown in panel A is disrupted, exposing the

RsmA-binding site. RsmA-binding then directly blocks the Shine Dalgarno (SD) sequence.

RsmA-binding is depicted as producing a smaller stem-loop structure (Gutiérrez et al.,

2005;Rife et al., 2005;Schubert et al., 2007), preventing ribosome access and thereby

inhibiting translation. *In vivo*, the Hfq- and RsmA-bound mRNAs are presumed to be

subsequently processed for degradation.

C. Simplified overview of the Gac/Rsm pathway that controls RsmA levels and pertinent

parts of its downstream regulon.

Fig. 1

Fig. 2

Fig. 3

A

B

Fig. 4

A

CGCGGUUUACUGGCACACUUCCUGAUCUGCCCGGUUGGGGGGAGCUCUCCGCUGAGCUCC

CGGGCCCUGGCAGGUCUCUCCAUAAGAACUC AAAAAAACA UCGCCUUGUACGUCAGGCAU

Hfq-site 1

Hfq-site 2   RsmA-binding site
A UUCUUUAUU CUUUC*GGGA*CUCGGGC*AUG*→
         Shine-   *Start*
         Dalgarno   *codon*

B dimeric molar excess
RsmA: 0   10× 14× 18×
U

C hexameric molar excess
Hfq: 0   1× 2× 3× 4× 6×
◁3
◁2
◁1
U

D
Hfq: 0   0   2× 2× 4× 4×
RsmA: 0 18× 0 18× 0 18×
3+
◁
2+
◁
1+
◁
U

Wild-type RsmA-binding mRNA
−UCUUUC*GGGA*CUCGG GC*AUG*−

E
Hfq: 0   0   2× 2× 4× 4×
RsmA: 0 18× 0 18× 0 18×
◁3
◁2
◁1
U

Mutant RsmA-binding site mRNA
−UCUUUC*GCCA*CUCGGGC*AUG*−

Fig. 5

Fig. S1

A

Hfq$_{P.\ aeruginosa}$ hexameric molar excess

Hfq$_{P.\ putida}$ hexameric molar excess

0   1×   2.5×   5×   1×   2.5×   5×

◁3
◁2
◁1

U

B

Hfq$_{WT}$ hexameric molar excess

distal mutant Hfq$_{Y25D}$ hexameric molar excess

proximal mutant Hfq$_{K56A}$ hexameric molar excess

0   1×   2.5×   5×   1×   2.5×   5×   1×   2.5×   5×

U

Fig. S2

A

KpnI–ApaI–XhoI–SalI–HindIII–EcoRV–EcoRI–PstI–SmaI–BamHI

B

C

D