Title: Comparing faster evolving *rplB* and *rpsC* versus SSU rRNA for improved microbial

community resolution

Authors: Jiarong Guo[a], James R. Cole[a], C. Titus Brown[b], James M. Tiedje[a]

Author affiliation:

[a]Center for Microbial Ecology, Michigan State University

[b]Department of Population Health and Reproduction, University of California, Davis

Corresponding author:

James M. Tiedje

tiedjej@msu.edu

Author contributions: J.G. performed the analyses under the supervision of J.M.T., C.T.B. and

J.R.C.. All also helped with the analysis approaches and writing of the manuscript.

Abstract:

Many conserved protein-coding core genes are single copy and evolve faster, and thus are more

resolving phylogenetic markers than the standard SSU rRNA gene but their use has been

precluded by the lack of universal primers. Recent advances in gene targeted assembly methods

for large shotgun metagenomes make their use feasible. To evaluate this approach, we compared

the variation of two single copy ribosomal protein genes, *rplB* and *rpsC,* with the SSU rRNA

gene for all completed bacterial genomes in NCBI RefSeq. As expected, among pairwise

comparisons of all species that belong to the same genus, 94.9% and 91.0% of the pairs

of *rplB* and *rpsC,* respectively, showed more variation than did their SSU rRNA gene sequences.

We used a gene-targeted assembler, Xander, to assemble *rplB* and *rpsC* from shotgun

metagenomic data from rhizosphere samples of three crops: corn (annual), and *Miscanthus* and

switchgrass (both perennials). Both protein-coding genes separated all three communities

whereas the SSU rRNA gene could only separate the annual from the two perennial communities

in ordination analyses. Furthermore, assembled *rplB* and *rpsC* yielded significantly higher

numbers of OTUs (alpha diversity) than the SSU rRNA gene. These results confirm these faster

evolving marker genes offer increased resolution of for comparative microbiome studies.

Introduction:

Shaped by 3.5 billion years of evolution, microorganisms are estimated to comprise up to one

trillion species and the majority of genetic diversity in the biosphere (Locey and Lennon 2016).

Our understanding of this diversity is limited by the magnitude of extend species, and that the

majority are yet to be cultured and their physiology or functions characterized. Since the

pioneering work of Carl Woese in the late 1970s, the small subunit (SSU) rRNA gene has been

the dominant marker used in microbial community structure analyses (Woese and Fox 1977;

Lane *et al.* 1985; Huse *et al.* 2008; Caporaso *et al.* 2012). Although it has advanced our

understanding of the microbial world, it does have important limitations, namely that it is highly

conserved and that there are usually multiple copies, and some with intra-genomic variations,

making this gene problematic for taxonomic identification at species and ecotypes levels and, in

some cases, incapable of reflecting community distinctions at ecologically meaningful levels

(Case *et al.* 2007; Wu and Eisen 2008; Roux *et al.* 2011).

With the accelerated accumulation of microbial genomes in recent years (Land *et al.* 2015),

whole genome-based comparison is now feasible and a more accurate method for species and

strain identification (Goris *et al.* 2007; Luo *et al.* 2011; Scortichini *et al.* 2013; Varghese *et al.*

2015; Rodriguez-R *et al.* 2018a, 2018b). However, whole genome-based comparison is relatively

expensive computationally, compared to marker genes, and it is not yet possible to obtain

genome sequences of many members of natural microbial communities due to the challenge of

metagenome assembly from complex environmental communities (Howe *et al.* 2014; Li *et al.*

2015; Olson *et al.* 2017). Hence, marker gene analysis remains useful. Single copy protein

coding housekeeping genes stand out as the best candidates as mark genes. First, their single

copy status provides more accurate species and strain counting, identification and OTU

(Operational Taxonomic Unit) clustering than the SSU rRNA gene. Second, they are present in

virtually all known members of the three domains of life. Third, protein coding genes evolve

faster than ribosomal RNA genes not only because rRNA genes are more conserved due to their

more critical role in ribosome function (compared to ribosomal proteins) (Carter *et al.* 2000), but

also because of the redundancy in the genetic code, especially at the third codon position (Case *et*

al. 2007).

Here, we evaluate two single copy protein coding genes, *rplB* (encoding 50S ribosomal large

subunit protein L2), and *rpsC* (encoding 30S ribosomal small subunit protein S3) as potential

housekeeping genes for phylogenetic markers for microbial community analyses. Earlier studies

showed the potential of protein coding genes over SSU rRNA genes as higher resolution

phylogenetic markers for microbial diversity analyses using both genomic data (111 genomes)

and metagenomic data (< 6 Gbp by Sanger sequencing) (Case *et al.* 2007; Roux *et al.* 2011). We

revisited this comparison with the now much larger data set - all completed bacterial genomes

(~4500 with one contig) and then tested the resolving power of these two genes versus the SSU

rRNA gene among different crop rhizospheres using large shotgun metagenomic data (~1TB).

The novelty of our analyses is 1) gene-targeted assembly can recover single copy protein coding

genes more efficiently and accurately than generic assembly tools from shotgun metagenomic

data (Wang *et al.* 2015) and 2) the use of *de novo* OTU-based diversity analyses, commonly used

in microbial diversity analyses, rather than just taxonomic identification that is limited by

reference databases (Case *et al.* 2007; Roux *et al.* 2011). Our analyses also have advantages over

existing tools such as phylOTU (Sharpton *et al.* 2011) and singleM

(https://github.com/wwood/singlem) that generate *de novo* OTUs from short reads since short

reads may not overlap and thus clustering could be unreliable.

87　Methods:

Bacterial genome assembly information from NCBI

(`ftp://ftp.ncbi.nlm.nih.gov/genomes/refseq/bacteria/assembly_summ`

`ary.txt`) was used to construct the link to download each genome based on the instructions

described in this link

(http://www.ncbi.nlm.nih.gov/genome/doc/ftpfaq/#allcomplete).

Command line "wget" was then used to retrieve the genome sequences with links obtained from

the above step.

For extracting genes from genomes, the SSU rRNA gene HMM (Hidden Markov Model)

from SSUsearch (Guo *et al.* 2015) was used to recover rRNA genes. Aligned *rplB* and *rpsC*

nucleotide sequences of the "training set" retrieved from the RDP FunGene database (Fish *et al.*

2013) were used to build the HMM models using hmmbuild command in HMMER (version

3.1b2) (Eddy 2009). The nhmmer command in HMMER was then used to identify SSU rRNA,

rplB and *rpsC* sequences from bacteria genomes obtained from NCBI using score cutoff (-T) of

60. Next, nhmmer hits of least 90% of the length of the HMM model were accepted as the target

gene. For the purpose of comparing SSU rRNA, *rplB* and *rpsC* gene distances, one copy of the

SSU rRNA gene was randomly picked from each genome. Pairwise comparison among gene

sequences was done using vsearch (version 1.1.3) with "--allpairs_global --acceptall --iddef 2"

(Rognes *et al.* 2016). Two species of environmental interest, *Rhizobium leguminosarum* and

Pseudomonas putida, and their genera and orders, were chosen for closer comparison of *rplB,*

rpsC, and SSU rRNA gene distances.

The shotgun data are from DNA from seven field replicates of rhizosphere samples of three

biofuel crops: corn (C) *Zea maize*, switchgrass (S) P*anicum virgatum*, and *Miscanthus* x gigantus

(M) that had been grown for five years. Shotgun sequence data (150 bp paired ends sequenced

from 275bp libraries) for the 21 samples were downloaded from the JGI web portal

(http://genome.jgi.doe.gov/); JGI Project IDs are listed in Table S1. Raw reads were quality

trimmed using fastq-mcf in EA-Utils (verison 1.04.662) (http://code.google.com/p/ea-utils) "-l 50

-q 30 -w 4 -k 0 -x 0 --max-ns 0 -X". Overlapping paired-end reads were merged by FLASH

(version 1.2.7) (Magoc and Salzberg 2011) with "-m 10 -M 120 -x 0.20 -r 140 -f 250 -s 25" as

described in (18).

SSU rRNA gene (*E.coli* position: 515 - 806, V4) amplicon data amplified from the same

DNA as shotgun sequence (forward primer: GTGCCAGCMGCCGCGGTAA, reverse primer:

GGACTACHVGGGTWTCTAAT) were also downloaded from JGI portal (JGI project ID:

1025756) and trimmed the same way as shotgun data (described above). Paired ends were joined

by FLASH (-m 10 -M 150 -x 0.08 -p 33 -r 200 -f 300 -s 25) (Magoc and Salzberg 2011) and

primer sequences were removed by cutadapt (-f fasta --discard-untrimmed) (Martin 2011). For

community analyses, the QIIME (version 1.8.0) open reference OTU picking method with a

default OTU clustering distance cutoff of 0.03 for de novo clustering step was used for clustering

and core_diversity_analyses. py was used for generating the OTU table with taxonomy

information (Kuczynski *et al.* 2012). SILVA reference database 108 was used for taxonomy

assignment and alignment. Each sample was subsampled to 57418 reads and singletons were

removed (details in https://github.com/jiarong/2016-rplB).

For SSU rRNA gene analyses with shotgun data, SSU rRNA gene fragments and those

aligned to a part of V4 region (*E. coli* position: 577 - 727) of each sample were identified using

the SSUsearch pipeline (version 0.8.2) (Guo *et al.* 2015) and clustered using RDP's McClust tool

(Cole *et al.* 2014) at a distance of 0.05 and minimal overlap of 25 bp and minimum read length of

100 bp, following the tutorial in SSUsearch (http://microbial-ecology-

protocols.readthedocs.io/en/latest/SSUsearch/overview.html).

Both *rplB* and *rpsC* sequences were assembled using Xander with

"MAX_JVM_HEAP=500G, FILTER_SIZE=40, K_SIZE=45, genes = *rplB* and *rpsC*,

MIN_LENGTH=150, THREADS=9" (Wang *et al.* 2015). The minimal length cutoff is set to 150

amino acids (450 bp in nucleotide). Sequence reads for each crop were assembled separately. The

assembled *rplB* or *rpsC* sequences (nucleotide and protein) from the three crops were pooled and

clustered using RDP's McClust tool (Cole *et al.* 2014). For each gene, a table of OTU counts of

each sample was made based on mean k-mer coverage of the representative sequence of each

OTU (provided in "*_coverage.txt" output file from Xander). An implementation of this pipeline

is publicly available at https://doi.org/10.5281/zenodo.1438073.

Further, the above OTU tables were loaded into R, singletons were removed, and all samples

were rarefied to the sample with the least total read number using "rrarefy" in vegan package

(Oksanen *et al.* 2015). Diversity analyses were done with the vegan package using function "rda"

for ordination with Bray-Curtis distance index, "diversity" for Shannon diversity, "estimateR"

for Chao1 species richness, and "specnumber" for OTU number, from the OTU (count) tables

(details in https://github.com/jiarong/2016-rplB). For alpha diversity comparisons among genes,

all samples were subsampled to 1200 sequences.

To assess how many potential target gene reads of *rplB* and *rpsC* were assembled by Xander,

we did a six-frame translation of the short reads (nucleotide sequences) into protein sequences by

transeq in EMBOSS tool (Rice, Longden and Bleasby 2000). We then searched HMMs against

the protein sequences and the hits with bit score > 40 (e-value < $6.2 * 10^{-6}$) were treated as reads

from the target gene. In addition, "*_match_reads.fa", a collection of reads that share a k-mer

(k=45) with assembled sequences, output from Xander, provided the reads assembled by Xander.

Then we compared the fold coverage of reads found by hmmsearch and reads used by Xander, by

estimating fold coverage of each read with median kmer coverage using khmer package (Brown

et al. 2012; Crusoe *et al.* 2015).

Results:

A total of 4,457 of complete bacteria genomes (genome as one sequence) were downloaded,

and 4,440 of them have all three genes. SSU rRNA gene copy number ranged from 1 to 16 with a

mean of 4, and *rplB* and *rpsC* are single copy in 99.9% of genomes (Table S2). When evaluating

intra-genomic variation among copies of SSU rRNA genes in completed genomes of *R.*

leguminosarum, *P. putida* and *E. coli*, *E. coli* had the largest variation with a minimum of 95.4%

identity (Fig. S1).

For the selected taxa, Rhizobiales, Pseudomonadales, *Rhizobium*, and *Pseudomonas*, *rplB*

and *rpsC* had similar variations and both had larger variation among the genomes than SSU

rRNA genes within their corresponding order (among genera), and genus (among species) (Fig. 1

and 2). When comparing all species of completed genomes that belong to the same genus, we

found SSU rRNA gene has an identity range of 63.2% to 100.0% and a median of 95.2%, *rplB*

has an identity range of 43.2% to 100.0% and a median of 87.2%, *rpsC* has an identity range of

46.0% to 100.0% and a median of 90.3%. Between *rplB* and SSU rRNA gene, 88,993 pairs

(94.9% of total) had larger variation in *rplB*, 3,573 pairs (3.8%) had larger variation in SSU

rRNA gene, and 1,167 pairs (1.2%) had the same variation (Fig. 3A); Between *rpsC* and SSU

rRNA gene, 77,885 pairs (91.0% of total) had larger variation in *rpsC*, 6,074 pairs (7.1%) had

larger variation in SSU rRNA gene, and 1,622 pairs (1.9%) had the same variation (Fig. 3B);

Between *rplB* and *rpsC*, 54,755 pairs (63.7%) had larger variation in *rplB*, 28,393 pairs (33.0%)

had larger variation in *rpsC* gene, and 2,808 pairs (3.3%) had the same variation for *rplB* and

rpsC (Fig. 3C).

We compared SSU rRNA genes (V4) with *rplB and rpsC* to test the ability of shotgun data to

resolve community differences among plant rhizospheres. We chose these two genes as they had

a suitable length for Xander assembly (~ 660bp for *rpsC* and 830bp for *rplB*), were long enough

for resolving power, and had HMMs that were both specific and sensitive for fragment recovery

due to their uniqueness in sequence as parts of the ribosome. Both have also been used as

phylogenetic markers in other shotgun metagenomic studies (Hug *et al.* 2013; Sharon *et al.*

2015). On average, 0.04% of total reads were identified as SSU rRNA gene fragments and

0.004% of total reads aligned to the 150 bp of V4 region of the gene with SSUsearch (Guo *et al.*

2015). Another 0.01% and 0.008% of total reads were identified as *rplB* and *rpsC,* respectively,

by Xander (Table S3). The assembled sequences (using a length cutoff of 450 bp) have lengths

ranging from 453 to 843bp with a median of 822bp for *rplB* and from 453 to 678bp with a

median of 648 bp for *rpsC*. To test the sensitivity of Xander, we found that the number of

potential *rplB* and *rpsC* reads assembled were 49.5% and 47.9%, respectively, of those defined

by hmmsearch with bit score cutoff of 40 (Table S4) and have much higher fold coverage than

the rest of reads in hmmsearch hits (excluding shared reads with Xander) (Figure S2).

Beta diversity analyses of all three genes showed that the rhizosphere communities of the

annual crop, corn, were different from those of the two perennial grasses, *Miscanthus* and

switchgrass, but only *rplB* and *rpsC* distinguished the communities of the two perennial grasses

(Fig. 4). This was true whether the analysis was at the nucleotide or protein level. The alpha

diversity of the corn rhizosphere communities was significantly lower than those of *Miscanthus*

and switchgrass rhizospheres by all three measures except for Chao1 index with *rpsC* and SSU

rRNA gene (Fig. 5). When comparing among genes, the numbers of OTUs from *rplB* and *rpsC*

were also significantly higher than from the SSU rRNA gene (Fig. 5). Since SSUSearch returns

shorter fragments (~100bp to 150bp) than Xander assembled genes (~ 450bp to 849bp), we also

evaluated whether the longer fragments of SSU rRNA from amplicon data (~ 250 to 300 bp),

could distinguish the two perennial grass communities, and they could not (Fig. 4).

Discussion:

We confirmed the advantages of *rplB* and *rpsC* over the SSU rRNA gene as a more resolving

phylogenetic marker using updated large genomic data (~4500 complete genomes) (Fig. 1, 2 and

3). We also demonstrated that *rplB* and *rpsC* can be assembled from large shotgun metagenomes

and showed that they provided higher community resolution by separating *Miscanthus* and

switchgrass rhizosphere samples while the SSU rRNA gene did not (Fig. 4). The two perennial

grasses would be expected to have more similar microbiome than the annual since the latter is re-

established each year while the fibrous perennial grass roots are more similar and not physically

disturbed annually and thus do not have full regrowth at a new random site each year.

In large genomic data analyses, *rplB* and *rpsC* show advantages in following three aspects:

First, SSU rRNA gene, a multiple copy gene, poses difficulties for interpreting species

abundance, while *rplB* and *rpsC* do not have the same issue as it is single copy genes in > 99.9%

of complete genomes (Table S2). Additionally, variations among multiple SSU copies can cause

multiple OTUs (sequence clusters) from the same species (Figure S1) and thus leads to

overestimation of species richness (31). Since a single copy of the *rplB* and *rpsC* genes is

contained in every cell in a community, the abundance of *rplB* and *rpsC* gene sequences provides

a reference for estimating the fraction of organisms possessing other genes and also average

genome size (Raes *et al.* 2007; Frank and Sørensen 2011; Nayfach and Pollard 2015; Vital,

Karch and Pieper 2017).

Second, *rplB* and *rpsC* are better able to differentiate closely related species based on their

lower sequence identities compared to the SSU rRNA gene in pairwise comparisons among

genomes (Fig. 1, 2, and 3). This is consistent with the crucial role SSU rRNA plays in translation

(ensuring translation accuracy) (Carter *et al.* 2000), also confirmed by another study showing

SSU rRNA genes (along with LSU rRNA genes, tRNA and ABC transporter genes) to be the

most conserved genes (Isenbarger *et al.* 2008).

Third, SSU rRNA genes in genomes are also more prone to assembly errors (chimera) than

single copy genes due to their higher overall nucleotide identity and the presence of highly

conserved regions interspersed in SSU rRNA genes (Miller 2013; Sharon *et al.* 2015; Yuan *et al.*

2015). Note that these erroneous sequences might be further collected by databases and used as

references for taxonomy, alignment, and chimera detection, and thus have an impact on common

microbial ecology analyses. Single copy protein coding genes, however, are less challenging to

assemble compared to the SSU rRNA gene since they have more unique sequences among

different taxa and no highly conserved regions. They do however, require enough sequence for a

sufficient number of gene assemblies (Hug *et al.* 2013; Sharon *et al.* 2015).

Additionally, this method provides for higher resolution community diversity analyses in

large shotgun metagenomes, leveraging a scalable gene targeted assembler, Xander. Assembly is

desirable for short read data to correctly identify the gene and provide enough length for

resolving power, a major objective in ecology studies. While this method avoids the primer bias

of amplicon sequencing (Farris and Olson 2007; Bergmann *et al.* 2011; Guo *et al.* 2015), it does

miss the rarer species because not enough fragments of the targeted genes from rarer members

are sequenced to allow their assembly, perhaps missing up to 50% of the cells of lowest

abundance (Table S4 and Fig. S2). The hmmsearch though could also have recovered some false

positives due to mistaken short-read identification and thus overestimated the total gene number

(Orellana, Rodriguez-R and Konstantinidis 2017). However, *rplB* and *rpsC* still yielded

significantly higher alpha diversity (Fig. 5) than the SSU rRNA gene despite missing rare

members. These protein-coding genes, although shorter than the full-length SSU rRNA gene, are

both more rapidly evolving and longer than produced with commonly used SSU rRNA gene

primers. Thus they reveal more diversity among abundant members than SSU rRNA gene, which

offsets and exceeds the diversity of the rare members that are not assembled, further confirming

their higher resolution. Additionally, rare members are commonly removed in amplicon-based

analyses, especially singletons, to remove spurious OTUs caused by sequencing error and to

improve computational efficiency. Rare valid members are unavoidably removed in this process

too, but this practice has been shown to have minimal impact on beta diversity analyses (Edgar

2013; Bokulich *et al.* 2013; Auer *et al.* 2017).

OTU distance cutoff could also impact community analyses in our comparison. We used a

distance cutoff of 0.05 for *rplB* and *rpsC* since that is the cutoff recommend (22) (and we used)

for SSUSearch, We used the standard, and more resolving 0.03 cutoff for the SSU rRNA gene

amplicon data, but it showed less community resolution than *rplB* and *rpsC* indicating that our

cutoff is appropriate and fairly evaluating the comparison (S1 text).

We did not attempt assembling the SSU rRNA gene because: i) as noted above, the

conserved regions results in a high proportion of miss-assembly with short reads, generating

chimeras (Miller 2013; Sharon *et al.* 2015; Yuan *et al.* 2015), ii) the existing tools, e.g. EMIRGE

and REAGO, have drawbacks. EMIRGE relies heavily on reference databases and does not

perform well when close relatives are absent. Further, EMIRGE was shown to have significant

false positives (Fan, McElroy and Thomas 2012; Miller 2013; Yuan *et al.* 2015). REAGO has

only been tested with low diversity, mock community data and does not report abundance

information (Yuan *et al.* 2015).

We chose two protein-coding genes to ensure that our results were not gene specific, and both

gave very similar results at both the nucleotide and protein levels. At least from extensive

completed genomes, more than 99% of these two genes are single copy, making quantitative

(ratio) comparisons with other genes consistent. For future use, *rplB* might have a slight

advantage over *rpsC* since it is longer, about 830 bp on average vs. 660 bp of *rpsC* (Fish *et al.*

2013), providing a bit more resolving power, which is consistent with results in genome

comparisons showing *rplB* has lower median sequence identity than *rpsC* (Fig 3).

Besides higher resolution in de novo OTU related analyses, it is of course possible to do

taxonomy related analyses by finding the best match to the assembled sequence of these marker

genes in reference databases and potentially finer taxonomic resolution than provided by SSU

rRNA. But, the reference databases of *rplB* and *rpsC* are mostly from sequenced genomes and

hence are very unbalanced and incomplete compared to SSU rRNA gene databases (Quast *et al.*

2013; Cole *et al.* 2014; Wang *et al.* 2015) so this use is not generally beneficial at this time.

Although the sequencing depth needed varies depending on community diversity, we provide

an estimate based on our rhizosphere soil samples as a guide. The reads from *rplB* are around

0.01% of total (Table S3). Assuming a fold coverage of 3000 of *rplB* for each sample, to be

comparable to 3000 amplicons in planning amplicon-based studies, one needs about 25 Gbp

(3000 * 830 / 0.01%) of shotgun metagenome (830 bp is the average gene length of *rplB*). The

major requirement for using this method beyond sufficient shotgun sequence depth is an access

to a high performance computer since large memory (> 250 Gb recommended for soil samples) is

needed for the deBruin graph (the most memory and cpu consuming step) to run Xander.

However, once one has the deBruin graph, it can also be used to assemble other genes of interest

(Wang *et al.* 2015).

Conclusion:

We demonstrated that *rplB* and *rpsC*, single copy protein coding genes can provide finer

resolution of taxa and hence better distinguish among communities than the more commonly

used SSU rRNA gene and also provide finer scale *de novo* OTU diversity analysis. This method

does require shotgun sequence of sufficient depth, so is currently more costly than amplicon

based analyses, but as sequencing costs decline, capacity and access increase, and read length and

genome reference databases grow, single copy protein coding genes such *rplB* and *rpsC* have the

potential to complement or even replace the SSU rRNA gene as a phylogenetic marker and better

reflect the ecology of microbial communities.

Acknowledgement:

We thank Ribosomal Database Project (RDP), Institute for Cyber-Enabled Research

(iCER) and High Performance Computing Center (HPCC) at Michigan State University for

technical support. Support for this research was provided by the U.S. Department of Energy,

Office of Science, Program of Biological and Environmental Research (Awards DE-FC02-

07ER64494 and DE-FG02-99ER62848), and by the National Science Foundation DBI-1356380,

DBI-1759892, and Long-term Ecological Research Program (DEB 1637653) at the Kellogg

Biological Station, and by Michigan State University AgBioResearch.

References:

Auer L, Mariadassou M, O'Donohue M *et al.* Analysis of large 16S rRNA Illumina data sets: Impact of singleton read filtering on microbial community description. *Mol Ecol Resour* 2017;**17**:e122–32.

Bergmann GT, Bates ST, Eilers KG *et al.* The under-recognized dominance of Verrucomicrobia in soil bacterial communities. *Soil Biol Biochem* 2011;**43**:1450–1455.

Bokulich NA, Subramanian S, Faith JJ *et al.* Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing. *Nat Methods* 2013;**10**:57–9.

Brown CT, Howe A, Zhang Q *et al.* A Reference-Free Algorithm for Computational Normalization of Shotgun Sequencing Data. *ArXiv12034802 Q-Bio* 2012.

Caporaso JG, Lauber CL, Walters WA *et al.* Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME J* 2012;**6**:1621–1624.

Carter AP, Clemons WM, Brodersen DE *et al.* Functional insights from the structure of the 30S ribosomal subunit and its interactions with antibiotics. *Nature* 2000;**407**:340–348.

Case RJ, Boucher Y, Dahllof I *et al.* Use of 16S rRNA and rpoB genes as molecular markers for microbial ecology studies. *Appl Env Microbiol* 2007;**73**:278–288.

Cole JR, Wang Q, Fish JA *et al.* Ribosomal Database Project: data and tools for high throughput rRNA analysis. *Nucleic Acids Res* 2014;**42**:D633–642.

Crusoe MR, Alameldin HF, Awad S *et al.* The khmer software package: enabling efficient nucleotide sequence analysis. *F1000Research* 2015, DOI: 10.12688/f1000research.6924.1.

Eddy SR. A new generation of homology search tools based on probabilistic inference. *Genome Inf* 2009;**23**:205–211.

Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods* 2013;**10**:996–8.

Fan L, McElroy K, Thomas T. Reconstruction of ribosomal RNA genes from metagenomic data. *PloS One* 2012;**7**:e39948.

Farris MH, Olson JB. Detection of Actinobacteria cultivated from environmental samples reveals bias in universal primers. *Lett Appl Microbiol* 2007;**45**:376–381.

Fish JA, Chai B, Wang Q *et al.* FunGene: the functional gene pipeline and repository. *Front Microbiol* 2013;**4**, DOI: 10.3389/fmicb.2013.00291.

Frank JA, Sørensen SJ. Quantitative Metagenomic Analyses Based on Average Genome Size Normalization. *Appl Environ Microbiol* 2011;**77**:2513–21.

Goris J, Konstantinidis KT, Klappenbach JA *et al.* DNA-DNA hybridization values and their
relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol*
2007;**57**:81–91.

Guo J, Cole JR, Zhang Q *et al.* Microbial community analysis with ribosomal gene fragments
from shotgun metagenomes. *Appl Environ Microbiol* 2015:AEM.02772-15.

Howe AC, Jansson JK, Malfatti SA *et al.* Tackling soil diversity with the assembly of large,
complex metagenomes. *Proc Natl Acad Sci* 2014;**111**:4904–9.

Hug LA, Castelle CJ, Wrighton KC *et al.* Community genomic analyses constrain the
distribution of metabolic traits across the Chloroflexi phylum and indicate roles in
sediment carbon cycling. *Microbiome* 2013;**1**:22.

Huse SM, Dethlefsen L, Huber JA *et al.* Exploring microbial diversity and taxonomy using SSU
rRNA hypervariable tag sequencing. *PLoS Genet* 2008;**4**:e1000255.

Isenbarger TA, Carr CE, Johnson SS *et al.* The most conserved genome segments for life
detection on Earth and other planets. *Orig Life Evol Biosph* 2008;**38**:517–533.

Kuczynski J, Stombaugh J, Walters WA *et al.* Using QIIME to analyze 16S rRNA gene
sequences from microbial communities. *Curr Protoc Microbiol* 2012;**Chapter 1**:Unit
1E.5.

Land M, Hauser L, Jun SR *et al.* Insights from 20 years of bacterial genome sequencing. *Funct*
Integr Genomics 2015;**15**:141–161.

Lane DJ, Pace B, Olsen GJ *et al.* Rapid determination of 16S ribosomal RNA sequences for
phylogenetic analyses. *Proc Natl Acad Sci USA* 1985;**82**:6955–6959.

Li D, Liu CM, Luo R *et al.* MEGAHIT: an ultra-fast single-node solution for large and complex
metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 2015;**31**:1674–
1676.

Locey KJ, Lennon JT. Scaling laws predict global microbial diversity. *Proc Natl Acad Sci USA*
2016;**113**:5970–5975.

Luo C, Walk ST, Gordon DM *et al.* Genome sequencing of environmental Escherichia coli
expands understanding of the ecology and speciation of the model bacterial species. *Proc*
Natl Acad Sci USA 2011;**108**:7200–7205.

Magoc T, Salzberg SL. FLASH: fast length adjustment of short reads to improve genome
assemblies. *Bioinformatics* 2011;**27**:2957–2963.

Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal 2011;**17**:10–2.

Miller CS. Assembling full-length rRNA genes from short-read metagenomic sequence datasets
using EMIRGE. *Meth Enzym* 2013;**531**:333–352.

Nayfach S, Pollard KS. Average genome size estimation improves comparative metagenomics
and sheds light on the functional ecology of the human microbiome. *Genome Biol*
2015;**16**, DOI: 10.1186/s13059-015-0611-7.

Oksanen J, Blanchet FG, Kindt R *et al. Vegan: Community Ecology Package.*, 2015.

Olson ND, Treangen TJ, Hill CM *et al.* Metagenomic assembly through the lens of validation:
recent advances in assessing and improving the quality of genomes assembled from
metagenomes. *Brief Bioinform* 2017, DOI: 10.1093/bib/bbx098.

Orellana LH, Rodriguez-R LM, Konstantinidis KT. ROCker: accurate detection and
quantification of target genes in short-read metagenomic data sets by modeling sliding-
window bitscores. *Nucleic Acids Res* 2017;**45**:e14.

Quast C, Pruesse E, Yilmaz P *et al.* The SILVA ribosomal RNA gene database project: improved
data processing and web-based tools. *Nucleic Acids Res* 2013;**41**:D590–596.

Raes J, Korbel JO, Lercher MJ *et al.* Prediction of effective genome size in metagenomic
samples. *Genome Biol* 2007;**8**:R10.

Rice P, Longden I, Bleasby A. EMBOSS: The European Molecular Biology Open Software
Suite. *Trends Genet* 2000;**16**:276–7.

Rodriguez-R LM, Castro JC, Kyrpides NC *et al.* How Much Do rRNA Gene Surveys
Underestimate Extant Bacterial Diversity? *Appl Environ Microbiol* 2018a;**84**:e00014-18.

Rodriguez-R LM, Gunturu S, Harvey WT *et al.* The Microbial Genomes Atlas (MiGA)
webserver: taxonomic and gene diversity analysis of Archaea and Bacteria at the whole
genome level. *Nucleic Acids Res* 2018b;**46**:W282–8.

Rognes T, Flouri T, Nichols B *et al.* VSEARCH: a versatile open source tool for metagenomics.
PeerJ 2016;**4**:e2584.

Roux S, Enault F, Bronner G *et al.* Comparison of 16S rRNA and protein-coding genes as
molecular markers for assessing microbial diversity (Bacteria and Archaea) in
ecosystems. *FEMS Microbiol Ecol* 2011;**78**:617–628.

Scortichini M, Marcelletti S, Ferrante P *et al.* A Genomic Redefinition of Pseudomonas avellanae
species. *PLOS ONE* 2013;**8**:e75794.

Sharon I, Kertesz M, Hug LA *et al.* Accurate, multi-kb reads resolve complex populations and
detect rare microorganisms. *Genome Res* 2015;**25**:534–43.

Sharpton TJ, Riesenfeld SJ, Kembel SW *et al.* PhylOTU: a high-throughput procedure quantifies
425   microbial community diversity and resolves novel taxa from metagenomic data. *PLoS*
426   *Comput Biol* 2011;**7**:e1001061.

427 Varghese NJ, Mukherjee S, Ivanova N *et al.* Microbial species delineation using whole genome
428   sequences. *Nucleic Acids Res* 2015;**43**:6761–6771.

429 Vital M, Karch A, Pieper DH. Colonic Butyrate-Producing Communities in Humans: an
430   Overview Using Omics Data. *mSystems* 2017;**2**, DOI: 10.1128/mSystems.00130-17.

431 Wang Q, Fish JA, Gilman M *et al.* Xander: employing a novel method for efficient gene-targeted
432   metagenomic assembly. *Microbiome* 2015;**3**:32.

433 Woese CR, Fox GE. Phylogenetic structure of the prokaryotic domain: the primary kingdoms.
434   *Proc Natl Acad Sci USA* 1977;**74**:5088–5090.

435 Wu M, Eisen JA. A simple, fast, and accurate method of phylogenomic inference. *Genome Biol*
436   2008;**9**:R151.

437 Yuan C, Lei J, Cole J *et al.* Reconstructing 16S rRNA genes in metagenomic data.
438   *Bioinformatics* 2015;**31**:35–43.

460 **Figures**

461
462

464   Figure 1: Pairwise comparisons among all genomes of Rhizobiales (panels A, C, E) and of all

465   *Rhizobium* (panels B, D, F) using the SSU rRNA gene, *rplB* and *rpsC*. SSU rRNA gene identities

466   are higher than *rplB* and *rpsC*, and *rplB* and *rpsC* have similar sequence identities in most

467   genome pairs. The dashed line is y = x. Data below y=x line indicate the gene on X axis is more

468   conserved. The dot size indicates the number of pairwise comparisons with those values.

489 Figure 2: Pairwise comparison among all genomes of Pseudomonadales (panels A, C, E) and of

490 all *Pseudomonas* (panels B, D, F) using the SSU rRNA gene, *rplB* and *rpsC*. SSU rRNA gene

491 identities are higher than *rplB* and *rpsC* in most genome pairs, and *rplB* and *rpsC* have similar

492 sequence identities. The dashed line is y = x. Data below y=x line indicate gene on X axis is

493 more conserved. The dot size indicates the number of pairwise comparisons with those values.

Figure 3: Pairwise comparison among all completed genomes of species in the same genus using

the SSU rRNA gene, *rplB,* and *rpsC*. SSU rRNA gene identities are larger than *rplB* in most

genomes. The diagonal dashed line is y = x and data below the line indicates the gene on X axis

is more conserved. The dot intensity is the number of comparisons with those values. Subplot A,

B, and C are comparisons of *rplB* and SSU rRNA gene (93,733 pairwise comparisons), *rpsC* and

SSU rRNA gene (85,581 pairwise comparisons), *rplB* and *rpsC* (85,956 pairwise comparisons).

Figure 4: Comparison of the SSU rRNA gene, *rpsC* and *rplB* in beta diversity analyses

(ordination) using large soil metagenome sequences from seven field replicates. All genes show

that the microbial community of the corn (C) rhizosphere is significantly different from those of

Miscanthus (M) and switchgrass (S) while *rplB* and *rpsC* at both the nucleotide (n) and protein

(p) levels separate microbial communities of *Miscanthus* and switchgrass. The SSU rRNA gene

does not separate *Miscanthus* and switchgrass with either shotgun (SSU.sg) or amplicon

(SSU.am) data. "**" indicates $p < 0.01$ in PERMANOVA test.

Figure 5: Comparison of the SSU rRNA gene, *rplB*, and *rpsC* in alpha diversity analyses (Chao1,

Shannon, and OTU number) using large soil metagenome sequence. The labels with suffixes

"_n" and "_p" means assembled nucleotide and protein sequences by Xander respectively; those

with suffixes "_am" and "_sg" are SSU rRNA gene from amplicon data and shogun data

respectively. All genes show that the microbial community of the corn (C) rhizosphere has

significantly less alpha diversity than those of *Miscanthus* (M) and switchgrass (S) except for

rpsC and SSU rRNA gene with Chao1 index ($p < 0.01$). Wilcoxon test was used to compare

SSU_sg against each of the other genes including rplB_n, rplB_p, rpsC_n, rpsC_p and SSU_am.

For Chao1 index, rplB_n and rpsC_n show significantly higher abundance than SSU_sg in

Miscanthus and switchgrass; For Shannon index and OTU number, rplB_n and rpsC_n show

significantly higher abundance than SSU_sg in all three crops ("***" is $p < 0.001$, '**' is $p <$

0.01, '*' is $p < 0.05$, "ns" is $p > 0.05$).